

The existence of time global solutions for tumor invasion models

RISEI KANO

Center for the Advancement of Higher Education,
 Faculty of Engineering, Kinki University
 Takayaumenobe 1, Higashihiroshimashi, Hiroshima 739-2116 JAPAN
 Email:kano@hiro.kindai.ac.jp

AKIO ITO

Center for the Advancement of Higher Education,
 Faculty of Engineering, Kinki University
 Takayaumenobe 1, Higashihiroshimashi, Hiroshima 739-2116 JAPAN
 Email:aito@hiro.kindai.ac.jp

1 Introduction

In this paper, we consider a tumor invasion model with constraint, which is the following systems:

$$(P) \left\{ \begin{array}{l} \frac{\partial n}{\partial t} = \nabla \cdot \{K_1(\cdot)\nabla n - \lambda n \nabla f\} + \mu n(1 - n - f) \quad \text{in } Q(T) := \Omega \times (0, T), \\ \frac{\partial f}{\partial t} = -\delta m f \quad \text{in } Q(T), \\ \frac{\partial m}{\partial t} = K_2(\cdot)\Delta m + C_1 n - C_2 m \quad \text{in } Q(T), \\ 0 \leq n + f \leq 1, m \geq 0, f \geq 0, n \geq 0 \quad \text{in } Q(T), \\ n = 0, \quad \text{in } \Sigma(T) := \Gamma \times (0, T), \\ \frac{\partial m}{\partial \mathbf{n}} = 0 \quad \text{in } \Sigma(T), \\ n(0) = n_0, f(0) = f_0, m(0) = m_0 \quad \text{in } \Omega, \end{array} \right.$$

where $0 < T < \infty$; Ω is bounded domain in $\mathbb{R}^N (N = 1, 2, 3)$ with a smooth boundary $\Gamma = \partial\Omega$; $K_n(\cdot)$ is a non-negative function on $(0, T)$; $\lambda(\cdot)$ is a non-negative function on $Q(T)$; K_m, μ, δ, C_1 and C_2 are positive constants. \mathbf{n} is the outer unit normal vector on Γ ; n_0, m_0 and f_0 are initial date. In this model, the unknown functions n, f and m describe the densities of solid tumor cells, the extracellular matrix (denoted by ECM) and the matrix degrading enzymes (denoted by MDE), respectively.

Remark. K_n and K_m are to express that diffusion rate of the tumor cells and MDE, respectively. Originally, they are determined by the state of the protein that exists there.

Therefore, the coefficients must be determined to be dependent on time and space. However, K_n is dependent upon only time.

2 Approach by quasi-variational inequality

First of all, we define the operators which satisfy the following properties:

1. For each $t \in [0, T]$ and $v \in L^2(Q(T))$, we consider the problem $(P)_m$:

$$(P)_m \begin{cases} \frac{\partial \hat{m}}{\partial t} = K_m \Delta \hat{m} + C_1 v - C_2 \hat{m} & \text{in } Q(T), \\ \nabla \hat{m} \cdot \mathbf{n} = 0 & \text{on } \Sigma(T), \\ \hat{m}(0) = m_0 & \text{in } \Omega. \end{cases}$$

Then, we denote by $\Lambda_1(t)$ is a solution operator on $L^2(0, T; L^2(\Omega))$ which assigns a unique solution of $(P)_m$ to v , namely, $\hat{m} = \Lambda_1(t)v$.

2. For each $t \in [0, T]$ and $w \in L^2(Q(T))$, we define a function $\Lambda_2(t)w$ by

$$[\Lambda_2(t)w](x, s) := f_0(x) \exp\left(-\delta \int_0^s w(x, \tau) ds\right), \quad \forall (x, s) \in Q(T).$$

Then, $\Lambda_2(t)$ is a solution operator which assigns a unique solution \hat{f} of the problem $(P)_f$ below to w :

$$(P)_f \begin{cases} \frac{\partial \hat{f}}{\partial t} = -\delta \hat{f} w & \text{in } Q(T), \\ \hat{f}(0) = f_0 & \text{in } \Omega. \end{cases}$$

3. For each $t \in [0, T]$ we put $\Lambda(t) := \Lambda_2(t) \circ \Lambda_1(t)$.

Using these operators, we give the definition of (P).

Definition 2.1 For each $t \in [0, T]$ a triplet $\{n, f, m\}$ is called a solution of (P) on $[0, t]$ if and only if the following properties are fulfilled:

(S1) $n \in W^{1,2}(0, t; L^2(\Omega)) \cap L^\infty(0, t; H_0^1(\Omega))$.

(S2) $m = \Lambda_1(t)n$, $f = \Lambda(t)n$.

(S3) $0 \leq n \leq 1 - f$ a.e. in $Q(T)$,

$$\int_0^t \int_\Omega \left(\frac{\partial n}{\partial s}(s) - \mu n(s)(1 - n(s) - f(s)) \right) (n(s) - v(s)) dx ds \\ + \int_0^t \int_\Omega \left(\lambda(s) \{n(s) \nabla f(s)\} + K_n(s) \nabla n(s) \right) \cdot \nabla (n(s) - v(s)) dx ds \leq 0,$$

for $\forall v \in L^2(0, t; H_0^1(\Omega))$ with $0 \leq v \leq 1 - f$ a.e. in $Q(T)$.

(S4) $n(0) = n_0$ a.e. in Ω .

3 An abstract existence result

In this section, we express about the existence of solution in abstract theory. we use the following notation. Let H be a real Hilbert space equipped with a usual norm $|\cdot|_H$ and an inner product $(\cdot, \cdot)_H$, and X be a real reflexive Banach space, and let X^* be a dual space of X . We assume that X is density and compact imbeded in H .

We consider a nonlinear evolution problem the following formulation

$$(CP): \frac{du}{dt}(t) + \partial\varphi^t(u; u(t)) \ni g(t), \quad 0 < t < T, \quad u(0) = u_0, \quad \text{in } H,$$

where $\partial\varphi^t(u; \cdot)$ is the subdifferential of convex function $\varphi^t(u; \cdot)$ on H , $u' = \frac{du}{dt}$ and $u_0 : [-\delta_0, 0] \rightarrow H$ and $f : (0, T) \rightarrow H$ are the initial and forcing functions, respectively. We define the following functional space and its norm; we put

$$\begin{aligned} \mathcal{V}(-\delta_0, t) &:= W^{1,2}(-\delta_0, t; H) \cap L^\infty(-\delta_0, t; X), \quad 0 \leq t \leq T. \\ |v|_{\mathcal{V}(-\delta_0, t)} &:= |v|_{L^\infty(-\delta_0, t; X)} + |v'|_{L^2(-\delta_0, t; H)}. \end{aligned}$$

This is a sort of functional differential equations generated by subdifferentials of $\varphi^t(v; \cdot)$ with a nonlocal dependence upon v . The objective of this paper is to specify a class of convex functions $\{\varphi^s(v; \cdot)\}_{0 \leq s \leq t}$ as well as its nonlocal dependence upon $v \in \mathcal{V}(-\delta_0, T)$ in order that above Cauchy problem admits at least one local or global in time solution u .

Definition 3.1(Mosco convergence) Let $\{\varphi_n\}$ be a sequence of proper, lower semi-continuous(l.s.c.), convex functions on X . Then $\{\varphi\}$ converges to a proper, l.s.c., convex function φ on X in the sense of Mosco, if the following two conditons (M1) and (M2) are satisfied:

(M1) Let $\{n_k\}$ be any subsequence of $\{n\}$. If $\{v_k\}$ is a sequence in X and $v \in X$ such that $v_k \rightarrow v$ weakly in X as $k \rightarrow \infty$, then

$$\liminf_{k \rightarrow \infty} \varphi_{n_k}(v_k) \geq \varphi(v).$$

(M2) For each $v \in D(\varphi)$, there is a sequence $\{v_n\}$ in X such that

$$v_n \rightarrow v \text{ in } X, \quad \varphi_n(v_n) \rightarrow \varphi(v) \text{ as } n \rightarrow \infty.$$

For $\forall v \in \mathcal{V}(-\delta_0, t)$ we are given a family $\{\varphi^s(v; \cdot)\}_{0 \leq s \leq t}$ such that

(Φ1) $\varphi^s(v; z)$ is proper, l.s.c., non-negative, convex in $z \in H$; $\varphi^s(v; z)$ is determined by the value of v on $(-\delta_0, s)$, namely $\varphi^s(v_1; z) = \varphi^s(v_2; z)$ wherever $v_1, v_2 \in \mathcal{V}(-\delta_0, t)$, $v_1 = v_2$ on $(-\delta_0, s)$.

(Φ2) $\varphi^s(v; z) \geq C_0|z|_X^p$, $0 \leq \forall s \leq t$, $\forall v \in \mathcal{V}(-\delta_0, t)$, where $2 \leq p < \infty$ and $C_0 > 0$ are constants.

(Φ3) If $0 \leq s_n \leq t \leq T$, $v_n \in \mathcal{V}(-\delta_0, t)$, $s_n \rightarrow s$ and $v_n \rightarrow v$ weakly in $W^{1,2}(-\delta_0, t; H)$ and weakly* in $L^\infty(-\delta_0, t; X)$, then $\varphi^{s_n}(v_n; \cdot) \rightarrow \varphi^s(v; \cdot)$ on H in the sense of Mosco.

Definition 3.2 $u_0 \in C([-\delta_0, 0]; H)$ and $f \in L^2(0, T; H)$. Then we say that u is a solution of the Cauchy problem

$$(CP) \begin{cases} u'(t) + \partial\varphi^t(u; u(t)) \ni f(t), & 0 < t < T, \text{ in } H, \\ u(t) = u_0(t), & -\delta_0 \leq t \leq 0, \text{ in } H, \end{cases}$$

if u satisfies that $u \in C([-\delta_0, T]; H)$, $u = u_0$ on $[-\delta_0, 0]$, $u \in W_{loc}^{1,2}((0, T]; H)$, $\varphi^{(\cdot)}(u; u(\cdot)) \in L^1(0, T)$ and $f(t) - u'(t) \in \partial\varphi^t(u; u(t))$ for a.e. $t \in (0, T)$.

Theorem 3.1 Let $0 < T < +\infty$, $0 < \delta_0 < +\infty$, $f \in L^2(0, T; H)$ and $u_0 \in \mathcal{V}(-\delta_0, 0)$ with $\varphi^0(u_0; u_0(0)) < +\infty$. Assume that for all $M > 0$ and $M \leq M^* := M^*(f, u_0, \varphi^0(u_0; u_0(0)))$, there are two bounded families $A_M := \{a; v \in \mathcal{V}(-\delta_0, T), |v|_{\mathcal{V}(-\delta_0, T)} \leq M\}$ of non-negative functions in $L^2(0, T)$ and $B_M := \{b; v \in \mathcal{V}(-\delta_0, T), |v|_{\mathcal{V}(-\delta_0, T)} \leq M\}$ of non-negative functions in $L^1(0, T)$ such that

(H1) for each $v \in \mathcal{V}(-\delta_0, T)$, $|v|_{\mathcal{V}(-\delta_0, T)} \leq M$, $v = u_0$ on $[-\delta_0, 0]$, there exist $a \in A_M$ and $b \in B_M$ with the following property: for each $s, t \in [0, T]$ with $s \leq t$ and $z \in D(\varphi^s(v; \cdot))$, there exists $\tilde{z} \in D(\varphi^t(v; \cdot))$ such that

$$\begin{cases} |\tilde{z} - z|_H \leq \int_s^t a(\tau) d\tau (1 + \varphi^s(v; z)^{\frac{1}{2}}), \\ \varphi^t(\tilde{z}) - \varphi^s(z) \leq \int_s^t b(\tau) d\tau (1 + \varphi^s(v; z)), \end{cases}$$

(H2) for all $\varepsilon > 0$, there exists $\delta_\varepsilon > 0$ such that

$$\int_t^{t+\delta_\varepsilon} (a(s)^2 + b(s)) ds < \varepsilon, \quad \forall t \in [0, T - \delta_\varepsilon], \quad \forall a \in A_M, \quad \forall b \in B_M.$$

Then, problem (CP) has at least one solution u on an interval $[0, T']$ with $0 < T' \leq T$ such that $u \in \mathcal{V}(-\delta_0, T')$ and $\sup_{0 \leq t \leq T'} \varphi^t(u; u(t)) < +\infty$.

The detail of of proof is referred to the paper [5].

4 Main result

4.1 Auxiliary equation

In this paper, we give some propositions which is the existence of solutions of auxiliary problem and its estimate. We can directly apply the theory established in [2] to derive Proposition 4.1. So, we omit its proof in this paper.

Proposition 4.1. (cf. [7]) For each $t \in [0, T]$, $v \in \mathcal{V}(-\delta_0, t)$ and $\hat{n} \in L^\infty(0, T; H_0^1(\Omega))$ the problem

$$(AP)_{t,v,\hat{n}} \begin{cases} n'(s) + \partial\varphi^s(v; n(s)) \ni G(s, \hat{n}(s), [\Lambda(t)\hat{n}](s)) & \text{in } L^2(\Omega), \quad \text{a.e. } s \in (0, t), \\ n(s) = n_0 & \text{in } L^2(\Omega), \quad \forall s \in [-\delta_0, 0]. \end{cases}$$

has a unique solution $n = n_{t,v,\hat{n}} \in W^{1,2}(0, t; L^2(\Omega)) \cap L^\infty(0, t; H_0^1(\Omega))$.

Moreover, there exists a constant $R_1 > 0$, which depends on $\|\kappa_n\|_{C[0,T]}$ and $\|\kappa'_n\|_{L^1(0,T)}$, such that

$$\|n'\|_{L^2(Q(T))}^2 + \sup_{0 \leq s \leq t} \varphi^s(v; n(s)) \leq R_1 \left(1 + \|n_0\|_{H_0^1(\Omega)}^2 + \|G(\hat{n}, \Lambda(t)\hat{n})\|_{L^2(Q(T))}^2 \right),$$

$$\forall t \in [0, T], \quad \forall v \in \mathcal{V}(-\delta_0, t).$$

Lemma 4.1. There exist a constant $R_2 > 0$ and a non-negative, continuous and strictly increasing function $R_3(\cdot)$ on $[0, T]$ with $R_3(0) = 0$ such that

$$\begin{aligned} \|\Lambda(t)\hat{n}\|_{L^\infty(0,t; H^3(\Omega))}^2 &\leq R_2 \left(1 + \|f_0\|_{H^3(\Omega)}^2 \right) \\ &+ R_3(t) \left(1 + \|m_0\|_{H^1(\Omega)}^2 + \|\hat{n}\|_{L^\infty(0,T; H_0^1(\Omega))}^2 \right)^2, \quad \forall t \in [0, T]. \end{aligned}$$

Lemma 4.2. There exist a constant $R_4 > 0$ and a continuous, non-negative and strictly increasing function $R_5(\cdot)$ on $[0, T]$ with $R_5(0) = 0$ such that

$$\begin{aligned} \|G(\hat{n}, \Lambda(t)\hat{n})\|_{L^2(Q(T))}^2 &\leq R_4 \|\mu_n\|_{L^2(Q(T))}^2 \\ &+ R_5(t) \left(\|f_0\|_{H^3(\Omega)}^2 + 1 \right) \left(1 + \|m_0\|_{H^2(\Omega)}^2 + \|\hat{n}\|_{L^\infty(0,T; H_0^1(\Omega))}^2 \right)^3, \quad \forall t \in [0, T]. \end{aligned}$$

For each $t \in [0, T]$ and $v \in \mathcal{V}(-\delta_0, t)$, we define the solution operator $S(t, v)$ which assigns a unique solution $S(t, v)\hat{n} := n$ of $(AP)_{t,v,\hat{n}}$ to each $\hat{n} \in L^\infty(-\delta, t; H_0^1)$. We can apply Schauder fixed point theorem, we see that the operator $S(t, v)$ has at least one fixed point. Then, we give the existence theorem as follows:

Proposition 4.2. There exist a positive constant M_1 and a time $T_0 := T_0(M_1) \in (0, T]$ such that for each $v \in \mathcal{V}(-\delta_0, T_0)$ the problem

$$(AP)_v \begin{cases} n'(t) + \partial\varphi^t(v; n(t)) \ni G(t, n(t), [\Lambda(T_0)n](t)) & \text{in } L^2(\Omega), \quad \text{a.e. } t \in (0, T_0), \\ n(t) = n_0 & \text{in } L^2(\Omega), \quad \forall t \in [-\delta_0, 0]. \end{cases}$$

has a unique solution $n_v \in W^{1,2}(0, T_0; L^2(\Omega)) \cap L^\infty(0, T_0; H_0^1(\Omega))$ satisfying

$$\|n'_v\|_{L^2(Q_{T_0})}^2 + \sup_{0 \leq t \leq T_0} \varphi^t(v; n_v(t)) \leq M_1. \quad (4.1)$$

Moreover, there exists a positive constant M_2 , which is independent of $v \in V(-\delta_0, T_0)$, such that

$$\begin{aligned} & \|[\Lambda_1(T_0)n_v]'\|_{L^2(0,T_0;H^1(\Omega))} + \|\Lambda_1(T_0)n_v\|_{L^\infty(0,T_0;H^2(\Omega))} \\ & + \|\Lambda_1(T_0)n_v\|_{L^2(0,T_0;H^3(\Omega))} + \|\Lambda(T_0)n_v\|_{L^\infty(0,T_0;H^3(\Omega))} \leq M_2. \end{aligned} \tag{4.2}$$

Proof We fix T_0 , which is the same number as in Lemma 4.1, and $v \in V(-\delta_0, T_0)$ throughout this argument. Let $\{\hat{n}_k\} \subset \mathcal{W}_{M_1}$ and $\hat{n} \in \mathcal{W}_{M_1}$ so that $\hat{n}_k \rightarrow \hat{n}$ in $C([0, T]; L^2(\Omega))$ as $k \rightarrow \infty$. Then, we see that $G(\hat{n}_k, \Lambda(T_0)\hat{n}_k) \rightarrow G(\hat{n}, \Lambda(T_0)\hat{n})$ weakly in $L^2(Q(T_0))$ as $k \rightarrow \infty$. By using the results, we derive $S(T_0, v)\hat{n}_k \rightarrow S(T_0, v)\hat{n}$ in $C([0, T_0]; L^2(\Omega))$, so, $\mathcal{S}(T_0, v)\hat{n}_k \rightarrow \mathcal{S}(T_0, v)\hat{n}$ in $C([0, T]; L^2(\Omega))$ as $k \rightarrow \infty$.

By applying Schauder fixed point theorem, we see that $\mathcal{S}(T_0, v)$ has at least one fixed point \bar{n} , i.e., $\mathcal{S}(T_0, v)\bar{n} = \bar{n}$, in \mathcal{W}_{M_1} . It is clear from the definition of $\mathcal{S}(T_0, v)$ that \bar{n} is a solution of $(AP)_v$ on $[0, T_0]$.

In the rest of this proof, we show the uniqueness of solutions of $(AP)_v$ on $[0, T_0]$. Let n_i ($i = 1, 2$) be solutions of $(AP)_v$ on $[0, T_0]$. For simplicity, we put $\theta_i := \Lambda(T_0)n_i$ and $\zeta_i := \Lambda_1(T_0)n_i$.

First of all, we note that ζ_i ($i = 1, 2$) satisfies the following system:

$$(\zeta_1 - \zeta_2)' - \kappa_m \Delta(\zeta_1 - \zeta_2) + C_3(\zeta_1 - \zeta_2) = C_2(n_1 - n_2) \quad \text{a.e. in } Q(T_0), \tag{4.3}$$

$$\nabla(\zeta_1 - \zeta_2) \cdot \nabla \mathbf{n} = 0 \quad \text{a.e. on } \Sigma_{T_0}, \tag{4.4}$$

$$(\zeta_1 - \zeta_2)(0) = 0 \quad \text{a.e. in } \Omega. \tag{4.5}$$

We multiply (4.3) by $\zeta_1 - \zeta_2$ and $\nabla(4.3)$ by $\nabla(\zeta_1 - \zeta_2)$. By integrating these resultants over $Q(T_0)$, it is easily seen that there exists a constant $K_{22} > 0$ such that

$$\begin{aligned} & \sup_{0 \leq s \leq t} \|\zeta_1(s) - \zeta_2(s)\|_{H^1(\Omega)}^2 + \int_0^t \|\zeta_1(s) - \zeta_2(s)\|_{H^2(\Omega)}^2 ds \\ & \leq K_1 \int_0^t \|n_1(s) - n_2(s)\|_{H_0^1(\Omega)}^2 ds, \quad \forall t \in [0, T_0]. \end{aligned} \tag{4.6}$$

In order to show the uniqueness of solutions of $(AP)_v$ on $[0, T_0]$, we have to estimate the term $(G(n_1, \theta_1) - G(n_2, \theta_2), n_1 - n_2)_{L^2(\Omega)}$ by the following ways.

(1) It is easily seen from (4.2) that for any $\varepsilon_1 > 0$ there exists a constant $K_2(\varepsilon) > 0$ such that the following inequality holds for a.e. $t \in (0, T_0)$:

$$\begin{aligned} & \int_{\Omega} |\nabla n_1(x, t) - \nabla n_2(x, t)| |\nabla \theta_1(x, t)| |n_1(x, t) - n_2(x, t)| dx \\ & \leq \varepsilon_1 \|n_1(t) - n_2(t)\|_{H_0^1(\Omega)}^2 + K_2(\varepsilon_1) \|n_1(t) - n_2(t)\|_{L^2(\Omega)}^2. \end{aligned}$$

(2) By using (4.1), (4.2) and (4.6), we see that for any $\varepsilon_2 > 0$ there exists a constant

$K_3(\varepsilon_2) > 0$ such that the following inequality holds for a.e. $t \in (0, T_0)$:

$$\begin{aligned}
& \int_{\Omega} |\nabla n_2(x, t)| |\nabla \theta_1(x, t) - \nabla \theta_2(x, t)| |n_1(x, t) - n_2(x, t)| dx \\
& \leq C_1 \int_{\Omega} |\nabla n_2(x, t)| |\nabla f_0(x)| |n_1(x, t) - n_2(x, t)| \left(\int_0^t |\zeta_1(x, s) - \zeta_2(x, s)| ds \right) dx \\
& \quad + C_1^2 \int_{\Omega} |\nabla n_2(x, t)| |n_1(x, t) - n_2(x, t)| \\
& \quad \quad \times \left(\int_0^t |\zeta_1(x, s) - \zeta_2(x, s)| ds \right) \left(\int_0^t |\nabla \zeta_1(x, s)| ds \right) dx \\
& \quad + C_1 \int_{\Omega} |\nabla n_2(x, t)| |n_1(x, t) - n_2(x, t)| \left(\int_0^t |\nabla \zeta_1(x, s) - \nabla \zeta_2(x, s)| ds \right) dx \\
& \leq C_1 \|\nabla f_0\|_{C(\bar{\Omega})} \|n_2(t)\|_{H_0^1(\Omega)} \|n_1(t) - n_2(t)\|_{L^4(\Omega)} \int_0^t \|\zeta_1(s) - \zeta_2(s)\|_{L^4(\Omega)} ds \\
& \quad + C_1^2 \|n_2(t)\|_{H_0^1(\Omega)} \int_0^t \|\nabla \zeta_1(s)\|_{C(\bar{\Omega})} ds \\
& \quad \quad \times \|n_1(t) - n_2(t)\|_{L^4(\Omega)} \int_0^t \|\zeta_1(s) - \zeta_2(s)\|_{L^4(\Omega)} ds \\
& \quad + C_1 \sqrt{T_0} \|n_2(t)\|_{H_0^1(\Omega)} \|n_1(t) - n_2(t)\|_{L^4(\Omega)} \left(\int_0^t \|\nabla \zeta_1(s) - \nabla \zeta_2(s)\|_{L^4(\Omega)}^2 ds \right)^{\frac{1}{2}} \\
& \leq \varepsilon_2 \|n_1(t) - n_2(t)\|_{H_0^1(\Omega)}^2 + K_3(\varepsilon_2) \int_0^t \|n_1(s) - n_2(s)\|_{H_0^1(\Omega)}^2 ds.
\end{aligned}$$

(3) It is easily seen that for any $\varepsilon_3 > 0$ there exists a constant $K_4(\varepsilon_3) > 0$ such that the following inequality holds for a.e. $t \in (0, T_0)$:

$$\begin{aligned}
& \int_{\Omega} |\Delta \theta_1(x, t)| |n_1(x, t) - n_2(x, t)|^2 dx \\
& \leq \|\Delta \theta_1(t)\|_{L^4(\Omega)} \|n_1(t) - n_2(t)\|_{L^4(\Omega)} \|n_1(t) - n_2(t)\|_{L^2(\Omega)} \\
& \leq \varepsilon_3 \|n_1(t) - n_2(t)\|_{H_0^1(\Omega)}^2 + K_4(\varepsilon_3) \|n_1(t) - n_2(t)\|_{L^2(\Omega)}^2.
\end{aligned}$$

(4) It is easily seen that for any $\varepsilon_4 > 0$ there exists a constant $K_5(\varepsilon_4) > 0$ such that the

following inequality holds for a.e. $t \in (0, T_0)$:

$$\begin{aligned}
& \int_{\Omega} |n_2(x, t)| |\Delta \theta_1(x, t) - \Delta \theta_2(x, t)| |n_1(x, t) - n_2(x, t)| dx \\
\leq & C_1 \int_{\Omega} |\Delta f_0(x)| |n_1(x, t) - n_2(x, t)| \left(\int_0^t |\zeta_1(x, s) - \zeta_2(x, s)| ds \right) dx \\
& + 2C_1^2 \int_{\Omega} |\nabla f_0(x)| |n_1(x, t) - n_2(x, t)| \left(\int_0^t |\nabla \zeta_1(x, s)| ds \right) \\
& \quad \times \left(\int_0^t |\zeta_1(x, s) - \zeta_2(x, s)| ds \right) dx \\
& + 2C_1 \int_{\Omega} |\nabla f_0(x)| |n_1(x, t) - n_2(x, t)| \left(\int_0^t |\nabla \zeta_1(x, s) - \nabla \zeta_2(x, s)| ds \right) dx \\
& + C_1^3 \int_{\Omega} |n_1(x, t) - n_2(x, t)| \left(\int_0^t |\nabla \zeta_1(x, s)| ds \right)^2 \left(\int_0^t |\zeta_1(x, s) - \zeta_2(x, s)| ds \right) dx \\
& + C_1^2 \int_{\Omega} |n_1(x, t) - n_2(x, t)| \left(\int_0^t |\nabla \zeta_1(x, s)| ds + \int_0^t |\nabla \zeta_2(x, s)| ds \right) \\
& \quad \times \left(\int_0^t |\nabla \zeta_1(x, s) - \nabla \zeta_2(x, s)| ds \right) dx \\
& + C_1^2 \int_{\Omega} |n_1(x, t) - n_2(x, t)| \left(\int_0^t |\Delta \zeta_1(x, s)| ds \right) \left(\int_0^t |\zeta_1(x, s) - \zeta_2(x, s)| ds \right) dx \\
& + C_1 \int_{\Omega} |n_1(x, t) - n_2(x, t)| \left(\int_0^t |\Delta \zeta_1(x, s) - \Delta \zeta_2(x, s)| ds \right) dx \\
\leq & C_1 \|\Delta f_0\|_{L^4(\Omega)} \|n_1(t) - n_2(t)\|_{L^4(\Omega)} \int_0^t \|\zeta_1(s) - \zeta_2(s)\|_{L^2(\Omega)} ds \\
& + 2C_1^2 \|\nabla f_0\|_{C(\bar{\Omega})} \int_0^t \|\zeta_1(s)\|_{H^1(\Omega)} ds \\
& \quad \times \|n_1(t) - n_2(t)\|_{L^2(\Omega)} \int_0^t \|\zeta_1(s) - \zeta_2(s)\|_{H^2(\Omega)} ds \\
& + 2C_1 \|\nabla f_0\|_{C(\bar{\Omega})} \|n_1(t) - n_2(t)\|_{L^2(\Omega)} \int_0^t \|\zeta_1(s) - \zeta_2(s)\|_{H^1(\Omega)} ds \\
& + C_1^3 \left(\int_0^t \|\nabla \zeta_1(s)\|_{C(\bar{\Omega})} ds \right) \|n_1(t) - n_2(t)\|_{L^2(\Omega)} \int_0^t \|\zeta_1(s) - \zeta_2(s)\|_{L^2(\Omega)} ds \\
& + C_1^2 \left(\int_0^t \|\nabla \zeta_1(s)\|_{C(\bar{\Omega})} ds + \int_0^t \|\nabla \zeta_2(s)\|_{C(\bar{\Omega})} ds \right) \\
& \quad \times \|n_1(t) - n_2(t)\|_{L^2(\Omega)} \int_0^t \|\zeta_1(s) - \zeta_2(s)\|_{H^1(\Omega)} ds
\end{aligned}$$

$$\begin{aligned}
& +C_1^2 \left(\int_0^t \|\zeta_1(s)\|_{H^2(\Omega)} ds \right) \|n_1(t) - n_2(t)\|_{L^2(\Omega)} \int_0^t \|\zeta_1(s) - \zeta_2(s)\|_{C(\bar{\Omega})} ds \\
& +C_1 \|n_1(t) - n_2(t)\|_{L^2(\Omega)} \int_0^t \|\zeta_1(s) - \zeta_2(s)\|_{H^2(\Omega)} ds \\
& \leq \varepsilon_4 \|n_1(t) - n_2(t)\|_{H_0^1(\Omega)}^2 + K_5(\varepsilon_4) \int_0^t \|n_1(s) - n_2(s)\|_{H_0^1(\Omega)}^2 ds.
\end{aligned}$$

(5) It is easily seen that there exists a constant $K_6 > 0$ such that the following inequality holds for a.e. $t \in (0, T)$:

$$\begin{aligned}
& \int_{\Omega} \mu_n(x, t) [n_1(x, t) \{1 - n_1(x, t) - \theta_1(x, t)\} - n_2(x, t) \{1 - n_2(x, t) - \theta_2(x, t)\}] \\
& \quad \times \{n_1(x, t) - n_2(x, t)\} dx \\
& \leq \|\mu_n(t)\|_{L^\infty(\Omega)} \left(4\|n_1(t) - n_2(t)\|_{L^2(\Omega)}^2 + \int_{\Omega} |\theta_1(t) - \theta_2(t)| |n_1(t) - n_2(t)| dx \right) \\
& \leq K_6 \|\mu_n(t)\|_{L^\infty(\Omega)} \left(\|n_1(t) - n_2(t)\|_{L^2(\Omega)}^2 + \int_0^t \|\zeta_1(s) - \zeta_2(s)\|_{L^2(\Omega)}^2 ds \right).
\end{aligned}$$

We see from (1)–(5) that there exist constants $K_i > 0$ ($i = 7, 8$) such that

$$\frac{d}{dt} \Psi(t) \leq K_7 (1 + \|\mu_n(t)\|_{L^\infty(\Omega)}) \Psi(t), \quad \text{a.e. } t \in (0, T_0), \quad (4.7)$$

where

$$\Psi(t) := \|n_1(t) - n_2(t)\|_{L^2(\Omega)}^2 + K_8 \int_0^t \|n_1(s) - n_2(s)\|_{H_0^1(\Omega)}^2 ds.$$

By applying Gronwall lemma, we derive $n_1(t) = n_2(t)$ in $L^2(\Omega)$ for all $t \in [0, T_0]$, i.e., the uniqueness of solution of $(AP)_v$ on $[0, T_0]$. \blacksquare

4.2 Local existence of solutions

In this section, we state our main theorem of the present paper, which gives the existence of time-local solutions of (P), and show its proof.

Theorem 4.1. *(P) has at least one solution $[n, f, m]$ on $[0, T_0]$, where T_0 is the same time as in Proposition 4.2.*

Throughout this section, let M_1 and T_0 be the same constants as in Proposition 4.1. In order to show Theorem 4.1, we define a non-empty, closed and convex subset $\mathcal{W}_{M_1}(T_0)$ of $C([0, T_0]; L^2(\Omega))$, and an operator \mathcal{L} from $\mathcal{W}_{M_1}(T_0)$ into itself by

$$\mathcal{W}_{M_1}(T_0) := \left\{ v \in \mathcal{V}_{T_0}^+ \mid \|v'\|_{L^2(Q(T_0))} + \sup_{0 \leq t \leq T_0} \varphi_0(v(t)) \leq M_1 \right\}$$

and

$$\mathcal{L}v := n_v, \quad \forall v \in \mathcal{W}_{M_1}(T_0),$$

respectively. Actually, it is easily seen from Proposition 4.1 that the operator \mathcal{L} is well-defined on $\mathcal{W}_{M_1}(T_0)$.

Now, we give the proof of Theorem 4.1 below.

Proof of Theorem 4.1. Let $\{v_k\} \subset \mathcal{W}_{M_1}(T_0)$ and $v \in \mathcal{W}_{M_1}(T_0)$ so that

$$v_k \longrightarrow v \quad \begin{cases} \text{in } C([0, T_0]; L^2(\Omega)), \\ \text{weakly in } W^{1,2}(0, T_0; L^2(\Omega)), \\ \text{*weakly in } L^\infty(0, T_0; H_0^1(\Omega)) \cap L^\infty(Q(T_0)). \end{cases}$$

For simplicity, for each $k \in \mathbf{N}$ we put $\bar{n}_k := \mathcal{L}v_k$, $m_k := \Lambda_1(T_0)\bar{n}_k$ and $f_k := \Lambda(T_0)\bar{n}_k$. Then, it is easily seen from the definition of $\mathcal{W}_{M_1}(T_0)$ that there exist a subsequence of $\{k\}$, which is denoted by the same notation $\{k\}$, and $\bar{n} \in \mathcal{W}_{M_1}(T_0)$ such that the following convergences hold:

$$\bar{n}_k \longrightarrow \bar{n} \quad \begin{cases} \text{in } C([0, T_0]; L^2(\Omega)), \\ \text{weakly in } W^{1,2}(0, T_0; L^2(\Omega)), \\ \text{*weakly in } L^\infty(0, T_0; H_0^1(\Omega)) \cap L^\infty(Q(T_0)). \end{cases} \tag{4.8}$$

By using the continuity property of $\Lambda_1(T_0)$, we see that the following convergences hold:

$$m_k \longrightarrow \Lambda_1(T_0)\bar{n} \quad \begin{cases} \text{in } C([0, T_0]; H^1(\Omega)) \cap L^2(0, T_0; H^2(\Omega)), \\ \text{weakly in } W^{1,2}(0, T_0; H^1(\Omega)) \cap L^2(0, T_0; H^3(\Omega)), \\ \text{*weakly in } L^\infty(0, T_0; H^2(\Omega)). \end{cases} \tag{4.9}$$

By repeating the similar argument, we see that the following convergence holds:

$$G(\bar{n}_k, f_k) \longrightarrow G(\bar{n}, \Lambda(T_0)\bar{n}) \quad \text{weakly in } L^2(Q(T_0)). \tag{4.10}$$

In the rest of this proof, we show that n is a solution of $(AP)_v$ on $[0, T_0]$.

For this, we let z any function in $L^2(0, T_0; H_0^1(\Omega))$ satisfying $0 \leq z \leq 1 - \Lambda(T_0)v$ a.e. in $Q(T_0)$ and put $z_k := \min\{z, 1 - \Lambda(T_0)v_k\}$. Since z_k satisfies $0 \leq z_k \leq 1 - \Lambda(T_0)v_k$ a.e. in $Q(T_0)$, it is easily seen that the following inequality holds:

$$\begin{aligned} & \int_0^{T_0} (\bar{n}'_k(t), \bar{n}_k(t) - z_k(t))dt + \int_0^{T_0} \int_\Omega \kappa_n(t) \nabla \bar{n}_k(x, t) \cdot \nabla (\bar{n}_k(x, t) - z_k(x, t)) dxdt \\ & \leq \int_0^{T_0} (G(\bar{n}_k(t), f_k(t)), \bar{n}_k(t) - z_k(t))dt. \end{aligned} \tag{4.11}$$

By taking $\lim_{k \rightarrow \infty}$ in (4.11) and using (4.8)–(4.10) with $z_k \longrightarrow z$ in $L^2(0, T_0; H_0^1(\Omega))$, we see that the following inequality holds:

$$\begin{aligned} & \int_0^{T_0} (\bar{n}'(t), \bar{n}(t) - z(t))dt + \int_0^{T_0} \int_\Omega \kappa_n(t) \nabla \bar{n}(x, t) \cdot \nabla (\bar{n}(x, t) - z(x, t)) dxdt \\ & \leq \int_0^{T_0} (G(\bar{n}(t), [\Lambda(T_0)\bar{n}](t)), \bar{n}(t) - z(t))dt, \end{aligned} \tag{4.12}$$

which implies that \bar{n} is a solution of $(AP)_v$ on $[0, T_0]$, i.e., $\bar{n} = \mathcal{L}v$. Hence, we see that the operator $\mathcal{L} : \mathcal{W}_{M_1}(T_0) \longrightarrow \mathcal{W}_{M_1}(T_0)$ is continuous with respect to the strong topology of $C([0, T_0]; L^2(\Omega))$.

By applying Schauder fixed point theorem, we see that \mathcal{L} has at least one fixed point, namely, there exists $n \in \mathcal{W}_{M_1}(T_0)$ such that $\mathcal{L}n = n$. It is clear from the definition of $\Lambda(T_0)$ and $\Lambda_1(T_0)$ that a triplet $[n, \Lambda(T_0)n, \Lambda_1(T_0)n]$ is a solution of (P) on $[0, T_0]$. ■

References

- [1] Zuzanna Szymańska, Jakub Urbański and Anna Marciniak-Czochra, Mathematical modelling of the influence of heat shock proteins on cancer invasion of tissue, *J.Math.Biol.* (2009), Springer-Verlag
- [2] Mark A.J. Chaplain and Alexander R.A. Anderson, Mathematical Modelling of Tissue Invasion, *Cancer Modelling and Simulation, Chapter 10, A CRC Press UK 23 Blades Court Deodar Road London SW15 2NU UK.*
- [3] Risei Kano, Nobuyuki Kenmochi, Yusuke Murase, Nonlinear evolution equations generated by subdifferentials with nonlocal constraints, *Banach Center Publication, Volume 86, (2009), pp 175-194.*
- [4] R. Kano, N. Kenmochi and Y. Murase, Existence theorems for elliptic quasi-variational inequalities in Banach spaces, to appear.
- [5] R. Kano, N. Kenmochi and Y. Murase, Parabolic quasi-variational inequalities with non-local constraints, to appear.
- [6] Risei Kano, Applications of abstract parabolic quasi-variational inequalities to obstacle problems, *Banach Center Publication, Volume 86, (2009), pp 163-174.*
- [7] N. Kenmochi, Solvability of nonlinear evolution equations with time-dependent constraints and applications, *Bull. Fac. Education, Chiba Univ.*, 30(1981), 1-87.
- [8] N. Kenmochi, *Monotonicity and Compactness Methods for Nonlinear Variational Inequalities*, Chapter 4 (pp.203-298) in *Handbook of Differential Equations, Stationary Partial Differential Equations, Vol.4*, Elsevier, Amsterdam-Boston-Heidelberg-London-New York-Oxford-Paris-SanDiego-San Francisco-Singapore-Sydney-Tokyo, 2007.
- [9] Y. Murase, Abstract quasi-variational inequalities of elliptic type and applications, *Banach Center Publications, Inst. Math. Polish Acad. Sci.*, 86 (2009), 235-246.