

OBSERVATIONS ON THE TAXONOMY AND DISTRIBUTION OF
CHAETOGNATHS OF THE NORTH PACIFIC¹⁾

TAKASI TOKIOKA

Seto Marine Biological Laboratory, Sirahama

With 35 Text-figures and 74 Tables

CONTENTS

	Page
Introduction	350
Material and methods	351
Taxonomy	352
1. <i>Sagitta ferox</i> f. <i>americana</i>	353
2. <i>Sagitta tenuis</i> and <i>Sagitta friderici</i>	358
3. <i>Sagitta serratodentata tasmanica</i>	364
4. <i>Sagitta neglecta</i> and its allies	372
5. <i>Sagitta neodecipiens</i>	377
6. <i>Sagitta bedoti</i> f. <i>minor</i>	381
7. <i>Sagitta hexaptera</i>	382
8. <i>Sagitta enflata</i> f. <i>gardineri</i>	383
9. <i>Sagitta lyra</i>	385
10. Other species of <i>Sagitta</i>	385
11. <i>Krohnitta subtilis</i> and <i>pacifica</i>	386
Distribution and faunistic notes	387
1. Lagoons of Baja California	387
2. The inlet waters along the Pacific coast of Japan	389
3. <i>Elegans</i> -water in the North Pacific	389
4. Mixing area of <i>elegans</i> -water and the warm water	389
5. The blue-green water along southern California	390
6. <i>Bedoti</i> -area	
I. Occurrence of <i>Sagitta bedoti</i> in the eastern Pacific	391
II. Occurrence of <i>Sagitta bedoti</i> at the western stations near Japan	392
III. Other available data concerning the occurrence of <i>S. bedoti</i>	395
IV. The identity of <i>S. bedoti</i> from various localities	413
V. General account of the distribution of <i>S. bedoti</i>	414
VI. Possible factors limiting the distribution of <i>S. bedoti</i>	417
VII. " <i>Bedoti</i> -water"	418
7. The far oceanic water of the Central North Pacific	419
a. Distribution of S/E in collections examined	422
b. Distribution of S/E in other parts of the Indo-Pacific based on available data	426
c. Distribution of H/E	442
d. Distribution of P/E	443

1) Contributions from the Seto Marine Biological Laboratory, No. 339.

e. Distribution of other prominent species	443
8. Differentiation found in the chaetognath fauna of the North Pacific and the relation between communities and water masses	444
Summary and conclusions	449
References	450
Appendix. Station data 1-3.....	454

Introduction

Although I had a chance to look through a part of vast collection of plankton samples kept at the Scripps Institution of Oceanography and to examine a considerable number of specimens of chaetognaths during my stay there in 1956—57, I have refrained from publishing papers on this animal group, because I knew at that time that Mr. R. BIERI had already accumulated a great many data on chaetognaths during his long stay at Scripps and had begun to prepare manuscripts for publication. Now, two of his papers (1957 and 1959) have been published on the Pacific chaetognaths. The second paper is especially very important in giving a comprehensive outline of the distribution of different species in the North Pacific. Previous works on Pacific chaetognaths are all covered in the second paper. Five subsequent papers on the Pacific chaetognaths have appeared, two by FAGETTI (1958 a, b) on material from Chilean waters and three by SUND (1959 a, b, c) on material from the Gulf of Alaska and eastern tropical waters.

My data shown in the tables appended at the end of this article appear to contribute only supplemental information to the distribution-charts of the respective species as given by BIERI (1959), but the results of closer morphological examinations on the material comprise some new findings. A new species and a new form was found in the material from the eastern tropical Pacific. The *tenuis-friderici* problem is discussed again on Pacific specimens, the range of morphological variations is reviewed in some species and a few morphological notes on some additional species are presented. In addition, the relative abundance of the four important species (*Sagitta enflata*, *S. serratodentata pacifica*, *S. hexaptera* and *Pterosagitta draco*) is defined by computing ratios between *S. enflata* and each of *serratodentata pacifica*, *hexaptera* and *P. draco*. This procedure seems to indicate a relation between the chaetognath fauna and some water-masses more clearly than previously shown. The distribution of *Sagitta bedoti* in the western Pacific and the Indian Ocean is reviewed in detail on the basis of present material and also on previous records published to date. This leads to the conclusion that a characteristic water mass of comparatively large extent, the *bedoti*-water, can be defined distinctly in the Indo-Pacific.

Before going further, I want to express my hearty thanks to the staff of the Scripps Institution of Oceanography for granting me the favour of a seat at the laboratory for a year, especially to Prof. M. W. JOHNSON for his kindness in providing facilities and for reading the manuscript; and also to the Rockefeller Foundation for a postdoctoral fellowship.

Material and Methods

The material examined consists of the following 326 plankton samples:

1. Transpac Expedition 1953: 130 samples collected from 50 m to the surface by means of a small net having a 17 cm diameter mouth; 19 samples collected by a Japanese survey ship off north eastern Honsyû, Japan were presented to the research party of the Expedition. These samples were taken vertically from 50 m to the surface by using KITAHARA's quantitative net, a small modified HENSEN's net. A sample of the North Pacific Stranger Expedition (1955) was examined and the result is given at the end of the table of the Transpac Expedition.
2. Midpac Expedition 1950: 57 samples taken by various gears and also from various depths.
3. Shellback Expedition 1952: 48 samples, one of every five stations, were examined. The samples were collected with a large net from about 300 m to the surface.
4. Equapac Expedition 1956: Five of many samples were examined; they were taken with a large net from 200-400 m to the surface.
5. The blue-green water survey: 54 samples taken with a small net, having a 17 cm diameter mouth; of these 35 were collected off San Diego, and 19 were collected March 12-13, 1956 in the area extending from San Diego to Long Beach.
6. Lagoon samples: Ten samples from Scammons lagoon, Oct. 9-12, 1955 and two samples from Manuela lagoon, Oct. 12, 1952. All these samples were collected by members of the Foraminifera Laboratory of the Scripps Institution of Oceanography under the direction of Prof. F. B. PHLEGER. I am very grateful to Prof. PHLEGER and the members of the Laboratory for their kindness in allowing me to examine these samples and to quote some of the hydrographical data in this paper.
7. In addition, the data obtained by examining the samples collected during the Palao-New Guinea cruises 1940 made by the former Japanese fisheries experimental station at Palao are given herewith.

In treating the samples collected with small nets, the whole of each sample was examined with a magnification of $30\times$. Every individual of each species was identified carefully and counted. In treating the large net samples, however, only one fourth of each sample was examined in this way. The crude data, thus obtained, are arranged in 8 tables appended at the end of this paper. In measuring the body length or the length of the caudal segment, the caudal fin was always included. When the teeth were counted, the anterior portion of the head was dissected off under the binocular, the dental area was mounted and examined under the microscope. The corona ciliata was observed by staining it with rose bengal; the existence of the intestinal diverticula was ascertained by making the specimens transparent with glycerine and examining them under the microscope. The arrangement of fin rays and the number of eggs were determined under the microscope. When it was necessary to measure some small distances, the objects were drawn on a sheet of paper by using a camera lucida and the measurements were made from these sketched

figures. In computing percentages, ratios and means, fractions less than 1/2 were disregarded and the rest were counted as units.

Taxonomy

In all, the following 30 forms were identified in the whole series of samples :

1. *Sagitta hexaptera* D'ORBIGNY
2. *Sagitta lyra* KROHN
3. *Sagitta enflata* GRASSI
4. *Sagitta enflata* f. *gardineri* DONCASTER
5. *Sagitta elegans* VERRILL
6. *Sagitta bipunctata* QUOY et GAIMARD
7. *Sagitta ferox* DONCASTER
8. *Sagitta ferox* f. *americana* nov.
9. *Sagitta robusta* DONCASTER
10. *Sagitta bedoti* BÉRANECK
11. *Sagitta pulchra* DONCASTER
12. *Sagitta tenuis* CONANT
13. *Sagitta friderici* RITTER-ZÁHONY
14. *Sagitta serratodentata pacifica* TOKIOKA
15. *Sagitta serratodentata pseudoserratodentata* TOKIOKA
16. *Sagitta serratodentata tasmanica* THOMSON
17. *Sagitta neglecta* AIDA
18. *Sagitta oceania* GRAY
19. *Sagitta regularis* AIDA
20. *Sagitta bedfordii* DONCASTER
21. *Sagitta crassa* f. *naikaiensis* TOKIOKA
22. *Sagitta minima* GRASSI
23. *Sagitta decipiens* FOWLER
24. *Sagitta neodecipiens* n. sp.
25. *Sagitta planctonis* STEINHAUS
26. *Sagitta macrocephala* FOWLER
27. *Pterosagitta draco* (KROHN)
28. *Eukrohnia hamata* (MÖBIUS)
29. *Krohnitta subtilis* (GRASSI)
30. *Krohnitta pacifica* (AIDA)

Most of these species have been described repeatedly and a key to the tropical eastern Pacific Ocean species has been published by SUND (1959b). Hence, only descriptions of two new forms and some notes on problematic forms or on the range of variations will be given together with a few morphological notes. Three subspecies of *Sagitta serratodentata* KROHN are printed in tables as species in order to save space.

1. *Sagitta ferox* f. *americana* nov.

(Figs. 1-3, Tables 1-4)

A medium-sized species attaining 8.3 mm in body length of which the tail segment constitutes most frequently 27-29%. General appearance of the body and the relative size of the head resemble closely those of the typical *S. ferox*, from which it is very difficult to separate even under considerably high magnification. The head is medium in size, not so large as in *S. robusta*; neck distinct, no constriction at the trunk-tail septum. The collarette is fairly conspicuous at the neck region and reaches to the anterior end of the ventral ganglion, diminishing in thickness posteriorly. The anterior fin begins at the posterior end of the ventral ganglion and without any rayless zone. The posterior fin is a little longer than the anterior and more than half of the fin lies on the tail segment, being widest at the level slightly posterior to the trunk-tail septum; rayless zone is practically absent, except for a very narrow rayless area in front of the aperture of the female genital organ in some individuals. The distance between the anterior and posterior fins is approximately one fourth of the posterior fin. Eye-pigment is rather large, slightly elongate and curved. The corona ciliata (Fig. 1) begins at the posterior margin of the brain in front of eyes, it is about one and a half times as long as the head, and both sides are slightly sinuous; this is slightly shorter than in *S. robusta*. The intestinal diverticula are present




Fig. 1. 1—*Sagitta ferox* f. *americana* n. f. Corona ciliata (SB 20).

and very distinct. Hooks are most frequently 7-8 in adults, though they fluctuate from 6 to 9. The anterior teeth are slender and only slightly fewer than the posterior teeth which are much thicker and usually 10-12 in number (fluctuate from 9 to 14) in mature individuals. Frequently the anterior teeth may equal, or in some cases, exceed the posterior teeth in number. This was observed only in mature specimens larger than 6.7 mm (in 12 of 22 specimens examined longer than 6.7 mm). The difference in number between the anterior and posterior teeth is larger in smaller immature individuals than in mature ones. Rows of the anterior teeth usually meet




Fig. 2. *Sagitta ferox* f. *americana* n.f. Seminal vesicles arranged in the order of maturity. $\times 125$; 2-4—(SB 20), ventral; 5-6—(SB 60), ventral; 7—(SB 20), ventral; 8—6.8 mm long individual (SB 210), 9—ruptured (SB 60), ventral; 10—ruptured (SB 60), dorsal.

in an acute angle, but in obtuse angles on some widely expanded heads. Both hooks and teeth are brownish. Fully mature ovaries reach near the neck. The seminal vesicle (Fig. 2) is situated very near the rear end of the posterior fin, being slightly apart from the base of the tail fin. The distance between the vesicle and the tail fin is, however, very small and also very variable. In some cases, it is practically indiscernible, while in others it may reach up to 24% of the length of the vesicle; thus the ratio, the distance between the vesicle and the tail fin/the vesicle length, fluctuates from 0 to 0.24, being 0.14 on an average of 12 measured individuals. In structure the vesicle consists of a glandular anterior portion and a voluminous posterior sperm sac which can be seen clearly when the specimen is examined in the light field under the microscope. But externally, the distinction between the anterior and posterior portions of the vesicle is indistinct. Thus, the situation and the outline of the vesicle resemble more closely those of *S. robusta* rather than those of *S. ferox*, although the fully mature vesicles are more voluminous than in *S. robusta*. Rupture occur at the antero-lateral side. The seminal vesicle of typical *ferox* usually touches both the posterior and tail fins (the distance between the vesicle and tail fin/the vesicle length may reach 0.12 in some cases) and consists of a very conspicuous anterior glandular portion and a posterior saccular portion. There is also a swelling of the frothy tissue just in front of the vesicle in perfectly preserved specimens (Fig. 3); this swelling is quite absent in the present form and also in *S. robusta*. The ovary reaches to the anterior end of the posterior fin in a 5.5 mm long individual, to the ventral ganglion in a 6.2 mm long individual and near the neck in a 6.4 mm long individual. Fully swollen seminal vesicles are found on a 6.2 mm long individual.

Remarks: The armature formulae of *S. robusta* are closely related to those of the present form, although the hooks are most frequently 6 (5-7) in mature *robusta*




Fig. 3. *Sagitta ferox* DONCASTER. Seminal vesicle of a 6.7 mm long individual. $\times 125$ (EQP H-3)

and the difference in number between the anterior and posterior teeth is slightly larger in *robusta* than in the present form; besides both hooks and teeth of *robusta* yellowish rather than brownish. The shape and situation of the seminal vesicle of *robusta* resemble closely those of the present form. In spite of these resemblances, these two forms can be separated from each other distinctly by the size of the head which is exceedingly larger in *S. robusta*. *Sagitta hispida* CONANT from the Caribbean Sea, especially some of the specimens described by SUAREZ (1955) from Cuban waters, is also closely related to the present form, the number of hooks and teeth

1 *Sag ferox f americana*

No	body length	tail seg. in %	hook	ant. teeth	post. teeth	maturity	loc
1	3.6mm	33.9	9-9	4-5	6-6	im.	SB75
2	4.2	31.3	9-9	1+*-5	8-8	im.	SB75
3	4.5	29.0	7-8	4-5	8-8	im.	SB68
4	5.5	27.1	7-7	8-8	10-10	med.	SB55
5	5.9	28.6	6-7	7-7	11-12	med.	SB55
6	5.9	27.8	7-7	7-7	10-10	med.	SB75
7	6.0	26.9	7-7	7-7	9-10	med.	SB64
8	6.2	29.5	7-7	9-9	10-12	m.	SB64
9	6.2	26.3	7-7	7-7	9-10	m.	SB210
10	6.4	30.6	6-7	8	9	m.	SB30
11	6.6	26.7	6-6	7-8	9-10	m.	SB55
12	6.7	29.1	7-7	10-11	10-10	m.	SB60
13	6.8	28.6	8-8	10-10	10-10	m.	SB210
14	6.8	26.7	7-7	12-13	10-10	med.	SB71
15	6.9	28.3	7-7	10-10	10-11	m.	SB60
16	7.0	26.9	7-7	12-12	12-13	m.	SB60
17	7.2	29.1	8-8	9-9	11-12	m.	SB64
18	7.3	28.6	7-8	9-7	9-11	m.	SB64
19	7.3	26.5	6-6	8-8	9-10	m.	SB71
20	7.3	26.5	7-8	9-11	10-10	m.	SB71
21	7.5	26.1	7-7	5+*-10	10-10	m.	SB64
22	7.5	26.1	7-7	3+*-9	5+*-10	med.	SB64
23	7.6	29.9	7-7	7	12	m.	SB30
24	7.7	27.7	7-7	9-9	11-12	m.	SB60
25	7.7	28.6	7-7	9-10	12-13	m.	SB71
26	7.7	27.7	8-8	10-14	11-11	m.	SB75
27	7.8	30.8	7-7	9	11	m.	SB30
28	7.8	27.5	7-7	10-11	10-10	m.	SB68
29	7.9	28.1	7-7	10-11	10-11	m.	SB68
30	7.9	25.6	8-8	9-9	10-10	m.	SB160
31	8.0	27.6	7-8	10-11	11-11	m.	SB68
32	8.2	28.6	6-7	8-8	10-10	m.	SB175
33	8.3	27.6	7-7	10-10	13-14	m.	SB175

Table 1. Armature formulae of *S. ferox f. americana* n.f.

* Some torn out. SB...Shellback Exp.

seems to fall in the range of the variations found in the present form, although the anterior teeth are always slightly fewer than the posterior ones. Mature *S. hispida* are, however, larger than mature specimens of the present form, the ovary does not reach beyond the frontal end of the anterior fin and the shape of the seminal vesicle resembles that of *S. bedoti* and is separated from the caudal fin by about half the length of the vesicle. Although the general appearance of *S. ferox* resembles that of the present form, the anterior teeth are much fewer by about one half than

2 *Sag. ferox*

No.	body length	tail seg. in %	hook	ant. teeth	post. teeth	maturity	loc
1	6.7mm	29.1	8—8	6—7	14—14	m.	EQPH3
? 2	7.8	28.3	8	6	14	im.	SB 30
3	8.5	30.8	9—9	6—7	14—14	m.	EQPH17

3 *Sag. robusta*

No	body length	tail seg. in %	hook	ant. teeth	post. teeth	maturity	loc
1	4.6mm	28.6	7—7	5—5	7—8	im.	SB 75
2	6.3	27.8	7—7	5—5	8—8	im.	SB 75
3	6.5	31.0	6—7	6—6	9—11	im.	SB 71
4	8.5	29.0	7—7	7—8	12—12	im.	SB 64
5	8.5	27.5	6—6	8—8	12—12	im.	SB 95
6	10.4	25.0	6—6	7—8	12—12	m.	SB 71
7	11.1	29.4	6—6	7—8	12—12	m.	SB 64
8	15.0	26.1	6—6	6—7	10—10	m.	SB 105

4 *Sag. robusta* collected by Paul N. Sund (Scot sta. 23)

No	body length	tail seg. in %	hook	ant. teeth	post. teeth	maturity
1	9.5mm	28.4	6—6	8	13	im
2	11.5	26.1	7—7	10	13	im
3	12.0	27.5	6—6	10	13	im
4	12.7	29.9	6—6	10	13	im

Tables 2-4. Armature formulae of *S. ferox* (2) and *S. robusta* (3-4).
EQP...Equapac Exp., SB...Shellback Exp.

the posterior teeth. The structure and situation of the seminal vesicle also differ slightly. Most probably, *S. robusta*, *hispida*, *ferox* and the present form have differentiated from a common ancestor and are very closely related. At present, I can not judge whether the present form is a distinct species or a variety or merely a form of any already known species. Here I can only treat the present form provisionally

as a local form specialised from *S. ferox* and restricted to the Shellback area*, since the general appearance of both forms resemble each other so closely. *Sagitta helenae* RITTER-ZAHONY conforms to the present form in that the anterior teeth are numerous and usually exceed the posterior ones, but the difference between the two is very distinct, because the former, *S. helenae*, is quite devoid of the intestinal diverticula.

2. *Sagitta tenuis* CONANT and *Sagitta friderici* RITTER-ZAHONY

(Figs. 4-5, Tables 5-8)

Specimens of *S. tenuis* from Scammons and Manuela lagoons (Fig. 4) are less than 10.7 mm in body length and the tail segment occupies about 30% of the total body length. The body is rather rigid and provided with many tactile setae distributed over the body surface and thus assumes a hispid appearance. Collarete is

5 *Sag tenuis* from Scammons Lagoon

No	body length	tail seg in %	hook	ant teeth	post. teeth	maturity	TC
1	3.8mm	31.0	8—9	3—4	6—6	im.	66.7
2	4.4	26.5	8—9	3—3	4—4	im.	83.3
3	5.3	32.1	8—8	4—4	7—7	med.	65.8
4	5.3	26.5	8—8	3—4	7—7		64.1
5	5.5	31.8	8—8	4—4	7—7	med.	63.4
6	5.7	34.7	8—8	4—4	6—7		63.3
7	6.0	32.6	8—8	4—4	8—8	m.	69.0
8	6.2	29.5	8—8	5—5	8—8	med.	71.8
9	6.2	30.5	8—8	5—5	9—9	med.	68.9
10	6.2	31.3	7—8	5—5	10—10	med.	60.9
11	6.4	32.3	8—8	5—5	9—9	med.	60.4
12	6.6	30.7	8—8	5—5	9—10	med.	66.7
13	7.0	30.6	8—8	5—5	10—10	m.	66.7
14	7.1	29.4	8—8	4—4	11—11	med.	73.3
15	7.2	25.5	8—9	5—5	9—9	med.	62.5
16	7.2	29.1	7—7	4—5	10—11	m.	70.0
17	7.4	26.3	7—8	5—5	11—11	m.	62.5
18	7.5	29.6	8—8	6—6	13—13	m.	84.4
19	7.5	30.4	7—7	5—5	10—10	m.	73.3
20	7.5	28.4	8—8	5—5	11—11	m.	70.0
21	7.8	30.1	8—8	5—6	12—12		70.2
22	7.8	29.2	7—8	5—5	11—11	m.	63.6
23	8.6	28.9	9—9	7—7	14—14	m.	66.9
24	10.7	29.0	8—8	5—5	12—12	m.	91.5

Table 5. Armature formulae of *S. tenuis* from Scammons lagoon.

* The area surveyed by the Shellback Expedition 1952, see Fig. 25.

distinct near the neck. The anterior fin begins usually at the posterior end of the ventral ganglion, but in larger individuals there is a space of about one half of the ganglion length between the frontal end of the fin and the posterior end of the ganglion. Fin rays are all arranged obliquely towards the posterior; rayless-zone absent. The posterior fin is nearly as long as the anterior and widest at the level posterior to the trunk-tail septum; fin rays are arranged as on the anterior fin, although those near the frontal end are vertical to the base of the fin or even arranged slightly obliquely towards the anterior. Eye-pigments roundish. The corona ciliata is about twice the head length, with two shallow sinuses along each side. The intestinal diverticula absent. The ovary extends slightly beyond the anterior end of the posterior fin at maturity. Ova are usually arranged in two, dorsal and ventral, rows in each ovary and mostly 14-20 in number in each, the maximum observed was 24 in the 10.7 mm long individual. The seminal vesicle is elongate, rather low and with the anterior glandular portion slightly swollen. It touches both the posterior and caudal fins. The ratio TC (anterior part of the posterior fin along the trunk/posterior part of the posterior fin along the caudal segment $\times 100$) varies from 60.4 to 91.5, being 69.1 on an average of 24 specimens. The above-mentioned features together with the armature formulae (Table 5) evidently refers the specimens to *Sagitta tenuis*. Rows of anterior teeth meet each other in acute angles.

Other chaetognaths found in the blue-green water along the southern Californian coast and in some samples of the Transpac and Shellback Expeditions, collected at stations near this blue-green water region, seem to belong to *Sagitta friderici*. They resemble closely *S. tenuis*, but differ from that species in the following points :-




Fig. 4. *Sagitta tenuis* CONANT.
A 8.6 mm long individual from Scammons Lagoon. 12—Dorsal side of the animal, $\times 20$; 13—Corona ciliata, 14—Seminal vesicle.

1) mature individuals are larger, attaining 12.8 mm in length; tail segment is slightly shorter, 26.2% on an average of 34 measured individuals; 2) both the anterior and posterior teeth are fewer than in *S. tenuis* of the same body length; and 3) TC value is much larger, varying from 76 to 141.7 and 106 on an average of 33 measured specimens.

In addition, there are a number of specimens from offshore waters in the Shell-

6

No.	body length	tail seg. in %	hook	ant. teeth	post. teeth	mat.	TC	loc.
1	3.7mm	28.1	9-9	2-2	3-3	im.	76.0	TP
2	4.2	30.8	9-9	3-3	4-4	im.	80.0	TP
3	4.4	26.9	9-9	2+*-3	5-5	im.	82.8	TP
4	4.9	28.9	8-8	2-3	4-4	im.	85.7	SD
5	5.1	28.2	8-8	3-?	5-5	im.	88.2	TP
6	5.1	30.4	8-9	2-3	5-5	im.	89.9	SB 5
7	5.2	27.5	8-9	3-3	4-4	im.	83.3	SD
8	5.3	28.0	9-9	2-3	5-5	im.	91.4	TP
9	5.3	26.8	8-9	3-3	7-7	im.	88.2	TP
10	5.5	27.1	9-9	2-3	5-5	im.	94.3	TP
11	5.7	28.4	7-8	3-3	5-5	im.	101.6	SB 5
12	5.8	29.2	9-9	2-3	6-7	im.	89.7	SB 5
13	5.9	27.8	8-9	1+*-3	7-7	im.		SB 5
14	6.0	25.8	9-9	2-2	5-5	im.	100.0	SD
15	6.3	25.7	6-7	3-4	7-8	im.	122.8	SB 5
16	6.5	27.0	7-8	5-5	7-7	im.	87.1	SB 5
17	6.5	30.0	6-7	5-5	8-8	im.	100.0	SB 5
18	6.5	26.0	7-7	4-4	7-7	im.	106.3	SB 5
19	6.6	27.4	7-8	3-3	7-7	im.	108.6	SB 5
20	6.7	24.6	9-9	2-3	6-6	im.	105.3	TP
21	7.1	26.6	7-7	5-5	9-9	im.	133.3	SB 85
22	7.2	23.4	8-9	3-3	5-5	im.	119.0	SD
23	7.3	23.2	8-8	4-4	8-8	im.	105.0	TP
24	7.5	26.1	10-10	3-4	8-8	im.	119.0	TP
25	7.5	26.1	8-9	4-4	7-7	im.	114.3	TP
26	9.4	23.6	7-9	4-5	8-9	med	134.0	SD
27	10.4	25.0	8-8	5-5	11-11	med	121.8	TP
28	10.6	23.9	8-8	4-5	9-10	med	120.4	SD
29	11.0	24.9	8-8	5-6	11-11	med	100.0	SD
30	11.5	23.2	8-8	6-6	13-13	med	125.0	SB 1
31	11.6	22.5	8-8	6-6	11-11	med	136.4	SB 1
32	11.6	24.2	8-9	6-6	12-12	med	114.1	SD
33	12.0	22.7	8-9	7-7	12-14	med	133.3	SB 1
34	12.8	20.8	8-8	4-4	11-12	med	141.7	SB 1

Table 6. Armature formulae of *S. friderici* from the blue-green water.
* Some torn out. TP...Transpac Exp., SB...Shellback Exp.,
SD...San Diego.

back samples, which superficially closely resemble *S. friderici*. Close examination reveals that they do not agree in the armature formulae with the typical *S. friderici* collected from the San Diego region. The anterior and posterior teeth are more abundant in these. Roughly speaking, these chaetognaths can be divided into two groups, one with 7-8 (or 9) hooks (group A) and the other with 5-6 hooks (group B). The armature formulae of the former (group A) resembles somewhat those of *S. tenuis*. The ratio TC is relatively small, varying from 80 to 91.1 and 86.3 on an average of eleven measured specimens, while in group B it is slightly higher, varying from 88.4 to 114.3 and 96.6 on an average of 13 measured specimens. When the

7 Shellback specimens allied to *S. friderici*

No.	body length	tail seg. in %	hook	ant. teeth	post teeth	mat.	T C	loc.
1	6.2 mm	29.5	7—7	6—6	10—10	im.	85.7	SB 112
2	6.4	25.5	7—9	6—6	14—14	im.		SB 109
3	7.2	27.3	6—6	7—7	14—14	med	88.4	SB 100
4	7.5	27.0	6—6	7—7	14—14	med.	97.5	SB 5
5	7.5	26.1	5—6	7—7	14—14	m.	95.0	SB 90
6	7.6	27.4	7—8	6—6	12—12	im.		SB 5
7	7.7	28.0	6—6	7—7	14—14	im.		SB 109
8	7.7	28.6	6—6	7—7	15—17	med.		SB 95
9	7.8	29.2	6—6	7—8	14—15	m.	100.0	SB 85
10	7.8	28.3	5—6	8—8	15—16	m	93.3	SB 90
11	7.9	26.2	6—6	6—6	13—13	im.	100.0	SB 105
12	8.1	25.6	6—6	7—7	16—16	m	107.5	SB 50
13	8.1	27.2	6—6	6—6	13—13	im.	93.0	SB 105
14	8.1	27.2	8—8	7—7	15—15	im.	86.0	SB 112
15	8.1	28.0	6—6	8—8	15—16	m		SB 85
16	8.3	29.1	6—6	7—7	15—15	med.	88.9	SB 105
17	8.3	25.8	7—7	4+*-6	12—13	im.	91.1	SB 105
18	8.5	26.9	6—6	7—7	14—14	m		SB 5
19	8.5	29.2	6—6	7—7	16—17	m	91.3	SB 105
20	8.7	29.1	6—6	6—7	13—14	im.	90.9	SB 105
21	8.8	26.5	6—6	7—8	16—16	m.	96.2	SB 1
22	8.8	29.6	6—6	7—7	15—16	im.		SB 90
23	9.4	28.5	6—6	8—9	18—19	im.	114.3	SB 109
24	10.0	26.6	8—8	8—9	16—18	im.	83.3	SB 118
25	11.1	24.1	8—8	5+*-9	17+*-19	m.	89.2	SB 109
26	11.4	27.4	8—8	9—9	17—18	med	88.7	SB 122
27	11.7	25.6	8—8	7—11	20—22	med	81.3	SB 109
28	12.0	26.1	8—8	6+*-11	21—23	m.	90.7	SB 109
29	13.0	25.0	8—8	10—10	21—22	m.	80.0	SB 109
30	13.2	26.1	8—8	10—11	24—24	m.	87.5	SB 109
31	13.8	25.5	7—8	10—10	25—25	med	85.7	SB 109

Table 7. Armature formulae of the Shellback specimens allied to *S. friderici*.

* Some torn out. SB—Shellback Exp.

ratios TC of all of the above-mentioned forms, of course including both typical *S. tenuis* and *S. friderici*, are plotted on a graph (Fig. 5) and the relative position of values in respective groups are compared, then the idea that group A with 7-8 hooks may be identified as *S. tenuis*, seems possible. Theoretically it is not impossible that *S. tenuis* propagated in some embayments might be drifted out, the attainment of sexual maturity in these might be much retarded by environmental changes so they might grow to such very large size as 13.8 mm, which is evidently an exceptional length for *S. tenuis*. The increase of the body length in waters of lower temperature is not rare in species of chaetognaths. The anterior and posterior teeth might increase




Fig. 5. Ratio TC (anterior part of the posterior fin along the trunk/posterior part of the posterior fin along the caudal segment $\times 100$) of *Sag. tenuis*, *Sag. friderici* and their allied forms from the Shellback area.

with body length. A flow of water out into the open sea from embayments in tropical area usually results in a remarkable drop in temperature. However, there remains still a gap of considerable width on the graph between the group of the typical *S. tenuis* and that of group A. *Sagitta bipunctata* also resembles superficially group A. But it has usually 8-9 (though fluctuating from 7 to 10) hooks and slightly fewer anterior and posterior teeth (Table 8), and also somewhat elongate eye-pigments. In the armature formulae given by SUND (1959b), the number of hooks seems to

be rather fewer than usual. The body is slightly stouter than in *friderici* and the collarette seems to be more conspicuous. For a detailed comparison between these two species, the comprehensive work done by FAURE (1952) is available. *Sagitta bedoti* is also superficially related closely to group A, but this species is most frequently provided with 7 hooks and its fins are marked with conspicuous rayless zones. A careful comparison was also made between group A and *Sagitta crassa* f. *naikaiensis*, although the locality of the latter is far removed from the Shellback area. However, the absence of intestinal diverticula in the former denies absolutely a close relationship between these two forms. The distance between the eyes is relatively large in both groups A and B, distance/head width $\times 100$ is 36.2-47.5 (42.3 on an average of 6 measured specimens) in 8.1-13.8 mm long individuals and this feature seems to increase the superficial similarity between forms A and B and *S. bedoti* or *S. crassa*.

8 *Sagitta bipunctata*

No.	body length	tail seg in %	hook	ant teeth	post. teeth	maturity	loc
1	3.9mm	33.3	9-9	3-3	7-7	im.	SB 75
2	3.9	30.0	9-9	5-5	8-8	im.	SB 60
3	6.8	27.6	8-9	7-7	14-15	im.	SB 75
4	7.2	27.3	8-9	5-6	11-11	im.	SB 25
5	8.1	28.0	9-9	6-6	12-12	med.	SB 64
6	9.2	26.2	9-10	7-7	3-15	med.	SB 60
7	9.5	29.5	10-10	7-8	14-15	m.	SB 71
8	9.8	26.7	9-9	6-7	15-16	m.	SB 71
9	9.8	27.3	9-9	8-8	15-16	m.	SB 55
10	9.9	26.1	10-10	8-8	14-15	m.	SB 60
11	9.9	26.1	8-8	7-9	14-15	m.	SB 160
12	10.1	27.7	10-10	9-9	14-15	m.	SB 75
13	11.1	25.9	10-10	7-7	16-17	m.	SB 175

Table 8. Armature formulae of *S. bipunctata*.
SB...Shellback Exp.

The ovary at its maximum length reaches to the middle of the anterior fin in group A and to the frontal end of the posterior fin in group B. Group B is very peculiar in having fewer hooks and therefore compared with *S. bedoti*, *S. decipiens* and *S. neodecipiens*, both of which are provided most frequently with only 6 hooks. However, the eye-pigments of these two species are elongate and much larger than that of group B. Moreover, the intestinal diverticula are present in these two species. On the other hand, 6 hooks, up to 8 anterior teeth and up to 20 posterior teeth are included in the range of variations found by previous authors in the armature formulae of *S. friderici*. It is evident that *S. friderici* shows a wide range of diversity in the armature formulae according to localities. Consequently it is not strange that some differences are found in the number of hooks and teeth between specimens of *S. friderici* collected in the San Diego region and those from the Shellback area.

And the Shellback area itself is very extensive and as a result it is quite possible that there exist different stocks of *S. friderici*, which might show respectively different armature formulae or TC ratios. Therefore it is not improbable that groups A and B represent only different stocks of *S. friderici*, both distinctly differing from the specimens occurring in the San Diego region, because the localities of the groups A and B are somewhat apart from each other. If the mature seminal vesicles were found in some specimens of these two groups, then the above-mentioned problem would be settled very easily and decidedly. Unfortunately, however, every specimen of groups A and B is devoid of seminal vesicles, hence, at present, only their probable systematic relationships are explained, together with *S. tenuis* and *S. friderici*. As to *S. tenuis* and *S. friderici*, I can only say that speciation in these two forms seems not to have evolved so far from a common ancestor, but I cannot determine whether the degree of speciation is at the level of species or variety. I am of the opinion that they are not ecological forms of a single species, because intermediate forms are not yet known from the waters showing intermediate characteristics between the *tenuis*-and *friderici*-waters.

3. *Sagitta serratodentata tasmanica* THOMSON

(Figs. 6-12, Tables 9-11)

THOMSON (1947): Commonwealth of Australia, Council for Scientific and Industrial Research Bulletin No. 222, pp. 15-17, 1 text-fig.

Sagitta tasmanica...FURNESTIN (1953 a): Bulletin de l'Institut Océanographique, No. 1025, 39 pp.

FURNESTIN (1953 b): Bulletin des Travaux publiés par la Station d'Aquiculture et de Pêche de Castiglione (Alger), Nouvelle série, No. 4, p. 25.

FAGETTI (1958): Revista de Biología Marina (Valparaiso), Vol. VIII, Nos. 1-3, pp. 46-48, text-fig. 7.

Sagitta sp. (*serratodentata* group)...BIERI (1959): Limnology and Oceanography, Vol. IV, No. 1, p. 18, text-fig. 22.

Many specimens up to 15 mm from the Shellback area and a single specimen from St. 1 of the Transpac Expedition. Generally speaking, the body appears to be a little more transparent than in *S. pacifica*; the hooks and the anterior and posterior teeth are yellowish, but stained more faintly than in *pacifica* and the serration on the hooks is not so conspicuous as in *pacifica*. The frontal end of the anterior fin begins at about the posterior end of the ventral ganglion. The anterior part of the posterior fin along the trunk is slightly longer than the posterior part along the caudal segment, the fin is broadest at the level posterior to the trunk-tail septum and is provided with a very narrow rayless zone along the trunk in front of the female genital aperture as in *pacifica*. The collarette is practically absent in all examined specimens. The eye-pigment (Figs. 6 and 10) is elongate, but relatively thicker than in *pacifica*. For instance, in a perfectly preserved mature specimens of *pacifica* from the Shellback station SB 50, length of eye-pigment/length of eye $\times 100$

was 23.4-24.0 and width of pigment/length of pigment $\times 100$ was 25.0-27.3, whereas the former was 27.9-34.6 (30.5 on an average of 4 measurements) and the latter was 32.6-50.0 (43.7 on an average of 4 measurements) in *tasmanica*. And this difference provides a means of separating immature specimens of *pacifica* and *tasmanica*. The ovary at maximum length reaches slightly beyond the middle of the anterior fin and contains up to 30 eggs arranged in two rows, but usually it ends near the anterior end of the posterior fin and contains less than 20 eggs. The seminal vesicle (Figs. 7-8) is situated near the rear end of the posterior fin, being fairly apart from the

9 *Sag. serratodentata tasmanica* (1)

No.	body length	tail seg. in %	hook	ant. teeth	post. teeth	maturity	loc
1	6.5mm.	25.0	6-7	6-6	15-16	m	SB 75
2	6.8	26.7	5-5	6-6	13-13	m	SB 68
3	7.0	26.9	6-6	5-6	12-13	m.	SB 71
4	7.2	23.6	5-6	6-6	13-13	med	SB 71
5	7.6	23.1	7-7	6-6	14-14	im.	SB 90
6	8.1	26.4	6-6	6-7	13-14	im.	SB 95
7	8.2	23.8	6-6	5-5	11-12	m.	SB 68
8	8.5	26.2	6-6	6-6	14-15	med.	SB 95
9	9.0	24.6	6-6	7-7	15-16	med.	SB 100
10	9.1	25.0	5-6	7-8	16-17	im.	SB 145
11	9.2	24.6	6-6	6-6	11-12	med	SB 132
12	9.3	23.1	5-5	7-7	15-16	m	SB 95
13	9.4	25.5	6-6	5-6	14-15	med.	SB 85
14	9.4	24.1	6-6	6-7	16-16	m	SB 112
15	9.8	23.3	6-6	7-7	14-15		SB 170
16	9.8	26.7	6-6	6-7	14-15	med.	SB 85
17	9.8	25.3	6-6	5-6	13-13	m	SB 90
18	9.9	25.7	6-6	7-7	15-15	im.	SB 50
19	9.9	21.1	6-6	7-7	15-16	m.	SB 85
20	9.9	23.5	6-6	7-7	15-16	m.	SB 180
21	10.1	25.6	6-6	6-7	13-14	im.	SB 90
22	10.1	23.1	6-7	7-7	13-13	med.	SB 137
23	10.2	23.6	6-6	7-7	16-17	m.	SB 35
24	10.3	24.1	6-6	7-7	17-18	m.	SB 50
25	10.3	25.2	6-6	7-7	16-17	im.	SB 35
26	10.4	22.5	6-6	6-6	15-15		SB 30
27	10.5	24.8	6-6	8-8	15-17	m.	SB 35
28	10.5	23.5	6-6	7-8	19-19	m.	SB 109
29	10.6	26.4	6-6	5-7	13-14	im.	SB 35
30	10.6	23.9	5-5	7-8	16-17	m.	SB 35
31	10.7	24.8	6-6	6-7	15-15	im.	SB 35
32	10.7	25.5	6-6	6-7	16-17	m.	SB 50

Table 9-1. Armature formulae of *S. serratodentata tasmanica*.
SB—Shellback Exp.

caudal fin, although in very early stages of the development (16 in Fig. 7), it may be separated considerably from both of the posterior and tail fins. It is elongate, and slightly swollen anteriorly in early stages, but becomes oval at maturity. A small prominence may be produced at maturity at the antero-lateral corner (21-26 in Fig. 7), where the glandular tissue is located, and the rupture occurs just around this prominence (27 in Fig. 7 and Fig. 8); evidently protandric. The intestinal diverticula absent. Rows of anterior teeth meet each other in an acute angle. The armature formulae of some individuals (for instance Nos. 3, 7, 11, 17, 22 and 45 in Table 9) from the Shellback area resemble those of typical *serratodentata atlantica*

9 *Sag. serratodentata tasmanica* (2)

No.	body length	tail seg. in %	hook	ant. teeth	post. teeth	maturity	loc.
33	10.8	21.7	5-6	6-6	12-14	m.	SB 80
34	10.8	24.1	6-6	6-7	15-15	m.	SB 35
35	10.9	25.6	6-6	7-7	15-15	m.	SB 35
36	10.9	24.0	5-5	7-7	15-17	m.	SB 115
37	11.0	24.7	6-6	7-7	15-15	im.	SB 35
38	11.1	23.5	5-6	7-7	14-14	med.	SB 130
39	11.1	22.4	7-7	7-7	14-15		SB 1
40	11.2	24.4	5-5	7-8	16-17	m.	SB 105
41	11.4	21.7	5-5	6-7	17-18	m.	SB 137
42	11.4	22.9	6-6	7-7	14-14	med.	SB 137
43	11.7	25.0	6-6	7-7	14-16	med.	SB 35
44	11.8	23.1	6-6	7-7	16-16	m.	SB 137
45	12.0	24.3	6-7	6-7	13-13	im.	SB 132
46	12.1	23.7	7-7	5-5	13-15	m.	SB 137
47	12.2	25.1	6-6	7-8	18-18	med.	SB 105
48	12.2	25.1	6-6	8-8	17-17	med.	SB 112
49	12.2	25.1	6-7	7-7	14-14	med.	SB 132
50	12.2	23.7	5-5	7-7	21-21	m.	SB 137
51	12.4	23.7	5-5	4+*-7	15-15	m.	SB 118
52	12.5	21.9	5-6	10-10	22-22	m.	SB 137
53	12.6	23.7	6-6	6-7	16-17	med.	SB 115
54	12.7	24.1	6-6	5-6	16-16	med.	SB 115
55	12.7	23.1	6-6	7-8	17-18	im.	SB 137
56	12.7	24.1	7-7	6-7	15-15	m.	SB 1
57	13.0	23.0	5-5	7-7	15-16	m.	SB 109
58	13.1	22.3	6-6	7-7	16-17	m.	SB 118
59	13.2	23.6	6-6	5-6	13-14	m.	SB 1
60	13.3	23.9	5-5	8-8	20-21	med.	SB 112
61	14.3	22.8	6-6	8-8	17-18	m.	SB 122
62	14.7	23.5	6-6	8-8	15-15	med.	SB 122
63	15.0	21.7	5-5	7-7	16-17	m.	SB 132

Table 9-2. Armature formulae of *S. serratodentata tasmanica*.

* Some torn out. SB...Shellback Exp.




Fig. 6. *Sagitta serratodentata tasmanica* THOMSON. Eyes of a 10.3 and 10.7 mm long individuals, magnified.

10 *Sag. serratodentata pacifica*

No	body length	tail seg. in %	hook	ant. teeth	post. teeth	maturity	loc.
1	8.4mm	26.4	7-7	6-6	13-13	m.	SB50
2	8.5	27.7	6-6	7-7	13-13	m.	SB50
3	8.6	26.3	7-7	7-7	12-12	m.	SB50
4	9.0	23.9	6-6	10-11	19-21	med.	SB64
5	9.4	24.1	7-7	10-10	18-18	med.	SB90
6	9.9	25.0	6-6	8-9	15-18	med.	SB90
7	10.1	25.8	6-6	10-11	19-20	m.	SB35
8	10.4	26.3	6-7	9-9	18-19	im.	SB35
9	10.6	25.2	6-6	9-10	20-21	m.	SB35
10	10.9	25.6	5-6	9-10	21-22	m.	SB35
11	11.0	25.4	6-7	8-9	20-20	m.	SB35
12	11.1	23.5	5-6	11-11	22-23	m.	TP
13	11.2	26.0	6-6	11-11	21-21	m.	SB35
14	11.4	26.1	6-6	10-10	18-18	m.	SB105

11 *Sag. serratodentata pseudoserratodentata*

No.	body length	tail seg. in %	hook	ant. teeth	post. teeth	maturity	loc
1	7.0mm	28.0	6-6	4-4	9-12	m.	SB215
2	7.6	27.4	6-6	5-5	11-11	m.	SB215
3	7.6	27.4	6-6	6-6	13-13	m.	SB215
4	7.7	28.0	6-6	6-6	12-13	m.	SB215
5	7.7	28.0	6-6	5-6	12-12	m.	SB215
6	8.2	27.8	6-7	7-7	14-14	med.	SB80
7	9.8	25.3	6-6	7-7	13-13	im.	SB50
8	9.8	24.0	6-6	6-7	12-13	im.	SB50
9	10.8	26.5	6-6	6-7	14-14	im.	SB50

Tables 10-11. Armature formulae of *S. serratodentata pacifica* (10) and *S. serratodentata pseudoserratodentata* (11). SB...Shellback Exp.




Fig. 7. *Sagitta serratodentata tasmanica* THOMSON. Seminal vesicles of various stages of development, arranged in the order of maturity, $\times 125$. 16—SB 132, 20—dorsal; 21, 22—SB 137; 23, 24—dorsal; 25, 26—right vesicles, SB 137; 27—ruptured.

and those of some others (for instance Nos. 28, 47, 50, 52 and 60 in the same table) are related closely to those of typical *pacifica*. Since some of these unusual specimens are provided with seminal vesicles of the typical *tasmanica*-form, it is unlikely that some specimens of *atlantica* and *pacifica* were mistaken for *tasmanica*. The number of posterior teeth is most frequently 13-17 (Fig. 9) and this is evidently an intermediate between those of *atlantica* and *pacifica*.

Remarks: For comparison with the present subspecies, some specimens of *pacifica* and *pseudoserratodentata* from the Shellback area were examined closely. In *pacifica*,




Fig. 8. *Sagitta serratodentata tasmanica* THOMSON. Seminal vesicles of a 14 mm long individual (tail seg. 24.7%, 6-6 $\frac{7-8}{18-18}$), $\times 125$. 28—left vesicle, just rupturing; 29—ruptured portion; 30—right vesicle, fully ruptured.

the collarete is usually distinct, though not voluminous. The ovary may reach anteriorly to about the ventral ganglion and contains up to 40 eggs at maximum. There are very many individuals in which the seminal vesicles projected out obliquely anteriorly. The armature formulae of 14 specimens with mature seminal vesicles are given in Table 10. In this table, the occurrence of three specimens at SB 50 with exceptionally few teeth (Nos. 1-3) is noteworthy. The teeth are so few that at first these three specimens were mistaken for *atlantica*. The structure of the seminal vesicle is, however, of the typical *pacifica*-form as shown in the figure (Fig. 10). Specimens Nos. 4 and 12 are striking in that the seminal vesicles have no




Fig. 9. *Sagitta serratodentata tasmanica* THOMSON. Fluctuation of the number of posterior teeth of specimens from the Shellback area.

teeth; very probably they are mal-formed individuals of *pacifica*, because the cephalic teeth are numerous enough for *pacifica* and the vesicles are projected out just as in typical *pacifica*. Some authors doubt the validity of *pseudoserratodentata*. Indeed, the armature formulae of this form conforms well to those of *atlantica*. But the shape of the mature seminal vesicle (Fig. 11) is quite different between the two, also the body length of mature specimens is considerably less in *pseudoserratodentata* than in *atlantica*. In the Shellback specimens, the ovaries reach reach the middle of the anterior fin and contain up to 20 eggs in each at maturity. The serration of

hooks in *pseudoserratodentata* is the least conspicuous of the three forms. A specimen from SB 130 has seminal vesicles which have each two prominences at the anterior end (Fig. 12) and hence at first this was considered to be *atlantica*, but the situation and the shape of the prominences differ slightly from those of typical *atlantica*, in which they are situated at the antero-lateral corner; rather the situation resembles that of a prominence of the seminal vesicle in *tasmanica*. Thus, it is possible that this specimen represents an unusual individual of *tasmanica*.

Sagitta selkirki FAGETTI from Chilean waters around the Juan Fernandez Archipelago resembles most closely *tasmanica* in the appearance of the seminal




Fig. 10. *Sagitta serratodentata pacifica* TOKIOKA. A 8.5 mm long individual (No. 2 of the Table) from SB 50. 32—eye, magnified; 34—left vesicle, $\times 125$; 33—immature right vesicle of a specimen from SB 30. $\times 125$.

vesicle, but the constant occurrence of seven hooks and fewer posterior teeth, besides the comparatively small body length of mature individuals, seem to be unique for this form. On the other hand, the dental formulae, which are related closely with those of *selkirki* as given by FAGETTI, can be seen at some places of Table 9 to show the armature formulae of *tasmanica* from the Shellback area. So, it is not impossible that *selkirki* represents a peculiar stock of *tasmanica* as do three specimens of *pacifica* from SB 50 (p. 370). Therefore, I hesitate to judge conclusively the validity of this species at present. Lastly, here arises a question as to whether the forms,




Fig. 11.

Fig. 12.

Fig. 11. *Sagitta serratodentata pseudoserratodentata* TOKIOKA. Seminal vesicles, $\times 125$. 35—7.7 mm long individual from SB 215, 36—right vesicle of a 7.6 mm long individual from SB 215, 37—ruptured left vesicle of the same specimen.

Fig. 12. ?*Sagitta serratodentata atlantica*. Left seminal vesicle of a specimen from SB 130. $\times 125$.

atlantica, *pacifica*, *tasmanica* and *pseudoserratodentata* should be ranked as species or subspecies. Since there are several distinct and important characteristics common to all of these forms, the serrated hooks, the remarkably elongate eye-pigment, the situation of seminal vesicles and the existence of a conspicuous swelling at the middle of the vas deferens besides the absence of intestinal diverticula and the general structure of the fins, the relation between these forms seems to be much closer than that found between other species. Hence, I am inclined to treat the forms as subspecies of a single species *Sagitta serratodentata*, though in some parts of this paper, especially in tables, I listed them as species in order to conserve space.

4. *Sagitta neglecta* AIDA and its allies

(Figs. 13-16, Tables 12-15)

(1) *Sagitta neglecta* AIDA (Fig. 13, Table 12): This species is extremely rare in the eastern Pacific as noted already by BIERI (1959) and SUND (1959b), though it occurs rather commonly in the western part of the North Pacific. Thus, it seems necessary to compare the specimens from both sides of the North Pacific and to ascertain the identity of the eastern specimens with the western. The armature formulae of the Shellback specimens given in Table 12 conform well to those of

12 *Sagitta neglecta*

No.	body length	tailseg. in %	hook	ant. teeth	post. teeth	maturity	loc.
1	4.6mm	29.6	8—8	5—5	11—11	im.	SB 60
2	4.8	29.7	7—8	5—6	12—13	im.	SB 55
3	5.6	31.4	7—7	5—6	14—15	im.	SB 55
4	5.9	30.0	7—7	5—6	13—14	im.	SB 55
5	6.0	31.5	8—8	6—6	13—14	im.	SB 80
6	6.0	29.0	8—8	5—5	14—15	im.	SB 170
7	6.2	30.2	7—7	6—7	15—15	im.	SB 55
8	6.2	34.4	7—7	6—7	15—16	im.	SB 60
9	6.5	34.0	6—7	7—7	16—17	im.	SB 60
10	6.5	33.0	7—7	6—6	14—14	med.	SB 60
11	6.8	33.3	7—8	6—7	16—16		SB 170
12	6.8	31.4	7—7	5—6	13—13	m.	SB 195
13	6.9	30.2	8—8	6—6	17—18	im.	SB 60
14	6.9	28.3	7—8	6—7	16—16		SB 170
15	7.2	30.9	7—8	2+*-7	17—17	m.	SB 109
16	7.3	32.1	6—6	6—6	17—17	m.	SB 68
17	7.5	32.2	8—8	4+*-6	16—17	m.	SB 68
* 18	7.5	28.7	0+*-3+*	5—6	7—10	med.	SB 50
19	7.7	29.4	7—7	7—7	17—17		SB 170
20	8.1	28.0	7—8	7—8	18—18	m.	SB 170

13 *Sagitta oceania*

No.	body length	tailseg. in %	hook	ant. teeth	post. teeth	maturity	loc.
1	4.3mm	31.8	7—7	4—5	11—12	m.	EQPS28
2	4.7	31.5	7—8	5—5	13—13	m.	EQPS28
3	4.8	32.4	7—7	4—5	12—12	m.	EQPS28

Tables 12-13. Armature formulae of *S. neglecta* (12) and *S. oceania* (13).

* Some torn out. SB—Shellback Exp.

typical forms from the western Pacific. The general structure of the body agrees closely in these two groups of specimens. For these reasons, *S. neglecta* undoubtedly occur also in the Shellback area. In specimens examined from this area, the ovary may reach to the anterior end of the ventral ganglion. Individual No. 18 in Table 12 (Fig. 13) is very peculiar in that the posterior teeth are much fewer and the

intestinal diverticula are not distinct. The anterior fin begins at the posterior end of the ventral ganglion, the posterior fin is slightly longer than the anterior, equally divided on the trunk and the caudal segment and broadest at the level posterior to the trunk-tail septum; the constriction of the body at this septum is not remarkable. Rayless zones were not found on either fin. A thin collarette is present around the neck. The ovary at medium maturity reaches to the front end of the posterior fin. The seminal vesicle (39) is situated near the rear end of the posterior fin; its shape




Fig. 13. A problematic specimen (No. 18 of the Table), treated here provisionally as *Sagitta neglecta* AIDA, from SB 50. 38—eyes, magnified; 39—seminal vesicles, dorsal. $\times 73$.




Fig. 14. *Sagitta oceania* GRAY. Head of a 4.3mm long individual from EQP S-28. $\times 125$.

resembles that of *Sagitta oceania*. Eye-pigment (38) oval, length of pigment/length of eye $\times 100$ is 39.4–43.8, width of pigment/length of pigment $\times 100$ is 53.8–60.7. The fewer number of hooks is evidently attributable to some injury. The armature formula, the situation of the seminal vesicle, and the indistinctness of the intestinal diverticula suggest a possible identity of this specimen with *Sagitta setosa* from the European waters. However, since the shape of the corona ciliata was not ascertained, and the anterior fin seems too long for *setosa* and moreover the locality is so far

removed from the European waters, one hesitates to treat the specimen as *setosa*. Rather, it is treated here provisionally as an unusual individual of *neglecta* pending further detail study of specimens in better state of preservation.

(2) *Sagitta oceania* GRAY (Figs. 14-15, Table 13): Six specimens up to 4.8 mm were found in the sample from EQP-S 28. The anterior fin begins at the posterior end of the ventral ganglion; the posterior fin is longer than the anterior, nearly equally divided on the trunk and the tail segment and broadest at the level posterior to the trunk-tail septum. Rayless zone absent in both fins, although rays are very sparse in a small part near the frontal end. The collarette is practically absent, eye-pigment (41) is roundish and very large. The ovary reaches slightly beyond the posterior end of the anterior fin, eggs are unequal in size. The seminal vesicles are situated nearer the posterior fin than the caudal fin; distance to posterior fin/distance to caudal fin is 0.32 to 0.70, being 0.48 on an average of 4 measurements. The distinction between *neglecta* and *oceania* is rather exacting as mentioned already in detail when *Sagitta lacunae* was described from the lagoon water of the Palao Islands (TOKIOKA, 1942). The absence of the collarette and the large eye-pigment in the present specimens seem to refer them to *oceania* rather than *neglecta*. However, it is noteworthy that these specimens have attained maturity at such a small body size.

(3) *Sagitta regularis* AIDA (Table 14): The armature formulae of the Shellback-specimens are closely related to those of the Eastropic Expedition specimens as given by SUND (1959b). This species seems to mature much more rapidly in the Shellback area than in the western part of the North Pacific, as indicated in individuals only 4 mm long having 5-8 mature oval eggs arranged in zigzag rows and in a 4.6 mm specimen in which the ovary reached to the ventral ganglion. The formulae of some small individuals resemble closely those of *Sagitta bedfordii*, but the apparent shape of the eye-pigment differs distinctly between *regularis* and *bedfordii*; it is roundish in the latter, while it is distinctly elongate in the former.





Fig. 15. *Sagitta oceania* GRAY.
A 4.8 mm long individual from
EQP S-28.

14

Sagitta regularis

No.	body length	tail seg. in %	hook	ant. teeth	post. teeth	maturity	loc.
1	3.3mm	37.3	10-10	2-2	2-3	im.	SB35
2	3.9	30.0	7-7	2-2	2-2	im.	SB75
3	4.0	35.5	8-9	3-3	4-4	m.	SB30
4	4.0	32.8	8-8	2-2	3-3	m.	SB30
5	4.5	33.3	8-8	3-3	5-5	m.	SB30
6	4.6	34.3	6-7	3-3	4-4	m.	SB80
7	4.6	31.4	7-8	3-?	3-4		SB40
8	4.9	36.0	7-7	2-3	5-5	m.	SB35
9	4.9	33.3	7-7	4-4	4-4		SB35
10	5.0	32.5	8-8	4-4	6-6	im.	SB100
11	5.1	34.2	7-8	3-4	6-6	m.	SB35
12	5.1	33.3	8-8	3-3	4-4	m.	SB30
13	5.2	34.0	7-7	3-4	5-6	med.	SB35
14	5.2	33.8	6-6	3-3	5-5	med.	SB35
15	5.2	33.8	8-8	3-4	4-5		SB35
16	5.2	33.8	7-7	3-3	3-3	med.	SB35
17	6.9	32.0	8-8	3-3	4-5	m.	SB112

Table 14. Armature formulae of *S. regularis*.
SB...Shallback Exp.



40

Fig. 16. *Sagitta crassa* f. *naikaiensis* TOKIOKA. A 6 mm long individual from the offshore waters of North Eastern Honshû, Japan. Seminal vesicles. $\times 125$

15

Sagitta crassa f. *naikaiensis*

No.	body length	tail seg. in %	hook	ant. teeth	post. teeth	maturity	JAPAN
1	4.0 mm	30.6	9—9	2—2	2—3	im.	off the north eastern Honsyu
2	4.0	29.0	7—9	4—4	6—6	im.	
3	4.9	29.3	8—8	4—4	7—7	im.	
4	5.4	27.7	8—9	4—5	7—8	im.	
5	5.5	29.8	7—8	4—5	8—8	med.	
6	6.0	30.4	9—9	5—5	9—9	m.	

Table 15. Armature formulae of *S. crassa* f. *naikaiensis*.

(4) *Sagitta crassa* f. *naikaiensis* TOKIOKA (Fig. 16, Table 15): To ascertain the identification of the specimens from the offshore waters off North-Eastern Honsyū, Japan, some individuals were dissected. The distinct collarette around the neck, the corona ciliata beginning at the level posterior to eyes, the large eye-pigment and the distinct but rather small intestinal diverticula together with the shape and situation of the seminal vesicles (Fig. 16) seem to ascertain the identity of these specimens with *crassa* f. *naikaiensis*. The ovary is limited to a range posterior to the frontal end of the posterior fin.

5. *Sagitta neodecapiens* n. sp.

(Figs. 17-20, Tables 16-17)

Length up to 13.2 mm in examined specimens. The tail segment occupies most frequently 28-29% of the whole body length. General appearance of the body (Fig. 17) resembles closely that of *S. decapiens*. The anterior fin begins at the posterior end of the ventral ganglion. The posterior fin is nearly as long as or very slightly shorter than the anterior and divided approximately equally on the trunk and the caudal segment, although sometimes the anterior half along the trunk is slightly longer than the posterior half along the tail segment; it is widest near the trunk-tail septum. In both fins, the thin anterior terminal portion is devoid of fin rays; and in the posterior fin, there is a narrow rayless-zone along the trunk anterior to the female genital opening. The collarette is indiscernible in any of the examined specimens; no constriction at the trunk-tail septum. The eye-pigment (Fig. 18) is rather small, elongate and somewhat resembles that of *S. bipunctata* rather than that of *S. decapiens*. The pigment is not dense as in *decapiens*, but somewhat shorter than in *decapiens*. Length of pigment/length of eye $\times 100$ is 28.6 to 44.1, being 40.2 on an average of seven measurements, and width of pigment/length of pigment $\times 100$ is 31.8 to 50.0 and 39.2 on an average of seven measurements. In *decapiens*, length of pigment/length of eye $\times 100$ is 49.3 to 61.2 in three measurements and 55.7 on an average, and width of pigment/length of pigment $\times 100$ is 23.1 to 39.7 and 29.9 on an average. Thus, generally speaking, the pigment seems shorter, but




Fig. 17. *Sagitta neodecipiens*
n. sp. A specimen from SB
1, dorsal.

relatively thicker than in *decipiens*. It is also shorter than that of *S. serratodentata tasmanica*. The armature formulae resembles closely those of *decipiens* as shown in Tables 16 and 17. Hooks are most frequently 6, not serrated; the anterior teeth (Fig. 19) are very short and triangular as in *decipiens* and rows meet each other usually in right or obtuse angles. The ovary at maximum length reaches anteriorly to the posterior end of the anterior fin in examined specimens. The seminal vesicles (Fig. 20) are situated approximately at the middle of the distance between the posterior and caudal fins. They are elongate and provided with a glandular portion at the anterior end. The rupture occurs along the whole lateral side at maturity. The intestinal diverticula are present as in *decipiens*.

Remarks: The present species resembles so closely *S. decipiens* that it is hardly possible to separate distinctly immature specimens of these two forms not yet having any trace of the seminal vesicles. The only apparent distinction lies in the situation of these vesicles. The difference found in size of the eye-pigment is too exacting to be practicable as a clue to differentiating these forms. In *decipiens*, the vesicles are situated near the base of the caudal fin, being far apart from the posterior fin; distance between the posterior fin and the vesicle/distance between the vesicle and the caudal fin is 4.83 to 8.75 and 7.21 on an average of four measurements. While in the present form, the ratio is 0.29 to 2.08 and 1.12 on an average of 33 measurements. However, the original description of *Sagitta decipiens* was made by FOWLER (1905) on specimens without seminal vesicles, consequently it is impossible to determine definitely whether the present form is identical with FOWLER's original specimens or not. But in order to prevent confusion, it seems reasonable to recognize the forms described by RITTER-ZÁHONY (1911) as being identical with FOWLER's specimens. Then the present specimens are separable distinctly from *decipiens* described by





Fig. 18. Eyes of 45—*Sagitta neodecipiens* n. sp. from SB 181 and 46—*Sagitta decipiens* FOWLER from SB 181, magnified.

16 *Sagitta neodecipiens*

No	body length	tail seg. in %	hook	ant. teeth	post. teeth	maturity	loc.
1	7.3mm	31.0	6—6	8—8	15—16	med.	SB112
2	7.4	29.8	6—6	7—7	15—15	im.	SB112
3	7.8	31.7	6—6	8—8	17—17	med.	SB112
4	8.1	28.0	6—6	7—7	15—15	med.	SB112
5	8.1	28.0	6—7	6—6	13—13	im.	SB122
6	8.1	28.0	6—6	7—8	12—14	med.	SB215
7	8.5	30.7	7—7	7—7	14—14	med.	SB145
8	8.7	28.4	6—6	8—8	14—14	med.	SB122
9	8.9	29.2	6—6	8—8	16—16	med.	SB122
10	9.0	29.7	6—6	10—10	17—18	m.	SB181
11	9.4	30.0	6—6	7—8	14—14	m.	SB150
12	9.6	29.7	6—6	9—9	19—19	m	SB122
13	10.4	31.3	6—6	9—10	17—18	im.	SB 1
14	10.5	28.6	6—6	8—9	18—18	im.	SB50
15	10.8	29.6	6—6	9—10	17—17	im.	SB50
16	10.9	28.6	6—6	7—7	13—14	im.	SB 1
17	11.1	26.3	6—6	8—9	15—16		SB 1
18	11.1	29.4	6—6	9—9	16—17	im.	SB50
19	11.1	28.7	6—6	8—8	15—16	im.	SB50
20	11.2	29.1	6—6	6—9	17—17	im.	SB 1
21	11.2	27.7	6—6	8—9	16—18	med.	SB 1
22	11.4	29.7	6—6	9—10	18—19	im.	SB50
23	11.8	28.2	6—6	9—9	15—16	med.	SB 1
24	12.1	27.4	6—6	9—9	16—17	im.	SB 1
25	12.7	28.2	6—6	8—8	15—16	med.	SB 1
26	13.2	28.6	6—6	9—9	16—16	med.	SB 1

Table 16. Armature formulae of *S. neodecipiens*.
SB—Shellback Exp.



44

Fig. 19. *Sagitta neodecipiens* n. sp. Anterior and posterior teeth of an individual from SB 1, magnified.

17 *Sagitta decipiens*

No	body length	tail seg. in %	hook	ant teeth	post. teeth	maturity	loc
1	3.6mm	32.7	7-7	4-4	6-6	im.	EQPS28
2	5.5	29.4	7-7	7-7	11-13	im.	EQPH17
3	6.2	25.0	7-7	5-5	10-10	im.	EQPH17
4	6.2	29.5	7-8	6-6	11-11	im.	EQPH17
5	7.5	28.7	6-7	7-8	13-14	im.	SB122
6	8.8	27.4	6-7	8-8	14-14	im.	SB122
7	8.9	27.0	6-6	8-8	14-16	m.	SB90
8	8.9	27.7	7-7	9-9	15-16	m.	SB100
9	9.0	28.1	6-6	8-9	17-18	im.	SB85
10	9.1	25.0	6-6	8-8	14-15	m.	SB180
11	9.4	27.6	6-6	9-9	16-17	im.	SB50
12	9.4	27.6	5-6	9-10	17-18		SB30
13	9.6	25.7	6-6	8-8	17-17	med.	SB75
14	9.6	25.2	6-6	9-9	18-20	med.	SB50
15	9.6	27.2	6-6	9-10	18-19	m.	SB50
16	10.1	27.1	6-6	9-9	17-18	m.	SB85
17	10.6	26.4	6-6	9-10	17-17	med.	SB50
18	10.7	27.4	5-6	10-10	20-20		SB30
19	10.7	29.1	5-6	9-9	16-17	med.	SB1
20	10.7	27.4	6-6	9-10	18-19	m.	SB50
21	10.7	26.8	6-6	9-9	19-20	med.	SB50
22	10.7	27.3	6-6	9-9	15-17	med.	SB85
23	10.7	22.4	6-6	8-8	16-17	m.	SB195
24	11.2	25.0	6-6	7-7	14-15	m.	SB1
25	11.4	28.4	6-6	7-8	15-15	med.	SB1
26	13.2	24.6	6-7	7-8	19-19	med.	SB181
27			7-7	7-7	14-15	med.	EQPH11

Table 17. Armature formulae of *S. decipiens*.
SB...Shellback Exp.




Fig. 20. Seminal vesicles of 47, 48—*Sagitta neodecipiens* n. sp. from SB 50, respectively 11.1 and 10.8 mm in length, and 49, 50—*Sagitta decipiens* FOWLER respectively from SB 181 and SB 195 (10.7 mm in length).

RITTER-ZÁHONY by the situation of the seminal vesicles. So I think it best to treat the present form as a separate species differentiated from *decipiens* and named "*neodecipiens*" because of its close resemblance to *decipiens*.

6. *Sagitta bedoti* BÉRANECK f. *minor* TOKIOKA

(Table 18)

The specimens occurring in the Shellback area are all small, less than 13.3 mm in length and the ovary at its maximum length reaches to the posterior end of the anterior fin. The anterior end of the intestine may be slightly swollen and becomes whitish and opaque in some specimens. These features together with the armature formulae seem to assign the Shellback specimens to f. *minor* found abundantly in the tropical waters in the western Pacific. Specimens from the Eastropic Expedition resemble wholly the Shellback individuals (SUND, 1959b). The distance between eyes is comparatively large, distance between eyes/width of head is 34.3–40.0 in examined specimens.

18							
Sagitta bedoti f. minor							
No	body length	tail seg. in %	hook	ant. teeth	post. teeth	maturity	loc.
1	5.9mm	22.2	6—7	6—7	20—20	im.	SB 75
2	6.1	25.5	7—7	8—8	17—18	im.	SB 60
3	7.2	27.3	7—7	6—7	20—20	im.	SB 95
4	7.3	26.5	6—7	9—9	23—23	im.	SB 60
5	7.8	25.0	7—7	10—10	20—21	im.	SB 55
6	7.9	25.4	8—8	10—10	25—27	im.	SB 55
7	7.9	24.6	7—7	10—10	24—25	im.	SB 64
8	8.2	25.4	7—7	7—8	20—20	im.	SB 100
9	8.4	24.8	7—7	10—10	24—24	im.	SB 75
10	8.6	24.8	6—7	9—11	29—30	im.	SB 60
11	8.7	28.4	7—7	9—9	23—24	med.	SB 90
12	9.1	28.6	7—7	9—9	24—24	med.	SB 105
13	9.2	26.1	6—7	10—10	29—30	med.	SB 55
14	9.2	25.4	6—6	11—11	28—30	im.	SB 60
15	9.3	26.5	7—7	10—11	25—26	im.	SB 55
16	9.5	24.7	7—7	9—10	22—22	med.	SB 80
17	9.8	26.0	7—7	11—11	22—24	med.	SB 80
18	9.8	22.7	8—8	9—10	25—26	im.	SB 155
19	10.1	23.2	7—7	12—12	26—26	med.	SB 64
20	10.3	24.8	7—7	12—12	26—27		SB 166
21	10.4	24.7	7—7	10—10	27—27		SB 166
22	10.4	23.8	8—8	10—10	22—22	med.	SB 105
23	10.6	24.5	7—7	10—10	24—25	med.	SB 68
24	10.9	25.0	7—7	8—9	24—24	med.	SB 155
25	10.9	24.0	6—8	12—12	29—30	im.	SB 71
26	11.1	23.5	7—7	10—10	25—26	im.	SB 68
27	12.0	24.3	7—8	11—11	24—27	med.	SB 68
28	12.0	23.9	7—7	11—11	29—29	med.	SB 68
29	12.1	23.7	7—7	11—11	29—29	im.	SB 155
30	13.3	24.4	7—7	10—10	26—28	m.	SB 109
31	13.3	26.3	7—7	10—11	26—26	m.	SB 109

Table 18. Armature formulae of *S. bedoti*.
SB...Shellback Exp.

7. *Sagitta hexaptera* D'ORBIGNY

(Fig. 21)

Many large specimens, 40–50 mm in length, were collected from the Shellback area, the largest of all was the 58 mm long individual from SB 195. These seem to be slightly larger than the Eastropic specimens (SUND, 1959b). As there are some specimens in which the seminal vesicles are quite mature and preserved perfectly, a typical one is shown here in Fig. 21. The situation of the vesicle varies considerably,




Fig. 21. *Sagitta hexaptera* D'ORBIGNY. Seminal vesicle of an individual from MP 16. $\times 125$.

although it is nearer the caudal fin than the posterior fin; distance between posterior fin and vesicle/distance between vesicle and caudal fin is 4.33 to 6.64, being 5.25 on an average of 5 measurements. The shape is elongate and slightly swollen in the posterior half, where the glandular tissue is found.

8. *Sagitta enflata* f. *gardineri* DONCASTER

(Table 19)

Sagitta gardineri...DONCASTER (1903): Chaetognatha, with a note on the variation and distribution of the group. The fauna and geography of the Maldivic and Laccadive Archipelagoes, Vol. I.

Sagitta inflata, gardineri-Typus...RITTER-ZÁHONY (1909): Denkschriften der Mathematisch-Naturwissenschaftlichen Klasse der Kaiserlichen Akademie der Wissenschaften Wien, Bd. LXXXIV, pp. 5-7.

There were found many large individuals of *S. enflata* in the Shellback area, especially at Stations SB 112 and SB 122. They attain 27 mm in body length; the ovary is long, generally reaching to the middle of the anterior fin at maturity and contains up to one hundred or more eggs in one organ; it may reach even to the frontal end of the anterior fin (No. 5 in the Table). Contrarily, the ordinary small forms (*minor*-Typus by RITTER-ZÁHONY) have much shorter ovaries that are limited within the area posterior to the frontal end of the posterior fin and containing up to 30 eggs. FAGETTI (1958a) also found such larger individuals with longer ovaries in the waters off Central-Northern Chile. She notes that the larger specimens are provided with more teeth than the smaller ones and this is true also in the Shellback individuals. However, it is rather a general feature that the teeth increase with body length. The armature formulae of both forms from the Shellback area seem to be continuous as shown in the table. Ten anterior teeth are recorded also on

19

No	body length	tailseg. in %	hook	ant. teeth	post. teeth	maturity	loc.
Sagitta enflata ordinary form							
1	12.7mm	19.0	9—10	7—8	12—13	m.	SB 112
2	13.2	19.0	9—9	7—8	11—12	m.	
3	13.7	19.0	10—10	7—8	12—13	m.	
4	17.2	17.0	10—10	8—8	14—15	m.	
Sagitta enflata large form							
5	25.4	18.0	10—10	10—11	16—16	m.	SB 112
6	25.7	18.0	10—10	9—9	16—17	m.	
7	26.0	18.0	9—9	9—9	7+*-15	m.	
8	26.5	18.0	9—10	10—10	14—15	m.	
9	27.0	18.0	10—10	10—11	15—16	m.	
10	27.0	17.0	9—10	9—9	15—15	m.	

Table 19. Armature formulae of *S. enflata*.

* Some torn out. SB...Shellback Exp.

Some larger individuals of the small form as indicated by SUND (1959b) on a 11.8 mm long specimen from the Eastropic Expedition and on 15.4–15.6 mm long specimens which were collected by HIDA in the middle part of the North Pacific. For these reasons, I hesitate to consider the number of teeth as a significant characteristic. At any rate, these large forms with longer ovaries can be safely identified as *Sagitta enflata* f. *gardineri*, first described by DONCASTER (1903) from the waters around the Maldiva and Laccadive Islands and later reported by RITTER-ZÁHONY (1909) from the Red Sea. It is not yet known from the Mediterranean Sea (GHIRARDELLI 1951, MASSUTI 1958). As it is hardly possible to distinguish young individuals of this form from those of the ordinary type, the exact range of distribution is not yet fully known. If the occurrence of this form is confined to some special water masses or if it is ascertained that there is no intermediate state between the longer and

shorter ovaries at maturity, then *gardineri* should be treated as a variety or even a distinct species.

9. *Sagitta lyra* KROHN

(Fig. 22)

The fully matured seminal vesicles of this species resemble closely that of *enflata*, nearly spherical in shape and rupture around the apex. The vesicles are situated near the rear end of the posterior fin, being widely separated from the caudal fin.




Fig. 22. *Sagitta lyra* KROHN. Seminal vesicles of a 32.5 mm long individual from SB 1, dorsal.

10. Other species of *Sagitta*

(Tables 20-22)

A few specimens of some other species were dissected to ascertain identification. Their armature formulae are given in small tables 20-22. Young individuals of *Sagitta pulchra* are sometimes provided with 7 hooks. Some specimens of *Sagitta minima* from the Shellback area seem to bear more teeth than the Eastropic speci-

mens (SUND, 1959b). The formulae of *planctonis* seem to fall in the range of variations generally admitted for this species.

20

Sagitta pulchra							
1	7.2	22.7	7—7	6—6	10—10	im.	MP 14
2	15.7	20.7	5—6	6—6	13—13	m.	TP
3			6—6	8—8	12—13	m.	EQPHII

21

Sagitta minima							
1	6.2	17.9	7—8	4—5	11—12	m.	SB 90

22

Sagitta planctonis							
No	body length	tail seg. in %	hook	ant. teeth	post. teeth	maturity	loc.
1	6.4mm	27.6	10—10	3—5	7—8	im.	SB 30
2	10.3	25.3	9—9	5—5	9—9	im.	SB170
3	19.8	26.2	9—9	8	14		SB 1
4	20.8	25.0	9—9	8	13		SB 1

Tables 20-22. Armature formulae of *S. pulchra* (20), *minima* (21) and *planctonis* (22). EQP...Equipac Exp., MP...Midpac Exp., SB...Shellback Exp., TP...Transpac Exp.

11. *Krohnitta subtilis* (GRASSI) and *K. pacifica* (AIDA)

(Fig. 23, Tables 23-24)

The mature seminal vesicles of *K. subtilis* are very elongate and touch both the posterior and the caudal fins; the rupture occurs at first at the antero-lateral corner of the vesicle (Fig. 23), but afterwards it extends to nearly the whole lateral

23

Krohnitta pacifica						
1	5.9	33.3	9—9	15—15	im.	SB 35
2	6.2	32.3	8—8	13—13	m	SB170
3	6.9	29.2	8—8	13—14	m.	SB 35
4	7.3	28.6	9—10	14—14	m.	SB 35
5	8.0	30.9	8—8	15—15	m	SB 35
6	8.0	33.3	9—10	15—15	m.	SB115
7	8.3	29.7	8—9	14—15	m	SB118

24

Krohnitta subtilis						
1	11.4	31.4	8—8	10—11	m.	SB170
2	12.0	32.4	7—7	11—12	m	SB170

Tables 23-24. Armature formulae of *K. pacifica* (23) and *subtilis* (24). SB...Shellback Exp.




Fig. 23. *Krohnitta subtilis* (GRASSI). Seminal vesicles of an individual from SB 44. $\times 125$.

side leaving only a small posterior part of the vesicle intact. The Shellback specimens of *K. pacifica* seem to be provided with slightly more hooks than the Eastropic specimens (SUND, 1959b).

Distribution and Faunistic Notes

1. LAGOONS OF BAJA CALIFORNIA

(Fig. 24, Tables 25-26)

1) *Scammons lagoon*. This lagoon is situated approximately at $27^{\circ}45' N \times 114^{\circ} 10' W$ (Fig. 24) and the shallow bottom is covered by eel grass. Ten plankton samples were taken with a small net at nine of the 17 stations in the lagoon. Only *S. tenuis* was found in six of the samples. Generally speaking, plankton samples from the outer stations (Sts. 10, 15 and 16) were characterized by a dominance of the copepod, *Paracalanus*, those from the middle part (Sts. 11 and 17) were densely populated by *Oithona* and the samples from the inner portion (Sts. 12-14) were




Fig. 24. Map of Scammons Lagoon, showing the sampling stations.

25- Stations	Stations						total
	10	11	15A	15B	16	17	
Body length							
< 3 mm	1	3	23	12	12	2	53
3.1 - 5 mm	3	7	14	20	13	15	72
5.1 - 7 mm	1	—	1	13	3	21	39
7.1 - 9 mm	—	—	4	2	2	10	18
9.1 - 10.7 mm	—	—	—	—	—	1	1
total	5	10	42	47	30	49	183
Salinity (‰)	36.3	36.7	34.7		37.6	38.8	

Table 25. Occurrence of *Sag. tenuis* in Scammons lagoon.

26	Stations	1	2	total
	Body length			
	3.1 - 5 mm	1	4	5
	5.1 - 7 mm	—	2	2
	total	1	6	7

Table 26. Occurrence of *Sag. tenuis* in Manuela lagoon.

crowded with *Acartia*. However, on some tidal phase these copepods may occur mingled with one another or *Acartia* may be very abundant at the outer-most station (St. 15) where the water flows in or out very violently. The distribution of *S. tenuis* seemed to be confined to the *Paracalanus-Oithona* area at the time when the collections were made; none was found in the inner portion (Sts. 1-4, 6, 8, 12-14) where the salinity reached 39.5-47.9‰. The highest salinity recorded at stations where this chaetognath was found was 38.8‰ (St. 17) and the maximum body length measured on the collected specimens at these stations was 10.7 mm.

2) *Manuela lagoon*. A few specimens of *S. tenuis* were found in two plankton samples collected with a small net. Very probably *S. tenuis* is the only chaetognath, which can live in such lagoons situated along the coast of Lower California and characterized by comparatively high salinity.

2. THE INLET WATERS ALONG THE PACIFIC COAST OF JAPAN

A single haul with a small net, 17 cm in the mouth diameter, was made from 0-17 m in the morning of Oct. 7 during the Transpac Expedition near the entrance to Yokosuka Harbour in Tōkyō Bay. Eighty chaetognaths were found in this sample, all belonging to *S. crassa* f. *naikaiensis* which is known as a plankton indicator of the inlet waters, being found very abundantly and commonly in embayments along the coast of Japanese Islands.

3. ELEGANS-WATER IN THE NORTH PACIFIC

(Appendix table 2)

Of the samples of the Transpac Expedition 1953, thirty-nine (Nos. TP 19-50, 53, 61, 63-66 and 73) contain only cold water species, *Sagitta elegans* and *Eukrohnia hamata*, or only the former. *E. hamata* was found only at two stations (TP 20 and 21) and only a single individual in each case.

4. MIXING AREA OF THE ELEGANS-WATER AND THE WARM WATER

(Appendix table 3)

With regards to samples from the surface water above 50 m, *Sagitta minima*

seems to be the commonest of five warm water species occurring throughout the whole mixing area. In the eastern part of the Pacific, *S. lyra* occurs rather regularly in the mixing area, while in the western half *S. inflata* joins the fauna and in more western part than 150° E, *S. bedoti* and *S. crassa* f. *naikaiensis* are added to the population. The occurrence of *S. lyra* has been reported by LEA (1955) along the west coast of the Queen Charlotte Islands, British Columbia, by SUND (1959) in the Gulf of Alaska, by TCHINDONOVA (1955) in the southern part of the Bering Sea, and by HIDA (1957) in the Transition Zone north of the Hawaiian Islands. The last author stated *Pterosagitta draco*, *S. hexaptera*, *S. serratodentata* and *S. bipunctata* decrease markedly in the transition zone, but on the contrary *S. lyra* increases there. MARUMO and others (1958) also mention that besides *S. elegans*, *S. bedoti*, *S. minima* and *E. hamata*, *S. lyra* occurred in the upper 80 m layer at 40° N, 150° E (p. 183). The higher concentration of *S. minima* near the northern boundary of the warm water mass in the North Pacific is clearly illustrated by BIERI (1959). The occurrence of *S. bedoti* and *S. crassa* f. *naikaiensis* in the western mixing area is evidently a natural result of the distribution of *bedoti*-water as is shown later in the section on *bedoti*-water, and of course this phenomenon is also observable in the Japan Sea.

5. THE BLUE-GREEN WATER ALONG SOUTHERN CALIFORNIA

(Table 27)

Nineteen plankton samples were collected March 12-13, 1956 in the coastal blue-green water along the Californian coast from San Diego to Long Beach. A small net, 17 cm in the mouth diameter, was towed from 20-40 m to the surface. Chaetognaths, represented by a single species *S. friderici*, were found in sixteen of these samples. Since small, young individuals were distributed widely in the surveyed area as shown in Table 27, it appears that this species propagates within this blue-green water. Of the 35 plankton samples collected off San Diego with a similar small net and stored at Scripps Institution, 27 contained *S. friderici* and some unidentifiable

27	Body length	Number of individual	Frequency of Occurrence
	1 - 4 mm	92	16/16
	5 - 8 mm	23	3/16
	9 - 12 mm	3	1/16
	13 - 16 mm	5	2/16
	17 - 19 mm	4	2/16
	total	127	

Table 27. Occurrence of *Sag. friderici* in the blue-green water along Souther California March 12-13, 1956.

juveniles or damaged specimens. Some cold water plankton animals, such as *Oikopleura labradoriensis* and *Fritillaria borealis* f. *typica*, were found rather commonly in this water, together with a neritic species such as *Oikopleura dioica*. Therefore, the blue-green water is apparently a mixture of northern cold subarctic water and coastal water, although no individual of *S. elegans* has ever been reported from the area. The details of the occurrence of *S. friderici* off California have been described by MICHAEL (1911), under the name *S. bipunctata*, and also referred by BIERI (1959).

6. BEDOTI-AREA

I. Occurrence of *S. bedoti* in the eastern Pacific (Fig. 25, Table 28)

The occurrence of *S. bedoti* in the eastern Pacific was first reported by BIERI (1957) from the waters off Peru. Later, the distribution of this species in the Shellback area was clearly illustrated in his second paper (1959). More recently the




Fig. 25. Distribution of *Sagitta bedoti* in the Shellback area.

28	Stations	Number of ind.	%	$\frac{\text{bed.}}{\text{enf.}}$	Stations	Number of ind.	%	$\frac{\text{bed.}}{\text{enf.}}$
	SB 20	1	*	*	SB 109	101	12	0.54
	SB 35	5	*	0.02	SB 155	30	2	0.06
	SB 44	6	*	0.02	SB 160	11	1	0.03
	SB 55	20	1	0.05	SB 166	215	19	0.52
	SB 60	38	4	0.19	SB 170	299	15	0.33
	SB 64	4	1	0.04	SB 175	27	2	0.08
	SB 68	137	22	0.77	SB 180	41	2	0.11
	SB 71	1	*	0.02	SB 181	7	1	0.02
	SB 75	12	3	0.17	SB 187	131	7	0.16
	SB 80	32	2	0.06	SB 195	12	3	0.14
	SB 90	1	*	*	SB 200	105	5	0.09
	SB 95	1	*	*	SB 215	1	*	*
	SB 100	15	1	0.02	SB 217	2	*	*
	SB 105	7	*	0.01				

* <1% or 0.01

Table 28. Occurrence of *S. bedoti* in the Shellback area.

presence of the species was reported by SUND (1959b and c) also in the area of the Eastropic Expedition. The species seems to be absent off northern-central Chile (FAGETTI; 1958a). Three additional papers dealing with the chaetognaths of the coastal waters of North America have appeared; namely MICHAEL (1911), LEA (1955) and SUND (1959a). In all of these reports *S. bedoti* was quite absent. I found this species in 27 of the 48 Shellback samples examined (Frequency of Occurrence 56%), of which the northernmost station was SB 217, the southernmost station SB 95 and the western-most station was SB 20. It was most abundant at stations along the coast between 15° N and 5° S, but decreases rather sharply in the offshore water. This species occurs in the Shellback area in rather large numbers at some stations, but yet the proportion of the species to the total chaetognaths is not high; it was 22% at SB 68, but most frequently it is less than 5%. The proportion of *S. bedoti* to *S. enflata* (*bedoti/enflata*) is especially low, never reaching 0.8 and mostly less than 0.5.

II. Occurrence of *S. bedoti* at the western stations near Japan (Fig. 26, Tables 29-32).

The Occurrence of this species in the waters around Japan has been reported repeatedly (AIDA 1897, as *S. bipunctata*; TOKIOKA 1939, 1940, 1951, 1954 and 1957; FURUHASHI 1953, BIERI 1959). Besides, there are several data reporting the occur-




Fig. 26. Distribution of *Sagitta bedoti* in the Japanese and adjacent waters.

29

Stations	Number of ind.	%	bed. enf.	Stations	Number of ind.	%	bed. enf.
TP 59A	6	27	∞	TP 76	60	29	2.61
TP 69	12	34	∞	TP 78	58	32	6.44
TP 70	3	19	1.50	TP 93	1	1	0.02
TP 71	3	11	∞	TP 95	2	2	0.08
TP 74	6	4	0.33	TP 99	2	1	0.04

Table 29. Occurrence of *S. bedoti* at the western stations of the Transpac Expedition near Japan.

30

Stations	Number of ind.	%	bed. enf.	Stations	Number of ind.	%	bed. enf.
J 1	16	6	0.73	J 8	1	33	∞
J 2	1	13	∞	J 9	4	25	∞
J 3	1	17	∞	J 10	2	25	2.00
J 5	2	9	2.00	J 11	9	23	9.00
J 6	1	3	0.33	J 12	1	3	0.50
J 7	18	16	3.00	J 19	17	24	1.42

Table 30. Occurrence of *S. bedoti* in the material collected by a Japanese survey ship in the offshore waters off North-Eastern Honsyū, Japan.

31

Percentage of <i>S. bedoti</i>	Eastern Pacific	Western Pacific
* - 5%	22	6
6 - 10%	1	2
11 - 20%	3	5
21 - 30%	1	6
31 - 40%	0	3
total	27	22

* less than 1%

Table 31. Comparison of the percentage of *S. bedoti* to the total chaetognaths between the eastern and western sides of the Pacific.

Eastern Pacific			Western Pacific		
*	5	27	*	0	7
0.01 - 0.09	13		0.01 - 0.09	3	
0.10 - 0.49	6		0.10 - 0.49	2	
0.50 - 0.99	3		0.50 - 0.99	2	
1.00 - 4.99	0	0	1.00 - 4.99	6	15
5.00 - 9.99	0		5.00 - 9.99	2	
10.00 - ∞	0		10.00 - ∞	7	
total	27		total	22	

Table 32. Frequency of occurrence of respective values of the ratio *bedoti/enflata* in the eastern and western sides of the Pacific.
* less than 0.01

rence of this species off the Pacific coast of middle Japan (UYENO and FURUHASHI 1955 and others) and off northeastern Japan (MARUMO 1955 and others). I found *S. bedoti* in 10 of the Transpac samples from the stations near Japan and in 12 of the 19 samples (Frequency of Occurrence 63%) collected off north-eastern Japan by a Japanese survey ship and presented to the Transpac Expedition of Scripps Institution. As the samples were collected with small net, only a small number of this species occurred in the samples. However, when the proportion of this species is referred to total chaetognaths higher percentages seem to occur more frequently in the western than in the eastern Pacific (Table 31); and this tendency is more pronounced when the values of *bedoti/enflata* are computed and compared with those in the eastern Pacific (Table 32). The northern-most latitude where this sagitta occurred was 44°06' N and the eastern-most limit of the distribution was 161°39' E,

III. Other available data concerning the occurrence of *S. bedoti*.

a) *The waters around Japan.*

i. Off north-eastern Honsyû, north to 40° N. (Table 33). The results of the examination of plankton samples collected by the Sôyô-Marû, a survey ship of the former Central Fisheries Experimental Station in Tôkyô, in the years 1934 and 1937-39 (TOKIOKA, 1957) show that *S. bedoti* occurred commonly in the mixing area of the cold Oyasio and the warm Kuroshio in the southern part of this area. Here this species constitutes high percentages of the population, 33-100%, the average being 84%. The northern-most station where *S. bedoti* occurred in this material was 42°05' N and the eastern-most station was 154°14' E.

ii. Off north-eastern Honsyû, south to 40° N. (Table 33). All of the seven samples mentioned in the preceding section, which were collected in this area, contained a large number of *S. bedoti*; its percentage was also very high ranging from 15 to 100%

33

Percentage	North to 40°N		South to 40°N	
* - 5%	—	1	—	2
6 - 10%	—		—	
11 - 20%	—		1	
21 - 30%	—		1	
31 - 40%	1		—	
41 - 50%	—		—	
51 - 60%	—	5	—	5
61 - 70%	—		1	
71 - 80%	—		—	
81 - 90%	2		—	
91 - 100%	3		4	
total	6		7	

Table 33. Occurrence of *S. bedoti* in the waters off North-Eastern Honsyû, Japan (Sôyô-Marû material).
* present, but less than 1%.

34

Percentage	off the South-Western Honsyû		Sagami Bay	
* - 5%	3	9	—	4
6 - 10%	2		—	
11 - 20%	1		2	
21 - 30%	2		1	
31 - 40%	—		1	
41 - 50%	1		—	
51 - 60%	—	3	2	2
61 - 70%	—		—	
71 - 80%	2		—	
81 - 90%	1		—	
total	12		6	

Table 34. Occurrence of *S. bedoti* in the waters off South-Western Honsyû, Japan (Sôyô-Marû material). * present, but less than 1%.

and 71% on an average. The greater part of this area is one of mixing between the Oyasio and the Kuro시오, and the abundance of *S. bedoti* along the boundary between these two water-masses in this area has also been reported in the oceanographic reports (Kaiyō-Hōkoku) published by the Meteorological Department of Japan (1956-1958). MARUMO (1955) stated that *S. bedoti* was comparatively abundant in the Kuro시오 and in the area where the Oyasio and the Kuro시오 mingle. This is clearly illustrated also by BIERI (1959).

35

Stations	Coastal Water					Proper part of Kuro시오					offshore water beyond Kuro시오		
	I ₁					I ₂	I ₃	I ₄		I ₅			
Aug. 1954	2					4	2	0		—			
Oct. 1954	0					0	0	0		0			
Mar. 1955	20					0	0	0		0			
Stations	I ₂ '					I ₃ '		I ₄ '					
Oct. 1954	0					4		0					
Stations	J ₁	J ₂	J ₃	J ₄	J ₅	J ₆	J ₇						
Aug.-Sept. 1954	4	1	4	1	0	0	0						
Oct. 1954	0	0	0	0	0	0	—						
Stations	K ₁			K ₂	K ₃	K ₄	K ₅	K ₆	K ₇				
May 1954	—			—	—	—	—	0	—				
Aug.-Sept. 1954	2			0	1		0	0	0				
Oct. 1954	8			0	0	0	0	0	—				
Stations	G ₁				G ₂	G ₃	G ₄	G ₅	G ₆	G ₇	G ₈		
May 1954	11				13	0	4	0	0	0	—		
Aug.-Sept. 1954	33				1	0	0	0	0	0	3		
Oct. 1954	4				4	0	0	0	8	—	—		
Mar. 1955	0				0	96	0	0	0	—	—		
Stations	H ₁	H ₂	H ₃	H ₄	H ₅	H ₆			H ₇				
Mar. 1955	16	8	32	16	0	0			0				
Stations	U ₁	U ₁₀	U ₁₁	U ₁₂	U ₁₃	U ₂	U ₃	U ₄	U ₇	U ₈	U ₉	U ₅	U ₆
Aug.-Sept. 1954	12	8	0	0	4	0	6	0	0	8	8	0	4

Table 35. Individual number of *S. bedoti* occurred at respective stations in the Kuro시오 area off the South-Western Honshū, Japan. (Data offered by the Kōbe Marine Observatory.)




Fig. 27. Occurrence of *Sagitta bedoti* in the Kuroshio region off the South Western Honsyū, Japan.

iii. Off south-western Honsyû. (Fig. 27, Tables 34-38). In 12 of the 13 samples of the Sôyô-Marû material (Frequency of Occurrence 92%), collected in this area, from 2 to 82% of the chaetognath population consisted of *S. bedoti* (31% on an average) (Table 34). Here, *S. bedoti* appeared to occur more abundantly in the eastern part of this area than in the western part near Kyûsyû.

Concerning the distribution of *S. bedoti* in the southern waters off Honsyû, UYENO and FURUHASHI (1955a and 1955b) showed that this species occurred in the coastal part of the Kuroshio and in the axial portion (the most strongly flowing portion) of the current, but it was quite absent in the counter current of the Kuroshio far beyond the axial portion. Prior to reading these papers, I asked to Mr. FURUHASHI of the Kôbe Marine Observatory to let me see the data to ascertain the details of the occurrence of this species. Very kindly, Mr. FURUHASHI sent me the detailed data and allowed me so generously to quote and publish them in my paper in his letter of

36

Table 36. Occurrence of *S. bedoti* in Tôkyô Bay (MURAKAMI, 1957).

* present, but less than 1%.

Outer part	May 15-17 1947	Sept. 11-13 1947	Nov. 19-22 1947	Dec. 6-7 1947	Mar. 17 1948
0	3	8	6	4	2
* - 5%	2	—	—	—	—
6 - 10%	2	—	2	—	—
11 - 20%	—	—	3	1	—
21 - 30%	1	—	1	1	—
31 - 40%	—	—	2	1	—
41 - 50%	—	—	3	—	—
51 - 60%	—	—	1	1	—
Occurrence	5	0	12	4	0
mean	10	—	30	33	—

Table 36-1

Central part	May 13-15 1947	Sept. 9-11 1947	Dec. 4-6 1947	Mar. 12-17 1948
0	15	15	10	13
* - 5%	—	—	3	2
6 - 10%	—	—	2	—
Occurrence	0	0	5	2
mean	—	—	4	2

Table 36-2

37

Inland Sea	Kii Channel	Osaka Bay	Harima Nada	Bingo Nada	Aki Nada	Suô Nada	Iyo Nada	Bungo Channel
Number of Samples	186	281	132	274	92	85	131	12
* - 5%	68	13	12	6	1	—	2	—
6 - 10%	17	3	3	—	—	—	1	1
11 - 20%	13	6	1	—	—	—	1	—
21 - 30%	5	1	—	—	—	0	1	—
31 - 40%	3	—	—	—	—	—	—	—
41 - 50%	1	—	1	—	—	—	1	1
51 - 60%	5	—	1	—	—	—	—	—
61 - 70%	5	—	—	—	—	—	1	—
71 - 80%	2	16	—	1	—	0	—	1
81 - 90%	—	—	—	—	—	—	—	—
91 - 100%	4	1	—	—	—	—	—	1
Occurrence	123	24	18	6	1	0	7	3
Frequency of Occurrence	66%	9	14	2	1	0	5	25
mean percentage	16%	11	10	1	2	0	24	50

Table 37. Occurrence of *S. bedoti* in the Inland Sea (MURAKAMI, 1957 and 1959).

* present, but less than 1%.

38

Osaka Bay 1938	
Number of Stations	36
* - 5%	11
6 - 10%	2
Occurrence	13
Frequency of Occurrence	36%
mean Percentage	3%

Table 38. Occurrence of *S. bedoti* in Osaka Bay (TOKIOKA, 1939).

* present, but less than 1%.

May 9, 1955. Moreover, he stated in his letter of July 6, 1955 that *S. bedoti* seemed to disappear in the waters 40-50 miles off Sionomisaki. All the samples were collected from 100m to the surface with the 0-net of the Kôbe Marine Observatory, constructed of No. 3 gauze and with the diameter of the mouth adjusted so as to cut 1/10m². *S. bedoti* occurred in 17 of 29 samples (Frequency of Occurrence 59%) obtained in the coastal waters, in 14 of 58 samples from the proper part of the Kuroshio (Frequency of Occurrence 24%) and in 2 of 9 samples from the offshore waters beyond the proper part of the Kuroshio (Frequency of Occurrence 22%) (Table 35). The average number of specimens per haul was 10.9 in the coastal water, 4.8 (excluding the unusual occurrence of 96 individuals at G₃ in March 1955) in the axial part of the Kuroshio and 3.5 in the offshore water beyond the Kuroshio. It is evident that the species is found abundantly

near the coast and diminishes markedly towards the open sea, although some small masses of the coastal water containing many *S. bedoti* may be carried, across the Kuroshio, into the more offshore waters beyond the current by some counter currents and contrarily some masses of the far offshore water containing few *S. bedoti* may drift just near the shore of the Pacific coast of middle Japan.

In addition to the coastal waters of this region of Japan, *S. bedoti* is found in the following localities :

A) Sagami bay (Table 34). The first record of *S. bedoti* in Japanese waters was in Misaki Harbour in Sagami Bay by AIDA (1897) under the name of *S. bipunctata*.

All of the six samples of the Sōyō-Maru material collected in this bay included this species which constituted from 13 to 53% (33% on an average) of the total chaetognaths. While I was working at the Mitsui Institute of Marine Biology in 1937-38, I very frequently collected chaetognaths from the surface water of Sagami and Suruga bays, near the laboratory of the Institute, by using KITAHARA's quantitative net. In this material, I found about 1300 *S. bedoti*, this was approximately 16% of all chaetognaths collected.

B) Tōkyō Bay (Table 36). MURAKAMI (1957) made a series of observations on the distribution of chaetognaths in this bay in the years 1947-48. When the bay is divided into three areas, an inner part inside a line connecting the estuaries of the River Rokugō and the River Yōrō, an outer part outside a line connecting the capes of Kannon-saki and Futtu-saki and a central part between these two, *S. bedoti* occurred most frequently (Frequency of Occurrence 48%) in the outer part and less frequently (Frequency of Occurrence 12%) in the central part, but it did not occur in any of the twenty-eight samples hauled in the inner part. The percentage of *S. bedoti* to the total chaetognaths ranged from 2 to 57 in the outer part, but from 1 to 8 in the middle portion. Higher percentages of the chaetognath population in the inner and middle portions are made up of *S. crassa* and its form *naikaiensis*.

C) Inland Sea and adjacent areas (Tables 37-38, see also Fig. 27). MURAKAMI (1957 and 1959) gives details of the occurrence of chaetognaths in vast plankton samples collected in the Inland Sea and adjacent areas in the years 1951-1957. His data on *S. bedoti* are arranged here in Table 37 so as to show its abundance in respective sections of the area under consideration. *S. bedoti* enters the Inland Sea from both Kii and Bungo Channels, but diminishes markedly towards the middle part of the Sea, where *S. crassa* and especially its forma *naikaiensis* are very abundant. In Kii-Channel the percentage of *S. bedoti* is very often much lowered by dense population of *S. enflata*. I made four hauls in Kii-Channel in 1938; *S. bedoti* occurred in three of these samples, its percentage varies from 3 to 20, 13 on an average (1939). In February 1954, a dense patch of chaetognaths appeared near the Seto Marine Biological Laboratory facing Kii-Channel, a small part of this patch was collected and 13 species were identified in 1112 specimens. Among these only five individuals of *S. bedoti* occurred (TOKIOKA, 1954a). In Ōsaka Bay, this species occurs

39

Southern Waters off Kyūsyū	TOKIOKA 1940		Material of Sōyō-Maru	
Number of Samples	16		6	
* - 5%	1	11	2	4
6 - 10%	3		—	
11 - 20%	4		—	
21 - 30%	2		—	
31 - 40%	1		2	
41 - 50%	—		—	
51 - 60%	1		1	
Occurrence	12		4	
Frequency of Occurrence	75%		67%	
mean Percentage	20%		20%	

Table 39. Occurrence of *S. bedoti* in the waters off Southern Kyūsyū, Japan.

* present, but less than 1%.

40

Western Waters off Kyūsyū	
Number of Samples	9
* - 5%	1
6 - 10%	—
11 - 20%	—
21 - 30%	—
31 - 40%	1
41 - 50%	—
51 - 60%	2
61 - 70%	1
71 - 80%	—
81 - 90%	1
91 - 100%	2
Occurrence	8
Frequency of Occurrence	89%
mean Percentage	62%

Table 40. Occurrence of *S. bedoti* in the waters off Western Kyūsyū, Japan. (TOKIOKA, 1940).

* present, but less than 1%.

more abundantly in the western half along Awazi Island. Of 36 samples hauled in 1938 in Ōsaka Bay by myself, *S. bedoti* occurred in 13 samples (Frequency of Occurrence 36%) and its percentage varied from 1 to 10. It occurred mostly near the south-western and north-western channels leading respectively to Kii-Channel and Harima-Nada (TOKIOKA, 1939). The species has never been found in any of the small embayments situated in the central part of the Inland Sea. For instance, none of the 31 plankton samples hauled in Kaida Bay in the years 1950-1951 nor 115 samples hauled in Kasaoka Bay in the years 1952-53 contained *S. bedoti* (MURAKAMI, 1957).

iv. The southern waters off Kyūsyū (Table 39). The data found in my previous paper (1940) and those of the Sōyō-Maru material (TOKIOKA, 1957) are available. The highest percentage was 59% found in the data of 1940. The density of *S. bedoti* seems to fluctuate rather violently in this area.

v. The western waters off Kyūsyū (Table 40). Only the data found in my paper of 1940 are available at present.

vi. The Korea Straits (Table 42). The data in my paper of 1940 show that *S. bedoti* is abundant in this area as well as in the Japan Sea.

vii. The Japan Sea (Table 41). The samples dealt with in my paper of 1940 were collected mostly along the eastern coast of Korea, not far from the land. All the data quoted in Table 41 are from samples collected at stations situated in the warm water region of the Japan Sea. At stations in the mixing area between the cold

41

The Japan Sea	TOKIOKA 1940	TOKIOKA 1951 Surface hauls	TOKIOKA 1951 50-350 m to 0 m	FURUHASHI 1953
Number of Samples	21	66	35	4
* - 5%	—	—	10	—
6 - 10%	—	—	10	1
11 - 20%	1	1	3	1
21 - 30%	2	1	2	2
31 - 40%	1	1	5	—
41 - 50%	1	—	2	—
51 - 60%	3	—	—	—
61 - 70%	2	—	—	—
71 - 80%	1	—	1	—
81 - 90%	4	—	—	—
91 - 100%	6	2	—	—
Occurrence	21	5	33	4
Frequency of Occurrence	100%	8%	94%	100%
mean Percentage	68%	56%	17%	18%

Table 41. Occurrence of *S. bedoti* in the Japan Sea.
* present, but less than 1%.

42

	The Yellow Sea	The Korea Straits
Number of Samples	19	32
* - 5%	3	—
6 - 10%	—	2
11 - 20%	—	2
21 - 30%	1	3
31 - 40%	—	2
41 - 50%	1	1
51 - 60%	—	2
61 - 70%	—	2
71 - 80%	—	1
81 - 90%	—	1
91 - 100%	7	13
Occurrence	12	29
Frequency of Occurrence	63%	91%
mean Percentage	62%	69%

Table 42. Occurrence of *S. bedoti* in the Yellow Sea and the Korea Straits (TOKIOKA, 1940).
* present, but less than 1%.

Liman Current and the warm Tusima Current, *S. bedoti* constitutes fairly high proportion of the chaetognaths, although its percentage is often lowered by the dense population of *S. minima*.

b) *The Yellow Sea* (Table 42)

According to the data of the material treated in my previous paper of 1940, *S. bedoti* seems to be very abundant at stations in this sea except for northern stations and excluding the samples gathered in the very surface layer, where the salinity of the water is rather low and only *S. crassa* is dominant. In Table 42, only the data of samples collected at stations situated south of the latitude of the point of Shantung peninsula in North China are treated. Both the frequency of occurrence and the mean percentage should be much raised if more stations in the northern part are omitted from consideration.

c) *The East China Sea* (Table 43).

S. bedoti was very abundant at stations around the Chu-San Islands near the estuary of the Yang-Tse River (TOKIOKA, 1940). In the East China Sea, it seems to be very common in the western half near the continent, but gradually decreases towards the east and is scarcely found near the Ryūkyū Islands. The Sōyō-Maru material con-

43	The East China Sea	Sōyō-Maru material		Around the Chu-san Islands	
	Number of samples	4		11	
	* - 5%	—	1	—	1
	6 - 10%	1		—	
	11 - 20%	—		—	
	21 - 30%	—		1	
	31 - 40%	—		—	
	41 - 50%	—		—	
	51 - 60%	—	1	—	10
	61 - 70%	—		1	
	71 - 80%	—		—	
	81 - 90%	—		—	
	91 - 100%	1		9	
	Occurrence	2		11	
	Frequency of Occurrence	50%		100%	
	mean Percentage	50%		88%	

Table 43. Occurrence of *S. bedoti* in the East China Sea.
* present, but less than 1%.

tains two samples from the Taiwan Straits, in both of which *S. bedoti* occupies very high percentage, respectively 54 and 21%. Mr. ENOMOTO of the Seikai Regional Fisheries Research Laboratory at Nagasaki stated in his letter of March 27, 1957 that this sagitta seemed to be the most important of all chaetognaths in the East China Sea, when he sent me several specimens of *S. bedoti* for identification, which were collected in June 1956 near the middle of that sea.

d) *The South China Sea* (Table 44)

In this area, *S. bedoti* maintains still higher percentages in the north-western portion (18-81%), but decreases greatly near Hainan Island (less than 3%). It is recorded also from the coastal waters of Annan (SERENE, 1937), but there it forms only a small part of the chaetognath fauna (HAMON, 1956). The minor form occurred in the Bay of Nhatrang-Cauda at a frequency of 6-76%, but it occupied only less than 1.7% of the whole population.

e) *The waters around the Philippines and adjacent waters* (Tables 45-46)

According to MICHAEL (1919), during the Philippine Expedition by the Albatross, *S. bedoti* occurred at only four stations in the Sulu Sea and adjacent waters. The frequency of occurrence was 9% of the total 46 sampling stations.

The south-eastern waters off Mindanao (Mindanao Current): During the Snellius

44

The South China Sea	
Number of samples	9
* - 5%	3
6 - 10%	—
11 - 20%	1
21 - 30%	1
31 - 40%	—
41 - 50%	—
51 - 60%	—
61 - 70%	—
71 - 80%	—
81 - 90%	1
Occurrence	6
Frequency of Occurrence	67%
mean Percentage	22%

Table 44. Occurrence of *S. bedoti* in the South China Sea (Sōyō-maru material).

* present, but less than 1%.

45	Sulu Sea	Number of Sampling Sts.	Frequency of Occurrence	mean Percentage
	SIBOGA-Exped.	5	60%	6.1 (1 station)
	SNELLIUS-Exped.	6	83%	4 (1~5%)

Table 45. Occurrence of *S. bedoti* in the Sulu Sea.

46	Mindanao Current	Number of Sampling Sts.	Frequency of Occurrence	mean Percentage
	SIBOGA-Exped.	2	50%	+
	SNELLIUS-Exped.	30	43%	2% (*.....2 1-5%...11)

Table 46. Occurrence of *S. bedoti* in the south-eastern waters off Mindanao.

* less than 1%.

Expedition, the frequency of occurrence of this species was higher in the southern part than in the northern part of this area.

f) *The waters around the Malay Archipelagoes* (Tables 47-51).

Since BÉRANECK'S (1895) original description of *S. bedoti* was published, stating that—"It is abundant in the bay of Amboina (p. 152)"—, there have been published three important paper: FOWLER'S report on the material of Siboga-Expedition (1906), OYE'S paper on the chaetognaths of the Java Sea (1918) and SCHILP'S extensive work on the material of the Snellius Expedition (1941). According to OYE'S data, *S. bedoti* seems to be more abundant in the western part than in the eastern part of the Java Sea.

g) *The waters adjacent to the above-mentioned area* (Tables 52 and 53).

The stations of the Snellius Expedition in the Arafura Sea were situated in the north-western part of the sea, while TOKIOKA'S material came from the south-western part of the sea. Besides, TOKIOKA (1942) states that *S. bedoti* was common in the plankton samples taken near the estuary of the River Liverpool, Arnhem Land.

h) *The Indian Ocean* (Fig. 28, Table 54).

The record of *Sagitta polyodon* by DONCASTER (1903) from the waters around the Maldive Islands is the first one dealing with the occurrence of *S. bedoti* in this ocean. There, this species was "found in fair abundance in both winter and spring (p. 213)". During the Deutsche Südpolar Expedition by the "Gauss" (RITTER-ZÁHONY,

47	Java Sea	SIBOGA 1906		OYE 1918		SNELLIUS 1941	
	Number of sampling stations	10		77		24	
	*	7	8	—	32	2	20
	1 - 5%	1		19		11	
	6 - 10%	—		6		2	
	11 - 15%	—		7		5	
	16 - 20%	—		4		1	
	21 - 30%	—	0	3	8	—	1
	31 - 40%	—		—		—	
	41 - 50%	—		1		—	
	Occurrence	8		40		21	
	Frequency of Occurrence	80%		52%		88%	
	mean Percentage	+		10%		6%	

Table 47. Occurrence of *S. bedoti* in the Java Sea.
* present, but less than 1%.

48	Celebes Ses	SIBOGA 1906		SNELLIUS 1941	
	Number of sampling stations	9		44	
	+ or *	4	7	1	29
	1 - 5%	2		19	
	6 - 10%	1		8	
	11 - 15%	—		1	
	16 - 20%	—	0	2	4
	21 - 30%	—		—	
	31 - 40%	—		2	
	Occurrence	7		33	
	Frequency of Occurrence	78%		75%	
	mean Percentage	3% (4 stations)		7%	

Table 48. Occurrence of *S. bedoti* in the Celebes Sea.
* less than 1%.

49		Banda Sea			Moluccas Archipelago				
		SIBOGA 1906		SNELLIUS 1941		SIBOGA 1906		SNELLIUS 1941	
	Number of sampling stations	15		68		23		53	
	+ or *	12	12	2	43	11	19	1	34
	1 - 5%	—		25		3		19	
	6 - 10%	—		12		4		10	
	11 - 15%	—		4		1		4	
	16 - 20%	1	1	4	9	—	0	—	4
	21 - 30%	—		2		—		2	
	31 - 40%	—		3		—		—	
	41 - 50%	—		—		—		1	
	100%	—		—		—		1	
Occurrence	13		52		19		38		
Frequency of Occurrence	87%		76%		83%		72%		
mean Percentage	+		8%		7% (8 stations)		7%		

Table 49. Occurrence of *S. bedoti* in the Banda Sea and the Moluccas Archipelago.
* less than 1%.

50	Flores Sea	SIBOGA 1906		SNELLIUS 1941	
	Number of sampling stations	11		42	
	+ or *	6	6	2	27
	1 - 5%	—		10	
	6 - 10%	—		8	
	11 - 15%	—		7	
	16 - 20%	—	0	1	2
	21 - 30%	—		1	
	Occurrence	6		29	
	Frequency of Occurrence	55%		69%	
	Mean Percentage	+		8%	

Table 50. Occurrence of *S. bedoti* in the Flores Sea.
* less than 1%.

51	Boni Bay in the Flores Sea	SNELLIUS 1941	
	Number of sampling stations	4	
	1 - 5%	1	2
	6 - 10%	1	
	11 - 15%	—	
	16 - 20%	1	2
	33%	1	
	Occurrence	4	
	Frequency of Occurrence	100%	
	Mean Percentage	16%	

Table 51. Occurrence of *S. bedoti* in Boni Bay, the Flores Sea.

52

	Timor Sea		Arafura Sea		
	SNELLIUS 1941	TOKIOKA 1956	SNELLIUS 1941	TOKIOKA 1956	
Number of sampling stations	12		18		
1 - 5%	3	6	3	9	
6 - 10%	3		5		4
11 - 15%	1		1		10
16 - 20%	1		0	3	7
21 - 30%	1	—		9	
31 - 40%	—	—		3	21
41 - 50%	—	—		—	20
Occurrence	9		12		
Frequency of Occurrence	75%		67%		
Mean Percentage	9%		9%		

Table 52. Occurrence of *S. bedoti* in the Timor Sea and the Arafura Sea.

53

Savu Sea	SNELLIUS 1941	SIBOGA 1906
Number of sampling stations	20	
1 - 5%	5	1
6 - 10%	3	
11 - 15%	3	
16%	1	
Occurrence	12	
Frequency of Occurrence	60%	
Mean Percentage	7%	

Table 53. Occurrence of *S. bedoti* in the Savu Sea.

54	Visakhapatnam Oct. 1952~Apr. 1953		Centra part of the Indian Ocean	
	10~15 miles off shore	more than 20 miles off shore	TOKIOKA 1956	
Number of sampling stations	13	21	Number of sampling stations	26
1 - 5%	1	—	1 - 5%	2
6 - 10%	—	—	6 - 10%	6
11 - 20%	—	1	11 - 15%	4
21 - 30%	1	—	16 - 20%	4
33%	—	1	21 - 30%	5
Occurrence	2	2	Occurrence	21
Frequency of Occurrence	15%	10%	Frequency of Occurrence	81%
Mean Percentage	15%	22%	Mean Percentage	15%

Table 54. Occurrence of *S. bedoti* in the Indian Ocean.

1911), *S. bedoti* was collected in the south-western part of the ocean from the surface water off Port Natal. On the opposite side in this ocean, it is reported from Sharks Bay on the western coast of Australia (RITTER-ZAHONY, 1910a). During the "Sea lark" Expedition in the western half of the Indian Ocean in 1905, *S. bedoti* occurred in only small numbers, the frequency of occurrence was also quite low, 12% (BURFIELD, 1926). GHIRARDELLI (1948) reported that this sagitta occurred in the waters along the coast of Somaliland at a frequency of 34% (of 44 stations) and the number of specimens throughout the whole samples constituted 4% of the total chaetognaths.

In the Bombay Harbour, this sagitta is the commonest species and is found there throughout the year in considerably large numbers (LELE and GAE, 1936). On the eastern side of India, the species was not found in coastal water of Madras (JOHN, 1933), but later its occurrence in this region was ascertained by SUBRAMANYAM (1940). SATYANARAYANA RAO (1958) has recently shown that *S. bedoti* is as common as *S. enflata* throughout the year in Lawson's Bay, Waltair, East India. It is most abundant in January-February and June-July, but it diminishes in the offshore water and it is quite absent there in February and October. Further, SATYANARAYANA RAO and GANAPATI (1958) discuss the occurrence of *S. bedoti* in the waters off Visakhapatnam Harbour south of Waltair as follows: the species is "very common in the inshore waters during most of the year; recorded rarely in the offshore waters, mostly surface dwelling (p. 150)". My examination (TOKIOKA, 1956) of 26 plankton samples collected in the years 1954-1955 in the central part of the Indian Ocean showed that this sagitta was one of the commonest species in this area. Also, I had an opportunity to examine three plankton samples from the north-eastern part of this ocean, near Nicobar, Nias and Bali-Lombok. In each of these samples, *S. bedoti* was found,




Fig. 28. Distribution of *Sagitta bedoti* in the Indian Ocean.

though only as 4, 3 and 9 percent of the whole population respectively. *S. bedoti* was not found by STEINHAUS (1896) in the southern part of the Indian Ocean nor was it found during the Gazelle Expedition (RITTER-ZÁHONY, 1909b). GRAY (1923) also failed to find this species in samples taken in the southern Indian Ocean. In the Red Sea, *S. bedoti* was not found by either Pola (RITTER-ZÁHONY, 1909a), Snellius, nor Cherso (GHIRARDELLI, 1948).

i) *The waters along the eastern coast of Australia.*

Although both JOHNSTON and TAYLOR (1919) and TOKIOKA (1940b) could not find this sagitta in the southern part of this area, THOMSON (1947) found it in the south-eastern waters of Australia and BURFIELD (1950) in the Great Barrier Reef region. In the south-eastern waters, it occurred at 25 stations, this constitutes 9.1% of all sampling stations. According to Fig. 4 in THOMSON's paper, it was "infrequent" in the area northward to 28°S, but "rare" in the area between 28°S to 35°S. The frequency of occurrence was 8.8% in the oceanic zone, while it was 9.2% in the neritic zone. In the Great Barrier Reef region, it occurred inside the reef at 25% of 48 stations and the number of specimens throughout the whole collection was 3.4% of the total chaetognaths, while outside the reef the frequency of occurrence was 23% of 13 stations, and the percentage of specimens of this sagitta to the total chaetognaths was about 0.3%. Thus, BURFIELD states as follows:—"When present, this was fairly abundant inside the reef, but was very scarce outside. This is considered to be warm water form of the Indian and Pacific Oceans, and found from the surface to 50 m".

j) *The far oceanic waters of the Pacific, excluding the coastal waters along the continents, the Japanese and Philippine Islands and the Malay Archipelagoes (Table 55).*

During examination of a lot of plankton samples collected in the waters around the Palao Islands of the western Caroline Islands in 1940, I found *S. bedoti* in the material, but in only very small numbers. It appears to be distributed in the counter Equatorial Current, the South Equatorial Current, and the New Guinea Coastal Current, but rarely found in the North Equatorial Current and in the waters around the Palao Islands. In lagoons of the Palao Islands, this species was not found during my observation for eight months (1940-41). Later, I examined two other collections of chaetognaths from the equatorial region of the Pacific; a small series collected in 1953 near the Phoenix and the Gilbert Islands in the central Pacific, and the material collected in 1954 by the "Syunkotu-Maru" during her cruise to survey the after effects of a series of experiments with atomic and hydrogen bombs at Bikini Atoll. In both cases, I could not find any *S. bedoti*. ROSE (1952) examined plankton samples from lagoons of the Tuamotu Islands, but found no *S. bedoti*. Both HIDA's paper (1957) based on the material collected by the Pacific Oceanic Fishery Investigations of Hawaii in the middle North Pacific and BIERI's comprehensive report (1959)

55

Near the Western Caroline Islands	Around the Palao Islands	North Equa- torial Current	Counter Equa- torial Current	South Equa- torial Current
Number of sampling stations	22	24	19	16
* 1 - 5%	no data	—	1	2
* 6 - 10%		—	4	1
* 11 - 15%		—	2	6
* 16 - 20%		—	—	1
* 21 - 30%		—	1	—
* 31 - 32%		—	1	1
Occurrence	2	1	14	13
Frequency of Occurrence	9%	4%	68%	88%
* mean Percentage	+	+	13%	13%

* Data obtained from the examination of 31 plankton samples hauled during the Oceanographic surveys of Palao~New Guinea in March~May 1940.

Table 55. Occurrence of *S. bedoti* in the material of Palao~New Guinea cruises.

based on the material from the Equapac and the Troll Expeditions and also other collections show that *S. bedoti* does not occur in the vast central area of the northern half of the Pacific.

k) *The Atlantic Ocean.*

The only record of *S. bedoti* from the Atlantic is presented by THIEL (1938) who recorded that only a single specimen of *S. bedoti* was found at Station 20 (50-100 m) near Cape Town during the German Meteor Expedition in the Atlantic Ocean. It is possible that *S. bedoti* might be carried into the Atlantic Ocean from the Indian Ocean by the southward Mozambique Current and then westward by the Agulhas Current along the southern coast of Africa, because this sagitta occurs near Port Natal as mentioned in section (h). This is very probable, because *S. neglecta* and *S. regularis*, both typical Indopacific warm water species, are also found in the Benguela Current off the west coast of the Union of South Africa, being carried there by the Agulhas Current (HEYDORN, 1959).

IV. The identity of *S. bedoti* from various localities.

One can be somewhat skeptical about the record of *S. bedoti* from the Atlantic Ocean, because it is not impossible that an imperfectly preserved immature specimen of *S. friderici* might be mistaken for *S. bedoti*. In some cases, the distinction between these two species can be made only on the presence (*bedoti*) or absence (*friderici*) of the rayless zones on the lateral fins. This character becomes easily obscured in some poorly preserved specimens.

Next, are the specimens of *S. bedoti* occurring in the eastern Pacific along the

western coast of the equatorial Central and South Americas identical with those from the waters around the Philippines and Malay Archipelagoes? Examining closely the specimens from the eastern Pacific, I could affirm that they were quite identical with the so-called forma *minor* of *S. bedoti* found abundantly in the waters of the western tropical Pacific, as already shown in the section on taxonomic descriptions.

Lastly, are the specimens of *S. bedoti* from the Japanese waters identical with those found in the southern seas? For instance, THOMSON (1947) states that my records of *Sagitta bedoti* from the waters around Korea in 1940 might be erroneous, because they are much larger than those found in the southern seas. However, I believe they are truly *bedoti*. All characteristics, except for the body length, are quite common to specimens from both Japanese waters and the southern seas. The difference in body length is somewhat remarkable (30 mm to about 15 mm), but we find many other examples of such remarkable difference in body length. For instance, *Sagitta robusta* matures at 10 mm in the southern seas, but it reaches 21 mm in the water of somewhat low temperature in the Japan Sea; *S. enflata* is usually 15 mm or thereabout when mature, but it may attain 30 mm in the waters off Peru. Moreover, *S. bedoti* of the East China Sea and the Yellow Sea and that of the waters around the Malay Archipelagoes can be intermixed by the southward current in winter and the northward flow in summer through the South China Sea, although the strong influx of the North Equatorial Current seems to lower the density of this sagitta in this sea and lessen intermixing on a large scale. As far as concern the specimens examined by myself, there is not any doubt, I believe, that only a single species *S. bedoti* is here under consideration at present.

V. General account of the distribution of *S. bedoti*.

My acquaintance with *S. bedoti* gained through studies carried on at the Mitui Institute of Marine Biology near Simoda, Izu (1938), the observations made in the eastern part of the Inland Sea, namely Ōsaka Bay and Kii-Channel (1939), and the examination of the chaetognaths material from the waters around western Japan (1940), and also through references accessible for me by that time, led me to believe that *S. bedoti* should be one of the common warm oceanic water species, probably being distributed in the Pacific as widely as many other warm water species. Hence, it appeared very strange to find that this sagitta was quite rare in the waters around the Palao Islands of the western Caroline Islands, when I examined, in 1940, a lot of plankton samples hauled in this area. However, when I had finished the examination of chaetognaths collected from the central equatorial Pacific, extending from the Gilbert and Phoenix Islands to the Marshall Islands, I was aware of the possibility that the scantiness of *S. bedoti* in far oceanic water of the Pacific was not a temporal phenomenon but it is a distinct general feature. Consequently, I paid attention to the currents flowing in the western North Pacific and in the equatorial region, especially to the relation between the surface waters of the North Equatorial Current

and the Kuroshio. The Pacific coast of Japan is said to be washed by the Kuroshio. Therefore, if the greater part of the surface water of the Kuroshio were purely an extension of the North Equatorial Current, then *S. bedoti* would not appear abundantly along the Pacific coast of Japan. Furthermore, the contamination of the sea water by the fission products emitted by experiments at Bikini would be observed near the coast more frequently and in higher degree than it actually was. Of course, there could be water masses of somewhat coastal character along some parts of the Pacific coast of Japan and they may maintain *S. bedoti* and lessen the direct effect of the Kuroshio. Moreover, there should be the lateral mixing of water along the left (west) side of the Kuroshio, between it and the coastal water along the continent. This mixing might be carried on to a much larger scale than has been supposed. It is possible that *S. bedoti* may here be available as an indicator of the degree of this mixing in some parts of the Kuroshio. To clear up these points, I started to collect data regarding the occurrence of *S. bedoti*, and to outline preliminarily the distribution of this species (TOKIOKA, 1957) from the plankton material collected by the Sôyô-Maru over a wide area extending from the northern Japan to the South China Sea. Now, with BIERI's comprehensive report (1959) on the distribution of chaetognaths in the Pacific together with personal observations on the large plankton collections present at Scripps Institution of Oceanography and with data from various other papers and personal communications all indicating an absence of *S. bedoti* in the mid Pacific, I believe the general features of the species can be discussed. Of course, the distribution will differ somewhat according to seasons. The differences are especially remarkable in the Indian Ocean and in some parts of the western Pacific, where the direction of currents may be quite changed with seasons. It is important to know if sampling stations and times were distributed evenly in a certain area, or to know whether or not this sagitta shows some peculiar vertical distribution. Concerning the latter point, so far as I am aware, this species can be considered an inhabitant of the surface water, probably above 100 m, most abundantly above 50 m. As to the seasonal variation in distribution and the even distribution of sampling stations and sampling times, I do not believe the data at hand are sufficient, but still I think they are sufficient to show the general outline of the distribution.

Sagitta bedoti is distributed widely in the Indian Ocean excluding the southern part of the ocean. In general, it is not abundant there, usually less than 10% of all chaetognaths, although it occurs commonly and abundantly in some embayments such as Bombay Harbour on the western coast and Lawson's Bay on the eastern coast of India. It occurs widely in the waters around the Malay Archipelagoes, but there also its percentage of the chaetognath fauna is usually less than 10%, excepting some cases in Boni Bay in the Flores Sea and in the Arafura Sea in some seasons. Most probably in the Indian Ocean and the waters around the Malay Archipelagoes, *S. bedoti* propagates in certain embayments or bays, out of which it drifts and is distributed widely in the ocean by the remarkable water movements found in these

seas. As the currents vary remarkably in the Indian Ocean and around the Malay Archipelagoes depending upon season, the abundance of *S. bedoti* in these areas also fluctuates much according to season. At times waters-masses of somewhat coastal character and containing many *S. bedoti* may drift out into the central part of the Indian Ocean and give rise to the records of higher percentages in the middle of the ocean. The population of this species in the Arafura Sea is continuous with that along the eastern coast of Australia. Since there is no report indicating dense population of the species in the coastal waters of the eastern Australia, the population of this sagitta in certain portion of this area seems to be supplied in part from the Arafura Sea at least in the season when the currents in the Arafura Sea flow towards the east.

S. bedoti occurs in the South China Sea and in Philippine waters, but only infrequently and in very small numbers; for instance, in the Philippine waters it is recorded only from the Sulu Sea. Possibly this is due to the effect of the strong influx of the North Equatorial Current which is quite devoid of this sagitta. The species increases towards the north along the continent and attains a remarkably dense population (nearly 100% of all chaetognaths and also at a frequency of 100% in all plankton hauls) in the coastal part of the East China Sea and in the southern part of the Yellow Sea. It is usually absent in the south-eastern part of the East China Sea near Okinawa, where the northward extension of the North Equatorial Current flows. However, it appears in the north-eastern part of the sea, where the water of the East China Sea is conducted out by the strong flow of the Kuroshio, and reaches a considerable density near the southern end of Kyûsyû Island. Its population near the southern end of Kyûsyû seems to fluctuate rather violently according to season and also according to hydrographical conditions changing from time to time. This species increases steadily in numbers from the western coast of Kyûsyû to the Korea Straits, where the waters of the East China Sea and the Yellow Sea are mixed in a considerable degree with the water of a branch of Kuroshio, Tusima Current. In the Japan Sea it may constitute 100% of the chaetognath fauna at some stations along the boundary between the Liman and Tusima Currents. It also increases steadily in numbers along the Pacific coast of Honsyû, the mainland of Japan, and attains its maximum in the waters off the north-eastern part of Honsyû along the boundary between the Oyashio and the Kuroshio. The distribution may extend eastwards to 161°39' E; then it disappears completely in the vast area of the Pacific. Along the Pacific coast of Honsyû Island, this sagitta occurs commonly in the water mass on the coastal side of the Kuroshio, but diminishes remarkably in the main stream of the Kuroshio and disappears in the far oceanic water mass beyond the Kuroshio. There, the Kuroshio seems to constitute a barrier between the Japanese coastal water mass which is evidently populated by various organisms from the East China Sea, and the far oceanic water.

The species occurs in an isolated small area at the eastern-most part of the equatorial Pacific. It is quite absent in the North Equatorial Current, the Counter

Equatorial Current, and the South Equatorial Current along their long course to the western Caroline Islands, where it appears again in insignificant numbers in the North Equatorial Current and increases in numbers towards the south. Probably *S. bedoti* is derived from the Sulu or Celebes Seas and carried eastward by the Mindanao Current and the Counter Equatorial Current. At first, I thought it possible that the population at the eastern-most part of the equatorial Pacific might be continuous with that in the south-eastern waters of Australia through the eastward current in the southern part of the South Pacific, but this is unlikely, because this sagitta decreases sharply at the southern stations in the Shellback area, quite disappears in the waters off Chile and at the offshore stations in the south-eastern waters of Australia.

VI. Possible factors limiting the distribution of *S. bedoti*.

At a glance, *S. bedoti* might be considered as an inlet or bay-form since it occurs abundantly in Bombay Harbour and Lawson's Bay of India and in Boni Bay in the Flores Sea. However, there is a genuine inlet-water chaetognath in the Indian waters, namely *Sagitta bombayensis*. Also in the Japanese waters, the inlet-waters are populated by *S. crassa*, its forma *naikaiensis* and *Sagitta delicata*. *S. bedoti* is absent in the waters where these species are dominant. Along the Californian coast, the inlet-waters or lagoons are populated by *S. tenuis*. Thus, *S. bedoti* cannot be considered as an inlet-water form. Can it then be a coastal-water form? However the occurrence of the species in abundance in the central part of the Indian Ocean and in the far offshore waters along the boundary between the cold Oyasio and the the warm Kuroshio in the western North Pacific, conflicts with this idea. Nevertheless, it is an obvious fact that this species is distributed along the coasts in very many places of the Indo-Pacific. The ranges of temperature and salinity in the areas where *S. bedoti* occurs are surprisingly wide. For instance, SATYANARAYANA RAO and GANAPATI give 26.6–29.5°C for temperature and 20.79–34.16‰ for salinity in the Indian waters; SUND (1959c) gives 9–27°C for temperature and 27.9–34.8‰ for salinity in the upper 300 meters at all stations inspected where this species occurred during the Eastrop Expedition; the lowest salinity in my data is 30.95‰ in the surface and 31.89‰ in the deeper layer in the Korea Straits and the lowest temperature is 11.5°C in the Yellow Sea. Thus, it is evident that *S. bedoti* is euryhaline and eurythermal and this seems to support the idea that it is a coastal-water form or a form which is fond of the mixed water. As to the latter point, actually most of the coastal waters where *S. bedoti* occurs in high numbers are found near areas of mixing or the boundaries between different waters. In the Yellow Sea and the East China Sea, the far oceanic water of the western North Pacific and the coastal water of somewhat lower salinity are mixed. In the Japan Sea, the cold Liman Current and the warm Tusima Current mingle with each other. Off the North-Eastern part of Honsyû Island of Japan, the cold Oyasio and the warm Kuroshio waters mix. In the eastern part of the equatorial Pacific, some upwelled water and the cold

Humboldt Current mingle with the tropical warm surface water. However, the biological conditions found in the mixing areas of different waters are favourable not only for *S. bedoti* but also for many other chaetognaths under certain circumstances; for instance in the Shellback area in the eastern equatorial Pacific many species of *Sagitta* thrive. On the other hand, in the central part of the Indian Ocean, where no remarkable mixing of different waters can be seen, a comparatively dense population of *S. bedoti* is maintained at least in some seasons. Why the same chaetognath which occurs in the central part of the Indian Ocean cannot survive in the mid Pacific Ocean is a moot question. The temperature-salinity relationships seem to be insufficient to explain the facts regarding this eurythermal and euryhaline chaetognath. Are there then some special chemical or biological factors (through food organisms, predators or competition) which differ between the Indian Ocean and the Pacific? So far as I am aware, there has not been as yet any definite answers to this question.

VII. "*Bedoti*-water".

Then, the following idea is considered possible that *S. bedoti* has specialized from a certain ancestor in the water mass which to-day contains this species in a considerable density and that the mixing of the water between this water mass and other water masses has been carried on in such a scale that the density of *S. bedoti* in the latter, dispersed there from the former, is always maintained below the lowest population level at which this sagitta can propagate. *S. bedoti*, when dispersed into far oceanic water might survive, but might be unable to reproduce below a certain population level, because self-fertilization is impossible. The lowest density at which the reproduction of the species can be maintained should differ specifically and it is conceivable that this limiting density might be comparatively high for *S. bedoti*. In other words, a comparatively high degree of stability can be expected in water masses containing *S. bedoti* in considerable numbers. In the East China Sea, the Yellow Sea and the middle part of the Japan Sea, the water is evidently moving, but it is considered to be circulating in a semi-closed system and thus considerable stability is maintained there. As regards propagation of various plankton organisms in areas of mixing, it is evident that a certain degree of stability of the mixed water is required for reproduction. Thus, it seems reasonable to separate the water masses containing *S. bedoti* from far oceanic water masses free from this species. It is proposed here to call the former "*bedoti*-water" because of this characteristic indicator. In the eastern Pacific, *S. friderici* shows a distribution somewhat like that of *S. bedoti* and actually occurs in the Shellback area together with *S. bedoti*. Besides these, *S. neodicipiens* and a peculiar form of *S. ferox* (f. *americana*) occur solely in this eastern *bedoti*-water. In the western North Pacific, the distribution of a calanoid copepod *Calanus lighti* is limited to far oceanic water and does not occur in the *bedoti*-water, some of the euphausiids distributed widely in the western North Pacific seem to be

slightly differentiated in some parts of the *bedoti*-water, the Japan Sea or in the East China Sea. All these facts seem to be attributable to the stability of the *bedoti*-water.

It might be possible to judge the grade of segregation of water-masses containing *S. bedoti* from the far oceanic water mass free from *S. bedoti* by measuring the density of this chaetognath. For instance, in the South China Sea, the low population of *S. bedoti* indicates a strong influx of the North Equatorial Current; near the southern coast of Kyûsyû, the Pacific coast of Honsyû and also in the Japan Sea the population of this species fluctuates considerably, probably according to the relative amount of influx of far oceanic water. In the Japan Sea, the average percentage of *S. bedoti* differs remarkably according to season, namely 30% in April, 16% in July and only 5% in September (1951-53). These data agree completely with the variation in strength of the Tusima Current and consequently the extent of far oceanic water.

As to the dense population of this chaetognath in the area of mixing between the Oyasio and the Kurosio off the north-eastern part of Honsyû of Japan, the distribution in this area cannot be considered as allo-genetic one, but rather auto-genetic distribution, because there occur very many young individuals (12 to 74%, 49% on an average in examined samples) referable to *S. bedoti*, and moreover it frequently constitutes 100% of the whole chaetognath population. Thus, the water in this area is considered to show a considerable degree of stability. I cannot judge whether this stability is attributable simply to the effect of SCHRÖDER'S law applicable along the northern border of the Kurosio flowing eastwards or maintained by complex horizontal or vertical circulations; the questions will be answered by further works by hydro-graphers. At present, it is also unknown where the eastern distal end of this *bedoti*-water goes. It might extend towards the north east, where it is cooled below the critical temperature for *S. bedoti*, or it might be dispersed in the western North Pacific Central Water Mass below the limiting density required for propagation of this species. The wide distribution of *S. bedoti* in the Indian Ocean might be attributable partly to the smallness of this ocean itself or the existence of complex currents in this ocean.

7. THE FAR OCEANIC WATER OF THE CENTRAL NORTH PACIFIC

The remaining vast far oceanic water mass of the Pacific is divided into northern and southern halves by the Equatorial Currents. As to the possible differences in chaetognath fauna between these two halves and between the Western North Central Pacific and the Eastern North Central Pacific Water Masses, my data are less extensive than those of BIERI (1959), hence I shall not repeat any comments on this problem. However, I think it is useful to record here only the proportions between the four most commonly occurring species, mentioned below, in far oceanic waters and speculate as to the possibility of any constant differences in the chaetognath fauna between the above-mentioned divisions of the far oceanic waters. BIERI (1959) showed only the distribution of individual per unit volume of water for respective

species in the Pacific. The various collections I have dealt with in this paper were collected by various kinds of gears and it is certain that even the same gear might not be used always in quite the same manner. However, the ratios among certain dominant species seem to be rather stable, because the effect of different gears and the manner in which they were used is minimized by this procedure. The errors which inevitably occur during the course of computing the individuals per unit volume of water might also be much reduced in this representation.

In Table 56, it is seen clearly that *S. enflata* is the commonest of the species collected in the warm oceanic waters of the North Pacific, next in order is *S. serratodentata pacifica*, *P. draco*, and *S. hexaptera*, as I previously noted on material from the waters around the Marshall Islands (1955). In the material of the Transpac Expedition, *S. minima* (F. O.*—39%, average percentage—30) and *S. regularis* (F. O.—26%, average percentage—9) were the important species; and in the Shellback material, *S. serratodentata tasmanica* (F. O.—98%, average percentage—8) occurred commonly and followed by *K. subtilis*, *K. pacifica*, *S. minima*, *S. regularis*, *S. lyra* and *S. decipiens*. Therefore, I calculated the three ratios *S. serratodentata pacifica*/*S. enflata* (S/E), *S. hexaptera*/*S. enflata* (H/E) and *P. draco*/*S. enflata* (P/E); and plotted the values of S/E in Figs. 29–31.

Before deducing any tendencies from the figures and tables, it is necessary to consider whether or not the greater part of the population of each of the above-mentioned four dominant species occurs in the surface water above 50 m. If it does not and if some of these species occur very abundantly in layers deeper than 50 m under some conditions or in some regions, then the comparison of ratios throughout the whole materials consisting of samples collected from various depths would be quite meaningless. Very fortunately, however, the dominant species, here selected, and also most of the common species in the warm water are concentrated in the upper 50 m layer, as mentioned by THIEL (1938) in the report of the German Atlantic Expedition by the "Meteor". HIDA (1957) mentions that "On Hugh M. Smith cruise 30 large catches, as number per unit of water strained, of Chaetognaths and Pteropods were taken in the shallow (0–40 m) hauls than in the deeper (0–140 m) hauls both during the day and at night". The tendency of chaetognaths to concentrate in the upper water seems to be as true in the Pacific as in the Atlantic. Hence, a comparison of ratios on world wide collections seems to be significant in a search for trends, although the values themselves in different samples collected from different depths cannot be estimated strictly on the same level. This point will be most pronounced in the case of *P. draco*, if this species is distributed in the Pacific in the same manner as in the Atlantic, where the centre of the distribution of this species is moved to the layer slightly deeper than in other three species.

The ratios calculated on the material from the Transpac Expedition, in which a small net (17 cm in diameter) was used and the catches were very small, fluctuate

* F. O....Frequency of Occurrence

56	TP	J	MP	SB	EQP	P-N
<i>Sag. hexaptera</i>	80	—	748	588	1300	70
<i>Sag. lyra</i>	4	—	—	146	—	—
<i>Sag. enflata</i>	644	828	1944	3300	4000	3395
<i>Sg. bipunctata</i>	299	65	19	25	200	—
<i>Sag. ferox</i>	18	117	8	—	60	264
<i>Sag. ferox</i> f. <i>americana</i>	—	—	—	186	—	—
<i>Sag. robusta</i>	75	—	63	48	100	—
<i>Sag. bedoti</i>	204	1008	—	224	—	897
<i>Sag. pulchra</i>	—	—	*	78	40	49
<i>Sag. friderici</i>	16	—	—	50	—	—
<i>Sag. pacifica</i>	2738	693	2520	2200	1700	1720
<i>Sag. tasmanica</i>	1	—	—	784	—	—
<i>Sag. pseudoserratodentata</i>	45	—	112	34	20	—
<i>Sag. neglecta</i>	12	39	24	42	*	416
<i>Sag. oceania</i>	—	—	—	—	20	—
<i>Sag. regularis</i>	279	1035	14	134	200	1806
<i>Sag. bedfordii</i>	—	—	—	—	—	20
<i>Sag. crassa</i> f. <i>naikaiensis</i>	18	104	—	—	—	—
<i>Sag. minima</i>	1200	1587	154	192	*	—
<i>Sag. decipiens</i>	*	—	—	65	200	—
<i>Sag. neodecipiens</i>	—	—	—	23	—	—
<i>Sag. planctonis</i>	—	—	*	*	—	—
<i>Sag. macrocephala</i>	—	—	6	—	—	—
<i>Pterosag. draco</i>	600	—	1470	864	1700	869
<i>Ktta. subtilis</i>	28	—	105	276	80	3
<i>Ktta. pacifica</i>	217	—	11	162	80	240
<i>Eukr. hamata</i>	—	—	8	*	—	—
Damaged specimens or juv.	3420	4576	2047	600	900	264

Table 56. Occurrence of warm water species in respective materials (samples from inlet waters excluded).

Numbers indicate the value *Frequency of Occurrence* × *mean percentage* in respective species. *...less than 1, TP...Transpac Expedition material, J...material collected by a Japanese survey ship and presented to the party of the Transpac Expedition, MP...Midpac Expedition material, SB...Shellback Expedition material, EQP...Equipac Expedition material, P-N...Material of the Palao~New Guinea cruises.

57	layers	<i>S. enflata</i>	<i>S. hexaptera</i>	<i>S. serratodentata</i>	<i>Pterosag. draco</i>
	0 - 50 m	26.8	5.6	42.6	4.57
	50 - 100 m	7.2	4.68	12.94	7.0
	100 - 200 m	1.05	1.36	4.0	1.4
	200 - 400 m	0.75	0.2	1.2	0.2
	400 - 600 m		0.02	0.08	0.05
	600 - 800 m			0.07	0.01
	800 - 1000 m			0.06	

Table 57. Vertical distribution of four important species in the Atlantic (THIEL, 1938).

rather violently, and consequently a comparison between these ratios and those of other catches collected by large nets from deeper waters is less significant. Here also, the efforts must be limited to seeking out trends. Throughout Figs. 29-31 and Tables 58-68, 70 and 73 the most remarkable trend may be noted in the distribution of the ratio between *S. serratodentata pacifica* and *S. enflata* (S/E).

a) **Distribution of S/E in collections examined** (Figs. 29-31, Tables 58-60).

The distribution of S/E shows a distinct tendency towards lower values in water masses inhabited by *S. bedoti* and higher in the *bedoti* free water masses. Generally speaking, in samples containing *S. bedoti* the values of S/E higher than 1.00 occur less frequently than the values lower than 1.00, while in samples containing no *S. bedoti* the values higher than 1.00 occur more frequently than the values lower than 1.00. The latter trend is, however, rather obscured in the Equapac Expedition material, the collection during the Palao-New Guinea cruises, and also in collections from the area within the distributional range of *S. bedoti*, such as the waters off north-eastern part of Japan and the Shellback area. Further, Fig. 29 shows that S/E is lower than 1.00 at most stations along the equatorial zone in the central Pacific, south of approximately 10°N. In other words, *S. enflata* is more abundant than *S. serratodentata pacifica* in the *bedoti*-water and also in the equatorial water mass in the far oceanic area of the Pacific, but decreases conspicuously and becomes less abundant than the latter in other parts of the far oceanic water of the Central North Pacific. The scarcity of *S. enflata* is especially marked in the eastern half of the Central North Pacific, east of 180°, and most pronounced in the north eastern quarter of the Central North Pacific shown in Fig. 29, where *S. enflata* is nearly absent. Even *S. serratodentata pacifica* diminishes prominently there and the greater part of the chaetognath fauna consists of *S. minima*. HIDA's paper (1957) on the chaetognath fauna in the central North Pacific also shows distinctly the scarcity of *S. enflata* throughout the subtropic and the transition zones extending from 145°W to 170°E. The boundary




Fig. 30. Distribution of S/E in the Japanese and adjacent waters.




Fig. 29. Distribution of S/E in the North Pacific.




Fig. 31 Distribution of S/E in the Shellback area.

58	Japanese N.E. Waters			Transpac Exp.			Palao New Guinea			Shellback Exp.			total		
	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E
0/0	3	2	3	2	1	2							5	3	5
0	7	2	7	7	3	4	18	2	1	1		1	33	7	13
* -0.10		2			1	1	1	1	1	8	1	6	9	5	8
0.11 — 0.50		2			1	2	1	9	14	15	12	13	16	24	29
0.51 — 1.00		1			1			4	3	3	11	7	3	17	10
1.01 — 2.00					1			3			2			6	
2.01 — 5.00									1		1			1	1
5.01 — 10.00															
> 10.01								1						1	
∞ (enflata 0)		1			1									2	
Number of samples	10			9			20			27			66		

	H/E	S/E	P/E
0	38	10	18
* -1.00	28	46	47
1.01-∞	0	10	1

Table 58. Occurrence of respective values of H/E, S/E and P/E in samples collected in the *bedoti*-water mass in materials examined. *...present, but less than 1%.

59

	Transpac Exp. incl. Stranger's samp.			Midpac Exp.			Shellback Exp.		
	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E
0/0	15	7	14	6	6	6			
0				11	2	10			1
* — 0.10				6	5	3	9	2	6
0.11 — 0.50				11	6	7	8	4	12
0.51 — 1.00				3	5	5	2	4	1
1.01 — 2.00				5	3	5	2	4	1
2.01 — 5.00				4	11	7		5	
5.01 — 10.00				1	3	2		1	
> 10.01					5	2		1	
∞ (enflata 0)	1	9	2	4	5	4			
Number of samples	16			51			21		

0	15	7	14	17	8	16			1
* — 1.00				20	16	15	19	10	19
1.01 — ∞	1	9	2	14	27	20	2	11	1

Table 59. Occurrence of respective values of H/E, S/E and P/E in samples collected in the *bedoti*-free water mass in the eastern half of the North Pacific in materials examined (east of 160°W). *...present, but less than 1%.

between the *bedoti*-water and the far oceanic water showing higher S/E and that between the latter and the equatorial water mass showing lower S/E are defined rather distinctly, but the distribution of S/E in other parts of the far oceanic water in the Central North Pacific does not show any clear discontinuous lines as far as the collections examined are concerned, although the above-mentioned trends are discernible.

b) Distribution of S/E in other parts of the Indo-Pacific based on available data.

i) *Near the Japanese waters.* In Tables 61-63 and 73 B are given values of S/E calculated from the following material: Sōyō-Maru (TOKIOKA, 1957); western Japan

60

	Palao New Guinea			Japanese N.E. Waters			Transpac Exp.			Midpac Exp.			Equapac Exp.		
	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E
0/0	1		1	2	1	2	21	5	14						
0	6	2	4	4	1	4	24	3	12						
* — 0.10	1	1	1						1					1	
0.11 — 0.50	1	2	2		1		3	2	8	3		1	4	2	4
0.51 — 1.00		1			1			3	3	1		4	1	1	1
1.01 — 2.00		2	1		1		2	7	2	1	1			1	
2.01 — 5.00								10	3		4				
5.01 — 10.00								3							
> 10.01								1							
∞ (enflata 0)		1			1		2	18	9						
Number of samples	9			6			52			5			5		

0	7	2	5	6	2	6	45	8	26						
* — 1.00	2	4	3		2		3	5	12	4		5	5	4	5
1.01 — ∞		3	1		2		4	39	14	1	5			1	

Table 60. Occurrence of respective values of H/E, S/E and P/E in samples collected in the *bedoti*-free water masses in the western half of the North Pacific in materials examined (west of 160°W). *...present, but less than 1%

(TOKIOKA, 1940); Kii-Channel connecting the eastern end of the Inland Sea with the open sea (MURAKAMI, 1959); small collection from the Japan Sea (FURUHASHI, 1953); New Yamato Bank in the Japan Sea (TOKIOKA, 1951); Sagami and Suruga Bays from May 1936 to April 1937 (TOKIOKA, 1954); and a dense patch of chaetognaths that appeared near the Seto Marine Biological Laboratory on Feb. 25, 1954 (TOKIOKA, 1954); together with some fragmentary data.

Tōkyō Bay—According to MURAKAMI's data (1959), in the 44 samples collected from the mouth of the bay, the following frequencies prevailed—*S. enflata* 47%, *S. hexaptera* 14%, *S. serratodentata pacifica* 5%, and *P. draco* 9%. The water of this region is evidently *bedoti*-water as mentioned previously (Frequency of Occurrence of *S. bedoti*—47%), and S/E is apparently lower than 1.00. *S. hexaptera* occurred in six samples, in the two of these the species exceeded *S. enflata* as 1:0 and 5:2.

SÔYÔ MARU 1934 1937—39	N.E. Japan (B) north to 40°N			*N.E. Japan (B) south to 40°N			South Western Honsyu						Sagami Bay bedoti water			Japan Sea Furuhashi 1953			N.Y. B.			
							bedoti water			no bedoti												
	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E	S/E
0/0	4	4	4	2	2	2	1	2	2										2	2	1	
0	2	2	2	2	2	5	7	6	8			1	1	3	2	2			2	1	2	
* — 0.10				1			1							1		1						4
0.11 — 0.50				1			1	3	2	1				2	1	3						4
0.51 — 1.00				1	2			1							2				1			
1.01 — 2.00					1										1							
2.01 — 5.00							1															
5.01 — 10.00																						
> 10.01																						
∞ (enflata 0)							1														1	
Number of samples	6			7			12			1			6			4			8			

* N.E.---- North Eastern (B)-----bedoti water															
0	6	6	6	4	4	7	8	8	10		1	1	3	2	2
* — 1.00				3	2		2	4	2	1			3	3	4
1.01 — ∞					1		2							1	

*---New Yamato Bank in the Japan Sea			
4	3	3	
	1		8
		1	

Table 61. Occurrence of respective values of H/E, S/E and P/E in plankton samples collected in the Japanese waters (1). *...present, but less than 1%.

SÔYÔ MARU 62 1934 1937—39	Waters off the Southern Kyusyu						East China Sea						South China Sea					
	bedoti water			no bedoti			bedoti water			no bedoti			bedoti water			no bedoti		
	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E
0/0										1		1						
0	2	1	4	1		2	2	2	2	1		1	5	3	6	3	3	3
* — 0.10		1												1				
0.11 — 0.50	1	1		1									1	1				
0.51 — 1.00	1										1							
1.01 — 2.00														1				
2.01 — 5.00		1			2													
5.01 — 10.00																		
> 10.01																		
∞ (enflata 0)											1							
Number of samples	4			2			2			2			6			3		

0	2	1	4	1		2	2	2	2	2		2	5	3	6	3	3	3
* — 1.00	2	2		1							1		1	2				
1.01 — ∞		1			2						1			1				

Table 62. Occurrence of respective values of H/E, S/E and P/E in plankton samples collected in the Japanese waters (2). *----present, but less than 1%.

TOKIOKA 63
1940

	Waters off the Western Kyusyu						Waters off the Southern Kyusyu					
	bedoti water			no bedoti			bedoti water			no bedoti		
	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E
0/0	4	1	3				1	1	1			
0	2	2	2				8	6	8	3	3	2
* — 0.10	1								3			2
0.11 — 0.50	1	1	2	1		1	3	3		1	1	
0.51 — 1.00		1			1			1				
1.01 — 2.00								1				
2.01 — 5.00												
5.01 — 10.00												
> 10.01												
∞ (enflata 0)		3	1									
Number of samples	8			1			12			4		

0	6	3	5				9	7	9	3	3	2
* — 1.00	2	2	2	1	1	1	3	4	3	1	1	2
1.01 — ∞		3	1					1				

Kii Suidō (Murakami 59)								
bedoti water			no bedoti					
H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E
22	22	21	27	26	27			
100	44	53	37	26	28			
	46	42		7	5			
	6	3		1	3			
	1	1		1	1			
	3	1						
				1				
				1				
		1		1				
122			64					

122	66	74	64	52	55
	53	46		9	9
	3	2		3	

Table 63. Occurrence of respective values of H/E, S/E and P/E in plankton samples collected in the Japanese waters (3). *...present, but less than 1%.

Bungo Channel—This is the channel connecting the south-western end of the Inland Sea and the open sea. Twelve samples were examined by MURAKAMI (1959). *S. enflata* occurred in four of them, but no *S. serratodentata pacifica*, *S. hexaptera*, nor *P. draco* were found in the material. This region is a mixing area between the *crassa*-water of the Inland Sea and the *bedoti*-water off south-western Honsyu.

The Japan Sea—In 32 samples treated by TOKIOKA (1940), *S. enflata* occurred at a frequency of 30%, while *S. hexaptera*, *S. serratodentata pacifica* and *P. draco* were quite absent. Of the material from the waters near the New Yamato Bank (TOKIOKA, 1951), only S/E of the vertical hauls are shown in Table 61. *S. hexaptera* was not found in the vertical hauls and only two specimens of *P. draco* occurred in a single sample. On the other hand, 13 of the 66 surface hauls contained *S. enflata*, but no *S. hexaptera*, *S. serratodentata pacifica* nor *P. draco* were represented in the material. The surface water of this region is evidently *bedoti*-water, but the water in deeper layers is mixed with *elegans*-water as in the region where FURUHASHI (1953) collected his samples (36°05.2' N–36°32' N×133°39.0' E–135°41.5' E). In these waters *S. minima* was very abundant as mentioned previously (p. 404). In 32 samples collected in the *crassa* or *bedoti*-water of the Korea Straits (TOKIOKA, 1940), *S. enflata* occurred at a frequency of 63% and *S. serratodentata pacifica* at a frequency of 6%, while both *S. hexaptera* and *P. draco* were absent.

The Yellow Sea—In twenty-six samples examined (TOKIOKA, 1940), no *S. enflata*, *S. hexaptera*, *S. serratodentata pacifica* nor *P. draco* were detected.

The East China Sea—In twelve samples collected in the characteristic *bedoti*-water near the Chu San Islands (TOKIOKA, 1940), *S. enflata* occurred at a frequency of 42%, but *S. hexaptera*, *S. serratodentata pacifica* or *P. draco* were quite absent.

All these data appear to indicate a tendency of *S. enflata* to be abundant while *S. serratodentata pacifica* is very scarce within the range of the distribution of *S. bedoti*, and this does not regard the presence or absence of *S. bedoti* in respective samples. However, in the areas where the *bedoti*-water is adjacent to far oceanic water or is mixing with the latter, some masses of various sizes of far oceanic water may be carried into the *bedoti*-water in various degree and consequently remarkably high values of S/E are sometimes met with as is shown in the case of the dense chaetognath patch that appeared at Seto.

ii) *The South China Sea and the Philippine waters* (Tables 62, 64 and 66). In the samples collected near Hainan Island (Table 62), S/E is lower than 1.00 in most cases. The value should be also very low near Nhatrang-Cauda, Viet Nam, because HAMON (1956) gives the following data: (A) In 58 samples...*S. enflata*—2644 individuals in the material, and occurred at a frequency of 59%; *S. hexaptera* did not occur in the material; *S. serratodentata pacifica*—only 4 individuals in the material, and occurred at a frequency of 2%; and *P. draco*—only 2 in the material, and occurred at a frequency of 2%. (B) In 17 samples collected by Dr. M. RANSON ...*S. enflata*—7796 individuals in the material, and occurred at a frequency of 100%;

64 Sn SNELLIUS Sb SIBOGA	Sulu Sea (Sn)						Sulu Sea (Sb)						Celebes Sea (Sn)					
	bedoti water			no bedoti			bedoti water			no bedoti			bedoti water			no bedoti		
	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E
0/0				1	1	1										14	11	13
0	6	1	6	4	1	4		1	1	1	1		35	3	17	21	7	19
* — 0.10		3	2				2	1	1			1	3	6	15		4	3
0.11 — 0.50		1			1								3	9	6	1	5	2
0.51 — 1.00	1	1												14	5	1	2	
1.01 — 2.00	1	2											1	5		1	2	
2.01 — 5.00					2								1	5	1		2	
5.01 — 10.00													1					
> 10.01														2			2	
∞ (enflata 0)																	3	1
Number of samples	8			5			2			1			44			38		

0	6	1	6	5	2	5		1	1	1	1		35	3	17	35	18	32
* — 1.00	1	5	2		1		2	1	1			1	6	29	26	2	11	5
1.01 — ∞	1	2			2								3	12	1	1	9	1

Table 64. Occurrence of respective values of H/E, S/E and P/E in plankton samples hauled in the waters around the Malay Archipelagoes (1). *...present, but less than 1%.

65
 Sn SNELLIUS
 Sb SIBOGA

	Celebes Sea (Sb)			Java Sea (Sn)						Flores Sea (Sn)					
	bedoti water			bedoti water			no bedoti			bedoti water			no bedoti		
	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E
0/0				1	1	1	4	1	3	1	1	1	6	5	6
0	2		3	20	2	7	13	7	10	40	6	27	24	11	17
* — 0.10	1	2	1	2	8	13	1	3		8	19	10		1	
0.11 — 0.50		2			7	4		1	3	4	9	13		5	6
0.51 — 1.00	1			2	5			3	1		6	1		4	
1.01 — 2.00					2				1		4			2	1
2.01 — 5.00								1			5	1			
5.01 — 10.00															
> 10.01											3			1	
∞ (enflata 0)								2						1	
Number of samples	4			25			18			53			30		

0	2		3	21	3	8	17	8	13	41	7	28	30	16	23
* — 1.00	2	4	1	4	20	17	1	7	4	12	34	24		10	6
1.01 — ∞					2			3	1		12	1		4	1

Table 65. Occurrence of respective values of H/E, S/E and P/E in plankton samples hauled in the water around the Malay Archipelagoes (2). *...present, but less than 1%.

66

Sn SNELLIUS	Moluccas Arch. (Sn)						Moluccas Arch. (Sb)						East South off Mindanao(Sn)					
	bedoti water			no bedoti			bedoti water			no bedoti			bedoti water			no bedoti		
	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E
0/0	1	1	1	10	2	7										11	4	12
0	36	5	15	24	6	17	4		2	1	1	1	10		10	16	3	14
* — 0.10	6	18	16		2	1	9	11	9	1		1		3	1		3	2
0.11 — 0.50	4	14	15	1	8	4	1	3	3				1	3	3		1	1
0.51 — 1.00	3	2	2		1	2					1			3		1	3	1
1.01 — 2.00		4	1		2								1	1		1	1	
2.01 — 5.00		5			1	1							1	1		1	3	
5.01 — 10.00					3								1	2			4	
> 10.01		1			2									1			1	
∞ (enflata 0)					8	3											7	
Number of samples	50			35			14			2			14			30		

0	37	6	16	34	8	24	4		2	1	1	1	10		10	27	7	26
* — 1.00	13	34	33	1	11	7	10	14	12	1	1	1	1	9	4	1	7	4
1.01 — ∞		10	1		16	4							3	5		2	16	

Table 66. Occurrence of respective values of H/E, S/E and P/E in plankton samples hauled in the waters around the Malay Archipelagoes (3). *...present, but less than 1%.

67 Sn SNELLIUS Sb SIBOGA	Boni Bay (Sn)						Banda Sea (Sn)						Banda Sea (Sb)					
	bedoti water			no bedoti			bedoti water			no bedoti			bedoti water			no bedoti		
	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E
0/0	1	1	1	1	1	1	3	1	2	9	4	9						4
0	3		1	1	1	1	63	10	36	28	9	23	2		1			1
* — 0.10		4	3				9	22	21		2	1	1	2	1	1		
0.11 — 0.50	1						4	23	15		5	2						
0.51 — 1.00							2	9	5	1	3	1			1			
1.01 — 2.00								8	2		5	1		1				1
2.01 — 5.00							1	7			3							
5.01 — 10.00											1							
> 10.01											1	1						
∞ (enflata 0)								2	1		5							
Number of samples	5			2			82			38			3			1		

0	4	1	2	2	2	2	66	11	38	37	13	32	2		1			1
* — 1.00	1	4	3				15	54	41	1	10	4	1	2	2	1		
1.01 — ∞							1	17	3		15	2		1				1

Table 67. Occurrence of respective values of H/E, S/E and P/E in plankton samples hauled in the waters around the Malay Archipelagoes (4). *...present, but less than 1%.

68

SNELLIUS

	Arafura Sea						Timor Sea						Savu Sea					
	bedoti water			no bedoti			bedoti water			no bedoti			bedoti water			no bedoti		
	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E
0/0				4		4										6	3	6
0	10	2	8	10	7	8	7		3	3	1	3	10	3	4	5	4	5
* — 0.10	1	6	6		1		2	5	5					3	3			
0.11 — 0.50	3	2			1	1		2	1				2	3	6			
0.51 — 1.00		1						1			1		1	2				
1.01 — 2.00		1			1			1						2				
2.01 — 5.00		1				1					1							
5.01 — 10.00		1															1	
> 10.01																		
∞ (enflata 0)					4												3	
Number of samples	14			14			9			3			13			11		

0	10	2	8	14	7	12	7		3	3	1	3	10	3	4	11	7	11
* — 1.00	4	9	6		2	1	2	8	6		1		3	8	9			
1.01 — ∞		3			5			1			1			2			4	

Table 68. Occurrence of respective values of H/E, S/E and P/E in plankton samples hauled in the waters around the Malay Archipelagoes (5). *...present, but less than 1%.

S. hexaptera did not occur in the material; *S. serratodentata pacifica*—18 individuals in the material, and occurred at a frequency of 18%; and *P. draco* did not occur in the material.

During the Albatross Expedition in the Philippine waters, chaetognaths were collected at 46 stations in the area extending 5°36'–21°31' N×117°53'–127°44' E. In this material, the four species under consideration occurred in the following proportion (MICHAEL, 1919): *S. enflata*—2800 individuals in the material, and occurred at a frequency of 85%; *S. hexaptera*—700 individuals in the material, and occurred at a frequency of 57%; *S. serratodentata*—100 individuals in the material, and occurred at a frequency of 15%; and *P. draco*—22 individuals in the material, and occurred at a frequency of 11%. It is evident that S/E must be very low in the Philippine waters.

iii) *The waters around the Malay Archipelagoes* (Tables 64–69 and 73 B). Many data from this area are divided into two groups, namely those from samples collected in the *bedoti*-water masses and those hauled in the water masses containing no *S. bedoti*. Throughout the Tables 64–68 and 73 B, it is easily noted that higher

69	<i>bedoti</i> water mass	non <i>bedoti</i> -water mass	
Sulu Sea (Sn.)	3.00	1.50	○
Sulu Sea (Sb.)	∞	∞	
Celebes S. (Sn.)	2.67	3.22	×
Celebes S. (Sb.)	∞	—	
Moluccas Arch. (Sn.)	4.00	1.19	○
Moluccas Arch. (Sb.)	∞	∞	
East south off Mindanao	1.80	0.88	○
Java Sea	11.50	5.00	○
Flores Sea	3.42	6.50	×
Boni Bay	∞	∞	
Banda Sea (Sn.)	3.82	1.53	○
Banda Sea (Sb.)	2.00	0.00	○
Arafura Sea	3.67	1.80	○
Timor Sea	8.00	2.00	○
Savu Sea	5.50	1.75	○

Table 69. The value of S/E lower than 1.00/S/E higher than 1.00 in various parts of the waters around the Malay Archipelagoes.

- ...the value higher in *bedoti*-water than in non-*bedoti*-water.
 ×...the value higher in non-*bedoti*-water than in *bedoti*-water.

70

	Central Indian Ocean						off Somaliland						near Aden			Red Sea		
	bedoti water			no bedoti			bedoti water			no bedoti			bedoti water			no bedoti		
	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E
%										1	1	1						
0	5		1	3		4	10	6	9	20	14	19		2	2	6	6	6
* — 0.10	8		4				1	1	2		2	1	2					1
0.11 — 0.50	6	1	12					2	1	1	2	1					1	
0.51 — 1.00	1	8	3	1	4		2	2			2							
1.01 — 2.00		3						1	1		1					1		
2.01 — 5.00		8						1										
5.01 — 10.00																		
> 10.01																		
∞ (enflata 0)																		
Number of samples	20			4			13			22			2			7		

0	5		1	3		4	10	6	9	21	15	20		2	2	6	6	6
* — 1.00	15	9	19	1	4		3	5	3	1	6	2	2				1	1
1.01 — ∞		11						2	1		1					1		

Table 70. Occurrence of respective values of H/E, S/E and P/E in plankton samples collected in the Indian Ocean and the adjacent waters. *...present, but less than 1%.

values of S/E seem to occur more frequently in the non-*bedoti*-water masses than in the *bedoti*-water masses. The value $\frac{\text{S/E lower than 1.00}}{\text{S/E higher than 1.00}}$ in various parts of the waters around the Malay Archipelagoes are given in Table 69. OYE (1918) examined samples collected at 77 stations in the Java Sea and showed the proportion of the four species as follows: *S. enflata* occurred at a frequency of 97% and constituted 31% of the whole collection of chaetognaths, *S. hexaptera* occurred at a frequency of 64% and constituted 5% of the whole collection of chaetognaths, while *S. serratodentata* did not occur and *P. draco* occurred at a frequency of only 1%. He states that the absence of *S. serratodentata* in the material is a result of only surface towing, while the species is probably a mesoplanktonic form. This is, however, not likely, because most samples of the Snellius material were also collected in the surface layer and yet they comprised a considerable number of *S. serratodentata*. However, his data can be safely accepted in support of the tendency of *S. serratodentata* to be very scarce in this area of *bedoti*-water as compared to the far oceanic waters. In 55 samples from the Arafura and Timor Seas, examined by TOKIOKA (1955b), *S. serratodentata pacifica* and *P. draco* were extremely scarce and *S. hexaptera* was quite absent. The water of the Arafura Sea was of the typical *bedoti*-water at that time.

iv) *The Indian Ocean* (Tables 70 and 73 B). Besides the data on the material from the central Indian Ocean (TOKIOKA, 1956), the material collected by the "Cherso" from the Red Sea and from the waters along Somaliland (GHIRARDELLI, 1948) and a small collection from the North Eastern part of the Ocean (Nicobar, Nias and Bali-Lombok) shown in Tables 70 and 73 B, there are the following fragmentary data:—

In a single sample collected near Ceylon (TOKIOKA, 1956), H/E was 0, S/E 6.5 and P/E was 2.5. In a single sample hauled in the Red Sea (300–0 m, in April 1929) by the Snellius (SCHILP, 1941), H/E was 4.6, S/E 0.4 and P/E was 0. It is somewhat strange that S/E is rather high even in the *bedoti*-water mass in the central part of the Indian Ocean. Since *S. serratodentata pacifica* can be used as an indicator of oceanic water in the waters off Visakhapatnam near Waltair (S. RAO and GANAPATI, 1958, p. 167) and BURFIELD (1926) shows that *S. serratodentata* is most abundant in the surface layer of this ocean, the above-mentioned phenomenon might indicate that the water of the surveyed area in the central Indian Ocean still retains in considerable degree, the characteristics of far oceanic water in spite of the existence of *S. bedoti*.

v) *Along the Eastern coast of Australia* (Tables 71–72). THOMSON (1947) gives the frequency of occurrence of the four species in the waters of south-eastern Australia as in Table 71. The rate of increase in the oceanic water is much higher in *S. pacifica* than in *S. enflata*, but probably *S. enflata* is more abundant than *S. pacifica* in both neritic and oceanic waters. BURFIELD (1950) shows the proportion of the four species

71 F.O.	at 161 neritic stations over the continent shelf		at 113 oceanic stations beyond the continent shelf	
	<i>Sag. enflata</i>	51.5%		54.0%
<i>Sag. hexaptera</i>	22.4%		58.4%	
<i>Sag. pacifica</i>	31.6%		46.0%	
<i>Pterosag. draco</i>	30.4%		62.8%	

Table 71. Distribution of four important species in the waters of the South-Eastern Australia (THOMSON, 1947).

72	inside the reefs		outside the reefs	
	<i>Sag. enflata</i>	110728 ind.		14735 ind.
<i>Sag. hexaptera</i>	243		837	
<i>Sag. pacifica</i>	3		64	
<i>Pterosag. draco</i>	5		74	

Table 72. Distribution of four important species in the Great Barrier Reef region (BURFIELD, 1950).

in the Great Barrier Reef region as above: S/E is clearly higher in the waters outside the reefs, although the dominance of *S. enflata* is still retained there. Both of these surveyed areas belong to *bedoti*-water.

vi) *Central part of the Pacific* (Table 72 A and B). The material of the Syunkotumaru (TOKIOKA, 1954) indicates distinctly that S/E is higher in the North Equatorial Current than in the Counter Equatorial and South Equatorial Currents. S/E is also lower than 1.00 in the material collected near the Phoenix and Gilbert Islands.

vii) *Off Chile* (Table 73 A). FAGETTI (1958) examined 220 samples collected from the waters off north and middle Chile. In her material, *S. hexaptera* occurred only once and *P. draco* did only five times, while *S. pacifica* occurred at a frequency of 36%. S/E seems to vary from time to time, but the dominance of *S. enflata* was maintained throughout the whole material.

All of the data, quoted here, indicates that *S. enflata* is very rich in the *bedoti*-water, while *S. serratodentata pacifica* increases towards the offshore oceanic waters and the S/E becomes higher in the far oceanic water, excepting the area along the equatorial zone. The influx of the North Equatorial Current retaining higher S/E flows into the waters around the Malay Archipelagoes, and consequently the values of S/E higher than 1.00 occur rather frequently in this area. The dominance of *S. enflata* in the Shellback area is maintained also in the waters off north to middle Chile, although the frequency of occurrence of S/E higher than 1.00 seems to be higher in the latter to some extent.

73 A

	Syunkotu maru 1954									off Chile S/E			
	N.Eq. Current			Count.Eq.Curr.			S.Eq. Current			'54 Aug.	'54 Dec.	'55 Mar.	'56 Jun.
	H/E	S/E	P/E	H/E	S/E	P/E	H/E	S/E	P/E				
0/0													
0	1	1	1							5	2	4	6
* — 0.10	1		1	2			4		2	1		3	5
0.11 — 0.50	4	2	3	4	4	6		3	2	8	8	7	7
0.51 — 1.00	5	4	5		3	1		1		6	4	2	1
1.01 — 2.00	3	3	5	1						1	4		3
2.01 — 5.00	2	6	2								5	2	1
5.01 — 10.00	1									1	1		
> 10.01		1										2	1
∞ (enflata 0)										1	1	2	2
Number of samples	17			7			4			23	25	22	26

0	1	1	1							5	2	4	6
* — 1.00	10	6	9	6	7	7	4	4	4	15	12	12	13
1.01 — ∞	6	10	7	1						3	11	6	7

Table 73. Occurrence of respective values of H/E, S/E and P/E in plankton samples came from various parts of the Pacific. *...present, but less than 1%.

B

	H/E	S/E	P/E	
Java Sea (Siboga)	0.22	0.07	0	B
Nicobar (Tokio'55)	0.10	0.30	0.90	B
Nias (Tok.'55)	0.12	1.12	1.00	B
Bali Lombok (Tok.'55)	0.05	0.58	0.49	B
Phoenix Gilbert (Tok.'54)	0.07	0.46	0.08	n.B
Sagami Bay (Tok.'54)	0.02	0.57	0.13	B 16%
Dense patch at Seto (Tok.'54)	0.02	1.06	2.97	B <1%

B Sag.bedoti present
n.B no bedoti

c) Distribution of H/E.

As the number of specimens of *S. hexaptera* collected during the Transpac Expedition with a small net from the surface water above 50 m is very small, it is quite impossible to establish any trends in the distribution of this species from the data of this expedition. As shown in Table 57 indicating the results of the German Atlantic Expedition by the "Meteor" the population of *S. hexaptera* is relatively large in deeper layers as compared with those of *S. enflata* and *S. serratodentata*. For instance, let us compare the two columns of Appendix Table 1 No. 4, which show the chaetognath populations of the two samples collected at the same station respectively by a small net from 50 m to surface and by a large net from 450 m to the surface. In the sample of *TP 131 from the deeper water, a considerable number of *S. hexaptera* were found and actually the number was larger than that of *S. enflata*. But the animals seem to easily avoid being caught when a smaller net is used or when the haul is limited to the surface layer. The same phenomena are seen also in the results of the Snellius Expedition, in which *S. hexaptera* occurred commonly in samples collected from the deep waters, but very scarcely in those collected by towing a small net in the surface water.

However, in examining the tables and data given in the preceding section (b, i-vii), it may be safely said that *S. hexaptera* shows a tendency to decrease in the *bedoti*-water and the adjacent waters, while increasing in the offshore far oceanic waters. The distribution of this species in a part of the far oceanic water of the Central North Pacific is shown by HIDA (1957), who illustrates very distinctly that the species is one of the most important chaetognaths in the subtropic and the transition zones between meridians 145°W-170°E and consequently H/E is enormously large in this area. TOKIOKA (1955) also stated that the proportion of *S. hexaptera* increases towards the north in a limited area in the central Pacific, covering the North Equatorial Current, the Counter Equatorial Current and a part of the South Equatorial Current. The raise of H/E towards the north in far oceanic waters cannot be considered to result only from the decrease of *S. enflata*; it can result by actual increase of *S. hexaptera*.

Looking through Tables 64-68 and 73 B, showing the proportional abundance of the four species near the Malay Archipelagoes, *S. hexaptera* and *P. draco* seem superficially to occur more frequently and abundantly in the *bedoti*-water masses than in the non-*bedoti*-water masses. But, this is rather questionable, because most of the larger samples collected from the deep waters are included in the column of the *bedoti*-water. And it is evident that the frequency of occurrence of *S. hexaptera* and *P. draco* is much higher in the larger samples from deep waters than in smaller samples collected by towing in the surface water, and the same applies to *S. bedoti*. Hence, the occurrence of the first two species together with *S. bedoti* should be more frequent than those two species alone in the region within the range of the distribution of *S. bedoti*.

d) Distribution of P/E.

The distribution of *P. draco* in the *bedoti*-water resembles that of *S. hexaptera*; it is rather scarce there, but increases considerably in far oceanic waters. In the far oceanic water of the Central North Pacific, the population of this species is larger in the southern half than in the northern half where it decreases very sharply and is scarcely found in the northern parts. HIDA's statement (1957) on the distribution of this species in the area, mentioned in the preceding section, shows clearly this trend; there, *P. draco* is abundant in the subtropic zone south to ca. 33°N, but decreases sharply in the transition zone.

e) Distribution of other prominent species.

i) *S. bipunctata*, HIDA (1957) includes this species as a very important form in the subtropic and transition zones, considering it comparable with *S. serratodentata*. Actually it occurred in considerable numbers at considerable frequencies in the Transpac Expedition collections as shown in Appendix Table 1, but it was much scarcer than *S. serratodentata pacifica*. It is not improbable that this species occurs abundantly in rather deep water and consequently is met with less frequently in the present material of the Transpac Expedition, which consists only of samples gathered from 50 m to the surface. BIERI (1959) say that the distribution of *S. californica* (= *S. bipunctata*) is antitropical, avoiding a narrow band in the equatorial region. The results of my examination on five of the samples of the Equapac Expedition show that *S. bipunctata* is not so rare in this region. It occurred also in considerable numbers in the material of the Syunkotu-Maru and also in the small sample from the Gilbert-Phoenix Islands, while it was quite absent in the materials of the Palao-New Guinea cruises. Probably, the population of *S. bipunctata* fluctuates rather violently in the equatorial region.

ii) *S. regularis*. According to the present data from the material examined, this species seems to occur more frequently and abundantly in the western half of the Central North Pacific than in the eastern half. It is found abundantly in waters adjacent to the *bedoti*-water in the western North Pacific, but this does not seem to suggest any special correlation between this species and *S. bedoti*, because the former is rather scarce in the *bedoti*-water of the Shellback area. In the central equatorial part of the Indian Ocean, the species occurs much more abundantly than does *S. hexaptera*.

iii) *S. minima*. The abundance of this species in or near the region of mixing between the *elegans*-water and the warm water has been repeatedly explained in previous sections. Its dominance is especially pronounced in the north-eastern part of the far oceanic water of the Central North Pacific, because *S. enflata* and even *S. serratodentata pacifica* are very scarce there. In the mixing area, it might easily be mistaken for young individuals of *S. elegans*, because the general external appearance is nearly the same between these two species. Very probably some slight errors are inevitable in counting these forms in the mixing area. However, the direction of

fin rays, which is vertical in *S. minima* rather than oblique as in *S. elegans*, can be used as the apparent characteristic in separating immature *S. minima* and young *S. elegans*. *S. minima* is also fairly common in the waters off North and Central Chile (FAGETTI, 1958). Sometimes it occurs in considerable density in the tropical waters as seen in the material from the central equatorial part of the Indian Ocean (TOKIOKA 1956a) and in the material of the Eastropic Expedition (SUND, 1959c), although it is still very difficult, at present, to explain clearly the significance of such distributions.

8. DIFFERENTIATION FOUND IN THE CHAETOGNATH FAUNA OF THE NORTH PACIFIC AND THE RELATION BETWEEN COMMUNITIES AND WATER MASSES

The contents of the foregoing sections may be summarized as follows:—

1) Chaetognath fauna in the embayments.

- i. *West side of the North Pacific*:- *Crassa*-water in the Inland Sea, Ise Bay, Tōkyō Bay and Mutu Bay of Honsyū, Japan, also in the coastal waters of Korea and in Kiauchau Bay of North China. *Delicata*-water in Maizuru Bay, Tanabe Bay, Katu-ura Bay and Ago Bay of Honsyū, Japan.
- ii. *East side of the North Pacific*:- *Tenuis*-water in lagoons and small bays or inlets along lower California.

2) *Bedoti*-water.

- i. *West side of the North Pacific*:- (1) In the mixing area between the *bedoti*-water and the *elegans*-water or between the former and the *crassa*-water in the Yellow Sea, respectively *S. elegans* and *S. bedoti* or *S. crassa* and *S. bedoti* represent the components of the fauna. In the former region *S. minima* and a small number of *S. enflata* may join the fauna at times.
(2) Then, there is a fairly wide area in the Yellow Sea and the East China Sea, where *S. bedoti* and *S. enflata* are the chief components.
(3) The greatest part of the *bedoti*-water is the mixing region between (2) and the warm oceanic water, where most of the warm oceanic species occur together with *S. bedoti*, although they are not so abundant as in the far oceanic water. Here, *S. enflata* predominates over other forms including *S. serratodentata pacifica*, *S. hexaptera* is very scarce. *P. draco* occurs in the southern part, but not in abundance. *S. regularis* is relatively common.
- ii. *East side of the North Pacific*:- The *bedoti*-water here is characterized as in (3) mentioned in the preceding paragraph. The predominance of *S. enflata* over *S. serratodentata pacifica* is maintained here also. *S. hexaptera* occurs in considerable numbers as does also *P. draco*, since the area is situated in the tropics. However, *S. regularis* is rather scarce. Other than these species, there occur also in this area *S. serratodentata tasmanica*, *S. neodecapiens*, *S. ferox* f. *americana* and *S. friderici*. These are quite unknown from the western *bedoti*-water and according to present data *S. neodecapiens* and *S. ferox* f. *americana* are strictly confined to this area. Thus, the difference between the western and eastern *bedoti*-water is somewhat remarkable.

6) *Pacifica*-dominant oceanic water.

In this vast area of far oceanic water, *S. serratodentata pacifica* predominates numerically over *S. enflata*. This area can be divided into a northern transition zone characterized by prominent increase in *S. minima* and *S. lyra* and a southern subtropical to tropical zone characterized by the abundance of *P. draco* as is shown by HIDA (1957). The northern part is much narrower than the southern part. At the same time, this area is divisible into western and eastern halves, although the distinction is not very clear in the distribution of chaetognaths. The frequent occurrence of numerous *S. regularis* and relatively large population of *S. enflata* contrasted with more scarce occurrence of *S. hexaptera* are characteristics of the fauna in the western half and the extreme scarceness of *S. enflata* and *S. regularis* and the comparatively abundant occurrence of *S. hexaptera* are the features characterizing the fauna in the eastern half. These are all deductions from only the available material that I have examined and hence it is not certain whether or not all these trends are absolutely of universal or constant features. BIERI (1959) says "the absence of *S. robusta* and *S. regularis*, with *S. pseudoserratodentata* present, would show that the sample came from the Eastern North Pacific Central Water". At least, the scantiness of *S. regularis* in the eastern half is common to both BIERI's and my statements.




Fig. 33. Differentiation found in the chaetognath fauna along the line through the Yellow Sea, East China Sea, waters off West-South Kyūsyū and extending to the North Pacific Central Water.

It is evident that (5), *enflata*-dominant Equatorial Water, corresponds to Pacific Equatorial Water and that (6), *pacifica*-dominant oceanic water, corresponds to North Pacific Central Water. The predominance of *S. serratodentata* over *S. enflata* seems to be a characteristic of far oceanic water, since in the South Atlantic *S. serratodentata* was about 1.7 times as numerous as *S. enflata* as a whole, when the individuals of each species in a cylinder of water, 1000 m deep and 50 m in diameter, were compared (THIEL, 1938). As to the approximate situation of the boundary between the western and eastern halves of the chaetognath fauna in the North Pacific Central Water, I cannot add anything to the comment by BIERI (1959), since the material I dealt with is much smaller than his. HIDA (1957) made a comparison of chaetognath faunas in the subarctic, transition, and subtropic zones along 180° and 157°30' W as found on




Fig. 34. Differentiation found in the chaetognath fauna along the line connecting 145°E x 40°N and 130°E x 0° (64) and along 180° meridian (65).

Hugh M. Smith Cruise 30. The slight narrowing of the transition zone along 157°30'W and the relative abundance of *S. minima* in the transition zone along 180° is shown in his Fig. 5, but these do not seem to indicate any distinct boundaries between the east and western Central Waters. Consequently, it is best, at present, to refrain from any attempts to correlate the Western and Eastern North Pacific Central Waters with the chaetognath fauna. The abun-

74

<i>Sag. serratodentata</i>	305
<i>Sag. enflata</i>	179
<i>Pterosag. draco</i>	66
<i>Sag. hexaptera</i>	59.3

Table 74. Proportion of four important species in the South Atlantic. (THIEL, 1938).




Fig. 35. Distribution of the *bedoti*-water and the *enflata*-dominant equatorial water plotted on the current chart from "The Ocean".

dance of *S. enflata* in the *bedoti*-water and the Equatorial water is significant in that both the *bedoti*-water and the Equatorial water hold denser population of not only chaetognaths but also of many other plankton organisms than does the North Pacific Central Water. *S. regularis* is abundant in the western half of both the North Pacific Central water and Equatorial water. It is very strange therefore that the species decreases greatly towards the east in spite of the existence of the eastward flowing Counter Equatorial Current. The degree of the lateral mixing along this current, which might induce dispersion or intake of various species of chaetognaths, must be studied more in detail in the future.

In the Indo-Pacific, *S. friderici* occurs only in the coastal waters along the western coast of the New World and the known occurrence of *S. tenuis* seems to confine it to the same region. These two species are reasonably considered as relicts of the Caribbean chaetognaths showing an ancient connexion between the Pacific and the Caribbean Sea, at which time they might have been carried into the Pacific by the westward Equatorial Current. But probably species characteristic to the Pacific, such as *S. bedoti* and *S. regularis*, could not penetrate into the Caribbean Sea against this current. At first, it appeared that *S. ferox* f. *americana* might be closely related with *S. hispida* or *S. helenae* in the Caribbean Sea, but this is not likely. The existence of more anterior teeth than posterior teeth in many individuals is merely a superficial coincidence between this and *S. helenae*; the existence of the distinct intestinal diverticula in *S. ferox* f. *americana* seems to be enough to separate it distinctly from *S. helenae* which lacks completely the intestinal diverticula. The distinction between this and *S. hispida* seems to be clear, though it is somewhat delicate as mentioned in the taxonomic part. Most probably this form differentiated from an ancestral *ferox*-group in the eastern *bedoti*-water as *S. neodecipiens* apparently did. *S. hispida* and *S. helenae* known commonly from the Caribbean Sea might have differentiated there after the connexion between that sea and the Pacific was closed and hence they do not occur in the Shellback area to-day. It is difficult to consider the western and eastern *bedoti*-waters in the Pacific as having developed independently of each other; long ago these two *bedoti*-waters might have been continuous with each other. The differentiation of *S. neodecipiens* and *S. ferox* f. *americana* in the eastern *bedoti*-water seems to have started after the continuity between the eastern and western *bedoti*-waters was cut off. The chaetognath fauna in the eastern *bedoti*-water is evidently nourished partly by the northward Humboldt Current, since there are found a considerable number of *S. serratodentata tasmanica* and a small number of a special form of *S. enflata* (f. *gardineri*) having longer ovaries. Both of these are known from the waters off North and Central Chile.

Summary and Conclusions

1. A part of the plankton material, 326 samples in all, collected by and stored at the Scripps Institution of Oceanography were examined and the data obtained

concerning chaetognaths are given in 7 appended tables. Data from samples of the Palao-New Guinea cruises (1940) made by a Japanese survey ship are also given in an appended table.

2. Thirty forms of chaetognaths were discriminated throughout the whole material, one of these is a new species and another is a new form. A comparison was made between Pacific specimens of *S. tenuis* and *S. friderici*. Some notes are also given on the morphology and the range of variations in certain other species.

3. The chaetognath faunas of several embayments, the blue-green water along the Californian coast and of the north Subarctic waters are covered briefly.

4. An extensive area characterized by the occurrence of *S. bedoti* is discerned as a water-mass, here called "*bedoti*-water", which is distinct from the far oceanic water of the North Pacific.

5. The ranges of temperature and salinity of the water containing *S. bedoti* are very wide, leading to the assumption that it is a eurythermal and relatively euryhaline form. The small amount of the interchange of water, under favourable conditions for the species, between the *bedoti*-water and the adjoining water masses is considered the most important factor confining the species to the *bedoti*-water. Very probably the number of *S. bedoti* that are diffused into the neighbouring water mass from the "*bedoti*-water" by interchanging of the water might be less than the minimum required by this species for successful reproduction.

6. The distributions of ratios between *S. enflata* and each of *serratodentata pacifica*, *hexaptera* and *P. draco* (S/E, H/E and P/E) were examined. And the relation between these distributions and water masses of the North Pacific was studied. The Equatorial Water Mass is characterized by lower S/E and the North Pacific Central Water Mass is remarkable in maintaining a higher S/E.

7. The distinction between "*bedoti*-water" and far oceanic water of the North Pacific Central Water Mass and that between the latter and the Equatorial Water Mass were made clear by the studies mentioned above in 6. But the boundary between the Western North Pacific Central and the Eastern North Pacific Central Water Masses was not found distinctly by these studies, although the density of some species seems to trend towards one of these two water masses.

REFERENCES

- AIDA, T. (1897): Chaetognaths of Misaki Harbor. Annot. Zool. Japon., Vol. I, pp. 13-21.
 BÉRANECK, ED. (1895): Les chétognathes de la baie d'Amboine. Revue Suisse de Zool., 3, pp. 137-159.
 BIERI, R. (1957): The chaetognath fauna off Peru in 1941. Pacif. Sci., Vol. XI, pp. 255-264.
 ——— (1959): The distribution of the plankton Chaetognatha in the Pacific and their relationship to the water masses. Limnol. Oceanogr., Vol. IV, No. 1, pp. 1-28.
 BURFIELD, S.T. and HARVEY, E.J.W. (1926): The Chaetognatha of the "Sealark" Expedition, Transact. Linn. Soc. London, Vol. XIX, Part 1, pp. 93-119, pls. 4-7.
 BURFIELD, S.T. (1950): Chaetognatha. Great Barrier Reef Exped., Sci. Rep., Vol. V, No. 8, pp. 459-473.

- DONCASTER, L. (1903): Chaetognatha, with a note on the variation and distribution of the group. The Fauna and Geography of the Maldive and Laccadive Archipelagoes, Vol. I, pp. 209-218.
- FAGETTI GUAITA, ELDA (1958a): Investigaciones sobre quetognatos colectados, especialmente, frente a la costa central y norte de Chile. Revista Biol. Mar. Valparaiso, Vol. VIII, Nos. 1-3, pp. 25-82.
- (1958b): Quetognato nuevo procedente del Archipelago de Juan Fernandez. Ibid., pp. 125-131.
- FAURE, M.L. (1952): Contribution a l'étude morphologique et biologique de deux chaetognathes des eaux atlantiques du Maroc: *Sagitta friderici* RITTER-ZÁHONY et *Sagitta bipunctata* QUOY et GAIMARD. Vie et Milieu, Tom. III, Fasc. 1, pp. 25-43.
- FOWLER, G.H. (1905): Biscayan plankton of H.M.S. Research. Pt. III Chaetognatha Transact. Linn. Soc. London, Ser. 2, Vol. X, pp. 55-87.
- (1906): The Chaetognatha of the Siboga Expedition. Siboga Exped. Monogr. 21, 86 pp., 3 pls., 6 charts.
- FURNESTIN, M.L. (1953a): Contribution à l'étude morphologique, biologique et systématique de *Sagitta serrato-dentata* KROHN des eaux atlantiques du Maroc. Bull. Inst. Océanogr. (Monaco), No. 1025, 39 pp.
- (1953b): Chaetognathes récoltés en Méditerranée par le "Président Théodore Tissier" aux Mois de Juin et Juillet 1950. Bull. d. Trav. publ. par Stat. Aquicult. & Pêche Castiglione (Alger), n.s. No. 4, 44 pp.
- FURUHASHI, K. (1953): On the vertical distribution of animal plankton in the Sea of Japan off San'in-District in summer of 1952. Publ. Seto Mar. Biol. Lab., Vol. III (1), pp. 61-74.
- GHIRARDELLI, E. (1948): Chaetognati raccolti nel Mar Rosso e nell'Oceano Indiano dalla nave "Cherso". Boll. Pesca, Piscic. Idrobiol. Vol. II (n.s.), Fasc. 2, pp. 3-20.
- (1951): Cicli di maturità sessuale nelle gonadi di *Sagitta inflata* GRASSI del Golfo di Napoli. Boll. Zool., Vol. XVIII, Nos. 4-6, pp. 149-162.
- GRAY, B.B. (1923): Notes on species of *Sagitta* collected during a voyage from England to Australia. Proc. Roy. Soc. Queensland, Vol. 34, pp. 171-180.
- HAMON, M. (1956): Chétognathes recueillis dans la Baie de Nha-Trang-Cauda (Viet-Nam). Bull. Mus. Hist. Nat., Paris, Tom. XXVIII, No. 5, pp. 466-473.
- HEYDORN, A.E.F. (1959): The Chaetognatha off the west coast of the Union of South Africa. Invest. Rep. Fish. Mar. Biol. Surv. S. Afr., No. 36, 56 pp., 2 pls.
- HIDA, T.S. (1957): Chaetognaths and pteropods as biological indicators in the North Pacific. Spec. Sci. Rep. U.S. Fish & Wildlife Serv., Fish. No. 215, 13 pp.
- JOHN, C.C. (1933): *Sagitta* of the Madras coast. Bull. Madras Govt. Mus., (Nat. Hist.) N.S., Vol. III, pp. 1-10.
- (1937): Seasonal variations in the distribution of *Sagitta* of the Madras coast. Rec. Indian Mus., Vol. 39, pp. 83-97.
- JOHNSTON, T.H. and TAYLOR, B.B. (1919): Notes on the Australian chaetognaths. Proc. Roy. Soc. Queensland, Vol. 31, pp. 28-41.
- KING, J.E. and HIDA, T.S. (1957): Zooplankton abundance in the Central Pacific. Fish. Bull. U.S. No. 118, pp. 365-395.
- LEA, H.E. (1955): The chaetognaths of western Canadian coast waters. J. Fish. Res. Bd. Canada, Vol. 12, No. 4, pp. 593-617.
- LELE, S.H. and GAE, P.B. (1936): Common sagittae of Bombay harbour. J. Univ. Bombay, Vol. 4, pp. 105-113.
- MARUMO, R. (1955): Results of the oceanographic observations made in the Eastern waters of Honsyû in Feb.-Mar. 1955. Ocean. Rep. Jap. Met. Ag., Vol. 4 (2), pp. 4-6. (in Japanese)
- MARUMO, R., KITOU, M. and OHWADA, M. (1958): Vertical distribution of plankton at 40°N, 150°E in the Oyashio water. Oceanogr Mag., Vol. 10, No. 2, pp. 179-184.

- MASSUTI, M. (1958): Estudio del crecimiento relativo de *Sagitta enflata* GRASSI del plancton de Castellon. Invest. Pesq., Barcelona, Tom. XIII, pp. 37-48.
- Meteorological Agency of Japan (1956-58): Results of the oceanographic observations made in the Eastern and Southern waters of Honsyû in the years 1956-57. Ocean. Rep. Jap. Met. Ag., Vol. 5, Nos. 2-4; Vol. 6, No. 4; Vol. 7, No. 1. (in Japanese)
- MICHAEL, E.L. (1911): Classification and vertical distribution of the Chaetognatha of the San Diego Region. Univ. Calif. Publ. Zool., Vol. VIII, pp. 21-186.
- (1919): Report on the Chaetognatha collected by the U.S. Fisheries Steamer "Albatross" during the Philippine Expedition, 1907-1910. U.S. Nat. Mus. Bull. 100, Vol. 1, Part 4, pp. 235-277.
- MURAKAMI, A. (1957): The occurrence of planktonic chaetognaths in the bay and Inland Sea regions (I). On the occurring conditions in Tôkyô Bay, and the central and western parts of Seto-Inland Sea. Suisangaku-Syûsei, Tôkyô Univ., pp. 357-384. (in Japanese)
- (1959): Marine biological study on the planktonic chaetognaths in the Seto Inland Sea. Bull. Naikai Reg. Fish. Lab. No. 12, pp. 1-186. (in Japanese)
- OYE, P. (1918): Untersuchungen über die Chaetognathen des Javameeres. Contr. Faune des Indes Néerlandais, Fasc. IV, 61 pp.
- PIERCE, E.L. (1951): The Chaetognatha of the west coast of Florida. Biol. Bull., Vol. 100, No. 3, pp. 206-228.
- (1953): The Chaetognatha over the continental shelf of North Carolina with attention to their relation to the hydrography of the area. Sears Found. J. Mar. Res., Vol. XII, No. 1, pp. 75-92.
- RITTER-ZÁHONY, R. (1909a): Expedition S.M. Schiff "Pola" in der Rote Meer nördliche und südliche Hälfte 1895/96-1897/98. XXII Zoologische Ergebnisse, Chaetognathen. Denkschrift. Math. Naturwiss. Kl. Kaiser. Akad. Wiss. Wien, Bd. 84, pp. 1-12.
- (1909b): Die Chaetognathen der Gazelle-Expedition. Zool. Anz., Bd. 34, pp. 787-793.
- (1910a): Chaetognatha. Die Fauna Südwest Australiens, Vol. 3, pp. 123-126.
- (1910b): Westindische Chätognathen. Zool. Jahrb., Suppl. 11, pp. 133-143.
- (1911): Revision der Chaetognathen. Deutsche Südpolar Exped., Bd. 13(5), pp. 1-71.
- SATYANARAYANA RAO, T.S. (1958): Studies on Chaetognatha in the Indian Seas, Part II. The Chaetognatha of the Lawson's Bay, Waltair. Andhra Univ. Mem. in Oceanogr., Vol. II, pp. 137-146.
- SATYANARAYANA RAO, T.S. and GANAPATI, P.N. (1958): Studies on Chaetognatha in the Indian Seas. Part III. Systematics and distribution in the waters off Visakhapatnam. Ibid., pp. 147-163.
- SCHILP, H. (1941): Biological results of the Snellius Expedition IX. The Chaetognatha of the Snellius Expedition, Temminckia, Vol. VI, pp. 1-99.
- STEINHAUS, O. (1896): Die Verbreitung der Chätognathen im südatlantischen und indischen Ozean. Inaug. Diss. Kiel., pp. 1-49.
- SUÁREZ CAABRO, J.A. (1955): Quetognatos de los mares Cubanos. Mem. Soc. Cubana Hist. Nat., Vol. 22, No. 2, pp. 125-180.
- *SUBRAMANYAM, M.K. (1940): *Sagitta bedoti* BÉRANECK in Madras plankton. Curr. Sci. (Bangalore), 9, pp. 379-380.
- SUND, P.N. (1959a): The distribution of Chaetognatha in the Gulf of Alaska in 1954 and 1956. J. Fish. Res. Bd. Canada, Vol. 16, No. 3, pp. 351-361.
- (1959b): A key to the Chaetognatha of the tropical eastern Pacific Ocean. Pacif. Sci. Vol. XIII, No. 3, pp. 269-285.
- SUND, P.N. and RENNERT, J.A. (1959c): The Chaetognatha of the Eastropic Expedition, with notes as to their possible value as indicators of hydrographic conditions. Inter-American Tropical Tuna Commission Bulletin, Vol. 3, No. 9, pp. 396-436, 27 text-figs., 1 table.
- TCHINDONOVA, YU. G. (1955): Chaetognatha of the Kurile-Kamchatka Trench. Trudi Inst.

- Okeanol., 12, pp. 298-310. (in Russian)
- THIEL, M.E. (1938): Die Chaetognathen-Bevölkerung des Südatlantischen Ozeans, *Wiss. Ergebn. Deutsch. Atl. Exped. "Meteor"*, D.A.E. "Meteor" 1925/27, Bd. XIII, pp. 1-110.
- THOMSON, J.M. (1947): The Chaetognatha of south-eastern Australia. Council Sci. Ind. Res. Australia, Bull. No. 222, 43 pp.
- ТОКИОКА, Т. (1939a): Chaetognaths collected chiefly from the bays of Sagami and Suruga, with some notes on the shape and structure of the seminal vesicle. *Rec. Oceanogr. Wks. Japan*, Vol. X, No. 2, pp. 123-150.
- (1939b): Chaetognaths and pelagic tunicates in Ōsaka Bay. *Sea and Sky*, Vol. 19, No. 6, pp. 152-160. (in Japanese)
- (1940a): The Chaetognath fauna of the waters of western Japan. *Rec. Oceanogr. Wks. Japan*, Vol. XII, No. 1, pp. 1-22.
- (1940b): A small collection of chaetognaths from the coast of New South Wales. *Rec. Aust. Mus.*, Vol. 20, No. 6, pp. 367-379.
- (1942): Systematic studies of the plankton organisms occurring in Iwayama Bay, Palao. III Chaetognaths from the Bay and adjacent waters. *Palao Tropical Biol. St. Studies*, Vol. II, No. 3, pp. 527-548.
- (1951): Pelagic tunicates and chaetognaths collected during the cruises to the New Yamato Bank in the Sea of Japan. *Publ. Seto Mar. Biol. Lab.*, Vol. II (1), pp. 1-25.
- (1954a): Droplets from the plankton net XIII. Emergence of a dense patch of chaetognaths near the Laboratory. *Ibid.*, Vol. III (3), pp. 359-360.
- (1954b): Droplets from the plankton net XVI. On a small collection of chaetognaths from the central Pacific. *Ibid.*, Vol. IV (1), pp. 99-102.
- (1955a): On some plankton animals collected by the Syunkotu-Marû in May-June 1954. I Chaetognatha. *Ibid.*, Vol. IV (2-3), pp. 219-225.
- (1955b): Droplets from the plankton net. XVII A small collection of chaetognaths and pelagic tunicates from the north eastern part of the Indian Ocean. *Ibid.*, Vol. V (1), pp. 75-78.
- (1955c): Notes on some chaetognaths from the Gulf of Mexico. *Bull. Mar. Sci. Gulf Caribb.*, Vol. 5, No. 1, pp. 52-65.
- (1956a): On chaetognaths and appendicularians collected in the central part of the Indian Ocean. *Publ. Seto Mar. Biol. Lab.*, Vol. V (2), pp. 197-202.
- (1956b): On chaetognaths and appendicularians collected by Mr. Z. Sagara in the Arafura Sea in May-August 1955. *Ibid.*, Vol. V (2), pp. 203-208.
- (1957): Chaetognaths collected by the Sōyō-marû in the years 1934 and 1937-39. *Ibid.*, Vol. VI (2), pp. 137-146.
- UYENO, F. and FURUHASHI, K. (1955a): Report of the oceanographic observations made in the southern waters of Honsyū in 1954. *Ocean. Rep. Jap. Met. Ag.*, Vol. 4 (1), pp. 62-69. (in Japanese)
- (1955b): Report of the oceanographic observations made in the southern waters of Honsyū in March 1955. *Ibid.*, Vol. 4 (2), pp. 20-21. (in Japanese)

(* referred through S. RAO's paper)

APPENDIX

Station data 1. For the small-net samples of the Transpac Expedition 1953.

Station Number	Date	Time	Haul Distance in m	Line Angle
TP 1	VII 24	14.50	50	
TP 3	25		50	
TP 4	26		50	40°
TP 6	27	13.35-14.00	50	50°
TP 7	28	1.30	50	0°
TP 8	29	7.00	50	20°
TP 9	30	5.30	50	40°
TP 10	31	4.00	50	30°
TP 11	31	18.00	50	20°
TP 12	VIII 2		50	30°
TP 13	2		50	
TP 14	3		50	60°
TP 15	3		50	
TP 16	4	15.00		
TP 17	5		50	
TP 18	5	23.30	50	60°
TP 52	IX 8	4.00	50	30°
TP 56	11	6.15	50	30°
TP 57A	11	17.00	50	
TP 58A	12	9.00	50	45°
TP 59A	12	18.30	50	45°
TP 67	18	4.00		40°
TP 68	18	14.30	50	
TP 69	19	4.00		60°
TP 70	19	6.45	50	
TP 71	20	11.30	50	
TP 72	21	3.45	50	
TP 74	22	3.00	50	
TP 76	28	21.20	50	
TP 78	30	5.00	50	
TP 79	30	16.30	50	
TP 80	X 1	4.00	50	
TP 80	3000 m wire out. suspended below ca. 1000 fathoms, mid water trawl.			
TP 81	1			
TP 82	2	13.00	50	
TP 83	3	8.00	50	
TP 84	3	16.30	50	
TP 85	4	19.00	50	
TP 89	15	19.30	50	
TP 92A	18	12.45	50	
TP 93	24	3.45		25°
TP 94	24	10.15	50	20°
TP 95	24	10.57-11.21	1	
TP 96	25	8.00	50	
TP 97	25	16.05	50	30°
TP 98	26	9.00	50	45°

TP 99	X 26	22.15	50	
TP 99	27	10.06-12.10	Surface	
TP 100	29	0.45	50	60°
TP 101	29	12.30	50	
TP 102	30	6.30	50	50°
TP 103	30	19.30	50	0°
TP 104	31	9.45	50	30°
TP 105	31	19.45	50	45°
TP 106	XI 1	14.45	50	45°
TP 107	2	1.20		40°
TP 108	2	12.30	50	45°
TP 109	3	1.00		15°
TP 111	3	22.15	50	30°
TP 112	4	10.30	50	0°
TP 113	5	3.30		35°
TP 114	5	19.30	50	45°
TP 114	5	19.47-20.48	Surface	
TP 115	6	6.15		75°
TP 116	7	5.45	50	30°
TP 117	7		50	
TP 118	8	10.30	50	60°
TP 119	8		50	
TP 120	9	9.30	50	
TP 121	10	0.15	50	15°
TP 122	10	13.30	50	30°
TP 123	11	0.30		15°
TP 124	11	14.00	50	45°
TP 125	12	4.00		45°
Midway, off NE Reef inner side	12	10.00	Surface	
TP 126	13	2.30	50	45°
TP 127	13	13.00	50	45°
TP 128	14	2.15	50	45°
TP 129	14	15.00	50	45°
TP 130	15	8.00	50	60°
TP 131	15	23.00		55°
TP 132	16	13.30	50	60°
TP 133	17	4.00	50	
TP 134	17	17.30	50	30°
TP 135	18	8.00	50	50°
TP 137	22	22.08-22.31	Surface	
TP 141	26	23.03-22.30	Surface	

Station data 2. For the plankton samples collected by a Japanese survey ship and presented to the Transpac Expedition.

Station Number	Position	Date	Haul Distance	Line Angle
J 1	40°50'N × 142°05'E	Sept. 28, 1953	50 m	60°
J 2	40°52'N × 142°57'E	28	50	16°
J 3	40°53'N × 143°53'E	28	50	16°
J 4	40°49.5'N × 145°13'E	28	50	36°
J 5	40°07'N × 145°09'E	29	50	46°

Station Number	Position	Date	Haul Distance	Line Angle
J 6	39°32'N × 145°05'E	Sept. 29	50 m	43°
J 7	39°37'N × 143°58'E	29	50	7°
J 8	39°31'N × 142°53'E	29	50	3°
J 9	39°28'N × 142°20'E	30	50	36°
J 10	38°52'N × 142°12'E	30	50	0°
J 11	38°10'N × 141°47'E	30	50	8°
J 12	38°12'N × 142°36'E	Oct. 3	50	20°
J 13	38°14.5'N × 144°03.5'E	4	50	12°
J 14	38°16'N × 145°00'E	4	50	14°
J 15	37°30'N × 145°05'E	4	50	10°
J 16	36°51'N × 145°06'E	4	50	14°
J 17	36°38'N × 143°37'E	4	50	3°
J 18	36°38'N × 142°39'E	5	55	34°
J 19	36°20'N × 141°12'E	5	55	52°

Station data 3. For the sampling stations during the Palao-New Guinea Cruises (March-May 1940).

Station Number	Position	Date
2	6°19'N × 134°27.7'E	Mar. 17, 1940
3	5°40'N × 134°26.4'E	18
4	5°01'N × 134°25'E	18
5	4°30'N × 134°31'E	18
6	3°58'N × 134°38'E	18
7	3°27'N × 134°37'E	19
8	2°58'N × 134°39'E	19
9	2°29'N × 134°35'E	19
10	2°00'N × 134°30'E	19
11	1°29'N × 134°21'E	20
12	0°58'N × 134°10'E	20
13	0°30'N × 134°16'E	20
15	0.33'S × 134°27'E	21
52	6°57.4'N × 134°27.5'E	May 20, 1940
53	6°30'N × 134°30'E	20
54	6°56.8'N × 134°33.1'E	21
55	5°23.8'N × 134°36.1'E	21
56	4°50.5'N × 134°29.1'E	21
57	4°30'N × 134°32.5'E	21
58	4°03'N × 134°33.8'E	21
59	3°35.5'N × 134°35.7'E	22
60	3°07.8'N × 134°37.4'E	22
61	2°30'N × 134°32.1'E	22
62	2°00.5'N × 134°30.2'E	22
63	1°31.1' × 134°28.4'E	23
64	1°02'N × 134°26.5'E	23
65	0°30.4'N × 134°24.1'E	23
66	0°02.5'N × 134°22'E	23
67	0°33'S × 134°21'E	24

1

(No. 1)

Stations	TP 1	TP 3	TP 4	TP 6	TP 7	TP 8	TP 9	TP 10	TP 11	TP 12	TP 13	TP 14	TP 15	TP 52	TP 59	TP 69	TP 70	TP 74	TP 76	TP 78	
<i>Sag. hexaptera</i>																					
<i>Sag. lyra</i>	1										1										
<i>Sag. enflata</i>																	2	18	23	9	
<i>Sag. bipunctata</i>				1																	
<i>Sag. ferox</i>																					
<i>Sag. robusta</i>																					
<i>Sag. bedoti</i>															6	12	3	6	60	58	
<i>Sag. friderici</i>	12																				
<i>Sag. pacifica</i>	6	1	3	7	1	8								1	1				3		
<i>Sag. tasmanica</i>	1																				
<i>Sag. pseudoserratodentata</i>							4														
<i>Sag. neglecta</i>																				1	2
<i>Sag. regularis</i>																	2	3	4	5	
<i>Sag. crassa f. naikaiensis</i>																3		1	2		
<i>Sag. minima</i>	12	8				1	6	4	6	8	61	166	3		3	7	2	44	21	35	
<i>Sag. decipiens</i>																					
<i>Pterosag. draco</i>																					
<i>Ktta. subtilis</i>																					
<i>Ktta. pacifica</i>																				1	
Damaged specimens or juv.	45	46	* 1	* 5		* 2	* 7	1		3	* 13	* 15	1	2	12	* 13	* 7	* 95	* 92	* 72	
total number	77	55	4	13	1	11	17	5	6	11	75	181	4	3	22	35	16	167	207	181	
number of species	5	2	1	2	1	2	2	1	1	1	2	1	1	1	3	3	4	5	8	5	

* juv.

(No. 2)

Stations	TP 79	TP 80	TP 80	TP 81	TP 82	TP 83	TP 84	TP 85	TP 89	TP 92	TP 93	TP 94	TP 95	TP 96	TP 97	TP 98	TP 99	TP 99	Entrance to Yokosuka	TP 100	
<i>Sag. hexaptera</i>			1		1															1	
<i>Sag. lyra</i>																					
<i>Sag. enflata</i>	8	1	4	6	8	10		1	8	24	58	7	24		3	3	2	52		4	
<i>Sag. bipunctata</i>				1	1		3				1		3	1	1			9			
<i>Sag. ferox</i>							1														
<i>Sag. robusta</i>	1			1	1	4			1				7		1	1		28			
<i>Sag. bedoti</i>											1		2					2			
<i>Sag. friderici</i>																					
<i>Sag. pacifica</i>	14	10	17	19	14	14	5	5	4		4	4	27	3	1	7	3	41		8	
<i>Sag. tasmanica</i>																					
<i>Sag. pseudoserratodentata</i>																					
<i>Sag. neglecta</i>	2									1				1							
<i>Sag. regularis</i>	8	2		5	8	12	2	1	1			1	15		3	4	1	7		4	
<i>Sag. crassa f. naikaiensis</i>									1										80		
<i>Sag. minima</i>	1				1				1		1									4	
<i>Sag. decipiens</i>																					
<i>Pterosag. draco</i>	2		2		1				2	1	1		4			1		15		4	
<i>Ktta. subtilis</i>			3																		
<i>Ktta. pacifica</i>	2	2		3	2	1			1	1			3	1	2	3	1	1		1	
Damaged specimens or juv.	16	3	11	7	13	10	7	6	* 27	* 68	* 92	18	21	11	12	1	10	20		8	
total number	54	18	38	42	50	51	18	13	46	95	158	30	106	17	23	20	17	175	80	34	
number of species	8	4	5	6	9	5	4	3	8	4	6	3	8	4	6	6	4	8	1	7	

Appendix Table 1. Occurrence of chaetognaths in samples of the Transpac Expedition. (Nos. 1-2)

Stations	TP 101	TP 102	TP 103	TP 104	TP 105	TP 106	TP 107	TP 108	TP 109	TP 111	TP 112	TP 113	TP 114	TP 115	TP 116	TP 117	TP 118	TP 119	TP 120	TP 121
<i>Sag. hexaptera</i>																		1		
<i>Sag. lyra</i>																				
<i>Sag. enflata</i>	4	4		4	2	2		1	5	1	1		1				1		1	
<i>Sag. bipunctata</i>	3	2	1	3														1		
<i>Sag. ferox</i>		1																		
<i>Sag. robusta</i>						1		1												
<i>Sag. bedoti</i>																				
<i>Sag. friderici</i>																				
<i>Sag. pacifica</i>	9	4	1	7	5	7	1	10	11	2			8	6	2	1	11	6	1	1
<i>Sag. tasmanica</i>																				
<i>Sag. pseudoserratodentata</i>																				
<i>Sag. neglecta</i>																				
<i>Sag. regularis</i>	1					2		1	3											
<i>Sag. crassa f. naikaiensis</i>																				
<i>Sag. minima</i>		1			2			1	6	3	14	5			3		1			
<i>Sag. decipiens</i>																				
<i>Pterosag. draco</i>	2	2				1		2	4	1						1	2			2
<i>Ktta. subtilis</i>					1															
<i>Ktta. pacifica</i>	2			2		2			4	1										
Damaged specimens or juv.	6	5	2	4	4	6		* 37	* 48	5	8	3			1	2	7	5	3	2
total number	27	19	4	20	14	21	1	53	81	13	23	8	9	6	6	4	22	13	5	5
number of species	6	6	2	4	4	6	1	6	6	5	2	1	2	1	2	2	4	3	2	2

* juv.

Stations	TP 122	TP 123	TP 124	TP 125	TP 126	TP 127	TP 128	TP 129	TP 130	TP 131	*TP 131	TP 132	TP 133	TP 134	TP 135	TP 137	TP 141	*St. 89	F. O.	Mean percentage
<i>Sag. hexaptera</i>				1							232				2		4		10	8
<i>Sag. lyra</i>											30								4	1
<i>Sag. enflata</i>											117			1	1	1			46	14
<i>Sag. bipunctata</i>	2	2									24								23	13
<i>Sag. ferox</i>																			3	6
<i>Sag. robusta</i>											6								15	5
<i>Sag. bedoti</i>																			12	17
<i>Sag. friderici</i>																			1	16
<i>Sag. pacifica</i>			6	2	4		2	2	5	1	401	1		4	15	21	4		74	37
<i>Sag. tasmanica</i>																			1	1
<i>Sag. pseudoserratodentata</i>											22					1	1		5	9
<i>Sag. neglecta</i>																			6	2
<i>Sag. regularis</i>				1															31	9
<i>Sag. crassa f. naikaiensis</i>																			6	* 3
<i>Sag. minima</i>											16								40	30
<i>Sag. decipiens</i>											4								1	*
<i>Pterosag. draco</i>		2	1		3		1			2	329	2	1	3	3	1	8		40	15
<i>Ktta. subtilis</i>											97								4	7
<i>Ktta. pacifica</i>											30		1	1	1				31	7
Damaged specimens or juv.	1		6	5	1	2	2	2	2	2	76	8	1	2	9	1	1		90	38
total number	3	4	13	9	8	2	5	4	7	5	1384	11	3	7	20	19	35	4		
number of species	1	2	2	3	2	—	2	1	1	2	12	2	2	3	5	4	4	1		

* 3 Yokosuka excluded.

* TP 131 large net hauled from 450m. * St. 89 Stranger: North Pacific 5509-2nd, 23.00.

Appendix Table 1 (continued). Occurrence of chaetognaths in samples of the Transpac Expedition. (Nos. 3-4)

(No. 1)

Stations	TP 19	TP 20	TP 21	TP 22	TP 23	TP 24	TP 25	TP 26	TP 27	TP 28	TP 29	TP 30	TP 31	TP 32	TP 33	TP 34	TP 35	TP 36	TP 37	TP 38	TP 39
Situation	48-14.2 N	49-29.5 N	50-25.0 N	51-21.3 N	52-10.4 N	53-15.0 N	53-32.5 N	54-00.0 N	54-00.0 N	53-59.5 N	54-00.0 N	53-57.7 N	53-15.6 N	52-29.3 N	52-19.9 N	53-10.3 N	54-05.5 N	54-00.9 N	53-59.0 N	53-59.1 N	54-03.1 N
	153-20.1 W	154-56.5 W	156-37.0 W	158-20.0 W	160-0.80 W	161-55.0 W	163-20.8 W	168-19.8 W	170-03.0 W	171-40.0 W	173-15.5 W	175-04.5 W	175-37.6 W	176-09.0 W	176-57.5 W	177-59.0 W	178-56.2 W	178-58.7 E	176-55.0 E	174-49.0 E	172-28.8 E
Date	Aug. 6	Aug. 7	Aug. 8	Aug. 8		Aug. 9	Aug. 10	Aug. 12	Aug. 13				Aug. 15	Aug. 15	Aug. 24	Aug. 25	Aug. 25	Aug. 26	Aug. 27	Aug. 28	Aug. 29
Time		9.00		12.00				11.30	10.30		6.45				20.00	10.00	18.30	10.30		13.30	3.15
Haul distance (m.)	0-50	0-50		0-50	*	0-50		0-50		0-50	0-50		0-50	0-50	0-50	0-50	0-50	0-50		0-50	0-50
Rope angle				45°			50°		40°				20°	0°	45°	60°	60°	60°	70°	60°	60°
<i>Sag. elegans</i>	2	6	5	10	23	9	2	6	11	4	8	2	9	2	3	22	21	82	26	17	15
<i>Eukr. hamata</i>		1	1																		
Damaged specimens or juv.																					
total number	2	7	6	10	23	9	2	6	11	4	8	2	9	2	3	22	21	82	26	17	15

* #20 Net used.

(No. 2)

Stations	TP 40	TP 41	TP 42	TP 43	TP 44	TP 45	TP 46	TP 47	TP 48	TP 49	TP 50	TP 51	TP 53	TP 55	TP 61	TP 63	TP 64	TP 65	TP 66	TP 73	
Situation	53-57.0 N	53-32.1 N	53-32.4 N	53-35.5 N	53-40.8 N	52-23.0 N	51-13.0 N	49-50.1 N	48-46.3 N	47-35.7 N	46-16.5 N	45-05.5 N	42-07.9 N	41-19.0 N	45-15.1 N	46-23.5 N	46-54.5 N	45-29.9 N	44-09.0 N	40-53.4 N	
	171-10.8 E	168-50.8 E	166-10.8 E	163-42.0 E	161-55.5 E	163-15.2 E	164-34.3 E	165-49.2 E	166-48.1 E	167-44.8 E	168-52.2 E	170-08.5 E	169-28.1 E	168-20.2 E	158-20.0 E	155-28.6 E	153-55.2 E	154-19.7 E	152-56.8 E	143-51.5 E	
Date	Aug. 30		Sep. 1	Sep. 1	Sep. 2	Sep. 3	Sep. 3	Sep. 4	Sep. 5	Sep. 6	Sep. 6	Sep. 7	Sep. 8	Sep. 10	Sep. 14	Sep. 15	Sep. 15	Sep. 16	Sep. 17	Sep. 21	
Time	12.45		6.00	23.00	5.00	8.15	19.00	17.30	11.00		14.45	11.15	15.25	6.45	9.00	5.45	20.45	10.30	9.00	4.30	
Haul distance (m.)	0-50		0-50		0-50		0-50	0-50	0-50		0-50	0-50	0-50	0-50	0-50	0-50	0-50	0-50	0-50	0-50	
Rope angle	45°	15°		40°		40°			40°		30°	30°		45°	60°	45°	10°			F.O.	
<i>Sag. elegans</i>	6	2	6	1	7	1	1	1	2	10	3		5		2	6	4	3	* 4	2	95
<i>Eukr. hamata</i>																					5
Damaged specimens or juv.						5		1			1	1	1	1							1
total number	6	2	6	1	7	6	1	2	2	10	4	1	6	1	2	6	4	3	4	3	

* One of the four is *Sag. minima*?Appendix Table 2. Chaetognaths occurred in the *elegans*-water during the Transpac Expedition. (Nos. 1-2)

	West of 170°E				West of 150°E				
TP 18	TP 56	TP 67	TP 68	TP 71	J 3	J 4	J 10	F. O.	p
1								30	
	1						1	20	
3	3	9	9	6	5	1	1	100	
				3	1		2	30	
				2			1	20	
		3	2	3		1	2	60	
4		1		14			1	60	
8	4	13	11	28	6	2	8		
2	2	2	2	4	2	2	5		

tognaths occurred in samples from the northern mixing area of the *elegans*-water and the warm water.

J 6	J 7	J 8	J 9	J 11	J 12	J 13	J 14	J 15	J 16	J 17	J 18	J 19	F. O.
3	6			1	2	2	2	2			3	12	69
							1						13
											1	2	13
1	18	1	4	9	1							17	63
1	1					1	2	3		2		1	63
	1				2								13
1	1				2	3	2	3	2		1	1	69
				1									13
15	2		6	3	9	3	1					9	69
16	84	2	6	26	28	13	15		1	3	2	30	88
37	113	3	16	40	44	22	23	8	3	5	7	72	
5	6	1	2	4	5	4	5	3	1	1	3	6	

Table 4. Chaetognaths occurred in samples from the North-Eastern waters of Japan.

Stations	MP 1			MP 2		MP 3								MP 5			MP 7		MP 8				MP 10								
	J 1	J 2	J 3	J 4	J 6	J 7	J 8	J 9	J 10	J 11	J 12	J 12	J 14	J 15	J 16	J 18	J 21	J 23	J 25	J 26	J 28	J 30	J 30	J 32	J 33	J 34	J 35	J 37	J 38	J 39	
<i>Sag. hexaptera</i>			3		2					1	2	5			1	2	47	4	23			1	37	2	62	12		7	5		
<i>Sag. enflata</i>	1	3		4	4	1		6	2	47	17	36	61	5		10	12	103	25	218		15	16	18	1	41	25	10	13	13	
<i>Sag. bipunctata</i>												1							1										1		
<i>Sag. ferox</i>																															
<i>Sag. robusta</i>												1				* 1							3					23	7	17	
<i>Sag. pulchra</i>																															
<i>Sag. pacifica</i>	* 106	* 134	71	5	5			27	45	2	3	34	51	11			1	5	2	74			5	65	1	30	101	54	30	31	
<i>Sag. pseudoserratodentata</i>	23	29	6									1	1	2															1		
<i>Sag. neglecta</i>																															
<i>Sag. regularis</i>								1																			1				
<i>Sag. minima</i>	60	95	7	3															1										1		
<i>Sag. planctonis</i>																															
<i>Sag. macrocephala</i>																															
<i>Pterosag. draco</i>			3					3		23	9	8	12	4		1		13	1	12				283	14	283	170	31	24	55	
<i>Ktta. subtilis</i>	2	1		2	4					1	1		12							1			4		2			1			
<i>Ktta. pacifica</i>																									1					1	
<i>Eukr. hamata</i>																				1										1	
Damaged specimens or juv.	16	16	31	3	17			7	7	1	3	20	34	3		25	22	16	11	60	3	3	8	10	6	* 47	10	12	46	6	
total number	208	278	121	17	32	1	0	44	54	75	33	102	177	25	0	37	38	184	44	390	3	18	30	420	24	466	319	130	131	129	
number of species	5	5	5	4	4	1	0	4	2	5	4	6	7	4	0	3	4	4	5	7	—	1	3	6	4	6	5	4	9	7	

* *pseudoserratodentata* juv. included?* or *ferox* f. *americana* juv.?

* juv.

Stations	MP 12					MP 14	MP 15	MP 16		MP 17		MP 18				MP 20				MP 35	MP 36		MP 41		F.O.	Mean percentage			
	J 40	J 41	J 42	J 43	J 44	J 46	J 47	J 48	J 50	J 51	J 52	J 55	J 56	J 57	J 57	J 62	J 64	J 64	J 65	J 66	J 67	J 69	J 70	J 71			J 73	J 74	
<i>Sag. hexaptera</i>	1	4			15	45	1	244	23			5	3	3	19	6	1	16	28	7	24	1	24	25	6	4	68	11	
<i>Sag. enflata</i>	22	12		3	49	384	19	30				1	2		4	2		14	29	7	21	4	20	26	20	12	81	24	
<i>Sag. bipunctata</i>	2							3										1		1		2		1	1	1	19	1	
<i>Sag. ferox</i>	2	1																									4	2	
<i>Sag. robusta</i>	16	2			15													2			2		2				21	3	
<i>Sag. pulchra</i>						1																					2	*	
<i>Sag. pacifica</i>	10	2		3	140	8	7	48	2	5		13	10	33	43	17	4	48	61	42	93	16	41	61	30	26	84	30	
<i>Sag. pseudoserratodentata</i>					1			1							7	4		1	4		3		1			1	28	4	
<i>Sag. neglecta</i>				1																				2			4	6	
<i>Sag. regularis</i>		1															1			2			2		1	2	14	1	
<i>Sag. minima</i>								1							1												14	11	
<i>Sag. planctonis</i>					1																						2	*	
<i>Sag. macrocephala</i>							1																				2	3	
<i>Pterosag. draco</i>	29	25	2	1	154	52		104	2	1		1	2		11	4		17	35	22	19	4	12	15	18	6	70	21	
<i>Ktta. subtilis</i>					2	1			2					1	2					2	1		2	2	2			35	3
<i>Ktta. pacifica</i>	1				1																2	2					11	1	
<i>Eukr. hamata</i>																											4	2	
Damaged specimens or juv.	52	15		3	36	125	2	39	4	1		1	5	35	34	12	3	64	11	52	37	15	34	49	38	25	89	23	
total number	135	62	2	11	414	616	30	470	33	7	0	21	22	72	120	46	8	164	168	135	202	44	138	181	114	77			
number of species	8	7	1	4	9	6	4	7	4	2	0	4	4	3	6	6	2	8	5	7	8	7	8	7	6	7			

Appendix Table 5. Occurrence of chaetognaths in samples of the Midpac Expedition. (Nos. 1-2)

Stations	SB 1	SB 5	SB 10	SB 15	SB 20	SB 25	SB 30	SB 35	SB 40	SB 44	SB 50	SB 55	SB 60	SB 64	SB 68	SB 71	SB 75	SB 80	SB 85	SB 90	SB 95	SB 100	SB 105	SB 109	SB 112
<i>Sag. hexaptera</i>	4	18	13	11	63	25	16	65	209	140	30	141	109	21	18	31	25	20	149	56	168	44	101	70	65
<i>Sag. lyra</i>	41	2			14	5	14	14	171	29	4	11						9	4	33	11	49	13	7	15
<i>Sag. enflata</i>	33	12	105	460	694	616	182	285	160	391	54	408	204	109	179	43	69	526	408	773	597	927	723	188	721
<i>Sag. bipunctata</i>	13		28		14	4						1	3	2		2	5								
<i>Sag. ferox</i> f. <i>americana</i>				1	4							3	70	65	69	58	69								
<i>Sag. robusta</i>				1	7	3				1		11	16	2		2	1			4	4	75	13		
<i>Sag. bedoti</i>					1			5		6		20	38	4	137	1	12	32		1	1	15	7	101	
<i>Sag. pulchra</i>								1			1	2	1		2			27	6		1	1	1		
<i>Sag. friderici</i>	101	23									1								3	7	1	1	8	36	3
<i>Sag. pacifica</i>	50	273	29	76	64	38	111	150	377	222	324	401	144	92	157	148	115	211	170	126	296	287	383	45	178
<i>Sag. tasmanica</i>	247	17	15	18	27	28	88	264	5	108	131	41		2	7	2	4	35	71	71	13	26	140	169	39
<i>Sag. pseudoserratodentata</i>	11	49	3								3							1							
<i>Sag. neglecta</i>											1	6	38	3	18	3	2	3		1					5
<i>Sag. regularis</i>				1			18	9	14	32	8	22	5	2			2	52	45	38	26	149	29	1	1
<i>Sag. minima</i>	137	9	2	5	15	2	17	5	8	31	15	7	3			2	1	39	35	10	10	5	11	23	27
<i>Sag. decipiens</i>	39					2	4	12	10	1	25	7					2	4	17	15	3	6	9	1	1
<i>Sag. neodecipiens</i>	20										13												1		9
<i>Sag. planctonis</i>	2						2		1																2
<i>Pterosag. draco</i>		5	17	28	213	290	27	120	16	203	22	209	172	10	4	13	13	81	182	93	404	326	178	12	18
<i>Kitta. subtilis</i>	8	44	22	3		23	51	53	68	51	40	51	2	3	1		2	42	40	41	10	112	84	48	31
<i>Kitta. pacifica</i>			1	5			10	5		5	4	11	3	15	17	11	1	24	1	1	20	7	4		
<i>Eukr. hamata</i>						1	2	4	22	6	2									7	4		3	3	
Damaged specimens or juv.	82	20	13	23	16	12	10	44	73	23	21	57	46	29	19	6	40	217	91	123	226	236	251	162	51
total number	788	472	248	632	1132	1049	552	1036	1134	1249	699	1409	854	359	628	322	363	1323	1222	1400	1795	2266	1959	871	1161
number of species	13	10	10	11	11	12	13	14	12	14	17	17	14	13	11	12	15	15	13	16	16	15	17	14	13

Appendix Table 6. Occurrence of chaetognaths in samples of the Shellback Expedition. (No. 1)

Stations	S B 115	S B 118	S B 122	S B 125	S B 130	S B 132	S B 137	S B 142	S B 145	S B 150	S B 155	S B 160	S B 166	S B 170	S B 175	S B 180	S B 181	S B 187	S B 195	S B 200	S B 210	S B 215	S B 217	F. O.	Mean percentage		
<i>Sag. hexaptera</i>	60	18	85	29	12	15	3	16	39	40	75	69	55	168	71	59	40	52	46	25	4		2	98	6		
<i>Sag. lyra</i>	5	7	2	9				1	10	16	28	11		2	26	62	5	2		1		1	4	73	2		
<i>Sag. enflata</i>	267	52	398	42	155	31	42	226	567	374	527	350	416	895	333	366	315	811	83	1118	116	429	433	100	33		
<i>Sag. bipunctata</i>												1			3					10					25	1	
<i>Sag. ferox f. americana</i>												16	1		19	1		4		13	101				31	6	
<i>Sag. robusta</i>								2	3	8	8	4			8	5	4		2	5					48	1	
<i>Sag. bedoti</i>												30	11	215	299	27	41	7	131	12	105		1	2	56	4	
<i>Sag. pulchra</i>												42	11	201	136	16	38	6	51		48					39	2
<i>Sag. friderici</i>		1	1																						25	2	
<i>Sag. pacifica</i>	635	117	34	80	156	70	92	116	382	308	400	403	106	186	235	324	234	318	31	238	156	78	171	100	22		
<i>Sag. tasmanica</i>	97	25	43	17	18	26	202	55	86	85	87	53	1	8	121	173	83	278	55	8	72	113	86	98	8		
<i>Sag. pseudoserratodentata</i>																				2		5	3		17	2	
<i>Sag. neglecta</i>											1	19	5	29	4		1	7	2	8	1				42	1	
<i>Sag. regularis</i>	3	1							2	30	4	15			10	7	59	3	8	17	2		1		67	2	
<i>Sag. minima</i>	7	1	6	3	4	3	11	8	4	1	32	1	5	12	19	14	36	3	10	1	6	53	183	96	2		
<i>Sag. decipiens</i>			15		3				5	2	20	10			10	14	5	6	6	10	1		15		65	1	
<i>Sag. neodecipiens</i>			3		1				1	1							1					1	3		23	1	
<i>Sag. planctonis</i>														1											10	*	
<i>Pterosag. draco</i>	77	9	1	6	11	1	43	218	219	59	208	308	21	34	283	140	94	98	71	483	16	1			96	9	
<i>Ktta. subtilis</i>	2	2	10	2	30		10	15	9	7	67	89	13	64	89	147	8	82	15	20		5	9		92	3	
<i>Ktta. pacifica</i>	1	1			2	5	60	186	85	9	63	30	76	26	71	10	13	26	2	12	63	7	4		81	2	
<i>Eukr. hamata</i>		1									16														25	*	
Damaged specimens or juv.	48	48	11	12	31	18	19	22	86	19	100	38	18	79	38	610	11	55	73	24	24	46	23	100	6		
total number	1202	283	609	200	423	169	482	865	1498	959	1708	1439	1133	1939	1383	2011	922	1927	416	2148	562	740	939				
number of species	10	12	11	8	10	7	8	10	13	13	16	17	12	13	17	15	16	15	13	18	11	11	13			* less than 0.5	

Appendix Table 6 (continued). Occurrence of chaetognaths in samples of the Shellback Expedition. (No. 2)

Stations	EQP S 28	EQP H 3	EQP H 7	EQP H 11	EQP H 17	F. O.	Mean percentage
<i>Sag. hexaptera</i>	109	29	41	62	57	100	13
<i>Sag. enflata</i>	688	56	180	431	348	100	40
<i>Sag. bipunctata</i>	12	6	1	1	32	100	2
<i>Sag. ferox</i>	1	2			6	60	1
<i>Sag. robusta</i>	5	3	5	9	9	100	1
<i>Sag. pulchra</i>				17	2	40	1
<i>Sag. pacifica</i>	203	60	66	231	35	100	17
<i>Sag. pseudoserratodentata</i>			3			20	1
<i>Sag. neglecta</i>	5					20	*
<i>Sag. oceania</i>	7					20	1
<i>Sag. regularis</i>	29	1	7	20	32	100	2
<i>Sag. minima</i>				2		20	*
<i>Sag. decipiens</i>	14	1	8	17	26	100	2
<i>Pterosag. draco</i>	73	21	81	204	315	100	17
<i>Kitta. subtilis</i>		1	5	1	2	80	1
<i>Kitta. pacifica</i>				12	15	40	2
Damaged specimens or juv.	99	9	113	63	29	100	9
total number	1245	189	510	1070	908		
number of species	11	10	10	12	12		

* less than 0.5

Appendix Table 7. Occurrence of chaetognaths in samples of the Equapac Expedition.

Stations	North Equatorial Current					Counter Equatorial Current											South Equatorial Current											F. O.	Mean percentage		
	2	3	4	52	53	5	6	7	8	54	55	56	57	58	59	60	9	10	11	12	13	15	61	62	63	64	65			66	67
<i>Sag. hexaptera</i>										3						1						1			2				14	5	
<i>Sag. enflata</i>	3		7	3	28	14	6	47	28	20	12	21	12	7	1	3	37	8	25	20	17	16	17	30	49	47	50	61	12	97	35
<i>Sag. ferox</i>				3	6					2			2								1				3			2	24	11	
<i>Sag. bedoti</i>						9	8	28	5	1	3	7	3		1		22	5	7	3		2	1	12		10	17	18	24	69	13
<i>Sag. pulchra</i>												5															4		7	7	
<i>Sag. pacifica</i>	1	1			2	6	1	4	32			5	9	12	12	5	25	12	5	6	11	21	10	10	6	5	7	20	8	86	20
<i>Sag. neglecta</i>					2	3			1	1							2		4	1	3	14	1		30	10	6	8	15	52	8
<i>Sag. regularis</i>	3	3	10		18	15	7	12	7	7	20	1	6		1		15	5	1	5	11		10	16	17	11	26	20	3	86	21
<i>Sag. bedfordii</i>																		1						3			1		10	2	
<i>Pterosag. draco</i>						5	1	5	7	1	4	11	2	1	3	6	8	5	9	8	4		10	9	3	8	14	11	6	79	11
<i>Ktta. subtilis</i>								1																					3	1	
<i>Ktta. pacifica</i>						7	2		6				1				1	1		2	1	8	1			2	1	6	6	48	5
Damaged specimens or juv.					18		1															29		13	1	2		1	24	11	
total number	7	4	17	6	74	59	26	97	86	35	39	50	35	20	18	15	110	37	51	45	48	91	50	93	111	95	121	140	76		
number of species	3	2	2	2	5	7	6	6	7	7	4	6	7	3	5	4	7	7	6	7	7	6	7	6	7	7	7	9	8		

Sts. 2-15 March, Sts. 52-67 May.

Appendix Table 8. Occurrence of chaetognaths in samples collected during the Palao—New Guinea cruises, 1940.