

Microfossils of the core sample GDP-11-15
from the Amami Plateau,
the northern margin of the Philippine Sea*

By

Akira NISHIMURA, Isao KONDA*¹, Kazumi MATSUOKA*²,
Shiro NISHIDA*³ and Terufumi OHNO

(Received Aug. 11, 1976)

Abstract

The micropaleontological studies of a gravity core taken from a depth of 1,830 m on the Amami Plateau, situated at the northern margin of the Philippine Sea, have revealed a presence of hiatus between the late Pliocene and the late Pleistocene sediments. Furthermore, the conspicuous difference in the water depths of the accumulation of sediments is recognized between the portions above and below the hiatus. Consequently, it is considered that the plateau submerged with the amount of about 1,000 m during the latest Pliocene and early Pleistocene.

Lastly two foraminiferal species are described: *Globocassidulina ikebei* and *Pleurostomella delicata*.

Introduction

Recently, the geological and paleontological informations of the Philippine Sea have rapidly increased in process of various research projects. Many results of micropaleontological investigations on sediment cores collected by the R/V Vema and the D/V Glomar Challenger in the northern Philippine Sea, have already been published (ELLIS, 1975; KOIZUMI, 1970, 1975a, b; TAKAYAMA, 1969, 1970; UJIE, 1975a, b, c; UJIE and MIURA, 1970, 1971; THE SHIPBOARD SCIENTIFIC PARTY, 1975; LING, 1975a, b).

Nearly the same time, the eleventh research cruise of Japanese Geodynamics Project (GDP-11 Cruise) was executed by the R/V Tokaidai-gaku-maru II in the northern Philippine Sea in August, 1974. During this cruise, continuous echo sounding, continuous seismic reflection profiling and frequent bottom sampling were carried

* Read at the 82nd Annual Meeting of the Geological Society of Japan.

*¹ Nara Senior High School.

*² Department of Biology, Faculty of Science, Osaka City University.

*³ Department of Earth Science, Nara University of Education.




Fig. 1. Map showing the station GDP-11-15, after MISAWA and HOSHIZAWA (1975).

out. The geological and paleontological results were reported at the Symposium "Geological Problems of the Philippine Sea Area" at the annual meeting of the Geological Society of Japan, Kyoto, April 3, 1975 and at other opportunities, and several synoptical reports have been presented (AOKI, 1975; AOKI *et al.*, 1975; KONDA, 1975; KONDA *et al.*, 1975; KONDA *et al.*, in press; MATSUDA *et al.*, 1975; RESEARCH MEMBERS OF GDP-11 CRUISE, 1975; SHIKI *et al.*, 1975; SHIKI *et al.*, 1975; SUWA and AOKI, 1975). Besides these papers, acoustic stratigraphy and submarine topography of the Amami Plateau and the Komahashi Seamount have been described in detail (MISAWA, 1975; MISAWA and HOSHIZAWA, 1975).

One of the most interesting samples obtained in this cruise is a sediment core taken by a gravity corer at the station GDP-11-15 (Latitude 28°06.2'N., Longitude 131°35.2'E., Water depth 1,830 m) on the westward flat of the major peak of the Amami Plateau (Fig. 1). The sample is composed of pale-creamy calcareous ooze and has a whole length of fifty centimeters. The biostratigraphical and paleoecological studies of this core suggest the abrupt submergence of the Amami Plateau.

The purposes of the present paper are 1) to establish the precise biostratigraphy of microfossils of this core, 2) to discuss the change of the environmental conditions inferred from the microfossils, and 3) to describe two new benthonic foraminiferal species.

Material and Method

After the sample was taken out from the inner tube of the core sampler, its lithology, color *etc.* were described immediately on the shipboard (Fig. 2). The whole of the core sample is exclusively composed of calcareous ooze. The ooze is gradually consolidated downward. The sharp change of the color is observed at about fifteen centimeters below the top (near the boundary between the sample GDP-11-15-(17) and (18)). According to the observation under a binocular microscope, the composition of the sample is mostly foraminiferal tests and fine matrix, and siliceous microfossils such as radiolaria are rarely found. Glassy fragments and mafic minerals are scattered in the upper part of the core.

Before the analysis of the microfossils, the sample was prepared by the following way.

i) The sample was cut longitudinally into two parts, and one part has been deposited in the Department of Geology and Mineralogy, Kyoto University, and the other one was used for microfossil analysis.

ii) The sample for microfossil analysis was further divided horizontally every one centimeter and these were numbered from the top to the bottom.

iii) Microfossil analysis was performed at stratigraphic interval of five centimeters, and at closer interval around the boundaries of microfossil assemblages.

iv) Each material was added with a small amount of distilled water and stirred up to be disintegrated.




Fig. 2. Sketch of the core sample GDP-11-15.

v) The top of the water was decanted out and was provided for the analysis of calcareous nannoplanktons. The residue was used for foraminiferal analysis.

Analysis of calcareous nannoplanktons; In order to obtain nannoplanktons, suspended water samples were centrifuged. For the observation of a scanning electronmicroscope (SEM), after mounted on a glass, the samples were coated with carbon and gold. Under the SEM with TV apparatus, more than 300 specimens of calcareous nannoplanktons were identified and counted at random. Nannoplanktons were analyzed for seventeen samples, GDP-11-15-(1), (6), (7), (8), (9), (10), (11), (16), (17), (18), (21), (26), (31), (36), (41), (46), and (50).

Foraminiferal analysis; The sample for foraminiferal analysis was mechanically split into moderate fractions. Analyses of planktonic and benthonic foraminifers were performed separately using different fractions. Planktonic and benthonic specimens were picked up from the sieved residue of a 115 and 200 mesh screen respectively. Actual number of specimens per grams was not calculated. Both planktonic and benthonic foraminifers were analyzed for thirteen samples, GDP-11-15-(1), (6), (11), (16), (17), (18), (21), (26), (31), (36), (41), (46), and (50).

Micropaleontology

1. Calcareous nannoplankton fossils

Well preserved calcareous nannoplanktons were abundantly contained throughout the present selected core samples. Identified taxa in the core sample are shown in Table 1 and Figure 3.

Based on the composition of these nannoplankton assemblages, the present nannoflora can be subdivided into three assemblages as follows in ascending order; the assemblage I is GDP-11-15-(50) to (18), the assemblage II GDP-11-15-(17) to (11), and the assemblage III GDP-11-15-(10) to (1).

The assemblage I (GDP-11-15-(50)~(18))

The assemblage I is composed of *Reticulofenestra reticulata*, *R. reticulata* var., *Pseudoemiliana lacunosa*, *Cyclococcolithina macintyreii*, *C. leptopora*, *Discoaster pentaradiatus*, *D. surculus*, *D. brouweri* (5 and 6 ray forms), and *D. variabilis*. This nannoplankton assemblage is characterized by the abundant occurrence of *Reticulofenestra reticulata* and *R. reticulata* var., and the presence of *Discoaster pentaradiatus*, *D. surculus* and *Pseudoemiliana lacunosa*.

The assemblage II (GDP-11-15-(17)~(11))

This assemblage consists of *Gephyrocapsa oceanica*, *G. caribbeanica*, *Ceratolithus cristatus*, *Cyclococcolithina leptopora*, *Helicopontosphaera kamptneri*, *Pseudoemiliana lacunosa*, and *Reticulofenestra reticulata*. Among the above mentioned species, *Gephyrocapsa oceanica* occupies about 60 percent of this nannoplankton population.

Table 1. Occurrences of calcareous nannoplanktons. (Frequencies in percent)

SPECIES	SAMPLE	G D P — 11 — 15																
		(1)	(6)	(7)	(8)	(9)	(10)	(11)	(16)	(17)	(18)	(21)	(26)	(31)	(36)	(41)	(46)	(50)
<i>Ceratolithus cristatus</i>		0.3	+	0.8	0.2		0.2	2.6	1.5	0.3	+					+		+
<i>C. rugosus</i>		+	+					+	+	0.3		0.3		+				
<i>Coccolithus adriaticus</i>										4.2		0.3						
<i>C. pelagicus</i>		+	+					+	0.3	0.9	2.8	1.9	1.2	1.6	0.9	2.3	1.1	1.1
<i>C. productus</i>		0.5	0.3		0.2	0.4	0.5	0.6				0.3					+	
<i>Cyclococcolithina leptopora</i>		11.7	14.7	12.5	11.7	8.9	13.6	15.6	12.6	9.0	10.8	6.4	6.1	6.4	7.6	10.3	9.6	9.2
<i>C. macintyreii</i>		1.3						1.7	1.5	2.2	2.3	3.6	4.9	3.9	5.4	4.3	4.1	4.9
<i>Cyclolithella annula</i>			0.3	0.7	0.2			0.3	+	+		0.6	2.3	0.9	0.6	0.6	0.6	0.9
<i>Discoaster browneri</i> (5 ray)		+	+					+	+	0.3	1.1	1.1	0.3	0.9	1.1	1.7	0.9	0.9
<i>D. browneri</i> (6 ray)									+	0.3	1.5	0.6	0.3	0.3	0.3	0.6	0.6	0.9
<i>D. browneri rutellus</i>									+		1.1	0.3	0.3	0.6	0.6	0.3	0.9	0.3
<i>D. challengeri</i>									+	+		0.3	+	+	+	+	+	+
<i>D. pentaradiatus</i>		+	+						+	+	2.2	4.0	6.9	8.1	7.0	4.3	6.6	2.6
<i>D. surculus</i>									+	+	0.3	1.7	1.4	0.3	0.3	2.3	0.6	0.6
<i>D. tamalis</i>			+		+				+		+	+	1.5	0.3	+	+	0.3	0.6
<i>D. variabilis</i>											+	+	+	+	+	0.3	+	0.6
<i>Emiliana hurleyi</i>		19.1	19.2	33.0	18.7	28.1	24.4	0.6										
<i>Gephyrocapsa caribbeanica</i>		0.5	0.3		0.7	1.8	0.7			3.1								
<i>G. ocaenica</i>		41.1	45.7	36.0	47.1	36.9	37.5	59.9	57.8	18.3	6.5							
<i>Helicopontosphaera hyalina</i>			0.6															
<i>H. kamptneri</i>		5.6	7.4	3.7	6.4	4.9	4.9	6.5	11.3	6.7	1.4	1.7	2.0	1.8	2.8	3.2	1.7	2.6
<i>H. sellii</i>															+		0.9	0.9
<i>H. wallichi</i>		0.8	0.3	0.3				0.2	0.3	+	+	+	0.3	0.6	0.3	0.6	+	0.6
<i>Isthmolithus fossilis</i>		0.3																
<i>Pontosphaera alboranensis</i>		+	+					0.3	+				+					
<i>P. discopora</i>			+				0.2	+	+	+		0.3	+	0.9	+	+	+	0.3
<i>P. multipora</i>			+															
<i>P. japonica</i>		0.3	+		0.5	0.2	1.2	0.3	+	+								0.6
<i>Pseudoemiliana lacunosa</i>				0.3	0.5	0.2		0.3	1.5	1.1	5.9	7.2	6.7	4.8	8.2	3.7	6.3	6.6
<i>Reticulofenestra reticulata</i>					0.2	0.2				39.1	55.1	52.4	42.4	50.0	40.3	50.3	50.0	54.9
<i>R. reticulata</i> var.												13.0	14.5	17.0	22.0	16.4	15.7	16.1
<i>Rhabdosphaera clavigera</i>		3.2	4.3	2.0	2.6	3.3	4.4	1.4	0.9	3.7	0.3			0.9				+
<i>Scyphosphaera pulcherima</i>												0.3					0.6	0.3
<i>Sphenolithus abies</i>										+								
<i>Syracosphaera pulchra</i>		1.0	1.7	0.8	0.7	1.3	1.6			0.3	+	1.4	0.6	1.5	0.6	0.9		0.6
<i>Thoracosphaera pelagica</i>		+	+							+								
<i>Umbellosphaera tenuis</i>		0.3	0.6			0.3		0.9		0.3								
<i>Umbilicosphaera mirabilis</i>		10.9	9.1	10.0	9.8	13.1	9.9	9.9	12.5	12.9	4.8	0.6	0.9	0.6				

Microfossils of the core sample GDP-11-15 from the Amami Plateau




Fig. 3. Distributions of some important microfossils in selected samples.

There is no species of *Discoaster* spp. except for ill-preserved species possibly derived from the lower part.

The assemblage III (GDP-11-115-(10)~(1))

The assemblage III is made of *Emiliana huxleyi*, *Gephyrocapsa oceanica*, *Ceratolithus cristatus*, *Cyclococcolithina leptopora*, *Helicopontosphaera kamptneri*, *Rhabdosphaera clavigera*, and *Umbilicosphaera mirabilis*. *Emiliana huxleyi* is a dominant species and occupies 18 to 30 percent of the assemblage.

2. Planktonic foraminifers

In the present work, 26 species of planktonic foraminifers were identified. The results of the analysis are shown in Table 2 and Figure 3.

Globigerina glutinata, *Globigerinoides ruber*, *G. sacculifer*, and *Globorotalia inflata* commonly occur throughout the core. *Globigerina pachyderma*, *Globigerinella siphonifera*, and *Orbulina universa* are constantly found in a small amount. The core faunas can be divided into two assemblages as follows in ascending order; the assemblage A is GDP-11-15-(50) to (18), and the assemblage B GDP-11-15-(17) to (1).

The assemblage A (GDP-11-15-(50)~(18))

Globigerinoides extremus, *Globoquadrina altispira*, *Globorotalia acostaensis*, *G. multicamerata*, and *G. tosaensis* characterize this assemblage. The coiling directions of *Globorotalia menardii* and *Pulleniatina obliquiloculata* are predominantly dextral. *Globorotalia truncatulinoides*, which seems to be contaminated, is found in this assemblage within 0.4 percent of the total specimens.

The assemblage B (GDP-11-15-(17)~(1))

Globoquadrina dutertrei and *Globorotalia truncatulinoides* are characteristic species. The species characterizing the assemblage A occur in this assemblage within 1 percent of the total specimens and these seem to be derived from the lower part. The predominant coiling direction of *Globorotalia menardii* is sinistral and that of *Pulleniatina obliquiloculata* dextral in this assemblage.

3. Benthonic foraminifers

44 genera, 120 species and 8 subspecies of benthonic foraminifers were identified. These faunas are mostly composed of calcareous forms, and arenaceous ones are scarcely found throughout the whole section. Most of specimens are assumed to be indigenous, because they are fresh in appearance and are in well state of preservation.

The species of more than one percent frequency are listed in Table 3, and the frequency diagram of dominant and depth-diagnostic species are also shown in Figure 3.

Table 2. Occurrences of planktonic foraminifers. (Frequencies in percent)

SPECIES	SAMPLE	G D P — 11 — 15													
		(1)	(6)	(11)	(16)	(17)	(18)	(21)	(26)	(31)	(36)	(41)	(46)	(50)	
<i>Candeina nitida</i> D'ORBIGNY										0.4					
<i>Globigerina pachyderma</i> (EHRENBERG)		0.9	1.2	3.1	3.1	3.6	3.0	1.8	3.5	0.9	1.7		0.9	2.3	
<i>G. quinqueloba</i> NATLAND		1.8		1.0	0.6	0.2		0.3			0.2				
<i>G. spp.</i>		47.3	29.0	31.9	34.0	26.7	38.7	36.2	41.2	41.9	24.3	45.1	31.3	29.1	
<i>Globigerinella siphonifera</i> (D'ORBIGNY)		0.9	1.2	0.5		1.4	0.5	0.5	1.0	2.6	2.1	0.7	0.4	0.3	
<i>Globigerinita glutinata</i> (EGGER)		7.3	4.4	3.1	8.6	0.2	9.9	8.4	8.2	7.0	7.6	6.9	7.5	3.0	
<i>Globigerinoides conglobatus</i> (BRADY)		0.9	0.4	0.3	1.2	0.5	1.0		0.4	0.4	0.4			0.5	
<i>G. extremus</i> BOLLI							0.7	3.1	1.6	1.7	3.0	1.5	0.4	0.8	
<i>G. ruber</i> (D'ORBIGNY)		8.2	17.1	14.6	8.0	13.7	8.4	10.2	10.1	4.4	10.6	4.7	10.6	11.0	
<i>G. sacculifer</i> (BRADY)		5.5	6.3	4.4	3.1	3.6	4.4	8.4	7.3	7.9	6.4	3.6	7.9	6.8	
<i>Globoquadrina altispira</i> (CUSHMAN et JARVIS)		0.5		0.3	0.6		1.0	1.3	1.5	0.4	0.1	1.1	1.3	2.5	
<i>G. dutertrei</i> (D'ORBIGNY)		3.2	5.6	5.7	2.5	4.1	2.2								
<i>Globorotalia acostaensis</i> BLOW								1.8	0.6		0.5			2.0	
<i>G. crassaformis</i> (GALLOWAY et WISSLER)		1.4	1.2	1.0		5.0	1.7	1.0	2.1	6.1	7.6	2.9	2.6	4.5	
<i>G. hirsuta</i> (D'ORBIGNY)						0.2									
<i>G. inflata</i> (D'ORBIGNY)		10.5	17.9	20.4	22.8	24.6	20.7	20.9	17.3	15.7	25.3	24.0	30.0	28.8	
<i>G. menardii</i> (D'ORBIGNY)		0.5	0.4	0.3	0.6	0.2	2.2	2.6	3.4	1.7	3.9	4.4	3.5	4.5	
<i>G. multicamerata</i> CUSHMAN et JARVIS											0.2	0.4			
<i>G. scitula</i> (BRADY)		0.9	2.4	2.1	0.6	2.3	0.7	0.3	0.5	2.2	2.4	1.5	1.8	1.0	
<i>G. tosaensis</i> TAKAYANAGI et SAITO		0.5					0.5	0.3	0.2	0.4	0.1	0.4			
<i>G. truncatulinoides</i> (D'ORBIGNY)		5.5	8.3	7.8	8.6	7.5	2.2	0.8	0.2	1.3	0.4	0.4	0.4	1.0	
<i>G. tumida</i> (BRADY)						0.2	0.2	0.8	0.4	1.3	0.7	1.1	0.4	0.8	
<i>Orbulina bilobata</i> (D'ORBIGNY)										0.4					
<i>O. suturalis</i> BRÖNNIMANN												0.4			
<i>O. universa</i> D'ORBIGNY		1.8	0.8	1.8	1.9	2.7	1.2	0.3	0.2	1.3	0.2	0.4		0.5	
<i>Pulleniatina obliquiloculata</i> (PARKER et JONES)		1.8	0.4	0.8	1.9	1.6		0.3	0.2	0.9	0.1	0.4	0.4	0.3	
<i>Sphaeroidinella dehiscens</i> (PARKER et JONES)				0.5			0.2	0.3	0.1		0.7		0.4	0.5	
unidentified specimens		0.9	3.6	0.3	1.9	1.6	0.5	0.8		0.9	1.4	0.4			
Total number of specimens		220	252	383	162	439	406	392	822	229	803	275	227	399	
Coiling direction															
<i>G. menardii</i>	number of specimens	20	20	20	20	20	20	20	28	20	31	20	20	20	
	dextral percent	90	100	100	95	85	40	5	4	10	6	0	0	5	
<i>P. obliquiloculata</i>	number of specimens	20	20	20	20	20	20	6	12	20	4	3	3	7	
	dextral percent	100	100	100	100	100	100	100	100	95	100	100	100	100	

Table 3. Occurrences of benthonic foraminifers.
(Frequencies in percent)
(Species of less than 1.0% frequencies are excluded in this list.)

SPECIES	SAMPLE	G D P — 11 — 15												
		(1)	(6)	(11)	(16)	(17)	(18)	(21)	(26)	(31)	(36)	(41)	(46)	(50)
<i>Bolivinita cf. minuta</i> (NATLAND)		4.7		3.8	2.3	3.1	0.8			0.3		0.1	0.9	
<i>Brisalina pacifica</i> (CUSHMAN et MCCULLOCH)				0.4		2.2								
<i>B. pussilla</i> (SCHWAGER)			1.3	1.3	1.0	0.4	0.4			0.3				
<i>Bulimina rostrata</i> BRADY		0.4	0.9	2.6	1.7	1.8	1.2		1.1	1.0	0.9	0.4	2.1	3.7
<i>B. translucens</i> PARKER					1.0	0.4	0.4							
<i>Carpenteria balaniformis</i> GRAY		2.2	0.9	0.4	1.7		0.4							
<i>Cassidulina carinata</i> SILVESTRI		1.7	4.3	2.6	2.0	3.5	0.8							
<i>C. neocarinata</i> THALMANN								1.7	1.1	1.3		0.1	0.9	
<i>C. sagamiensis</i> ASANO		0.9	0.4	1.3	0.3	0.4	1.7	3.0	5.0	5.1	6.0	5.8	6.4	9.7
<i>Civcidoides bradyi</i> (TRAUTH)		1.7	0.9	0.9	1.3	1.3	0.4			0.4	0.3	0.4		
<i>C. pseudoungerianus</i> (CUSHMAN)		3.0	1.3	0.4	1.3	0.9			0.4	0.3		0.5	0.4	0.7
<i>C. wuellerstorfi</i> (SCHWAGER)		1.7	1.7	1.7	3.0					0.7	0.4	0.4		
<i>Ehrenbergina pacifica</i> CUSHMAN		4.7	5.6	2.1	0.7	1.8	0.4		0.4					
<i>E. undulata</i> PARKER		7.3	7.3	9.8	7.3	7.9	3.7	0.4		1.0				
<i>Epistominella exigua</i> (BRADY)		0.9	0.4	1.3	0.7	3.1								
<i>E. pulchella</i> FUSEJIMA et MARUHASI		0.4	2.2	2.1	0.3									
<i>Fissurina formosa</i> (SCHWAGER)					1.0									
<i>F. orbignyana</i> (SEGUENZA)		0.9	0.9	0.4	0.3		1.2				0.4		0.4	
<i>Gavelinopsis translucens</i> (PHLEGER et PARKER)		0.4	2.6	1.7		2.2	0.4	2.2	1.8	1.0		0.9	2.1	0.7
<i>Globocassidulina depressa</i> (ASANO et NAKAMURA)		3.4	2.6	6.0	2.3	2.2	0.8							
<i>G. elegans boscensis</i> (KUWANO)		1.3	5.6	1.3	6.6	1.8	2.5	0.9	1.1	1.3		0.9	0.4	0.7
<i>G. ikebei</i> KONDA sp. nov.			2.2		1.0		0.4				0.4			0.7
<i>G. pacifica</i> (CUSHMAN)		9.1	6.0	3.8	0.7	0.4	0.8	1.3	3.5	1.3	3.4	0.6	7.3	1.5
<i>G. subglobosa</i> (BRADY)		14.2	9.9	10.7	6.6	7.9	6.6	5.6	7.4	7.7	6.4	7.8	4.7	5.2
<i>Gyroidina orbicularis</i> (D'ORBIGNY)				5.1	4.0	3.5	1.2			1.7	0.4	0.8	0.4	0.7
<i>Laticarinina halophora</i> (STACHE)		1.3	0.4	0.4	0.3							0.1		
<i>Melonis affinis</i> (REUSS)		1.3	0.4		1.3	0.9								
<i>Oridosalis imbonatus</i> (REUSS)		1.3	1.7	1.7	1.7	1.8	0.4	1.3	1.4	1.0	2.6	1.3	3.4	2.2
<i>Orthomorphina challengeriana</i> (THALMANN)									0.4	2.0	0.4	0.8		1.5
<i>Osangularia bengalensis</i> (SCHWAGER)		0.4	0.4	0.4		0.4	0.4	0.9	0.4		0.4	0.8	1.3	2.2

Microfossils of the core sample GDP-11-15 from the Amami Plateau

Table 3. (continued)

SPECIES	SAMPLE	G D P — 11 — 15												
		(1)	(6)	(11)	(16)	(17)	(18)	(21)	(26)	(31)	(36)	(41)	(46)	(50)
<i>Parafissurina ovalis</i> TODD			0.4	0.4	1.0			0.4	0.4	0.3		0.4		
<i>Pleurostomella acuminata</i> CUSHMAN		0.4				0.4	1.7	2.2	2.5	3.3	4.3	3.6	4.7	1.5
<i>P. brevis</i> SCHWAGER										0.3		0.9	1.3	1.5
<i>P. elliptica</i> GALLOWAY et HEMINWAY			0.4	0.4			2.5	4.8	3.5	3.7	4.3	4.0	5.5	3.7
<i>P. delicata</i> KONDA sp. nov.								0.9		1.4		0.3	0.9	
<i>P. praegerontica</i> CUSHMAN et STAINFORTH							2.1	0.4		0.7	0.4	1.6	1.3	
<i>P. rapana recens</i> DERVIEK												0.1	0.4	1.5
<i>Pullenia apertula</i> CUSHMAN		0.4	0.4	1.7		0.4				0.3				
<i>P. bulloides</i> (D'ORBIGNY)		2.2	1.7	0.9	1.0	1.8	1.2	0.4	1.8	1.0	1.3	1.6	1.7	3.7
<i>P. quinqueloba</i> (REUSS)		0.4	0.4		0.7	0.4	0.8	0.4	0.4	1.0		1.0	0.4	
<i>Pyrgo murrhina</i> (SCHWAGER)		0.4	0.4	1.7	1.3	0.9	1.7	0.9	0.4	0.7	0.4	0.8		
<i>Sigmotilina edwardsi</i> (SCHLUMBERGER)			2.2	1.3	0.3	0.4		0.4		0.9	0.8	1.3		
<i>Siphonodossaria consobrina</i> (D'ORBIGNY)			0.4			2.2	9.1	12.6	16.0	13.1	14.5	13.1	9.4	5.2
<i>S. oinomikadoi</i> (ISHIZAKI)		0.4					4.1	3.0			7.3	3.8	1.7	7.5
<i>Siphonovigerina ampullacea</i> (BRADY)		6.5	3.4	3.0	10.6	10.1	9.5	6.5	5.0	9.4	5.6	4.0	3.4	2.2
<i>Sphaeroidina bulloides</i> D'ORBIGNY			0.9		1.7	0.9	0.4			0.3		0.3		
<i>Stilostomella bradyi</i> (CUSHMAN)										2.0	0.9	0.3	0.4	
<i>S. hayasakai</i> (ISHIZAKI)					0.3		0.4	1.3		0.3		0.3	0.4	
<i>S. hyugaensis</i> (ISHIZAKI)									1.1				0.4	
<i>S. japonica</i> (ISHIZAKI)		0.4			0.3		4.6	8.7	8.9	14.1	5.1	5.6	5.5	4.5
<i>S. ketiensiensis</i> (ISHIZAKI)					0.7			1.7				0.9	0.4	1.5
<i>S. lepidula</i> (SCHWAGER)		0.9	0.9	0.4	1.0	4.0	6.6	8.7	5.7	4.0	6.4	7.1	8.9	11.2
<i>Trifarina angulosa</i> (WILLIAMSON)						0.4	2.9	5.2	4.6	1.7	1.3	1.9	3.8	3.0
<i>T. bradyi</i> CUSHMAN							0.4	1.3	1.8	1.0	2.1	0.3	0.4	
<i>Uvigerina aubertiana</i> D'ORBIGNY		1.3	1.7	1.3	2.0	1.8	1.7		0.4			0.1		0.7
<i>U. dirupta</i> TODD		1.7	3.9	2.1	4.6	1.3	0.8	0.4						
<i>U. proboscides</i> SCHWAGER		0.9		1.3	0.3	0.4		0.4		0.3				
Total number of specimens		232	232	234	302	227	241	231	282	297	234	773	235	134

Throughout the whole section, *Globocassidulina subglobosa* and *Siphonodossaria amplacea* are found with 8 and 6 percent on the average respectively, accompanied with *Globocassidulina pacifica*. However, the faunal assemblages of this core are divided into two types; namely, the assemblage α is GDP-11-15-(50) to (18), and the assemblage β GDP-11-15-(17) to (1).

The assemblage α (GDP-11-15-(50)~(18))

This assemblage is characterized by common occurrence of *Siphonodossaria consobrina*, *S. oinomikadoi*, *Stilostomella japonica*, *S. lepidula*, *Pleurostomella acuminata*, *P. elliptica*, *Cassidulina sagamiensis*, and *Trifarina angulosa* in order of abundance. Furthermore, *Cassidulina neocarinata*, *Orthomorphina anbigua*, *O. challengeriana*, *O. cf. obituensis*, *Pleurostomella praegerontica*, *P. brevis*, *P. rapana recens*, *Stilostomella bradyi*, *S. hayasakai*, and *Trifarina bradyi* are contained only in this assemblage, though each frequency of them is relatively low.

The assemblage β (GDP-11-15-(17)~(1))

In this assemblage, species characterizing the assemblage α as mentioned above are completely replaced by *Ehrenbergina undulata*, *E. pacifica*, *Globocassidulina depressa*, and *Cassidulina carinata*. Besides these species, rare but continuous occurrence of *Bolivina cf. minuta*, *Brizalina pussilla*, *Laticarinina halophora*, *Melonis affinis*, *Uvigerina dirupta*, and species of *Epistominella* spp. is remarkable.

Table 4. Relationship among the assemblages of calcareous nannoplanktons, planktonic and benthonic foraminifers.

Sample No.		Nannoplankton fossils		Foraminifers			Age
				planktonic		benthonic	
GDP 11-15	(1)	III	NN 21	B	N. 22	β	late pleistocené
	(10)				N. 23		
	(11)	II	NN 20				
	(17)						
	(18)	I	NN 16	A	N. 21	α	
(50)							

4. Other microfossils.

Organic-walled microfossils such as dinoflagellates, acritarchs, pollen grains and spores were not found in these selected core samples. A small amount of siliceous microfossils of radiolaria, sponge spicules, diatoms and ebridians were contained in several samples; GDP-11-15-(36), (26), (18), (17), and (16).

The relationship among the assemblages of calcareous nannoplanktons, planktonic and benthonic foraminifers is shown in Figure 3 and Table 4.

Conclusion

On the basis of nannoplanktons and planktonic foraminifers, the geological age of the present core is considered as follows.

The nannoplankton flora in this core is subdivided into three assemblages as mentioned above.

The assemblage I, judging from the coexistence of *Discoaster pentaradiatus* and *D. surculus*, belongs to the *Discoaster surculus* Zone, NN 16 (MARTINI and WORSELY, 1970) indicated by the interval from the last occurrence of *Reticulofenestra pseudumbilica* to that of *Discoaster surculus*. In the range chart of calcareous nannoplankton datum indicators presented by MARTINI (1971), the first appearance of *Pseudoemiliania lacunosa* is recorded in the middle part of the *Discoaster surculus* Zone. A lack of *Reticulofenestra pseudumbilica* and a few but continuous occurrence of *Pseudoemiliania lacunosa* from the lowest part in this core suggest that this assemblage may occupy the middle to upper part of the *Discoaster surculus* Zone.

Rare and sporadic occurrence of *Discoaster brouweri*, *D. pentaradiatus*, *D. tamalis* and *Pseudoemiliania lacunosa* of the assemblage II, suggests that these nannoplankton fossils might be contaminated and/or derived from the lower part. Based on the presence of *Gephyrocapsa oceanica* and the lack of *Emiliania huxleyi*, this assemblage is assigned to the *Gephyrocapsa oceanica* Zone, NN 20.

Abundant occurrence of *Emiliania huxleyi* in the assemblage III indicates that this assemblage belongs to the *Emiliania huxleyi* Zone, NN 21.

Accordingly, the *Discoaster pentaradiatus* Zone, NN 17, the *Discoaster brouweri* Zone, NN 18, and the *Pseudoemiliania lacunosa* Zone, NN 19, are lacking in this core.

NISHIDA (1976) noticed that *Gephyrocapsa protohuxleyi* is a valuable marker species in the middle to late Pleistocene in and around Japan. But there is no specimen of *G. protohuxleyi* in the assemblage II or III either. Consequently, it might be possible that sediments are lacking just below the appearance of *Emiliania huxleyi*.

As for the planktonic foraminifers, the assemblage A is thought to be of Zone N. 21 proposed by BLOW (1969), judging from the presence of *Globorotalia tosaensis* and *Globoquadrina altispira*. The assemblage B represents Zone N. 22 to N. 23 because of the frequent occurrence of *Globorotalia truncatulinoides*. The change of coiling direction of *Globorotalia menardii* supports this assignment. Furthermore, it is noticeable that specimens of *Pulleniatina obliquiloculata* are mostly dextral throughout the core. In other deep sea cores (Lamont Core V21-98; DSDP Leg 7 Site 62, 63, 64, 65) taken from the Philippine Sea and the western equatorial Pacific, the sinistral coiling *Pulleniatina obliquiloculata* have been predominantly recognized within Zone N. 21 and the lower part of Zone 22 (BRÖNNIMAN and RESIG, 1971; TAKAYAMA, 1973; UJIE and MIURA, 1970, 1971). In comparison with these results, it is considered that the upper part of Zone N. 21 and the lower part of Zone N. 22 are lacking in the present core, corresponding to the result from the study of the nannoplanktons.

Thus, it is concluded that there is a hiatus between the upper Pliocene and the upper Pleistocene corresponding to a time gap of about 2 million years in this core.

On the benthonic foraminiferal fauna, a remarkable break is observed between the late Pliocene assemblage α and the late Pleistocene assemblage β .

In comparison with the depth distribution of Recent benthonic foraminiferal assemblage in the Philippine and adjacent seas (CUSHMAN, 1921; FUSEJIMA *et al.*, 1943; POLSKI, 1959; YABE and HANZAWA, 1925), the assemblage α is considered to be of the upper bathyal zone, perhaps at depth of several hundred meters, under a warm water condition. Whereas, the assemblage β is regarded to be of the present depth. Therefore, the locality of the core is considered to have been under the bathyal condition in the late Pliocene. Subsequently, the sea bottom subsided relatively with the amount of about 1,000 m and in the late Pleistocene, this area had come to be at the present water depth.

In the Philippine Sea, some biostratigraphic gaps and paleoenvironmental changes have been discovered in the core sections during Leg 31 of Deep Sea Drilling Project (ELLIS, 1975; THE SHIPBOARD SCIENTIFIC PARTY, 1975; UJIE, 1975b, c). According to UJIE (1975c), the disconformity in the early to middle Miocene at Site 292 might have been caused by submarine erosion, and thereby reflected in lithologic change. In the present core, the fairly distinct lithologic change is observed between the portions above and below the hiatus. However, we cannot prove sufficiently the submarine erosion only with that change. Besides, KOIZUMI (1975a) recognized four cold-climate epochs and the corresponding sea level lowering after the Jaramillo Event on the basis of diatom distribution in the Lamont Core V21-80 taken from the Shikoku Basin.

However, nothing has been known on the hiatus and the environmental change

during the period from the late Pliocene to the late Pleistocene in the northern Philippine Sea. The results reported here are the first which elucidate them during that period.

We have not sufficient data to discuss the origin of environmental change. Late Cenozoic sea level changes and/or crustal movements in the Philippine Sea and adjacent land areas have been discussed by HOSHINO (1975), KIZAKI and TAKAYASU (1976), IKEBE (1976), and OKUDA *et al.* (1976) from various points of view. More data of the similar kind to this paper should be necessary at various stations in the Philippine Sea and other areas to prove the geologic events. Therefore, benthonic fauna of depth indicators would be as significant as planktonic fossils of age indicators in the future investigation of the micropaleontology of deep sea cores.

Acknowledgements

We are greatly indebted to Captain Magoshichi SATO and the crew members of the Tokaidai-gaku-maru II for contribution to the survey by their excellent operation and navigation. Our grateful thanks are due to Dr. Tsunemasa SHIKI (chief scientist of the GDP-11 Cruise) and Dr. Takao TOKUOKA of Kyoto University, and Professor Hitoshi AOKI and Dr. Yoshifumi MISAWA of Tokai University for many helpful suggestions on the geology of the Amami Plateau.

Deep gratitude is expressed to Dr. Manzo CHUJI, Director of the Osaka Museum of Natural History, for his valuable advice on the description of new foraminiferal species, and to Professor Keiji NAKAZAWA for his help to complete this paper.

Finally, we wish to acknowledge our indebtedness to students of Tokai University who cooperated in the work on board.

Description of New Species

by Isao KONDA

Family CASSIDULINIDAE D'ORBIGNY, 1839

Genus *Globocassidulina* VOLOSHINOVA, 1960

Globocassidulina ikebei n. sp.

(Pl. 5, Figs. 1, 2, 3; Pl. 6, Figs. 1a, b, c, 2a, b, c)

Description: Test free, small, subglobular in outline, periphery broadly rounded, slightly lobulated; wall smooth; chambers few, inflated, three pairs composing adult test, alternate chambers extending to the umbilicus on one side with only a small triangular portion visible on the opposite side; sutures flush with surface in early portion, later very slightly depressed; aperture large and broad, but nearly covered by a large crescent lip.

Dimension: Length of holotype 0.390 mm; breadth 0.341 mm.

Holotype: Department of Geology and Mineralogy, Faculty of Science, Kyoto University, Coll. Cat. No. JC1011, from the sample GDP-11-15-(6), picked up from 5.0–6.0 cm depth under the surface in the gravity core GDP-11-15, Lat. 28°06.2'N., Long. 131°35.2'E., at a water depth 1,830 m.

Remarks: This new species resembles *Cassidulina caudriae* described by CUSHMAN and STEINFORTH (1945) from the Oligocene Cipero Marl Formation of Trinidad, West Indies. However, compared with the figured holotype of *C. caudriae*, this new species is distinguished from *C. caudriae* by its much smaller size and its different aperture nearly covered by its large crescent lip. The tribal name of this new species is dedicated to Professor Emeritus Nobuo IKEBE of Osaka City University in recognition of his work on the Japanese Neogene biostratigraphy and chronology.

Occurrence: Rare in core samples GDP-11-15-(50), (36), (18), (16), and (6).

Famly PLEUROSOMELLIDAE REUSS, 1860

Genus *Pleuromella* REUSS, 1860

Pleuromella delicata n. sp.

(Pl. 5, Figs. 8a, b; Pl. 6, Figs. 3a, b, c, 4)

Description: Test free, small, elongate-oval in outline, nearly twice as long as broad, circular in cross section; wall thin, smooth, translucent; chambers few, each chamber covering major portion of preceeding one, reaching nearly to the base of test; sutures slightly depressed, indistinct in early stage; aperture small, with two small teeth on one side, and internal tube.

Dimension: Length of holotype 0.415 mm; breadth 0.220 mm.

Holotype: Department of Geology and Mineralogy, Faculty of Science, Kyoto University, Coll. Cat. No. JC1012, from the sample GDP-11-15-(21), picked up from 20.0–21.0 cm depth under the surface in the gravity core GDP-11-15, Lat. 28°06.2'N., Long. 131°35.2'E., at a water depth 1,830 m.

Remarks: This new species can be easily distinguished from other species of the genus in having a elongate-oval test. It is also characterized by a few chambers reaching nearly to the base of test.

Occurrence: Rare in core samples GDP-11-15-(50), (41), (31), and (21).

References

- AOKI, H. (1975): Volcanic rocks and others dredged in the GDP-11 cruise. In NAKAZAWA, K. *et al.* (eds.), *Geological problems of the Philippine Sea*, p. 87. (in Japanese)
- AOKI, H., KIM, Y., ISHIKAWA, M. and EGAWA, R. (1975): Petrological results in the GDP cruise. *Mar. Sci.*, 7 (7), pp. 28–33. (in Japanese with English abstract)
- BLOW, W.H. (1969): Late Middle Eocene to Recent planktonic foraminiferal biostratigraphy.

- In BRÖNNIMANN, P. and RENZ, H.H. (eds.), *Proc. Int. Conf. Planktonic Microfossils, Ist., Geneva, 1967*, 1, pp. 199–421, pls. 1–54.
- BRÖNNIMANN, P. and RESIG, J. (1971): A Neogene globigerinacean biochronologic timescale of the Southwestern Pacific. *Initial Rept. DSDP.*, 7, pp. 1235–1470.
- CUSHMAN, J.A. (1921): Foraminifera of the Philippine and adjacent seas. *U.S. Nat. Mus. Bull.*, 4, pp. 1–480, pls. 1–100.
- CUSHMAN, J.A. and STEINFORTH, R.M. (1945): The Foraminifera of the Cipero marl formation of Trinidad, British West Indies. *Cushman Lab. Foram. Res., Spec. Pub.*, 14, pp. 3–75, pls. 1–16.
- CUSHMAN, J.A., TODD, R. and POST, R.V. (1954): Recent Foraminifera of the Marshal Island. *U. S. Geol. Surv., Prof. Pap.*, 26;H, pp. 319–384, pls. 82–93.
- ELLIS, C.H. (1975): Calcareous nannofossil biostratigraphy—Leg 31, DSDP. *Initial Rept. DSDP.*, 31, pp. 655–676.
- FUSEJIMA, R., MARUHASHI, M. and KITAZAKI, U. (1943): The interrelation between foraminiferal assemblage and environmental conditions in the adjacent sea of the Philippine. *Mis. Rept. Res. Inst. Natur. Resources*, 1, pp. 23–40. (in Japanese)
- GALLOWAY, J.J. and HEMINWAY, C.E. (1941): The Tertiary Foraminifera of Porto Rico. *New York Acad. Sci., Sci. Surv. Porto Rico and Virgin Island*, 3, pt. 4, pp. 275–491, pls. 1–35.
- GORDON, W.A. (1961): Planktonic foraminifera and correlation of the middle Tertiary rocks of Puerto Rico. *Micropaleontology*, 7, pp. 451–460.
- HOSHINO, M. (1975): *Eustacy in relation to orogenic stage*. Tokai Univ. Press., Tokyo, 416p.
- IKEBE, N. (1976): Neogene geohistory of Japan in relation to the geohistory of the Philippine Sea. *Mar. Sci.*, 8 (3), pp. 24–29. (in Japanese with English abstract)
- KIZAKI, K. and TAKAYASU, K. (1976): Outline of geohistory of the Ryukyu Arc. *ibid.*, 8 (1), pp. 50–56. (in Japanese with English abstract)
- KOIZUMI, I. (1970): Diatom thanatocoenoses in deep sea cores from the Philippine Sea. *Jour. Mar. Geol.*, 6 (2), pp. 67–69. (in Japanese with English abstract)
- KOIZUMI, I. (1975a): Diatom flustule assemblages in the Philippine Sea. In NAKAZAWA, K. et al. (eds.), *Geological problems of the Philippine Sea*, pp. 46–52. (in Japanese)
- KOIZUMI, I. (1975b): Neogene diatoms from the western margin of the Pacific Ocean, Leg 31, Deep Sea Drilling Project. *Initial Rept. DSDP.*, 31, pp. 779–819, 5pls.
- KONDA, I. (1975): Some paleontological results and problematic subjects on GDP research cruise. *Mar. Sci.*, 7 (7), pp. 33–38. (in Japanese with English abstract)
- KONDA, I., HARADA, K., KITAZATO, H., MATSUOKA, K., NISHIDA, S., NISHIMURA, A., OHNO, T. and TAKAYAMA, T. (1975): Some paleontological results of the GDP-1, 8, 11 cruise. In NAKAZAWA, K. et al. (eds.), *Geological problems of the Philippine Sea*, pp. 91–98, 2pls. (in Japanese)
- KONDA, I., MATSUOKA, K., NISHIMURA, A. and OHNO, T.: Nummulites boninensis Hanzawa from the Amami Plateau in the northern margin of the Philippine Sea. *Trans. Proc. Paleont. Soc. Japan, N. S.* (in press)
- LING, H.Y. (1975a): Radiolaria: Leg 31 of the Deep Sea Drilling Project. *Initial Rept. DSDP.*, 31, pp. 703–761, 13 pls.
- LING, H.Y. (1975b): Silicoflagellates and Ebridians from Leg 31. *ibid.*, 31, pp. 763–777, 2 pls.
- MARTINI, E. (1971): Standard Tertiary and Quaternary calcareous nannoplankton zonation. In FARINACCI, A. (ed.), *Proc. II Plankton. Conf. Roma, 1970*, 2, pp. 739–785, incl. 4 pls.
- MARTINI, E. and WORSLEY, T. (1970): Standard calcareous nannoplankton zonation. *Nature*, 225, pp. 289–290.
- MATSUDA, J., SAITO, K. and ZASHU, S. (1975): K-Ar ages and Sr isotopic study of the igneous rock fragments in the manganese nodules dredged from the Amami Plateau, the Western

- Philippine Sea. In NAKAZAWA, K. *et al.* (eds.), *Geological problems of the Philippine Sea*, pp. 99–101. (in Japanese)
- MISAWA, Y. (1975): Morphological and geological results of the GDP-11 cruise in the Philippine Sea. *Mar. Sci.*, 7 (7), pp. 38–46. (in Japanese with English abstract)
- MISAWA, Y. and HOSHIZAWA, M. (1975): Sub-marine topography and acoustic survey during the GDP-11 cruise. In NAKAZAWA, K. *et al.* (eds.), *Geological problems of the Philippine Sea*, pp. 75–86. (in Japanese)
- NISHIDA, S. (1976): Late Cenozoic calcareous nannoplankton biostratigraphy in Japan. *Abstract of Pap., 1st CPNS*, pp. 159–162.
- OKUDA, Y., INOUE, E., ISHIIHARA, T., KINOSHITA, Y., TAMAKI, K., JOSHIMA, M. and ISHIBASHI, K. (1976): Submarine geology of the Nankai Trough and its peripheral area. *Mar. Sci.*, 8 (3), pp. 48–56. (in Japanese with English abstract)
- PARKER, F.L. (1964): Foraminifera from the experimental Mohole Drilling near Guadalupe Island, Mexico. *Jour. Paleont.*, 38 (4), pp. 617–636, pls. 97–102.
- POLSKI, W. (1959): Foraminiferal biofacies off the north Asiatic coast. *ibid.*, 33 (4), pp. 569–587, pl. 78.
- RESEARCH MEMBERS OF GDP-11 CRUISE (1975): Nummulites and pebbles of hornblende-tonalite and other igneous rocks, collected at the Amami Plateau. *Jour. Geol. Soc. Japan*, 81 (4), pp. 269–271. (in Japanese)
- SHIKI, T., AOKI, H. and MISAWA, Y. (1975): Geological results on the recent studies of the Philippine Sea—with special reference to GDP-8, 11. *Mar. Sci.*, 7 (7), pp. 22–28. (in Japanese with English abstract)
- SHIKI, T., TOKUOKA, T., AOKI, H., MISAWA, Y., KONDA, I. and NISHIDA, S. (1975): GDP cruise in the Philippine Sea with special reference to the bottom sampling in GDP-8 and 11. In NAKAZAWA, K. *et al.* (eds.), *Geological problems of the Philippine Sea*, pp. 67–74. (in Japanese)
- SUWA, K. and AOKI, H. (1975): Plutonic rocks from the Komahashi Seamount and the Amami Plateau. *ibid.*, pp. 88–89. (in Japanese)
- TAKAYAMA, T. (1969): Discoasters from the Lamont Core V21-98 (Preliminary reports of the Philippine Sea Core, Part 2). *Bull. Nat. Sci. Mus. Tokyo*, 12 (2), pp. 431–450.
- TAKAYAMA, T. (1970): The Pliocene-Pleistocene boundary in the Lamont Core V21-98 and at le Castella, southern Italy. *Jour. Mar. Geol.*, 6 (2), pp. 70–77.
- TAKAYAMA, T. (1973): On the distribution of calcareous nannoplankton in the youngest Cenozoic of Japan. *Mem. Geol. Soc. Japan*, 8, pp. 45–64. (in Japanese with English abstract)
- TODD, R. (1964): Planktonic Foraminifera from deep-sea cores off Eniwetok Atoll. *U. S. Geol. Surv., Prof. Pap.*, 260-cc pp. 1067–1100, pls. 289–295.
- THE SHIPBOARD SCIENTIFIC PARTY (1975): Site 296. *Initial Rept. DSDP.*, 31, pp. 191–223.
- UJIE, H. (1975a): Recent micropaleontological studies concerned to the Philippine Sea—especially to JOIDES. In NAKAZAWA, K. *et al.* (eds.), *Geological problems of the Philippine Sea*, pp. 43–45. (in Japanese)
- UJIE, H. (1975b): Micropaleontology of the Philippine Sea region. *Mar. Sci.*, 7 (7), pp. 13–22. (in Japanese with English abstract)
- UJIE, H. (1975c): Planktonic foraminiferal biostratigraphy in the western Philippine Sea, Leg 31 of DSDP. *Initial Rept. DSDP.*, 31, pp. 677–691, 3 pls.
- UJIE, H. and MIURA, M. (1970): Planktonic Foraminifera from the Laomnt core V21-98, Philippine Sea. *Jour. Mar. Geol.*, 6 (2), pp. 93–98.
- UJIE, H. and MIURA, M. (1971): Planktonic foraminiferal analysis of a calcareous ooze core from the Philippine Sea. In FARINACCI, A. (ed.), *Proc II Plankt. Conf. Roma*, 1970, 2, pp. 1231–1249, 3 pls.
- YABE, H. and HANZAWA, S. (1925): Globigerina ooze from the sea lying south of Okinawa-jima (the Riukiu Islands). *Japan Jour. Geol. Geogr.*, 4, (1–2), pp. 47–54.

Explanations of Plates

Plate 3

(Scanning photomicrographs of planktonic foraminifers)

Figs. 1a, b: *Globorotalia truncatulinoides* (D'ORBIGNY),
from the sample GDP-11-15-(11), $\times 60$.

Figs. 2a, b: *Globorotalia acostaensis* BLOW,
from the sample GDP-11-15-(21), $\times 90$.

Figs. 3a, b: *Globorotalia tosaensis* TAKAYANAGI et SAITO,
from the sample GDP-11-15-(18), $\times 100$.

Figs. 4, 5: *Globoquadrina altispira* (CUSHMAN et JARVIS),
4; from the sample GDP-11-15-(41), $\times 75$.
5; from the sample GDP-11-15-(50), $\times 75$.

Figs. 6a, b: *Globigerinoides extremus* BOLLI,
from the sample GDP-11-15-(21), $\times 100$.

Figs. 7a, b: *Globorotalia multicamerata* (CUSHMAN et JARVIS),
from the sample GDP-11-15-(41), $\times 40$.

Plate 4

(Scanning photomicrographs of benthonic foraminifers)

Figs. 1, 2: *Ehrenbergina undulata* PARKER,
from the sample GDP-11-15-(16), $\times 75$.

Figs. 3, 4: *Cassidulina neocarinata* THALMANN,
3; from the sample GDP-11-15-(21), $\times 100$.
4; from the sample GDP-11-15-(26), $\times 100$.

Fig. 5: *Stilostomella lepidula* (SCHWAGER),
from the sample GDP-11-15-(50), $\times 45$.

Figs. 6, 7: *Stilostomella ketienziensis* (ISHIZAKI),
from the sample GDP-11-15-(41), $\times 40$.

Figs. 8, 9: *Ehrenbergina pacifica* CUSHMAN,
8; from the sample GDP-11-15-(6), $\times 90$.
9; from the sample GDP-11-15-(1), $\times 85$.

Fig. 10: *Stilostomella japonica* (ISHIZAKI),
from the sample GDP-11-15-(31), $\times 40$.

Fig. 11: *Orthomorphina challengerina* (THALMANN),
from the sample GDP-11-15-(31), $\times 50$.

Fig. 12: *Trifarina angulosa* (WILLIAMSON),
from the sample GDP-11-15-(46), $\times 100$.

Fig. 13: *Trifarina bradyi* CUSHMAN,
from the sample GDP-11-15-(26), $\times 100$.

Fig. 14: *Uvigerina auberiana* D'ORBIGNY,
from the sample GDP-11-15-(16), $\times 75$.

Fig. 15: *Uvigerina dirupta* TODD,
from the sample GDP-11-15-(16), $\times 50$.

Figs. 16, 17: *Siphonodosaria consobrina* (D'ORBIGNY),
16; from the sample GDP-11-15-(46), $\times 50$.
17; from the sample GDP-11-15-(46), $\times 30$.

Plate 5

(Scanning photomicrographs of benthonic foraminifers)

Figs. 1, 2, 3: *Globocassidulina ikebei* KONDA, n. sp., paratypes,

1; from the sample GDP-11-15-(16), $\times 90$.

2; from the sample GDP-11-15-(36), $\times 90$.

3; from the sample GDP-11-15-(6), $\times 120$.

Figs. 4, 5: *Cassidulina sagamiensis* ASANO *et* NAKAMURA,

4; from the sample GDP-11-15-(31), $\times 100$.

5; from the sample GDP-11-15-(18), $\times 90$.

Figs. 6, 7: *Globocassidulina depressa* ASANO *et* NAKAMURA,

from the sample GDP-11-15-(16), $\times 200$.

Figs. 8a, b: *Pleurostomella delicata* KONDA, n. sp., paratype,

from the GDP-11-15-(31), a; $\times 150$, b; $\times 500$.

Figs. 9, 10: *Cassidulina carinata* SILVESTRI,

9; from the sample GDP-11-15-(16), $\times 100$.

10; from the sample GDP-11-15-(6), $\times 125$.

Fig. 11: *Laticarinina halophora* (STACHE),

from the sample GDP-11-15-(1), $\times 50$.

Plate 6

Figs. 1a, b, c: *Globocassidulina ikebei* KONDA, n. sp., holotype,

from the sample GDP-11-15-(6), $\times 110$, a, c; side view b; edge view.

Figs. 2a, b, c: *Globocassidulina ikebei* KONDA, n. sp., paratype,

from the sample GDP-11-15-(16), $\times 110$, a, c; side view, b; edge view.

Figs. 3a, b, c: *Pleurostomella delicata* KONDA, n. sp., holotype,

from the sample GDP-11-15-(21), $\times 110$, a, c; side view, b; edge view.

Fig. 4: *Pleurostomella delicata* KONDA, n. sp., paratype,

from the sample GDP-11-15-(31), $\times 110$, aperture side.

