

Title	Effectiveness of simplified 15-min refresher BLS training program: a randomized controlled trial.
Author(s)	Nishiyama, Chika; Iwami, Taku; Murakami, Yukiko; Kitamura, Tetsuhisa; Okamoto, Yoshio; Marukawa, Seishiro; Sakamoto, Tetsuya; Kawamura, Takashi
Citation	Resuscitation (2015), 90: 56-60
Issue Date	2015-05
URL	http://hdl.handle.net/2433/199884
Right	© 2015 Elsevier Ireland Ltd. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International http://creativecommons.org/licenses/by-nc-nd/4.0/ . NOTICE: this is the author's version of a work that was accepted for publication in "Resuscitation". Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in [Resuscitation, Volume 90, May 2015, Pages 56–60, doi:10.1016/j.resuscitation.2015.02.015]; The full-text file will be made open to the public on 1 June 2016 in accordance with publisher's 'Terms and Conditions for Self-Archiving'.
Type	Journal Article
Textversion	author

1 **Effectiveness of simplified 15-min refresher BLS training program: a randomized**
2 **controlled trial**

3

4 **Chika Nishiyama^a, Taku Iwami^b, Yukiko Murakami^b, Tetsuhisa Kitamura^c,**
5 **Yoshio Okamoto^d, Seishiro Marukawa^e, Tetsuya Sakamoto^f, Takashi Kawamura^b**

6

7 ^a Department of Critical Care Nursing, Kyoto University Graduate School of Human
8 Health Science, 53 Shogoin Kawahara-cho, Sakyo-ku, Kyoto 606-8507, Japan.

9 ^b Kyoto University Health Service, Yoshida-Honmachi, Sakyo-ku, Kyoto 606-8501,
10 Japan.

11 ^c Division of Environmental Medicine and Population Sciences, Department of Social
12 and Environmental Medicine, Graduate School of Medicine, Osaka University, 2-5
13 Yamada-oka, Suita, Osaka 565-0871, Japan.

14 ^d Kagawa Prefectural Central Hospital, 1-2-1 Asahimachi, Takamatsu City, Kagawa
15 760-8557, Japan.

16 ^e Iseikai Hospital, 6-2-25 Sugawara, Higashi Yodogawa-ku, Osaka 533-0022, Japan.

17 ^f Department of Emergency Medicine, Teikyo University School of Medicine, 2-11-1
18 Kaga, Itabashi-ku, Tokyo 173-8605, Japan.

19 **Word count of the paper:** 2665 words

20 **Word count of the abstract:** 241 words

21 **Number of figures:** 1

22 **Number of tables:** 2

23

24 Corresponding author:

25 Taku Iwami, MD, PhD

26 Kyoto University Health Service, Yoshida Honmachi, Sakyo-ku, Kyoto 606-8501,

27 Japan.

28 Phone: +81-75-753-2401, Fax: +81-75-753-2424

29 E-mail address: iwamit@e-mail.jp

30

31 *Keywords:*

32 Basic life support (BLS)

33 Bystander CPR

34 Cardiac arrest

35 Cardiopulmonary resuscitation (CPR)

36 Retraining

- 37 Chest compression
- 38 Education
- 39 Randomized controlled trial
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54

55 **A B S T R A C T**

56 *Objectives:* To evaluate the long-term effectiveness of 15-min refresher basic life
57 support (BLS) training following 45-min chest compression-only BLS training.

58 *Methods:* After the 45-min chest compression-only BLS training, the participants were
59 randomly assigned to either the refresher BLS training group, which received a 15-min
60 refresher training 6 months after the initial training (refresher training group), or to the
61 control group, which did not receive refresher training. Participants' resuscitation skills
62 were evaluated by a 2-min case-based scenario test one year after the initial training.

63 The primary outcome measure was the number of appropriate chest compressions
64 during a 2-min test period.

65 *Results:* 140 participants were enrolled and 112 of them completed this study. The
66 number of appropriate chest compressions performed during the 2-min test period was
67 significantly greater in the refresher training group (68.9 ± 72.3) than in the control
68 group (36.3 ± 50.8 , $p = 0.009$). Time without chest compressions was significantly
69 shorter in the refresher training group (16.1 ± 2.1 seconds versus 26.9 ± 3.7 seconds, $p <$
70 0.001). There were no significant differences in time to chest compression (29.6 ± 16.7
71 seconds versus 34.4 ± 17.8 seconds, $p = 0.172$) and AED use between the groups.

72 *Conclusions:* A short-time refresher BLS training program 6 months after the initial
73 training can help trainees retain chest compression skills for up to one year. Repeated
74 BLS training, even if very short, would be adopted to keep acquired CPR quality
75 optimal. (UMIN-CTR UMIN 000004101)

76

77

78

79

80

81

82

83

84

85

86

87

88

89

90

91 **1. Introduction**

92 Out-of-hospital sudden cardiac arrest (OHCA) continues to be a leading cause of
93 death in many regions of the world.¹⁻³ Bystander cardiopulmonary resuscitation (CPR)
94 can double or triple the likelihood of survival after OHCA.^{1,4,5} Chest compressions are
95 an especially critical component of CPR.¹ Animal and clinical investigations have
96 suggested that continuous chest compressions without ventilation are as effective as
97 chest compression plus rescue breathing resuscitation for most cases of cardiac arrests⁶⁻
98 ⁸ and even more effective in some types of cardiac arrests.⁹⁻¹¹ Chest compression-only
99 CPR is attractive not only because it can provide a greater number of chest
100 compressions without interruption, but also because it is simpler and easier to learn and
101 perform than conventional CPR.¹²⁻¹⁴

102 Despite intensive efforts to train the general public in CPR and usage of an
103 automated external defibrillator (AED), the proportions of bystander CPR and AED use
104 generally remain unacceptably low.^{1,2,15,16} Because this complex psychomotor task is
105 difficult to learn and perform,^{6,13,17} basic life support (BLS) training for laypersons
106 should be more focused on essential skills and performed repeatedly. We have
107 demonstrated that our 45-min chest compression-only BLS training makes it possible

108 for the general public to perform more appropriate chest compressions than the 3-hour
109 conventional BLS program.¹⁸

110 However, it is well known that CPR skills and knowledge rapidly deteriorate over
111 time.¹ Current American Heart Association (AHA) BLS training courses certify trainees
112 for two years, but the optimal retraining time for maintaining these skills has not been
113 established.¹ The purpose of this study is to evaluate the skill retention of one year after
114 the 45-min chest compression-only BLS training between those who had or did not have
115 the 15-min refresher BLS training 6 months after the initial training.

116

117 **2. Methods**

118 *2.1 Study design*

119 This study was designed as a randomized controlled trial and was conducted
120 between April 2010 and March 2012.

121

122 *2.2 Participants recruitment and randomization*

123 The employees and students of Kyoto University aged 18 years or older were
124 recruited for this study via billboard advertisements and the Kyoto University web-site
125 as well as by word of mouth from the participants themselves. Participants were offered

126 \$20 as an incentive to be evaluated at one year after the initial chest-compression BLS
127 training. Health care professionals, medical/co-medical students and paramedics, and
128 those whom the program director considered unsuitable for resuscitation training (e.g.,
129 those with obvious physical and mental disabilities) were excluded.

130 Participants were randomly assigned to either the refresher training group or the
131 control group using permuted blocks after stratification by sex and age (age < 40 or ≥40
132 years). A computer-generated randomization list was provided by an independent
133 biostatistician. The allocation was concealed from all participants and instructors until
134 the completion of the initial chest compression-only BLS training.

135

136 *2.3 Intervention*

137 For members of both the refresher training group and the control group, the 45-min
138 chest compression-only BLS training consisting of chest compressions, and AED use
139 was carried out using a digital video disk (DVD) instructional material and a compact
140 personal resuscitation manikin named Mr.PUSH[®]. This personal training manikin
141 makes a sound when a trainee provides chest compressions with appropriate pressure to
142 reach 5cm based on clinical data.¹⁹ Effectiveness of the 45-min chest compression-only
143 BLS training program was previously evaluated.¹⁸ Mr.PUSH[®] was designed to train

144 people to perform chest compression. Both the DVD and Mr.PUSH[®] were developed
145 by the Osaka Life Support Association, a non-profit organization which is one of the
146 most active bodies dedicated to spreading BLS training in communities in Japan
147 (<http://osakalifesupport.jp/osakalsa/>). The main instructor of the training program
148 conducted the course with the DVD with supporting instructors assigned to every 20
149 participants. Each supporting instructors observed and assisted 20 trainees whether they
150 followed up the main instructor and DVD instruction or not. If not, the supporting
151 instructors supported them to keep up. A total of 20-100 participants underwent the
152 training at the same time (http://osakalifesupport.jp/push_e/index.html).

153 The refresher training group members were given an additional 15-min refresher
154 training program 6 months after the initial BLS training. The refresher training program
155 included 5 items: (1) Overview (1 min), (2) emergency call and cardiac arrest
156 recognition (3 min), (3) chest compressions (5 min), (4) AED use (5 min), and (5)
157 questions and answers (1 min). To standardize the quality of training program and
158 instructors, the 15-min refresher training program was carried out based on DVD-based
159 training program. The instructor was only allowed to give brief feedback for basic skills
160 such as hand position or compression depth. The control group members did not attend
161 the 15-min refresher training program.

162 A total of 20 physicians, nurses and emergency medical technicians, all of whom
163 were instructors of the Immediate Cardiac Life Support (ICLS) course certified by the
164 Japanese Association for Acute Medicine (JAAM), were specifically trained for this
165 study to maintain the quality of this training program.

166

167 *2.4 Outcome measures*

168 One year after their initial 45-min chest compression-only BLS training,
169 resuscitation skills were evaluated using a case-based scenario. In this test, each
170 participant was called individually into the testing room and provided the following
171 scenario: “Imagine that you are at a department store. Suddenly a man collapses in front
172 of you. You are the only person around. Do whatever you can do to help this man.”
173 After presentation of the scenario, we evaluated their CPR skills including initial
174 assessment, call for 119 (the emergency call number in Japan), call for an AED, and
175 chest compressions. After the CPR evaluation, AED was brought to the participant by
176 the instructor and participants were encouraged to use it. The Laerdal[®] PC Skill
177 Reporting software[™] (Laerdal Medical, Stavanger, Norway) automatically recorded
178 CPR performance variables for each subject.

179 The primary outcome measure was the number of appropriate chest compressions

180 during a 2-min test period at the evaluation test. An appropriate chest compression was
181 defined as one with a depth of over 5.0 cm, correct hand position, and completely
182 recoiling according to the Japanese CPR guidelines.⁵

183 The secondary outcome measures included the number of total chest compressions,
184 the proportion of appropriate chest compressions, and time without chest compressions
185 during the 2-min test period. Time from starting the presentation to first chest
186 compression and time from arriving at AED beside the participant to the first
187 defibrillation were measured. Calls for 119 and AED, switching-on the AED, correct
188 positioning of defibrillator pads, and assuring safety of the victim were also assessed by
189 the instructors using the check list.

190

191 *2.5 Statistical methods*

192 The sample size was calculated for the number of appropriate chest compressions
193 one year after the training based on previous reports,^{12,20} and was assumed to be 81
194 times in the refresher training group and 48 times in the control group. Under the
195 condition of an alpha error of 5% and a power of 80%, 53 subjects were needed per
196 group. Projecting a 10% dropout, the sample size was estimated to be 120 subjects in
197 total.

198 Analyses were performed on an intention-to-treat basis, but participants who were
199 absent from the one-year-later evaluation test were not included in the analyses
200 regardless of the participation to the refresher training. The data were compared across
201 groups using chi-square test for categorical variables and Student's *t*-test for continuous
202 variables. An analysis of covariance was conducted to adjust for sex and age. Analyses
203 were performed using SPSS Ver.21J (IBM SPSS, Armonk, NY). A two-tailed value of
204 $p < 0.05$ was considered statistically significant.

205

206 *2.6 Ethical considerations*

207 All procedures were conducted according to the Declaration of Helsinki. The
208 participants submitted written informed consent prior to participation. This study was
209 approved by the Ethics Committees of Kyoto University Graduate School of Medicine
210 (registration number E999).

211

212 **3. Results**

213 *3.1 Flow and baseline characteristics*

214 In total, 140 participants were enrolled in this study, and 71 and 69 were assigned to
215 the refresher training group and the control group, respectively. Among those assigned

216 to the refresher training group, 62 participants actually underwent the 15-min refresher
217 training. One year later, 57 (80.3%) in the refresher training group and 55 (79.7%) in
218 the control group were completely evaluated for their BLS skills (Fig. 1). The
219 demographic data of the two groups are summarized in Table 1. There were no
220 significant differences in sex ratio, age, previous CPR training, experience of actual
221 CPR, and family history of sudden cardiac arrest between the groups. Although nine
222 participants who did not attend the refresher training program in the refresher training
223 group, their demographic data was not different from that of those who completed the
224 study. Nineteen participants who did not attend the one-year-later evaluation test also
225 did not have significantly different demographic data from those who completed the
226 study.

227

228 *3.2 Performance of BLS skills one year after training*

229 Table 2 shows the participants' activation of emergency medical services (EMS)
230 and their chest compression skills one year after the training. Among those whose CPR
231 skills were tested, 55 (96.5%) in the refresher training group tried CPR, as opposed to
232 52 (94.5%) in the control group. The number of participants placing a 119 call was 46
233 (83.6%) in the refresher training group and 39 (75.0%) in the control group ($p=0.343$).

234 Fifty-two (94.5%) of those in the refresher training groups called for AED, while 43
235 (82.7%) in the control groups did so ($p=0.027$).

236 The number of total chest compressions was also significantly greater in the
237 refresher training group than in the control group (182.0 ± 41.7 versus 142.0 ± 59.1 , $p <$
238 0.001). The number of appropriate chest compressions performed during the 2-min test
239 period was significantly greater (68.9 ± 72.3) in the refresher training group than in the
240 control group (36.3 ± 50.8 , $p = 0.009$). The proportion of appropriate chest
241 compressions was significantly greater in the refresher training group than in the control
242 group ($38.3\% \pm 37.6\%$ versus $27.5\% \pm 34.4\%$, $p = 0.009$). Time without chest
243 compressions was significantly shorter in the refresher training group (16.1 ± 2.1
244 seconds versus 26.9 ± 3.7 seconds, $p < 0.001$). However there were no significant
245 differences in time to first chest compression between the two groups (29.6 ± 16.7
246 seconds versus 34.4 ± 17.8 seconds, $p = 0.172$).

247 All participants attempted to use an AED. Specific of AED use including turning-on,
248 pad-positioning, and area clearing were not significantly different between the groups.

249

250 **4. Discussion**

251 This is a randomized control trial to show the effectiveness of a very short refresher
252 BLS training on longer-term CPR and AED skill retention in the general public. One
253 year after the training, the number of total and appropriate chest compressions were
254 both significantly greater in the refresher training group compared to the control group.

255 Our novel refresher BLS training was only 15 minutes in duration and focused
256 entirely on the chest compressions and AED use, using the handy personal manikin. It
257 is well accepted that hands-on training is effective for reinforcing the quality of
258 CPR.^{21,22} However, limited resources and time have been a barrier to more hands-on
259 practice with a manikin,²³ leading to poor skill acquisition.²⁴ This is very important
260 because a shorter length refresher training could be better accepted by both instructors
261 and participants. Focusing on chest compressions and AED use and the use of a
262 personal manikin could make this short training program more efficient and effective.

263 This study successfully suggested that a refresher BLS training program would help
264 the general public retain their chest compression skills for up to one year. Previous
265 studies suggested that CPR skills declined faster than expected,^{25, 26} and that a repeated
266 training course can obviously maintain CPR performance for a long time.^{22,27} The CPR
267 Guidelines recommend that skill performance should be reinforced more often with an
268 interval of at most 12 to 24 months to maintain the quality of the rescuer's CPR. A

269 previous study suggested that, in order to minimize skill decay, the recommended
270 interval for refresher training should not be longer than seven months.²⁸ In addition,
271 Wollard M et al. showed that frequent short intervals refreshers improved not only skills
272 but also confidence.²⁹ Whilst Bohn A et al. reported that annual resuscitation training
273 provided by trained teachers are effective and adequate in children aged 10 years.³⁰ The
274 optimal timing for reassessment or reinforcement should be future discussed.¹

275 Although this study demonstrated that a 15-min short refresher BLS training could
276 maintain CPR and AED skills, CPR qualities including the number of appropriate chest
277 compressions, chest compressions with appropriate depth, and chest compressions with
278 correct hand position were not sufficient compared with the guideline-recommended
279 levels regardless of the group. This suggests the need for more frequent and effective
280 CPR training. The current program, which was scheduled and instructor-led, might still
281 be inconvenient for both trainees and instructors, even though the refresher training
282 course was short. Different approaches to skill retention, such as poster retraining with a
283 manikin,^{31,32} self-instruction using voice assistance with a manikin,^{22,33} and mobile
284 phone-based reminding have been attempted.³⁴ Further study initiatives with simpler,
285 more cost-effective, unscheduled, and autonomous refresher training courses would be
286 needed to improve CPR skill retention. In the next study, we plan to evaluate the

287 effectiveness of a self-learning refresher training program.

288 In terms of AED operations, the participants generally used an AED well
289 irrespective of refresher training. Our previous study also showed that 70% of the
290 untrained citizens who were provided with a self-learning video before attending a BLS
291 training course attempted to use an AED.¹⁴ Gundry et al. reported that AEDs were
292 safely and successfully operated by sixth-year pupils, who performed defibrillation in
293 90 seconds in a simulated resuscitation.³⁵ The operation of AED is rather simple and
294 there is almost no skill deterioration.^{29,36} In addition, the voice prompts of the AED can
295 support lay rescuers to use it at the actual emergency scene.³⁷⁻³⁹ However, the
296 proportion of those who called for AED was significantly higher and the time to
297 defibrillation tended to be shorter in the refresher group. In addition, we have to take
298 into account the fact that we encouraged participant to use an AED in this study.
299 Considering the clinically important rolls of AEDs and simplicity to train how to use an
300 AED, AED training should be included in the refresher training program, even though
301 the training program is short.

302 Our study has some limitations. First, the resuscitation skills were evaluated by a
303 case-based scenario test, and resuscitation performance in the real setting where lay
304 persons would easily panic was unknown. Second, data on further long-term retention

305 and the effects of repetitive training were lacking. We are planning a study to evaluate
306 the trainees' BLS skills for regular refresher training programs (such as every 90 days).
307 Third, there might be bias by the allocation because we did not blind training
308 assignment. Moreover, to ensure the effectiveness of this refresher training program, we
309 are planning to evaluate it in the real world, measuring the proportion of bystander CPR
310 and survival after OHCA.

311

312 **5. Conclusion**

313 A short-time refresher BLS training program would make it possible to retain chest
314 compressions skills for up to one year. Repeated BLS training would be adopted to keep
315 the acquired CPR quality optimal. Further study is warranted to identify whether
316 refresher training delivered with intervals shorter than 6 months may have benefits for
317 chest compression and AED skill retention.

318

319 **Conflict of interest**

320 Taku Iwami is one of the developers of Mr.PUSH[®] and is an executive director of the
321 non-profit organization Osaka Life Support Association, but has no financial conflict of
322 interest to declare. The rest of the authors also have no conflict of interest to declare.

323 **Role of funding source**

324 This study was supported by a Grant-in-Aid for Health and Labour Sciences Research
325 Grants (H24-Shinkin-001) from the Ministry of Health, Labour and Welfare.

326

327 **Acknowledgements**

328 We gratefully acknowledge Masaaki Matsumoto, Katsuharu Hirai, Keiji Akatsuka,
329 Katsuo Ogura, Seiji Kasatani, Nobuyuki Iwai, Yasuyuki Shinkai, and Yuji Yamazaki
330 for instructing in the CPR training program. We also thank all the faculty members of
331 Kyoto University School of Public Health for their helpful comments on the study's
332 design and data analyses, both of which were critical to the study's success.

333

334

335

336

337

338

339

340

341 **References**

- 342 1. 2010 American Heart Association Guidelines for Cardiopulmonary Resuscitation
343 and Emergency Cardiovascular Care. *Circulation* 2010;112 (Supp 3):S640-S946.
- 344 2. Rea TD, Eisenberg MS, Becker LJ, Murray JA, Hearne T. Temporal trends in
345 sudden cardiac arrest: a 25-year emergency medical services perspective.
346 *Circulation* 2003;107:2780-5.
- 347 3. Kitamura T, Iwami T, Kawamura T, Nagao K, Tanaka H, Hiraide A. Nationwide
348 public-access defibrillation in Japan. *N Engl J Med* 2010;11:994-1004.
- 349 4. European Resuscitation Council guidelines for resuscitation 2010. *Resuscitation*
350 2010;81:1219-2451.
- 351 5. Japan Resuscitation Council. *2010 Japanese Guidelines for Emergency Care and*
352 *Cardiopulmonary Resuscitation*. Tokyo, Japan: Health Shuppansha;2011.
- 353 6. Berg RA, Kern KB, Sanders AB, Otto CW, Hilwig RW, Ewy GA. Bystander
354 cardiopulmonary resuscitation. Is ventilation necessary? *Circulation* 1993;88:1907-
355 15.
- 356 7. Iwami T, Kawamura T, Hiraide A, et al. Effectiveness of Bystander-Initiated Chest
357 compression only Resuscitation for Patients with Out-of-Hospital Cardiac Arrest.
358 *Circulation* 2007;116:2900-7.

- 359 8. Hallstrom A, Cobb L, Johnson E, Copass M. Cardiopulmonary resuscitation by
360 chest compression alone or with mouth-to-mouth ventilation. *N Engl J Med*
361 2000;342:1546-53.
- 362 9. Iwami T, Kitamura T, Kawamura T, et al. Chest Compression-Only
363 Cardiopulmonary Resuscitation for Out-of-Hospital Cardiac Arrest With Public-
364 Access Defibrillation: A Nationwide Cohort Study. *Circulation* 2012;126:2844-51.
- 365 10. Hüpfel M, Selig HF, Nagele P. Chest-compression-only versus standard
366 cardiopulmonary resuscitation: a meta-analysis. *Lancet* 2010;376:1552-7.
- 367 11. Bobrow BJ, Spaite DW, Berg RA, et al. Chest compression-only CPR by lay
368 rescuers and survival from out-of-hospital cardiac arrest. *JAMA* 2010;304:1447-54.
- 369 12. Nishiyama C, Iwami T, Kawamura T, et al. Effectiveness of simplified chest
370 compression-only CPR training for the general public: a randomized controlled trial.
371 *Resuscitation* 2008;79:90-6.
- 372 13. Heidenreich JW, Sanders AB, Higdon TA, Kern KB, Berg RA, Ewy GA.
373 Uninterrupted chest compression CPR is easier to perform and remember than
374 standard CPR. *Resuscitation* 2004;63:123-30.

- 375 14. Nishiyama C, Iwami T, Kawamura T, et al. Effectiveness of simplified chest
376 compression-only CPR training program with or without preparatory self-learning
377 video: A randomized controlled trial. *Resuscitation* 2009;80:1164-8.
- 378 15. Iwami T, Nichol G, Hiraide A, et al. Continuous improvements of “chain of
379 survival” increased survival after out-of-hospital cardiac arrests: a large-scale
380 population-based study. *Circulation* 2009;119:728-34.
- 381 16. Herlitz J, Andersson E, Bang A, et al. Experiences from treatment of out-of-hospital
382 cardiac arrest during 17 years in Goteborg. *Eur Heart J* 2000;21:1251-8.
- 383 17. Chamberlain DA, Hazinski MF. Education in resuscitation: an ILCOR symposium:
384 Utstein Abbey: Stavanger, Norway: June 22-24, 2001. *Circulation* 2003;108:2575-
385 94.
- 386 18. Nishiyama C, Iwami T, Kitamura K, et al. Long-term retention of cardiopulmonary
387 resuscitation skills after shortened chest compression-only cardiopulmonary
388 resuscitation training and conventional cardiopulmonary resuscitation training: a
389 randomized controlled trial. *Acad Emerg Med* 2014;21:47-54.
- 390 19. Retzer E, Yuen TC, Cua JL, Edelson DP. Sex Disparities in Chest Compression
391 Quality. *Circulation* 2011; 124: A290.

- 392 20. Andresen D, Arntz HR, Gräfling W, et al. Public access resuscitation program
393 including defibrillator training for laypersons: a randomized trial to evaluate the
394 impact of training course duration. *Resuscitation* 2008;76:419-24.
- 395 21. Bobrow BJ, Vadeboncoeur TF, Spaite DW, et al. The effectiveness of ultrabrief and
396 brief educational videos for training lay responders in hands-only cardiopulmonary
397 resuscitation: implications for the future of citizen cardiopulmonary resuscitation
398 training. *Circ Cardiovasc Qual Outcomes* 2011;4:220-6.
- 399 22. Niles D, Sutton RM, Donoghue A, et al. "Rolling Refreshers": a novel approach to
400 maintain CPR psychomotor skill competence. *Resuscitation* 2009;80:909-12.
- 401 23. Issenberg SB, McGaghie WC, Hart IR, et al. Simulation technology for health care
402 professional skills training and assessment. *JAMA* 1999;282:861-6.
- 403 24. Kaye W, Rallis SF, Mancini ME, et al. The problem of poor retention of
404 cardiopulmonary resuscitation skills may lie with the instructor, not the learner or
405 the curriculum. *Resuscitation* 1991;21:67-87.
- 406 25. Moser DK, Coleman S. Recommendations for improving cardiopulmonary
407 resuscitation skills retention. *Heart Lung* 1992;21:372-80.
- 408 26. Anderson GS, Gaetz M, Statz C, Kin B. CPR skill retention of first aid attendants
409 within the workplace. *Prehosp Disaster Med* 2012;27:312-8.

- 410 27. Jeffrey D. Karpicke and Henry L. Roediger III. The critical importance of retrieval
411 for learning. *Science* 2008;319: 966-8.
- 412 28. Woollard M, Whitfield R, Newcombe RG, Colquhoun M, Vetter N, Chamberlain D.
413 Optimal refresher training intervals for AED and CPR skills: a randomised
414 controlled trial. *Resuscitation* 2006;71:237-47.
- 415 29. Woollard M, Whitfield R, Smith A, et al. Skill acquisition and retention in
416 automated external defibrillator (AED) use and CPR by lay responders: a
417 prospective study. *Resuscitation* 2004;60:17-28.
- 418 30. Bohn A, Van Aken HK, Möllhoff T, et al. Teaching resuscitation in schools: annual
419 tuition by trained teachers is effective starting at age 10. A four-year prospective
420 cohort study. *Resuscitation*. 2012;83:619-25.
- 421 31. de Vries W, Schelvis M, Rustemeijer I, Bierens JJ. Self-training in the use of
422 automated external defibrillators: the same results for less money. *Resuscitation*
423 2008;76:76-82.
- 424 32. de Vries W, Bierens JJ. Instructor retraining and poster retraining are equally
425 effective for the retention of BLS and AED skills of lifeguards. *Eur J Emerg Med*
426 2010;17:150-7.

- 427 33. Monsieus KG, De Regge M, Schelfout S, et al. Efficacy of a self-learning station
428 for basic life support refresher training in a hospital: a randomized controlled trial.
429 Eur J Emerg Med 2012;19:214-9.
- 430 34. Ahn JY, Cho GC, Shon YD, Park SM, Kang KH. Effect of a reminder video using
431 a mobile phone on the retention of CPR and AED skills in lay responders.
432 Resuscitation 2011;82:1543-7.
- 433 35. Gundry JW, Comess KA, DeRook FA, Jorgenson D, Bardy GH. Comparison of
434 naive sixth-grade children with trained professionals in the use of an automated
435 external defibrillator. Circulation 1999;100:1703-7.
- 436 36. Cummins RO, Schubach JA, Litwin PE, Hearne TR. Training lay persons to use
437 automatic external defibrillators: success of initial training and one-year retention of
438 skills. Am J Emerg Med 1989;7:143-9.
- 439 37. Callejas S, Barry A, Demertsidis E, Jorgenson D, Becker LB. Human factors impact
440 successful lay person automated external defibrillator use during simulated cardiac
441 arrest. Crit Care Med 2004;32:S406-13.
- 442 38. Beckers S, Fries M, Bickenbach J, Derwall M, Kuhlen R, Rossaint R. Minimal
443 instructions improve the performance of laypersons in the use of semiautomatic and
444 automatic external defibrillators. Crit Care 2005;9:R110-6.

445 39. Mosesso VN Jr, Shapiro AH, Stein K, Burkett K, Wang H. Effects of AED device
446 features on performance by untrained laypersons. Resuscitation 2009;80:1285-9.

447

448

449

450

451

452

453

454

455

456

457

458

459

460

461

462

463 **Legend to figure**

464 **Fig. 1.** Participants flow.

Fig. 1. Participant flow.




Table 1 Baseline characteristics of participants.

	Refresher training group (n=71)	Control group (n=69)	<i>p</i> -value
Men, n (%)	53 (74.6)	49 (71.0)	0.705
Age, yr, means \pm SD	37.3 \pm 13.8	38.4 \pm 14.4	0.650
Previous CPR training, n (%)	28 (39.4)	32 (46.4)	0.495
Experience of actual CPR, n (%)	1 (1.4)	5 (7.2)	0.113
Family history of sudden cardiac death, n (%)	5 (7.0)	8 (11.6)	0.396

CPR denotes cardiopulmonary resuscitation; SD, standard deviation.

Table 2 BLS performance one year after training.

	Refresher training group (n=57)	Control group (n=55)	<i>p</i> - value
Activation of EMS, n (%)*			
Call for help (119)	46 (83.6)	39 (75.0)	0.343
Call for AED	52 (94.5)	43 (82.7)	0.027
Chest compressions during 2 minutes test period*			
Total chest compressions, n, means \pm SD	182.0 \pm 41.7	142.0 \pm 59.1	<0.001
Appropriate chest compressions, n, means \pm SD	68.9 \pm 72.3	36.3 \pm 50.8	0.009
Chest compressions with appropriate depth, n, means \pm SD	121.9 \pm 79.7	87.7 \pm 71.9	0.025
Chest compressions with correct hand position, n, means \pm SD	101.7 \pm 80.3	65.9 \pm 60.8	0.006
Chest compressions with appropriate recoil, n, means \pm SD	179.7 \pm 41.6	140.9 \pm 58.1	<0.001
Proportion of appropriate chest compressions, %, means \pm SD [†]	38.3 \pm 37.6	27.5 \pm 34.4	0.009
Resuscitation time course, sec, means \pm SD*			
Time to chest compression	29.6 \pm 16.7	34.4 \pm 17.8	0.172
Time without chest compression during 2 min-test period	16.1 \pm 2.1	26.9 \pm 3.7	<0.001
AED operations			
Turn on the AED first, n (%)	36 (63.2)	36 (65.5)	0.845
Correct positioning of defibrillator pad, n (%)	53 (93.0)	49 (89.1)	0.524
Clear self and area, n (%)	49 (86.0)	41 (74.5)	0.209
Time to first defibrillation, sec, means \pm SD	86.5 \pm 15.1	93.5 \pm 27.6	0.103

CPR denotes cardiopulmonary resuscitation; EMS, emergency medical services; 119, emergency call number in Japan; AED, automated external defibrillator; SD, standard deviation.

P-values were derived by analysis of covariance adjusting for sex and age for continuous variables.

* Data are available for those with chest compressions (n=55 (96.5%) in the refresher training group; n=52 (94.5%) in the control group, p=0.676).

[†] Proportion of appropriate chest compressions over total number of chest compressions.