

THE AKE PRINCIPLE MEETS CLASSIFICATION THEORY

JUNGUK LEE

ABSTRACT. Using a type-amalgamation style criterion of NATP, we give a proof of the AKE principle for NATP in the case of equicharacteristic 0.

In [1], Ahn and Kim introduced a new dividing line, called the *antichain tree property* (ATP), and they provide a nice dichotomy for SOP_1 , that is, a complete theory T is SOP_1 if and only if it is TP_2 or ATP as like unstability is equal to IP or TP, and TP is equal to TP_1 or TP_2 . So, NATP is located in the map of the universe in Model Theory as follows:

$$\begin{array}{ccccccc}
 \text{DLO} & \longrightarrow & \text{NIP} & \longrightarrow & \text{NTP}_2 & \longrightarrow & \text{NATP} \\
 & & \uparrow & & \uparrow & & \uparrow \\
 & & \text{Stable} & \longrightarrow & \text{Simple} & \longrightarrow & \text{NSOP}_1 \longrightarrow \text{NSOP}_2 \longrightarrow \dots
 \end{array}$$

On the other hand, one of most fundamental principles in model theory of valued fields is the *Ax-Kochen-Ershov* principle (in short, the AKE principle), roughly saying that elementary properties of unramified henselian valued fields of characteristic 0 is determined by elementary properties of its residue field and value group. In [4, 8], Ax and Kochen, and independently Ershov showed that two henselian valued fields of equicharacteristic 0 are elementary equivalent if and only if their residue fields and value groups are elementary equivalent.

In [2], we studied several basic properties of NATP, for example, witness of NATP in a single variable and the equivalence ATP and k -ATP, and provided several algebraic examples of NATP theories. Most of all, we show that the Ax-Kochen-Ershov principle for NATP in the case of equicharacteristic 0. Namely, a given henselian valued field of

The author is supported by KAIST Advanced Institute for Science-X fellowship. This work was supported by the Research Institute for Mathematical Sciences, an International Joint Usage/Research Center located in Kyoto University.

equicharacteristic 0 is NATP if and only if its residue field is NATP. In this note, we aim to give a new proof of the AKE principle for NATP in the case of equicharacteristic 0, given in [2, Theorem 3.27], via a type-amalgamation criterion of NATP ([2, Theorem 3.27]).

Theorem 0.1. [7, Théorème 8][5, Theorem 7.6][2, Theorem 3.27] *Let $\mathcal{K} = (K, \Gamma, k)$ be a henselian valued field of equicharacteristic 0 in the Denef-Pas language, where K is a valued field, Γ is a value group, and k is a residue field.*

- (1) (Delon) $\text{Th}(\mathcal{K})$ is NIP if and only if $\text{Th}(k)$ is NIP.
- (2) (Chernikov) $\text{Th}(\mathcal{K})$ is NTP_2 if and only if $\text{Th}(k)$ is NTP_2 .
- (3) (Ahn-Kim-L.) $\text{Th}(\mathcal{K})$ is NATP if and only if $\text{Th}(k)$ is NATP.

Also, using similar methods, we can prove the AKE principle for NATP in the case of mixed characteristic in [10], and we can find several examples having NATP in [3], for example, a random parametrization of DLO is NATP, which is TP_2 and SOP.

1. PRELIMINARIES

1.1. Antichain tree property. We briefly recall several notions and facts around the antichain tree property from [2].

Notation 1.1. Let κ and λ be cardinals.

- (1) By κ^λ , we mean the set of all functions from λ to κ .
- (2) By $\kappa^{<\lambda}$, we mean $\bigcup_{i < \lambda} \kappa^i$ and call it a tree.

Let $\eta, \nu \in \kappa^{<\lambda}$.

- (3) By $\eta \trianglelefteq \nu$, we mean $\eta \subseteq \nu$. If $\eta \trianglelefteq \nu$ or $\nu \trianglelefteq \eta$, then we say η and ν are comparable.
- (4) By $\eta \perp \nu$, we mean that $\eta \not\trianglelefteq \nu$ and $\nu \not\trianglelefteq \eta$. We say η and ν are incomparable if $\eta \perp \nu$.
- (5) By $\eta \wedge \nu$, we mean the maximal $\xi \in \kappa^{<\lambda}$ such that $\xi \trianglelefteq \eta$ and $\xi \trianglelefteq \nu$.
- (6) By $l(\eta)$, we mean the domain of η .
- (7) By $\eta <_{\text{lex}} \nu$, we mean that either $\eta \trianglelefteq \nu$, or $\eta \perp \nu$ and $\eta(l(\eta \wedge \nu)) < \nu(l(\eta \wedge \nu))$.
- (8) By $\eta \widehat{\ } \nu$, we mean $\eta \cup \{(l(\eta) + i, \nu(i)) : i < l(\nu)\}$.
- (9) We say a subset X of $\kappa^{<\lambda}$ is an *antichain* if the elements of X are pairwise incomparable.

We use a language $\mathcal{L}_0 := \{\trianglelefteq, <_{\text{lex}}, \wedge\}$ for a tree with a natural interpretation in Notation 1.1. We say that an antichain A of $\kappa^{<\lambda}$ is *universal* if any finite antichain of $\kappa^{<\lambda}$ is embedded into A as \mathcal{L}_0 -structures.

Let T be a complete theory in a language \mathcal{L} , and let M be a monster model of T . For a tuple a of elements in M and a parameter A , we write $\text{tp}_{\mathcal{L}}(a/A)$ for the complete type of a over A and write $\text{qftp}_{\mathcal{L}}(a/A)$ for the quantifier free type of a over A . If there is no confusion, we omit the subscription \mathcal{L} . Let $(a_{\eta})_{\eta \in \kappa < \lambda}$ be a tree-indexed tuple of parameters from M . For $\bar{\eta} = (\eta_0, \dots, \eta_m)$, we denote $(a_{\eta_0}, \dots, a_{\eta_m})$ by $a_{\bar{\eta}}$. We say that $(a_{\eta})_{\eta \in \kappa < \lambda}$ is *strongly indiscernible* if for tuples $\bar{\eta}, \bar{\nu}$ of elements in $\kappa < \lambda$,

$$\text{qftp}_{\mathcal{L}_0}(\bar{\eta}) = \text{qftp}_{\mathcal{L}_0}(\bar{\nu}) \Rightarrow \text{tp}_{\mathcal{L}}(a_{\bar{\eta}}) = \text{tp}_{\mathcal{L}}(a_{\bar{\nu}}).$$

Definition 1.2. [2, Definition 1.1, Definition 3.19] Let $\varphi(x; y)$ be a \mathcal{L} -formula. For a positive integer $k \geq 2$, we say that $\varphi(x; y)$ has the *k-antichain tree property (k-ATP)* if there is a tree of parameters $(a_{\eta})_{\eta \in 2 < \omega}$ such that

- For any antichain $A \subset 2 < \omega$, $\{\varphi(x; a_{\eta}) : \eta \in A\}$ is consistent.
- For any comparable distinct elements $\eta_0, \eta_1 \in 2 < \omega$,

$$\{\varphi(x; a_{\eta_0}), \varphi(x; a_{\eta_1})\}$$

is inconsistent.

The *antichain tree property (ATP)* means the 2-antichain tree property. We say that T has *k-ATP* if there is a formula having *k-ATP*. We write *NATP* for non *ATP*.

Fact 1.3. [2, Theorem 3.17, Lemma 3.20]

- (1) *If there exists a witness of ATP, then there exists a witness of ATP in a single free variable.*
- (2) *T is ATP if and only if T has k-ATP for some $k \geq 2$. Moreover, if a formula $\varphi(x; y)$ has k-ATP for some $k \geq 2$, then a formula of the form*

$$\varphi(x; y_0) \wedge \dots \wedge \varphi(x; y_m)$$

has ATP for some $m \geq 1$.

We recall a type-amalgamation style criterion for *NATP*.

Fact 1.4. [2, Theorem 3.27] *Let κ and κ' be cardinals such that $2^{|T|} < \kappa < \kappa'$ with $\text{cf}(\kappa) = \kappa$. The following are equivalent.*

- (1) *T is NATP.*
- (2) *For any strongly indiscernible tree $(a_{\eta})_{\eta \in 2 < \kappa'}$ with $|a_{\emptyset}| \leq |T|$ and b with $|b| = 1$, there are some $\rho \in 2^{\kappa}$ and b' such that*
 - (a) *$(a_{\rho \cap 0^i})_{i < \kappa'}$ is indiscernible over b' , and*
 - (b) *$b \equiv_{a_{\rho}} b'$.*

- (3) For any strongly indiscernible tree $(a_\eta)_{\eta \in 2^{<\kappa'}}$ with $|a_\emptyset| \leq |T|$ and b with $|b| = 1$, there are some $\rho \in 2^\kappa$ such that for $p(x; a_\rho) = \text{tp}(b/a_\rho)$,

$$\bigcup_{i < \kappa'} p(x; a_{\rho \smallfrown 0^i})$$

is consistent.

We finish this subsection with a fact on the existence on a ‘homogeneous’ universal antichain.

Fact 1.5. [2, Corollary 3.23(b)] *Let $(a_\eta)_{\eta \in 2^{<\kappa'}}$ be a strongly indiscernible tree and b be a tuple of elements in M for some cardinals κ and κ' such that $2^{|T|+|a_\emptyset|+|b|} < \kappa < \kappa'$ and $\text{cf}(\kappa) = \kappa$. Then, there is a universal antichain $S \subset 2^\kappa$ such that*

$$a_\eta \equiv_b a_{\eta'}$$

for all $\eta, \eta' \in S$.

1.2. Denef-Pas language. Let $\mathcal{K} = (K, \Gamma, k, \nu : K \rightarrow \Gamma \cup \{\infty\}, \text{ac} : K \rightarrow k)$ be a henselian valued field of characteristic $(0, 0)$ in the **Denef-Pas language** $\mathcal{L}_{Pas} = \mathcal{L}_K \cup \mathcal{L}_{\Gamma, \infty} \cup \mathcal{L}_k \cup \{\nu, \text{ac}\}$ where $\nu : K \rightarrow \Gamma \cup \{\infty\}$ is a valuation map and $\text{ac} : K \rightarrow k$ is an angular component map.

We recall Delon’s description of types in a single variable on the valued field sort (c.f. [7, 11]). Let $\mathcal{M} = (M, \dots)$ be a monster model of $\text{Th}(\mathcal{K})$. Take $x \in M \setminus K$, and define

$$I_K(x) := \{\gamma \in \Gamma : \exists a \in K (\gamma = \nu(x - a))\}.$$

Then, there are three possibilities of $I_K(x)$:

- (Immediate) $I_K(x) = \{\nu(x - a) : a \in K\}$ and does not have a maximal element.
- (Residual) $I_K(x) = \{\nu(x - a) : a \in K\}$ and has a maximal element.
- (Valuational) $I_K(x) \neq \{\nu(x - a) : a \in K\}$.

Fact 1.6. [7, Théorème 5][11, Theorem 2.1.1] *There are three possibilities of $\text{tp}(x/K)$:*

- (1) (Immediate) *Let (a_ρ, γ_ρ) be a sequence indexed by a well-ordered set such that $a_\rho \in K$, $\gamma_\rho = \nu(x - a_\rho)$, and (γ_ρ) is cofinal in $I_K(x)$. Then,*

$$\{\nu(x - a_\rho) = \gamma_\rho\} \models \text{tp}(x/K).$$

- (2) (Residual) *There are $a \in K$ and $\gamma \in \Gamma$ such that $\nu(x - a) = \gamma$, $\text{ac}(x - a) \notin k$, and*

$$\{\nu(x - a) = \gamma\} \cup \text{tp}(\text{ac}(x - a)/k) \models \text{tp}(x/K).$$

- (3) (Valuational) There is $a \in K$ such that $\nu(x - a) \notin \Gamma$ and
 $\text{tp}(\nu(x - a)/\Gamma) \cup \text{tp}(\text{ac}(x - a)/k) \models \text{tp}(x/\mathcal{K})$.

2. PROOF OF THEOREM 0.1(3)

In this section, we aim to give a new proof of Theorem 0.1(3) using a type-amalgamation style criterion for NATP (Fact 1.4(3)) and a description of type in a single variable on the valued field sort (Fact 1.6).

Let $\mathcal{K} = (K, \Gamma, k, \nu : K \rightarrow \Gamma, \text{ac} : K \rightarrow k)$ be a henselian valued field of characteristic $(0, 0)$, which is saturated enough, in the Denef-Pas language. We use x, y, z, \dots for tuples of variables on K , $x^\Gamma, y^\Gamma, z^\Gamma, \dots$ for tuples of variables on Γ , and x^k, y^k, z^k, \dots for tuples of variables on k . We first recall the following fact.

Fact 2.1. [2, Lemma 4.27] *Suppose $\text{Th}(k)$ is NATP. Let*

$$\varphi(x; yy^\Gamma y^k) \equiv \chi(\nu(x - y); y^\Gamma) \wedge \rho(\text{ac}(x - y); y^k)$$

where $|x| = |y| = 1$, $\chi \in \mathcal{L}_{\Gamma, \infty}$ and $\rho \in \mathcal{L}_k$. Then, $\varphi(x; yy^\Gamma y^k)$ does not witness ATP. Moreover, by Fact 1.3(2), it does not witness k -ATP for all $k \geq 2$.

From now on, we assume that $\text{Th}(k)$ is NATP. By [9], $\text{Th}(\Gamma)$ is automatically NATP. Note that k and Γ are stably embedded (c.f. [6, Lemma 2.3]).

Let κ and κ' be cardinals such that $2^{\aleph_0} < \kappa < \kappa'$ with $cf(\kappa) = \kappa$. Take a strongly indiscernible tree $(a_\eta)_{\eta \in 2^{<\kappa'}}$ with $|a_\eta|$ countable and let b a tuple of element in \mathcal{K} of length 1. Without loss of generality, we may assume that each $a_\eta = (K_\eta, \Gamma_\eta, k_\eta)$ is a elementary substructure of \mathcal{K} , where K_η is a valued field, Γ_ρ is a value group, and k_ρ is a residue field. Suppose $b \in \Gamma$. Since Γ is stably embedded, $\text{tp}(b/a_\eta) = \text{tp}(b/\Gamma_\eta)$ for each η . Since $\text{Th}(\Gamma)$ is NATP, by Fact 1.4(3), there is $\rho \in 2^{<\kappa'}$ such that for $p(x; a_\rho) = \text{tp}(b/a_\rho)$,

$$\bigcup_{i < \kappa'} p(x; a_{\rho \smallfrown 0^i})$$

is consistent. By similar way, the same thing holds for the case that $b \in k$. So, we may assume that $b \in K$.

By Fact 1.5, there is a universal antichain $S \subset 2^\kappa$ such that $a_\eta \equiv_b a_{\eta'}$ for all $\eta, \eta' \in S$. Take $\rho \in S$ arbitrary. Put $p(x; a_\rho) = \text{tp}(b/a_\rho)$ and put $q(x) := \bigcup_{i < \kappa'} p(x; a_{\rho \smallfrown 0^i})$.

Claim 2.2. $q(x)$ is consistent.

Proof. . If $b \in K_\rho$, then by strong indiscernibility, $b \in K_\eta$ for all $\eta \in S$ and so $b \models q$. Now, we assume that $b \notin K_\rho$. Suppose q is not consistent. Then, by compactness, strong indiscernibility, and Fact 1.6, there are

- a formula of the form:

$$\varphi(x; yy^\Gamma y^k) \equiv \chi(\nu(x - y); y^\Gamma) \wedge \rho(\text{ac}(x - y); y^k),$$

- for a positive integer n and for each $i < n$,

$$c_i \in a_{\rho \smallfrown 0^i}, c_i^\Gamma \subset \Gamma_{\rho \smallfrown 0^i}, c_i^k \subset k_{\rho \smallfrown 0^i}$$

such that

$$\bigwedge_{i < n} \varphi(x; c_i c_i^\Gamma c_i^k)$$

is inconsistent. Since $b \models \bigcup_{\eta \in S} p(x; a_\eta)$ and S is a universal antichain, the formula $\varphi(x; yy^\Gamma y^k)$ witness n -ATP, a contradiction to Fact 2.1. Thus, $q(x)$ is consistent. \square

Now, by Fact 1.4, $\text{Th}(\mathcal{K})$ is NATP.

REFERENCES

- [1] J. Ahn and J. Kim, SOP₁, SOP₂, and antichain tree property, submitted.
- [2] J. Ahn, J. Kim, and J. Lee, On the antichain tree property, accepted in J. Math. Log.
- [3] J. Ahn, J. Kim, H. Lee, and J. Lee, Preservation of NATP, in preparation.
- [4] J. Ax and S. Kochen, Diophantine problems over local fields. III. Decidable fields, *Ann. of Math.*, **83** (1966), 437-456.
- [5] A. Chernikov, Theories without the tree property of the second kind, *Ann. Pure. Appl. Logic*, **165** (2014), 695-723.
- [6] A. Chernikov and M. Hils, Valued difference fields and NTP₂, *Israel J. Math.*, **204** (2014), 299-327.
- [7] F. Delon, Types sur $\mathbb{C}((X))$, Study Group on Stable Theories (Bruno Poizat), Second year: 1978/79, Exp. No. 5, 29 pp., Secrétariat Math., Paris, 1981.
- [8] Y. L. Ershov, On the elementary theory of maximal normed fields, *Dokl. Akad. Nauk SSSR*, **165** (1965), 21-23.
- [9] Y. Gurevich and P.H. Scmitt, The theory of ordered abelian groups does not have the independence property, *Trans. Amer. Math. Soc.*, **284** (1984), 171-182.
- [10] J. Lee, The AKE principle meets NATP, in preparation.
- [11] P. Sinclair, Relationships between Model Theory and Valuations of Fields, PhD thesis, McMaster University, 2018.

KAIST, DEPARTMENT OF MATHEMATICAL SCIENCES, 291 DAEHAK-RO YUSEONG-GU, 34141, DAEJEON, SOUTH KOREA

Email address: ljwhayo@kaist.ac.kr