

# Mathematical reflections on locality

Sylvie Paycha  
University of Potsdam  
joint work with Li Guo and Bin Zhang

Women in Mathematics, RIMS, Kyoto, 8th September  
2022

## I. The principle of locality revisited

The principle of **locality** states that an object is influenced directly only by its immediate surroundings.

Thus, one can **separate** events located in different regions of space-time and should be able to measure them **independently**.


### Our aim

- Propose a **mathematical framework** which encompasses the main features of the **locality principle** in QFT;
- use this framework to carry out **renormalisation** (**evaluate meromorphic germs at their poles**) in accordance with the **locality principle**.

## Causal separation

### Light cone, past and future

In the **Minkowski** space  $(\mathbb{R}^d, g)$ , where  $g(x, y) = -x_0y_0 + \sum_{j=1}^{d-1} x_jy_j$  is the **Lorentzian** scalar product, there is a notion of "**past**" and "**future**":



(picture downloaded from Wikipedia)

Two sets  $S_1$  and  $S_2$  are **causally separated** ( $S_1 \parallel S_2$ ) if and only if  $S_i$  **does not lie in the future** of  $S_j$  for  $i \neq j$ .

## Locality in axiomatic QFT

The Wightman field  $\varphi : \mathcal{S}(\mathbb{R}^d) \rightarrow \mathcal{O}(H)$  obeys the **locality** axiom

$$\text{Supp}(f_1) \parallel \text{Supp}(f_2) \implies [\varphi(f_1), \varphi(f_2)] = 0. \quad (1)$$

The (relative) **scattering matrix**  $S_f$  satisfies the **locality** condition

$$\begin{aligned} \text{Supp}(f_1) \parallel \text{Supp}(f_2) &\implies S_f(f_1 + f_2) = S_f(f_1) S_f(f_2) \\ &\implies [S_f(f_1), S_f(f_2)] = 0. \end{aligned} \quad (2)$$

## Mathematical interpretation

We introduce two **binary relations**

- on sets:

$$O_1 \top O_2 \Leftrightarrow [O_1, O_2] = 0, \quad (3)$$

- on test functions:

$$f_1 \top f_2 \Leftrightarrow \text{Supp}(f_1) \parallel \text{Supp}(f_2). \quad (4)$$

Interpretation of (1) as a **locality map** (see later)

$$f_1 \top f_2 \implies \varphi(f_1) \top \varphi(f_2). \quad (5)$$

Interpretation of (2) as a **locality morphism** (see later)

$$f_1 \top f_2 \implies S_f(f_1 + f_2) = S_f(f_1) S_f(f_2). \quad (6)$$

## II. Locality as a symmetric binary relation

### Algebraic locality

#### Definition of locality

A **locality set** is a couple  $(X, \top)$  where  $X$  is a set and  $\top \subseteq X \times X$  is a **symmetric relation** on  $X$ , called **locality relation** (or **independence relation**) of the locality set.

$$x_1 \top x_2 \iff (x_1, x_2) \in \top, \quad \forall x_1, x_2 \in X.$$

#### First examples of locality

- $X \top Y \iff X \cap Y = \emptyset$  on subsets  $X, Y$  of a set  $Z$ .
- $X \top Y \iff X \perp Y$  on subsets  $X, Y$  of an euclidean vector space  $V$ .

#### (almost-)Separation of supports

Let  $U \subset \mathbb{R}^n$  be an open subset and  $\epsilon \geq 0$ . Two functions  $\phi, \psi \in \mathcal{D}(U)$  are **independent** i.e.,  $\phi \top \psi$  whenever  $d(\text{Supp}(\phi), \text{Supp}(\psi)) > \epsilon$ .

For  $\epsilon = 0$ , this amounts to **disjointness of supports**, otherwise to  **$\epsilon$ -separation of supports**.

## Further examples

### Probability theory: independence of events

Given a probability space  $\mathcal{P} := (\Omega, \Sigma, P)$  and two events  $A, B \in \Sigma$ :

$$A \top B \iff P(A \cap B) = P(A)P(B).$$

### Geometry: transversal manifolds

Given two submanifolds  $L_1$  and  $L_2$  of a manifold  $M$ :

$$L_1 \top L_2 \iff L_1 \pitchfork L_2 \iff T_x L_1 + T_x L_2 = T_x M \quad \forall x \in L_1 \cap L_2.$$

### Number theory: coprime numbers

Given two positive integers  $m, n$  in  $\mathbb{N}$ :

$$m \top n \iff m \wedge n = 1.$$

## Partial products

- **Locality set:**  $(X, \top)$ ,
- **Polar set:**  $U^\top := \{x \in X, x \top u \quad \forall u \in U\}$  for  $U \subseteq X$ ;
- **Graph** of the **locality** relation:  $\top = \{(x_1, x_2) \in X^2, x_1 \top x_2\}$ ;
- **Partial product:**  $m_X : X \times X \supset \top \rightarrow X$  i.e.  $m_X(\top) \subset X$ .

### $(X, m_X, \top)$ locality semi-group

**semi-group** condition:  $\forall U \subseteq X, m_X((U^\top \times U^\top) \cap \top) \subseteq U^\top$   
or equivalently

$$(x_1 \top u_1 \text{ and } x_2 \top u_2 \quad \forall u_1, u_2 \in U) \implies (m_X(x_1, x_2) \top w \quad \forall w \in U).$$

### Counterexample

Equip  $\mathbb{R}$  with the **locality** relation  $x \top y \iff x + y \notin \mathbb{Z}$ .

$(\mathbb{R}, \top, +)$  is **NOT** a **locality semi-group**: for  $U = \{1/3\}$  we have  $(1/3, 1/3) \in (U^\top \times U^\top) \cap \top$  but  $1/3 + 1/3 = 2/3 \notin U^\top$

## Locality category

### Locality structures

- set  $X \rightsquigarrow$  **locality set**  $(X, \mathbb{T})$ ;
- semi-group  $(X, m_X) \rightsquigarrow$  **locality semi-group**  $(X, m_X, \mathbb{T})$ ;
- vector space  $(V, +, \cdot) \rightsquigarrow$  **locality vector space**  $(V, +, \cdot, \mathbb{T})$   
( $U \subset V \implies U^\mathbb{T}$  **vector space**);
- algebra  $(A, +, \cdot, m_A) \rightsquigarrow$  **locality algebra**  $(A, +, \cdot, m_A, \mathbb{T})$ .

### Locality morphisms: $f : (X, \mathbb{T}_X) \rightarrow (Y, \mathbb{T}_Y)$

- **locality map**:  
 $(f \times f)(\mathbb{T}_X) \subset \mathbb{T}_Y$  or equivalently  $x_1 \mathbb{T}_X x_2 \implies f(x_1) \mathbb{T}_Y f(x_2)$ ;
- **locality semi-group morphism**  $f : (X, m_X, \mathbb{T}_X) \rightarrow (Y, m_Y, \mathbb{T}_Y)$ :  
 $f$  is a **locality map** such that  
 $x_1 \mathbb{T}_X x_2 \implies f(m_X(x_1, x_2)) = m_Y(f(x_1), f(x_2))$ .  
.....

## III. Evaluating meromorphic germs at poles in QFT

## Functions of several variables in QFT

### Speer's analytic renormalisation [JMP 1967] revisited

Eugene Speer considers **Feynman amplitudes** given by the coefficients of the **perturbation-series expansion** of the  $S$  matrix in a Lagrangian field theory (with non zero mass).

### Excerpt of Speer's article

*In this paper we apply a method of defining **divergent quantities** which was originated by Riesz and has been used in various contexts by many authors. [...] We find it necessary to consider functions of **several complex variables**  $z_1, \dots, z_k$ , one associated with **each line** of the Feynman graph. The main difficulty is the extension of the above [Riesz's] treatment of poles to the **more complicated singularities** which occur in **several complex variables**...*

## Brain teaser

(We assume the poles are at zero)

Speer shows [Theorem 1] that the divergent expressions lie in the **filtered algebra**  $\mathcal{M}^{\text{Feyn}}(\mathbb{C}^\infty) := \bigcup_{k=1}^\infty \mathcal{M}^{\text{Feyn}}(\mathbb{C}^k)$  consisting of **Feynman functions**  $f : \mathbb{C}^k \rightarrow \mathbb{C}$ ,

$$f = \frac{h(z_1, \dots, z_k)}{L_1^{s_1} \dots L_m^{s_m}}, \quad L_i = \sum_{j \in J_i} z_j, \quad J_i \subset \{1, \dots, k\}, \quad h \text{ holom. at zero}$$

### Questions:

- ① How to **evaluate**  $f$  consistently at the **poles**  $z_1 = \dots = z_k = 0$ ?
- ② What freedom of choice do we have for the **evaluator**?

### Evaluating a fraction with a linear pole at zero

$$f(z_1, z_2) = \frac{z_1 - z_2}{z_1 + z_2} \Big|_{z_1=0, z_2=0} = \begin{cases} 1? \\ 0? \\ 10000? \end{cases}$$

## Speer's generalised evaluators

They consist of a family  $\mathcal{E} = \{\mathcal{E}_k, k \in \mathbb{N}\}$  of linear forms  $\mathcal{E}_k : \mathcal{M}^{\text{Feyn}}(\mathbb{C}^k) \rightarrow \mathbb{C}$ , compatible with the filtration, which fulfill the following conditions

- 1 (extend  $\text{ev}_0$ )  $\mathcal{E}$  is the ordinary evaluation  $\text{ev}_0$  at zero on holom. germs;
- 2 (partial multiplicativity)  $\mathcal{E}(f_1 \cdot f_2) = \mathcal{E}(f_1) \cdot \mathcal{E}(f_2)$  if  $f_1$  and  $f_2$  depend on different sets (later called independent) of variables  $z_i$ ;
- 3  $\mathcal{E}$  is invariant under permutations of the variables  $\mathcal{E}_k \circ \sigma^* = \mathcal{E}_k$  for any  $\sigma \in \Sigma_k$ , with  $\sigma^* f(z_1, \dots, z_k) := f(z_{\sigma(1)}, \dots, z_{\sigma(k)})$ ;
- 4 (continuity) If  $f_n(\vec{z}_k) \cdot L_1^{s_1} \dots L_m^{s_m} \xrightarrow[n \rightarrow \infty]{\text{uniformly}} g(\vec{z}_k)$  as holomorphic germs, then  $\mathcal{E}_k(f_n) \xrightarrow[n \rightarrow \infty]{} \mathcal{E}_k(\lim_{n \rightarrow \infty} f_n)$ .

**Drawback:** Speer's approach depends on the choice of coordinates  $z_1, \dots, z_k, \dots$ .

### IV. Locality on meromorphic germs comes to the rescue



## Back to the **locality** principle in QFT

We consider  $\mathcal{M} := \mathcal{M}(\mathbb{C}^\infty) := \bigcup_{k=1}^{\infty} \mathcal{M}(\mathbb{C}^k)$  consisting of **meromorphic** functions/germs  $f : \mathbb{C}^k \rightarrow \mathbb{C}$  with **linear poles** at zero,

$$f = \frac{h(z_1, \dots, z_k)}{L_1^{s_1} \dots L_m^{s_m}}, \quad L_i \text{ linear in } z_1, \dots, z_k, \quad h \text{ holom. at zero}$$

**Aim:** evaluate meromorphic germs at poles according to the **principle of locality**: "two events separated in space can be **measured independently**"

**Principle of locality:** factorisation on **independent** events

$$\underbrace{a \text{ and } b}_{\in \mathcal{A}} \text{ independent} \quad \xRightarrow{\text{factorisation}} \quad \text{Meas} \quad \underbrace{(a \vee b)}_{\text{concatenation}} = \text{Meas}(a) \cdot \text{Meas}(b).$$

- We shall later equip  $\mathcal{M}$  with a **locality** relation  $\top$ ;

**Principle of locality revisited:** **locality evaluators**

$f \top g \implies \mathcal{E}(f \cdot g) = \mathcal{E}(f) \mathcal{E}(g)$  for two **meromorphic germs**  $f$  and  $g$  in an appropriate subalgebra  $\mathcal{M}^\bullet$  of  $\mathcal{M}$ .

## Locality on/independence of meromorphic germs

**Meromorphic germs with linear poles**

- $\mathcal{M}(\mathbb{C}^k) \ni f = \frac{h(\ell_1, \dots, \ell_m)}{L_1^{s_1} \dots L_m^{s_m}}$ ,  $h$  holomorphic germ,  $s_i \in \mathbb{Z}_{\geq 0}$ ,
- $\ell_j : \mathbb{C}^k \rightarrow \mathbb{C}$ ,  $L_j : \mathbb{C}^k \rightarrow \mathbb{C}$  **linear forms** with real coefficients (lie in  $\mathcal{L}(\mathbb{C}^k)$ ).

**Locality on meromorphic germs:** orthogonality

- **Dependence** set  $\text{Dep}(f) := \langle \ell_1, \dots, \ell_m, L_1, \dots, L_n \rangle$ .
- $Q$  inner product on  $\mathbb{R}^k$  induces one on  $\mathcal{L}(\mathbb{C}^k)$
- $f_1 \perp^Q f_2 \iff \text{Dep}(f_1) \perp^Q \text{Dep}(f_2)$ .
- **polar germs:**  $\mathcal{M}_-^{\bullet Q}(\mathbb{C}^k) \ni f \leftarrow h \perp^Q L_i$  for all  $i = 1, \dots, n$ .
- **Theorem:** (L. Guo, S.-P., B. Zhang/ N. Berline, M. Vergne 2015)  
 $\mathcal{M}^\bullet(\mathbb{C}^k) = \mathcal{M}_+(\mathbb{C}^k) \oplus^Q \mathcal{M}_-^{\bullet Q}(\mathbb{C}^k)$

## Where we stand

### Data

- $(\mathcal{M}^\bullet, \perp^Q)$  an (locality) algebra of meromorphic germs at zero with a prescribed type of poles (e.g. Chen  $\subset$  Speer  $\subset$  Feynman);
- $\mathcal{M}_+ \subset \mathcal{M}^\bullet$  the algebra of holomorphic germs at zero;
- the evaluation at zero:  $ev_0 : \mathcal{M}_+ \rightarrow \mathbb{C}$ ;
- the Galois group  $Gal^Q(\mathcal{M}^\bullet/\mathcal{M}_+)$  of (locality) isomorphisms of  $(\mathcal{M}^\bullet, \perp^Q)$ ;
- $\mathcal{M}_-^Q$  is generated by polar germs  $f = \frac{h}{g}$  with  $h \perp^Q g$ .

### Orthogonal projection

$\perp^Q$  induces a splitting

$$\mathcal{M}^\bullet = \mathcal{M}_+ \oplus^Q \mathcal{M}_-^Q \quad \text{and} \quad \pi_+^Q : \mathcal{M}^\bullet \rightarrow \mathcal{M}_+$$

## Theorem [Guo, S.P., Zhang 2022]

### Definition

A locality evaluator at zero  $\mathcal{E} : \mathcal{M}^\bullet \rightarrow \mathbb{C}$  is a linear form which i) extends the ordinary evaluation  $ev_0$  at zero and ii) factorises on independent germs (or is a locality character):

$$f_1 \perp^Q f_2 \implies \mathcal{E}(f_1) \perp^Q \mathcal{E}(f_2).$$




Example: Minimal subtraction scheme:






$$\mathcal{E}^{MS} : \mathcal{M}^\bullet \xrightarrow{\pi_+^Q} \mathcal{M}_+ \xrightarrow{ev_0} \mathbb{C} \text{ is a locality evaluator.}$$

### Theorem

Given an inner product  $Q$ , a locality evaluator at zero  $\mathcal{E} : \mathcal{M}^\bullet \rightarrow \mathbb{C}$  is of the form:  $\mathcal{E} = \underbrace{ev_0 \circ \pi_+^Q}_{\mathcal{E}^{MS}} \circ \underbrace{T_{\mathcal{E}}}_{Gal^Q(\mathcal{M}^\bullet/\mathcal{M}_+)}$ .

THANK YOU FOR YOUR ATTENTION!

-  P. Clavier, L. Guo, B. Zhang and S. P., An algebraic formulation of the locality principle in renormalisation, *European Journal of Mathematics*, Volume 5 (2019) 356-394
-  P. Clavier, L. Guo, B. Zhang and S. P., Renormalisation via locality morphisms, *Revista Colombiana de Matemáticas*, Volume 53 (2019) 113-141
-  P. Clavier, L. Guo, B. Zhang and S. P., Renormalisation and locality: branched zeta values, in "Algebraic Combinatorics, Resurgence, Moulds and Applications (Carma)" Vol. 2 ,Eds. F. Chapoton, F. Fauvet, C. Malvenuto, J.-Y. Thibon, *Irma Lectures in Mathematics and Theoretical Physics* **32**, *European Math. Soc.* (2020) 85–132
-  P. Clavier, L. Guo, B. Zhang and S. P., Locality and renormalisation: universal properties and integrals on trees, *Journal of Mathematical Physics* **61**, 022301 (2020)

-  L. Guo, B. Zhang and S. P., Renormalisation and the Euler-Maclaurin formula on cones, *Duke Math J.*, **166** (3) (2017) 537–571.
-  L. Guo, B. Zhang and S. P., A conical approach to Laurent expansions for multivariate meromorphic germs with linear poles, *Pacific Journal of Mathematics* 307 (2020) 159–196.
-  L. Guo, B. Zhang and S. P., Galois groups of meromorphic germs and multiparameter renormalisation (2022) (Preprint)
-  L. Guo, B. Zhang and S. P., Mathematical reflections on locality (2022) (Preprint)
-  R. Dahmen, A. Schmeding and S. P., A topological splitting of the space of meromorphic germs in several variables and continuous evaluators, arXiv:2206.13993 (2022)