

A New Species of *Rhacophorus* from Eastern Thailand (Anura: Rhacophoridae)

Masafumi Matsui^{1*} and Somsak Panha²

¹Graduate School of Human and Environmental Studies, Kyoto University, Sakyo-ku,
Kyoto 606-8501, Japan

²Department of Biology, Faculty of Science, Chulalongkorn University,
Bangkok 10330, Thailand

A new tree frog of the genus *Rhacophorus* is described on the basis of specimens collected from Kalasin and Roi Et Provinces, eastern Thailand. It can be distinguished from all other congeners by the combination of: moderate body size (about 38 mm in males and 44 mm in females); brownish dorsum with irregular, dark marking; third finger webbed broadly to base of disk or less on outer side, and fourth finger to distal subarticular tubercle or base of disk; and no dermal appendage on snout, vent, or heel. It is currently known only around the type locality, where habitats suitable for anurans are limited.

Key words: biogeography, classification, *Polypedates*, Southeast Asia, taxonomy

INTRODUCTION

The tree frog genus *Rhacophorus* Kuhl and Van Hasselt, 1822 (*sensu lato*) occurs in tropical to temperate regions of the Old World (Duellman, 1999). Liem (1970) split larger-sized rhacophorid members into two genera, *Rhacophorus* and *Polypedates* Tschudi, 1838, on the basis of a phenetic analysis of morphological characters; however, more than three decades after Liem's (1970) revision, classification of the Old World treefrogs is still unstable (cf. Wilkinson *et al.*, 2002; Matsui and Orlov, 2004). Some authors combine the two genera and admit only *Rhacophorus* (e.g., Tian and Jiang, 1986; Dubois, 1987; Fei *et al.*, 2004), while others recognize the validity of two genera (e.g., Frost, 1985; Jiang *et al.*, 1987; Zhao and Adler, 1993; Inger *et al.*, 1999; Maeda and Matsui, 1999; Malkmus *et al.*, 2002). This conflict mainly derives from ambiguous morphological and ecological distinctions between the two genera, due mainly to inadequate species sampling for comparisons, and some Chinese *Polypedates* species with a green dorsum require taxonomic reassessment (Matsui and Wu, 1994). Even so, Liem (1970) could differentiate the two lineages by some osteological characters in the specimens examined.

Setting these taxonomic problems aside, the genus *Rhacophorus* (*sensu stricto*) includes about 60 species (Frost, 2004), of which quite a few have been added recently from regions of Southeast Asia already well surveyed (e.g., Manthey and Steirof, 1998; Inger *et al.*, 1999; Matsui, 2000; Harvey *et al.*, 2002). This situation holds for Thailand, where recent extensive field surveys have resulted in the discovery of new taxa or separation of cryptic taxa in frogs of different lineages (Matsui *et al.*, 1996, 1998,

1999, 2005). During our survey of eastern Thailand in the autumn of 1996, we collected several specimens of a *Rhacophorus* distinctly different from its congeners hitherto known from Thailand (Taylor, 1962). These specimens, which resemble some Bornean or Sumatran members of this genus, but are easily distinguished morphologically from them, are described below as a new species.

MATERIALS AND METHODS

We conducted fieldwork in the Phu Sri Tan Wildlife Sanctuary, Kalasin Province, and Phu Pha Namtip Non-hunting Area, Roi Et Province, eastern Thailand, from 24–27 October 1996. Specimens were fixed in 10% formalin and later preserved in 70% ethanol. We took 18 body measurements (Table 1) to the nearest 0.1 mm with dial calipers under a binocular dissecting microscope: snout-vent length (SVL); head length (HL), from tip of snout to hind border of angle of jaw (not measured parallel with the median line); snout length (SL); eye length (EL); tympanum-eye distance (T-EL); tympanum diameter (TD); head width (HW); internarial distance (IND); interorbital distance (IOD); upper eyelid width (UEW); forelimb length (FLL); third finger disk width (3FDW); hindlimb length (HLL); tibia length (TL); foot length (FL) from proximal end of inner metatarsal tubercle to tip of fourth toe; fourth toe disk width (4TDW); first toe length (1TL) from distal end of inner metatarsal tubercle to tip of first toe; inner metatarsal tubercle length (IMTL). These measurements mainly followed the definitions by Matsui (1984). We also prepared radiographs to examine gross osteology. Finally, we examined museum specimens, including some types, of many described taxa of the genus *Rhacophorus* (*sensu stricto*), for comparisons (see Appendix 1 in Matsui and Orlov, 2004).

SYSTEMATICS

***Rhacophorus jarujini* sp. nov.**
(Figs. 1, 2)

Diagnosis. The new species is assigned to *Rhacophorus* (*sensu stricto*) by: vertebrae procoelous; frontoparietal lacking parieto-squamosal arch; distal end of terminal phalanx Y-shaped. It is distinguished from all other congeners

* Corresponding author. Phone: +81-75-753-6846;
Fax : +81-75-753-6846;
E-mail: fumi@zoo.zool.kyoto-u.ac.jp

Table 1. Measurements of 18 characters in *Rhacophorus jarujini*. SVL (means \pm SD, in mm) and medians of ratios (R) of other characters to SVL, followed by ranges in parenthesis. See text for character abbreviations.

		SVL	RHL	RSL	REL	RT-EL	RTD
Males	(N=4)	37.6 \pm 2.9 (33.7–40.0)	37.2 (36.4–38.6)	17.0 (15.6–17.2)	14.2 (13.5–15.7)	1.5 (1.2–1.9)	7.9 (7.5–8.8)
Females	(N=5)	43.6 \pm 1.6 (41.5–46.1)	36.2 (34.1–37.0)	16.1 (15.0–17.1)	14 (12.5–15.2)	1.6 (1.0–2.3)	7.8 (6.3–8.1)
		RHW	RIND	RIOD	RUEW	RFLL	R3FDW
Males	(N=4)	37.2 (36.8–39.8)	10.1 (9.8–10.5)	11.9 (10.5–12.8)	9.3 (8.3–10.8)	67.0 (63.2–70.3)	7.3 (6.6–7.7)
Females	(N=5)	37.1 (34.5–37.6)	8.9 (8.3–10.3)	12.4 (10.8–12.9)	8.7 (8.3–9.9)	68.0 (56.0–69.7)	6.9 (6.5–6.9)
		R4TDW	RHLL	RTL	RFL	R1TL	RITML
Males	(N=4)	4.9 (4.3–5.9)	163.4 (154.5–172.7)	52.8 (49.0–56.7)	44.6 (41.7–47.8)	12.8 (11.0–14.5)	4.4 (3.5–4.5)
Females	(N=5)	5.0 (4.6–5.5)	167.8 (147.3–178.9)	54.4 (46.6–57.2)	47.9 (40.1–49.0)	13.7 (10.4–14.2)	4.6 (4.1–5.1)




Fig. 1. Male paratype of *Rhacophorus jarujini* (KUHE 22287, SVL=33.7 mm); (A) dorsal and (B) ventral views. Scale bar indicates 20 mm.

by the combination of (1) moderate size, with females from 41.5–46.1 mm and males from 33.7–40.0 mm in SVL, (2) dorsum brownish rather than greenish, with irregular dark marking, (3) third finger webbed broadly to base of disk or




Fig. 2. Ventral views of (A) hand and (B) foot of male holotype (CUZM (A) 5251). Scale bar indicates 5 mm.

less on outer side, and fourth finger to distal subarticular tubercle or base of disk, and (4) no dermal appendage on snout, vent, or heel.

Etymology. The specific name is dedicated to Mr. Jarujin Nabhitabhata of National Science Museum of Thailand, who is actively elucidating faunal diversity in Thailand.

Holotype. CUZM (Chulalongkorn University, Zoological Museum) (A) 5251, a male from Phu Sri Tan Wildlife Sanctuary (104°10' E, 16°30' N, 500 m a.s.l.), Kalasin Province, Thailand. Collected on 26 October 1996 by M. Matsui.

Paratypes. CUZM (A) 5252–5253, KUHE (Kyoto University, Human and Environmental Studies), 23038, three females; KUHE 23037, a male; and CUZM (A) 5254, a juvenile; all the same data as for holotype. KUHE 22280, 23045, two females, and KUHE 22287, 22288, two males, from Phu Pha Namtip Non-hunting area, Roi Et Province, collected on 25 October 1996 by M. Matsui.

Description of holotype (measurements in mm). SVL 39.6; head about as long (14.4) as broad (14.6); snout (6.8) longer than eye (5.5), rounded dorsally and angular in profile, slightly pointed at tip and projecting slightly over lower jaw; canthus blunt; lores slightly oblique, slightly concave; nostril nearer to tip of snout than to eye; internarial distance

(4.0) narrower than interorbital (4.8); latter wider than eyelid (3.8); eye diameter (5.4) larger than eye-nostril (4.2); tympanum distinct, length (3.5) more than three-fifths eye diameter and separated from eye by one-seventh the tympanum diameter (0.5); vomerine teeth in short, nearly horizontal groups beginning near anterior corners of choanae, groups separated by more than the length of one group; a longitudinal opening into median subgular vocal sac on both sides of mouth floor.

First finger shorter than second, length of first (3.9, measured from distal edge of inner palmar tubercle) shorter than diameter of eye; fourth finger longer than second; tips of fingers dilated into large disks, that of third finger (2.6) narrower than tympanum; broad web (Fig. 2A) reaching base of subarticular tubercle of first finger, proximal margin of tubercle on inner edge of second finger, and to bases of disk on outer edge of second, proximal margin of distal tubercle on inner edge of third, distal margin of distal tubercle on outer edge of third, and distal margin of distal tubercle on fourth finger; weak supernumerary tubercles on metacarpals; a distinct inner and a round, indistinct outer palmar tubercle.

Hindlimb (61.2) about 2.3 times length of forelimb (26.1); tibia not long (19.4), heels slightly overlapping when limbs are held at right angles to body; tibiotarsal articulation of adpressed limb reaching anterior rim of eye; foot (17.0) shorter than tibia; tips of toes expanded into disks smaller than those of fingers (disk diameter of fourth toe 1.7); all toes webbed to disks (Fig. 2B); subarticular tubercles indistinct; a small inner metatarsal tubercle, length (1.4) less than one-third the length of first toe (4.8), but no outer metatarsal tubercle.

Dorsum smooth, free of skull; an oblique fold from eye above tympanum, ending at above arm insertion; skin of throat smooth, abdomen coarsely granular; a weak fringe of skin from outer edge of fourth finger to elbow; hindlimb smooth, except for very narrow fringes of skin along inner edge of first toe and outer edge of fifth; no dermal appendages or large tubercles at vent and heel; nuptial pad absent.

Color in life. Dorsum light brown; a narrow darker brown bar crossing upper eyelids; back with irregular, darker brown crossbands; sides of face same shade of brown as dorsal surface of head; lips unpatterned; abdomen without markings, anteriorly cream and posteriorly yellow; ventral surfaces of thigh, hand, and foot light orange; iris yellow; a white line on supracloacal ridge; a dark brown marking around vent; limbs with dark crossbars, three bands crossing shank and two crossing antebrachium; posterior thigh orange-brown, without any definable pattern; webbing reddish-orange with a light brown streak.

Osteology. Vertebral column procoelous; frontoparietal lacking parieto-squamosal arch; distal end of the terminal phalanx Y-shaped.

Variation. The average snout-vent length of five females ($\text{mean} \pm \text{SD} = 43.6 \pm 1.6$ mm) is significantly larger than that of four males (37.6 ± 2.9 mm; Student's t-test, $p < 0.01$), but no sexual difference was found in other measurements relative to SVL (Mann-Whitney U test, $p > 0.05$), possibly due to small sample size. A juvenile had a SVL of 26.8 mm. Length of hindlimb varies slightly, and tibiotarsal articulation of adpressed limb reaches middle of eye in one, anterior rim of

eye in four, between eye and nostril in three, and nostril in one individuals. Hand webbing is slightly better developed in some individuals than in the holotype, with broad web reaching base of disk on outer edge of third, and on inner edge of fourth finger. Individuals are nearly uniform in coloration and pattern of dorsal marking.

Range. Known only from the type locality, Phu Sri Tan Wildlife Sanctuary, Kalasin Province, and Phu Pha Namtip Non-hunting Area, Roi Et Province, eastern Thailand.

Natural history. In Phu Sri Tan, *R. jarujini* was found at night on a dried, rocky streambed (width <5 m), and it jumped into small pools when alerted. In Phu Pha Namtip, it was found at night perching on the leaves of trees (<2 m) along the bank of a small pond. No tadpoles or eggs were found in the pond, and calling males were absent in late October. All females had small ovaries, indicating a post-breeding condition. Frogs found associated with *R. jarujini* included *Occidozyga lima* (Gravenhorst, 1829), *O. martensi* (Peters, 1867), *Rana nigrovittata* (Blyth, 1856), *Fejervarya limnocharis* (Gravenhorst, 1829), *Limnonectes gyldenstolpei* (Andersson, 1916), *Polypedates leucomystax* (Gravenhorst, 1829), *Microhyla fissipes* Boulenger, 1884, *M. berdmorei* (Blyth, 1856), *M. heymonsi* Vogt, 1911, and *M. pulchra* (Hallowell, 1861).

Comparisons. Although the distinction between *Rhacophorus* and *Polypedates* is under debate, as noted above, *R. jarujini* n. sp. is assigned on the basis of osteology to *Rhacophorus* (having procoelous vertebral column and lacking parieto-squamosal arch of the frontoparietal) rather than to *Polypedates* (having diplasicoelous vertebral column and mostly having parieto-squamosal arch; Liem, 1970). Therefore, we compare with the new species only species listed as *Rhacophorus* by Frost (2004).

The dorsal ground color of *R. jarujini* is brown, which differentiates it from the following species that have a dorsum with the ground color green: *R. arboreus* (Okada and Kawano, 1924) and *R. schlegelii* (Günther, 1859) from the Japanese Main Islands; *R. owstoni* (Stejneger, 1907) and *R. viridis* (Hallowell, 1861) from the Ryukyus; *R. arvalis* Lue, Lai, and Chen, 1995, *R. aurativentris* Lue, Lai, and Chen, 1994, *R. moltrechti* Boulenger, 1908, *R. prasianthus* Mou, Risch, and Lue, 1983, and *R. taipeianus* Liang and Wang, 1978 from Taiwan; *R. dennysii* Blanford, 1881 from China; *R. feae* Boulenger, 1893 from China, Vietnam, Thailand, and Myanmar; *R. maximus* Günther, 1859 from China, Thailand, Nepal, and India; *R. calcaneus* Smith, 1924 and *R. duboisi* Ohler, Marquis, Swan, and Grosjean, 2000 from Vietnam; *R. promianus* Smith, 1924 (including *R. tunkui* Kiew, 1987) from Thailand and Malaysia; *R. taroensis* Smith, 1940 from Myanmar; *R. malabaricus* Jerdon, 1870 and *R. pseudomalabaricus* Vasudevan and Dutta, 2000 from India; *R. nigropalmatus* Boulenger, 1895 from India, Thailand, Malaysia, Sumatra and Borneo; *R. reinwardtii* (Schlegel, 1840) from Sumatra, Java, Malaysia, and China; *R. dulitensis* Boulenger, 1892 from Sumatra and Borneo; *R. achantharrhena* Harvey, Pemberton, and Smith, 2002 from Sumatra; *R. kajau* Dring, 1984 from Borneo; *R. edentulus* Mueller, 1894, *R. georgii* Roux, 1904, and *R. monticola* Boulenger, 1896 from Sulawesi.

Rhacophorus jarujini n. sp. is distinguished from some species having skin modifications: *R. angulirostris* Ahl, 1927

from Borneo and Sumatra; *R. translineatus* Wu, 1977 from China; and *R. appendiculatus* (Günther, 1859) from Malaysia to Sumatra, Borneo, and the Philippines have a markedly pointed snout, and the last two species have an anal flap like *R. rhodopus* Liu and Hu, 1959 from China and *R. namdaphaensis* Sarkar and Sanyal, 1985 from India. *Rhacophorus appendiculatus* also has a crenulated fringe along the tarsus, as do *R. bisacculus* Taylor, 1962 from Thailand and India and *R. exechopygus* Inger, Orlov, and Darevsky, 1999 from Vietnam. The following species all have a dermal flap or projections at the heel: *Rhacophorus baluensis* Inger, 1954 and *R. gauni* (Inger, 1966) from Borneo; *R. verrucopus* Huang, 1983 from China; *R. bipunctatus* Ahl, 1927 from India, Tibet, Myanmar, Thailand, and Malaysia; *R. barisani* Harvey, Pemberton, and Smith, 2002 and *R. catamitus* Harvey, Pemberton, and Smith, 2002 from Sumatra; *R. marginifer* (Schlegel, 1837) from Java; *R. pardalis* Guenther, 1859 from Malaysia, Sumatra, Borneo, and the Philippines; *R. hoanglienensis* Orlov, Lathrop, Murphy, and Ho, 2001 and *R. orlovi* Ziegler and Köhler, 2001 from Vietnam; and *R. namdaphaensis*, *R. rhodopus*, and *R. translineatus*. In having smooth dorsal skin, *R. jarujini* differs from species with a tuberculate or rugose dorsum (*R. calcadensis* Ahl, 1927 and *R. tuberculatus* Anderson, 1871 from India; *R. verrucosus* Boulenger, 1893 from Vietnam, Cambodia, Myanmar, and India; *R. everetti* Boulenger, 1894 from the Philippines and Borneo).

In *R. jarujini*, a broad web on the fourth finger reaches the distal subarticular tubercle or base of the disk. In some species, webs between the fingers are less developed than in *R. jarujini*, and webbing is limited at most to the base of the subarticular tubercle on the fourth finger (*R. baliogaster* Inger, Orlov, and Darevsky, 1999 from Vietnam; *R. poecilonotus* Boulenger, 1920 and *R. modestus* Boulenger, 1920 from Sumatra; *R. everetti*; *R. bisacculus*). In contrast, other species have hand webbing better developed than in *R. jarujini*. The web reaches the base of the disk on both sides of the third finger as a broad sheet in *R. annamensis* Smith, 1924 from Vietnam (including *R. notater* Smith, 1924; Matsui, 2005); *R. robinsonii* Boulenger, 1903 from Thailand and Malaysia; and *R. harrisoni* Inger and Haile, 1959, *R. rufipes* Inger, 1966, and *R. fasciatus* Boulenger, 1895 from Borneo. In *R. jarujini*, only the outer side of the third finger is sometimes fully webbed.

Although the degree of web development on the fourth finger is similar, *R. bimaculatus* (Peters, 1867) from the Philippines and *R. cyanopunctatus* Manthey and Steiof, 1998 from Thailand, Malaysia, Borneo, and Sumatra have a dark mask on the sides of the head, and blue spots on the flanks. *Rhacophorus turpes* Smith, 1940 from Myanmar has a much narrower head than *R. jarujini*, and the dorsum is immaculate or scattered with dots. *Rhacophorus lateralis* (Boulenger, 1883) from India has a series of dorsolateral markings and much less developed toe webbing.

DISCUSSION

Dubois (1987) divided members of *Rhacophorus* (*sensu lato*) into two subgenera, *Leptomantis* and *Rhacophorus*, and further subdivided the latter subgenus into several species groups. Among about 60 species of *Rhacophorus* mentioned above, *R. jarujini* is superficially similar to members

of the *fasciatus* group (*R. fasciatus* and *R. harrisoni*) or the *pardalis* group (*R. annamensis* [including *R. notater*; see above], *R. pardalis*, and *R. robinsonii*). Although Dubois (1987) noted that his classification was chiefly based on larval morphology, data are lacking for many species. Thus, the reasons for Dubois' (1987) placement of the abovementioned species in the *pardalis* group is unclear. Similarly, because no larvae are yet known for *R. jarujini*, its classification remains uncertain. Species of the *pardalis* group, as well as *R. catamitus* from Sumatra described more recently (Harvey *et al.*, 2002), are especially similar to *R. jarujini* in dorsal color pattern. Dorsal coloration, however, must be adaptive in its cryptic function and is expected to exhibit convergence in many different lineages of Old World treefrogs.

Rhacophorus jarujini was found in the center of the Thai-Lao Dry Plateau (Inger, 1999), where the native vegetation scarcely remains because of the paucity of large forest reserves or conservation areas. Due partly to this habitat modification and partly to poor sampling to date, the number of anuran species known from the plateau is small. In addition, the scarcity of mountain streams in this region seems to restrict the distribution of anuran species to those that breed in ponds or develop directly. Inger (1999) listed 35 species from the plateau, of which only two (*Chirixalus hansae* [Cochran, 1927] and *R. bisacculus*) were considered endemic to the region; *R. bisacculus* has now been reported from outside Thailand (Frost, 2004). Interestingly, both these species are rhacophorids. Inger (1999: 457) listed as endemic another rhacophorid, *Philautus parvulus* (Boulenger, 1893), but this species is actually not endemic to the region (Inger, 1999: 480). Thus, only *C. hansae* and *R. jarujini* can presently be regarded as endemic to the Thai-Lao Dry Plateau. A more intensive survey in this region would likely detect additional undescribed species, as indicated by the discovery of *R. jarujini*, but this should be done quickly, because the modification of amphibian habitats is in progress.

ACKNOWLEDGMENTS

We thank J. Nabhitabhata, H. Ota, K. Araya, M. Toda, and S.-L. Cheng for help in the field, T. Hikida for help in statistical analyses, and H. Ota, Y. Shibata, I. Das, and U. Manthey for providing literature or information. The National Research Council of Thailand and the Royal Forest Department of Thailand granted permission for fieldwork in Thailand. We thank S. Tunhikorn for help in obtaining permission. We are grateful to C. McCarthy and B. T. Clarke (BM), M. S. Hoogmoed (RMNH), N. Ananjeva and N. Orlov (ZIL), R. Günther and U. Manthey (ZMB), A. Haenggi (NHMB), G. Doria (MSNG), and A. Dubois and A. Ohler (MNHN) for allowing MM to examine specimens under their care. K. Nishikawa and T. Shimada prepared radiographs and photographs. This study was supported by grants under The Monbusho International Scientific Research Program (Field Research, Nos. 06041066, 08041144, and 10041166) to MM.

REFERENCES

- Dubois A (1987) *Miscellanea taxinomica* [sic] batrachologica (I). *Alytes* 5: 7–95
- Duellman WE (Ed) (1999) *Patterns of Distribution of Amphibians: A Global Perspective*, Johns Hopkins University Press, Baltimore
- Fei L, Ye CY, Jiang JP, Xi F, Huang, YZ (2004) *An Illustrated Key to Chinese Amphibians*. Sichuan Publishing Group and Sichuan

- Publishing House of Science and Technology, Chengdu
- Frost DR (Ed) (1985) Amphibian Species of the World: A Taxonomic and Geographic Reference. Allen Press Inc. and Association of Systematics Collections, Lawrence
- Frost DR (2004) Amphibian species of the world: an online reference, Version 3.0 (22 August, 2004). American Museum of Natural History, New York, Electronic Database accessible at <http://research.amnh.org/herpetology/amphibia/index.html>.
- Harvey BH, Pemberton AJ, Smith EN (2002) New and poorly known parachuting frogs (Rhacophoridae: *Rhacophorus*) from Sumatra and Java. *Herpetol Monogr* 16: 46–92
- Inger RF (1999) Distribution of amphibians in Southern Asia and adjacent islands. In "Patterns of Distribution of Amphibians: A Global Perspective" Ed by WE Duellman, Johns Hopkins University Press, Baltimore, pp 445–482
- Inger RF, Orlov N, Darevsky I (1999) Frogs of Vietnam: A report on new collections. *Fieldiana Zool NS* 92: 1–46
- Jiang SP, Hu SQ, Zhao EM (1987) The approach of the phylogenetic relationship and the supraspecific classification of 14 Chinese species of tree frogs (Rhacophoridae). *Acta Herpetol Sinica NS* 6: 27–42
- Liem SS (1970) The morphology, systematics and evolution of the Old World tree frogs (Rhacophoridae and Hyperoliidae). *Fieldiana Zool* 57: 1–145
- Maeda N, Matsui M (1999) Frogs and Toads of Japan, Rev Ed, Bunichi Sogo Shuppan, Tokyo
- Malkmus R., Manthey U, Vogel G, Hoffman P, Kosuch J (2002) Amphibians and Reptiles of Mount Kinabalu (North Borneo). ARG Gantner Verlag Kommanditgesellschaft, Ruggell
- Manthey U, Steiof C (1998) *Rhacophorus cyanopunctatus* sp. n. (Anura: Rhacophoridae), ein neuer Flugfrosch von der Malaiischen Halbinsel, Sumatra und Borneo. *Sauria* 20: 37–42
- Matsui M (1984) Morphometric variation analyses and revision of the Japanese toads (Genus *Bufo*, Bufonidae). *Contrib Biol Lab Kyoto Univ* 26: 209–428
- Matsui M (2000) Batrachology of Japan and adjacent regions -a systematic review. *Comp Bioch Physiol B* 126: 247–256
- Matsui M (2005) Taxonomic relationships of two species of *Rhacophorus* from Vietnam (Amphibia, Rhacophoridae). *Cur Herpetol* 24: 91–93
- Matsui M, Orlov N (2004) A new species of *Chirixalus* from Vietnam (Anura: Rhacophoridae). *Zool Sci* 21: 671–676
- Matsui M, Wu GF (1994) Acoustic characteristics of treefrogs from Sichuan, China, with comments on systematic relationship of *Polypedates* and *Rhacophorus* (Anura, Rhacophoridae). *Zool Sci* 11: 485–490
- Matsui M, Chan-Ard T, Nabhitabhata J (1996) Distinct specific status of *Kalophrynxus pleurostigma interlineatus* (Anura, Microhylidae). *Copeia* 1996: 440–445
- Matsui M, Nabhitabhata J, Panha S (1998) A new *Ansonia* from northern Thailand (Anura, Bufonidae). *Herpetologica* 54: 448–454
- Matsui M, Nabhitabhata J, Panha S (1999) On *Leptobrachium* from Thailand with a description of a new species (Anura: Pelobatidae). *Jpn J Herpetol* 18: 19–29
- Matsui M, Khonsue W, Nabhitabhata J (2005) A new *Ansonia* from Isthmus of Kra, Thailand (Amphibia, Anura, Bufonidae). *Zool Sci* 22: 809–814
- Taylor, EH (1962) The amphibian fauna of Thailand. *Univ Kansas Sci Bull* 43: 265–599
- Tian WW, Jiang YM (Eds) (1986) Identification Manual of Chinese Amphibians and Reptiles. Science Press, Beijing
- Wilkinson JA, Drewes RC, Tatum OL (2002) A molecular phylogenetic analysis of the family Rhacophoridae with an emphasis on the Asian and African genera. *Molec Phylogenet Evol* 24: 265–273
- Zhao EM, Adler K (1993) Herpetology of China. SSAR, Oxford, Ohio

(Received November 10, 2005 / Accepted December 21, 2005)