

## Photochemical Reaction of 1,4-Naphthoquinone with Olefins

Kazuhiro MARUYAMA\*, Tetsuo OTSUKI\*\*, Akio TAKUWA\*\* and Saburo KAKO\*\*

*Received May 30, 1972*

Photochemical reactions of 1,4-naphthoquinone with various types of olefins are examined. 1,4-Naphthoquinhydrone is produced in the reaction of 1,4-naphthoquinone with olefins having an abstractable allylic hydrogen atom. With the another types of olefins 1,4-naphthoquinone undergoes cycloaddition reaction to yield cyclobutene-type compounds (I) or cyclobutane-type compounds (II).

### I. INTRODUCTION

Recently photochemical reactions have been the attractive field in the synthetic organic chemistry.<sup>1)</sup> A variety of cyclobutane can easily be prepared by the application of photochemical reactions. Competitive formation of cyclobutanes and oxetans was reported when *p*-benzoquinones were irradiated in the presence of olefins.<sup>2)</sup> Photochemical behaviors of 1,4-naphthoquinone have not been elucidated with the exceptions of its dimerization reaction,<sup>3)</sup> and its reactions with indene, isocoumarin and benzofurane to give cyclobutane-type compounds.<sup>4)</sup>

Olefins examined in this experiment are  $\alpha$ -pinene, cyclopentene, cyclohexene, cyclooctene, acenaphthylene, indene and styrene.

### II. RESULTS AND DISCUSSIONS


In the photochemical reaction of 1,4-naphthoquinone with  $\alpha$ -pinene, cyclopentene and cyclohexene, 1,4-naphthoquinone is readily reduced to 1,4-naphthoquinhydrone. 1,4-Naphthoquinhydrone may be resulted from 1,4-naphthosemiquinone radical, which is produced through the hydrogen abstraction of photochemically excited 1,4-naphthoquinone. These processes might be analogous to those of photochemical reactions of *p*-benzoquinones with hydrogen donors.<sup>5)</sup> Thus,  $\alpha$ -pinene, cyclopentene and cyclohexene react with 1,4-naphthoquinone as hydrogen donors rather than olefins, since these olefins have the reactive hydrogen atom.

Another types of olefins, which seem to have no reactive hydrogen atom, add to 1,4-naphthoquinone to give cycloaddition compounds by irradiation. The photo-addition compounds are classified into two groups; cyclobutene-type compounds (I) and cyclobutane-type compounds (II). Cyclooctene and acenaphthylene give cyclobutene-type compounds in the photochemical reaction with 1,4-naphthoquinone (see (Ia) and (Ib))


\* 丸山和博: Department of Chemistry, Faculty of Science, Kyoto University, Kyoto.

\*\* 大槻哲夫, 宅和暁男, 加古三郎: Department of Chemistry, College of Liberal Arts and Science, Kyoto University, Kyoto.





Photochemical Reaction of 1,4-Naphthoquinone with Olefins




( I )




( II )



and indene and styrene give cyclobutane-type compounds (see (IIa) and (IIb)). Factors differentiating between two types of photo-addition compounds, *i.e.* (I) and (II), remain to be clarified. However, cyclobutane-type compounds (II) might be the precursor at the earlier stage of the reaction since the photo-dimer (III) of 1,4-naphthoquinone as a primary reaction product is converted to 1,4,5,8-tetrahydroxy-2,3; 6,7-dibenzobiphenylene (IV) by the treatment with alkali.<sup>3a)</sup> Concurrent formation of photo-addition com-




( III )




( IV )

pound (IIa) and 1,4-naphthoquinhydrone is observed in the reaction of 1,4-naphthoquinone with indene, presumably because indene has the reactive benzylic hydrogen atoms, which are comparable in their nature to allylic hydrogen atoms. Although cyclooctene is a higher analogue of cyclohexene, it gives a photo-addition compound of type (Ia). Molecular model reveals that cyclooctene has no active allylic hydrogen atoms owing to the deformation from the normal bond angles. Hence, cyclooctene reacts with 1,4-naphthoquinone to yield (Ia)-type photo-addition compound, but not 1,4-naphthoquinhydrone.

The principal course of the photochemical reaction is illustrated as follows;





At the first stage of the reaction, 1,4-naphthoquinone is excited through the  $n-\pi^*$  excitation of the carbonyl group by irradiation. In the presence of hydrogen donors photochemically excited 1,4-naphthoquinone (characterized by "\*" in Scheme) abstracts hydrogen atom to form 1,4-naphthosemiquinone radical. The lack of reactive hydrogen atoms should cause 1,4-naphthoquinone to add to olefinic double bonds. Considering these results, the abstraction of allylic hydrogen atoms by excited 1,4-naphthoquinone might proceed more easily than the other processes such as cycloaddition with olefins and dimerization of itself.<sup>6)</sup>

### III. EXPERIMENTAL

**General procedures:** 1,4-naphthoquinone was used after sublimation of commercially available guaranteed grade reagent (mp. 124.0–125.0°C). 10 mmol of 1,4-naphthoquinone and 20 mmol of olefins were dissolved into 20 ml of benzene with an exception of cyclooctene. In the reaction of 1,4-naphthoquinone with cyclooctene carbon tetrachloride was used as solvent. The solution was poured into a usual glass tube, sealed and then irradiated for 20–40 hours by high pressure Hg lamp (400 W). The reaction vessel set apart by 5 cm long from the light source was immersed in water. The reaction products were isolated by column-chromatography on florisol and purified by recrystallization.

#### Physical properties of products;

(Ia); brownish white solid after washing with acetone and ether repeatedly. mp 303.0–303.5°C (decom.). IR;  $\nu_{\text{OH}}$ : 3510  $\text{cm}^{-1}$  (sharp), 3350  $\text{cm}^{-1}$  (broad) (as KBr disk). NMR;  $\tau$ : 1.50–2.70 (4H, aromatic-H, multiplet),  $\tau$ : 5.40–6.35

Photochemical Reaction of 1,4-Naphthoquinone with Olefins

- (2H, methine-H, multiplet), 7.65–8.88 (12H, aliphatic-H, broad multiplet) (in the solution of  $\text{CF}_3\text{COOH-CDCl}_3$ ).
- (Ib); yellowish white needles after recrystallization from chloroform. mp 296.0–298.0°C (decom.). IR;  $\nu_{\text{OH}}$ : 3255  $\text{cm}^{-1}$  (broad) (as KBr disk). NMR;  $\tau$ : 2.20–2.58 (10H, aromatic-H, multiplet),  $\tau$ : 5.92 (2H, methine-H, singlet) (in the solution of  $\text{CDCl}_3$ ).
- (IIa); white plates after recrystallization from benzene. mp 180.5–181.5°C IR;  $\nu_{\text{C=O}}$ : 1670  $\text{cm}^{-1}$  (as KBr disk). NMR;  $\tau$ : 1.65–2.30 (4H, aromatic-H due to 1,4-naphthoquinone rest, multiplet),  $\tau$ : 2.65 (4H, aromatic-H due to indene rest, broad singlet),  $\tau$ : 5.90–6.17, 6.58–6.95 (6H, methine-H and methylene-H, multiplet) (in the solution of  $\text{CDCl}_3$ ).
- (IIb); white plates after recrystallization from benzene and petroleum ether. mp 66.0–67.0°C. IR;  $\nu_{\text{C=O}}$ : 1670  $\text{cm}^{-1}$  (as KBr disk). NMR;  $\tau$ : 1.65–2.30 (4H, aromatic-H due to 1,4-naphthoquinone rest, multiplet),  $\tau$ : 2.68 (5H, aromatic-H due to styrene rest, singlet),  $\tau$ : 6.18–6.75, 7.12–7.35 (5H, methine-H and methylene-H, multiplet) (in the solution of  $\text{CDCl}_3$ ).

REFERENCES

- (1) A. Schönberg, G. O. Schenk and O. A. Neumüller, *Preparative Organic Photochemistry*, Springer-Verlag, Berlin-Heidelberg-New York, 1968.
- (2) D. Bryce-Smith and A. Gilbert, *J. Chem. Soc.*, (1964) 87.
- (3) a) J. M. Bruce, *ibid.*, (1962) 2782.  
b) J. Rennert, S. Japar and M. Guttman, *Photochem. and Photobiol.*, **6**, 485 (1967).  
c) J. Dekker, P. J. van Vuuren and D. P. Venter, *J. Org. Chem.*, **33**, 464 (1968).
- (4) C. H. Krauch and S. Farid, *Tetrahedron Lett.*, (1966) 4783.
- (5) K. Maruyama, T. Otsuki and A. Takuwa, *Chem. Lett.*, (1972) 131.
- (6) This could be supported by the measurement of the reacting system using CIDNP methods.