

Table I. Palladium-Catalyzed Decomposition of **1**^a.

Solvent	Temp., °C	Reaction Time, hr	Conv., % ^b	Yield of Product, % ^c			
				2	3	4	5
Benzene	40	1.5	100	43	26	—	26
Benzene	reflux	0.5	100	43	29	2	27
Benzene ^d	reflux	0.25	100	45	29	3	22
Ethanol ^e	reflux	0.5	100	45	19	—	27
Diethylamine	reflux	2.0	96	42	15	—	22
<i>n</i> -Butylamine	reflux	2.0	84	29	11	—	0.4

^a Method A. ^b Consumed **1**. ^c Based on **1** used. ^d Method B. ^e 99.5%.










Table II. Effect of Hydrogen-Acceptors.

A	Method	Conv., % ^a	Yield of Products, % ^b			
			2	3	5	AH ₂
PhCH=CH ₂	B	87	61	2	26	40
<i>cis</i> -PhCH=CHPh	B	100	56	6	29	32
<i>trans</i> -PhCH=CHPh	B	100	63	7	23	33
CH ₃ COCH=CHPh	A	100	52	3	30	10
CH ₃ COCH=CHPh	B	100	61	2	27	25
HOC(CH ₃) ₂ C≡CC(CH ₃) ₂ OH	A	100	63	9	23	35(2) ^c
HOC(CH ₃) ₂ C≡CC(CH ₃) ₂ OH	B	100	76	0	22	57(3) ^c
PhC≡CPh	A	99	69	0	24	49(4) ^c
PhC≡CPh	B	97	75	0	24	59(4) ^c

^a Consumed **1**. ^b Based on **1** used. ^c Numbers out of and in parentheses are yields of *cis*- and *trans*-isomers, respectively.

Table III. Palladium-Catalyzed Decomposition of **1** in the Presence of Amine^a.

Azide	Amine ^b	A	Method	Conv., % ^c	Yield of Products, % ^d									
					2	9	3	8	5	10	11	12	Total ^e	AH ₂
1	8	—	A	100	41	—	30	—	16	10	2	1	29	—
1	8	H ^f	A	99	58	—	23	—	15	13	2	1	31	19
1	8	H ^f	B	93	57	—	24	—	14	12	2	1	29	21
13	3	—	A	100	—	44	—	30	1	3	9	19	32	—
13	3	H ^f	A	98	—	51	—	20	1	2	12	19	34	15
13	3	H ^f	B	90	—	51	—	19	1	2	11	17	31	18

^a **8**, Cl--CH₂NH₂; **9**, Cl--CN; **10**, -CH=NCH₂--Cl; **11**, Cl--CH=NCH₂-; **12**, Cl--CH=NCH₂--Cl; **13**, Cl--CH₂N₃.

^b 0.5 mmol. ^c Consumed azide. ^d Based on azide used. ^e Total yield of Schiff bases. ^f H, 2,5-Dimethyl-3-hexyne-2,5-diol.


employed to avoid unnecessary bimolecular reaction of **1**, did not alter the yields of products. The results are summarized in Table I.

Since, formally, the present reaction is composed of the disproportionation of the

intermediate benzyl nitrene and successive side-processes,⁴⁻⁶ the yield of **2** is expected not to exceed 50% under the present condition. In the presence of a suitable hydrogen-acceptor, however, the yield of **2** may be improved at the sacrifice of the yields of amine-products. Among many compounds investigated, those listed in Table II appeared to be good hydrogen-acceptors. Mono-substituted acetylenes and some olefins such as 1,1-dicyano-2-phenylethylene, 1,1-diacetyl-2-phenylethylene, and *trans*-1,2-dicyanoethylene prevented the decomposition of **1** almost completely. Table II also shows that the high-dilution technique is effective when the reaction is run in the presence of a hydrogen-acceptor. As is shown in Table III, the coexistence of an amine in the reaction mixture causes large decrease of the effect of the hydrogen-acceptor.


DISCUSSION

There is little doubt that a complex, **6**,⁷ might be formed initially by the interaction between **1** and palladium metal.⁸ The retardation of the reaction by the presence



of mono-substituted acetylenes or electron-deficient olefins may be accounted for by strong interaction between these compounds and palladium preventing the azide from coordination.

The successive processes may be obvious:




The fact that an acetylenic compound used for a hydrogen-acceptor is reduced to the *cis*-olefin predominantly supports the process in Eq. 4, which is the catalytic hydrogenation of the acceptor.⁹ The reaction in Eq. 6 is obvious from the results shown in Table III, and that in Eq. 7 is known.⁶ Thus, an excellent hydrogen-acceptor is the compound which has larger susceptibility to catalytic hydrogenation than **1** but has less facility of coordination to palladium (0) than **1**.

It is important to emphasize that, as is shown in Tables I and II, the yield of **5** remains constant (~25%) whole under a variety of reaction conditions, where the yield of **3** changes widely (0-30%). The results cannot be accounted for by reactions in Eqs. 3-7 by themselves, where **3** is proposed to be the precursor of **5**. Instead, it is reasonable to expect another process for the appearance of **5**.

Juday and Adkins proposed the intermediate formation of a dimer of the imine to account for the formation of a Schiff base from Raney nickel hydrogenation of aromatic nitrile.⁶ Under the present condition, the results are best interpreted by assuming a

trimer of **6**, which is in equilibrium with the monomer, for a precursor of **5** (Eqs. 8 and 9).



Let us assume $K=(3 \times 2)^3/1=216$ (mol/l)³ for the equilibrium in Eq. 8, then the maximum yields of **2**, **3**, and **5** are calculated to be 44, 33, and 22%, respectively, in the absence of a hydrogen-acceptor and those of **2**, **AH**₂, and **5** to be 78, 67, and 22%, respectively, in the presence of a hydrogen-acceptor, **A**. The calculated values are in good agreement with those observed experimentally. Of course, the value of the equilibrium constant assumed above has no real meaning because the rates of reactions from **6** and **7** have to be taken into account. The value, instead, suggests that the contribution of **6** is about twice as important as that of **7**.

Consequently, in order to improve the yield of **2** more than 80% one has to look for a condition which shifts the equilibrium toward right-hand side and the research is in progress to this end.

EXPERIMENTAL

Materials. Benzyl azide (bp 86–87°C/20 mmHg (lit.¹⁰) 82.5°C/16.5 mmHg), n_D^{14} 1.5407), *p*-chlorobenzyl azide (bp 80–82°C/1.5 mmHg, n_D^{15} 1.5600),¹¹) and diphenylacetylene (mp 60–61°C (lit.¹²) mp 60–61°C) were prepared according to the literature, respectively. Solvents were purified by distillation prior to the use. Other reagents were obtained from commercial sources and used without further purification.

Procedures. Method A. A mixture of benzyl azide (2.0 mmol) and palladium black (10 mg) in benzene (4.0 ml) was refluxed for 30 min or stirred for 1.5 hr at 40°C under an atmosphere of nitrogen. It was necessary to exclude air from the system when the reaction was carried out without refluxing, otherwise no reaction took place even after 12 hr.

To the reaction mixture were added 200 mg of biphenyl as an internal standard, 4 ml of ether, and 10 ml of water. The ether extract was subjected to VPC analyses on Yanagimoto G 1800 with a column of Silicon DC-200 (1m, 150°C, H₂ 0.4 kg/cm²) or BDS (1m, 120°C, H₂ 0.2 kg/cm²).

Method B. Into a refluxing mixture of palladium black (10 mg) and benzene (4.0 ml) was infused benzyl azide (2.0 mmol) over a period of 2 hr. The whole mixture was refluxed for additional 15 min and the products were analysed as described above.

In some experiments 100 mg of palladium on charcoal was employed in place of palladium black yielding similar results. Ethanol could be substituted for benzene, but amines were not good solvents for the reaction (Table I).

Identification of Products. In a typical run 3.3 g (0.02 mol) of *p*-chlorobenzyl azide was added to a refluxing mixture of 2,5-dimethyl-3-hexyne-2,5-diol (2.8 g, 0.02 mol) and palladium black (0.1 g) in benzene (20 ml) over a period of 1 hr and the whole mixture was refluxed for additional 10 min. The reaction mixture was filtered and the filtrate was washed with four 25 ml portions of water. The organic layer was dried over CaCl₂ and the solvent was evaporated *in vacuo*. The residue was distilled at a bath temperature

of 95°C under a pressure of 20 mmHg.


The distillate was recrystallized from cyclohexane giving 1.6 g (60% yield) of *p*-chlorobenzonitrile: mp 91–93°C (lit.¹³) mp 89–91°C.

The unvolatile substance was recrystallized from *n*-hexane giving 0.5 g (20% yield) of *N*-(*p*-chlorobenzyliden)-*p*-chlorobenzylamine: mp 58–60°C (lit.¹⁴) mp 62–65°C.

The water layers were combined and extracted with 50 ml of ether. The ether layer was washed with three 25 ml portions of water and dried over CaCl₂. The solvent was evaporated *in vacuo* and the residue was recrystallized from benzene yielding 1.0 g (40% yield) of *cis*-2,5-dimethyl-3-hexene-2,5-diol: mp 68–69°C (lit.¹⁵) mp 68–69°C.

REFERENCES

- (1) M. E. C. Biffin, J. Miller, and D. B. Paul, "Introduction of the Azide Group," in "The Chemistry of the Azide Group," S. Patai, Ed., Interscience Pub., New York, N. Y., 1971, Chapter 2, pp 57–190.
- (2) T. Sheradsky, "Azides as Synthetic Starting Materials," Chapter 6, pp 331–395 in Ref. 1.
- (3) H. Hayashi and S. Oka, *Bull. Inst. Chem. Res., Kyoto Univ.*, **52** (3), 514 (1974).
- (4) G. Brieger and T. J. Nestruck, *Chem. Rev.*, **74**, 567 (1974).
- (5) R. Juday and H. Adkins, *J. Amer. Chem. Soc.*, **77**, 4559 (1955).
- (6) N. Yoshimura, I. Moritani, T. Shimamura, and S. Murahashi, *ibid.*, **95**, 3038 (1973).
- (7) We do not specify the structure of the complex. It may be represented by one of **6a–6c** or by equilibrium(s) between two or three of them.



- (8) (a) G. L'abbé, *Chem. Rev.*, **69**, 345 (1969); (b) H. Kwart and A. A. Kahn, *J. Amer. Chem. Soc.*, **89**, 1950, 1951 (1967).
- (9) (a) R. L. Burwell, Jr., *Chem. Rev.*, **57**, 895 (1957); (b) E. F. Meyer and R. L. Burwell, Jr., *J. Amer. Chem. Soc.*, **85**, 2877 (1963).
- (10) T. Curtius and G. Ehrhart, *Chem. Ber.*, **55**, 1559 (1922).
- (11) A. J. Hubert, *Bull. Soc. Chim. Belg.*, **79**, 195 (1970); *Chem. Abstr.*, **73**, 3865d (1970).
- (12) L. I. Smith and M. M. Falkof, *Org. Synth.*, Col. Vol. **3**, 350 (1955).
- (13) C. R. Hauser and G. Vermillion, *J. Amer. Chem. Soc.*, **63**, 1224 (1941).
- (14) Y. Sasaki and F. P. Olsen, *Can. J. Chem.*, **49**, 271 (1971).
- (15) I. N. Nazarov, L. D. Bergel'son, L. P. Badenkova, and B. V. Lopatin, *Zh. Obshch. Khim.*, **28**, 1132 (1958); *Chem. Abstr.*, **52**, 19925f (1958).