TITLE:

ELLIPTIC BOUNDARY VALUE PROBLEMS IN THE SPACE OF DISTRIBUTIONS(Microlocal Geometry)

AUTHOR(S):

ANDRONIKOF, E.; TOSE, N.

CITATION:

ANDRONIKOF, E. ...[et al]. ELLIPTIC BOUNDARY VALUE PROBLEMS IN THE SPACE OF DISTRIBUTIONS(Microlocal Geometry). 数理解析研究所講究録 1993, 845: 13-18

ISSUE DATE:

1993-06

URL:

<http://hdl.handle.net/2433/83623>

RIGHT:

ELLIPTIC BOUNDARY VALUE PROBLEMS IN THE SPACE OF DISTRIBUTIONS

E. ANDRONIKOF (UNIV. PARIS XIII)

AND

N. TOSE (KEIO UNIV.)

戸澤 洋之 (慶応大)

Introduction

Elliptic boundary value problems have their own long history. For the general system they were, however, first clearly formulated microlocally by M. Kashiwara and T. Kawai [K-K]. Their theorem has enjoyed many applications, for example, to solvability of operators of simple characteristics, hypoelliptic operators, and tangential Cauchy-Riemann systems. The theorem does not give, however, much information if we restrict ourselves in the space of distributions. This note aims at giving an analogous theorem of Kashiwara-Kawai type in case function spaces are tempered. See Theorem 3 in Section 1 for the main theorem. By this theorem, we can obtain many application to distribution boundary values of holomorphic functions (e.g. M. Uchida[U]). The result of this note was obtained while the second author was staying in Univ. de Paris VI and Univ. Paris XIII.

1. Main theorem

Let M be a real analytic manifold of dimension n with a complex neighborhood X . Let $\mathcal{M}$ be a coherent $\mathcal{D}_X$ module on X and assume that $\mathcal{M}$ is elliptic on M , *i.e.*

$$(1) \quad \text{char}(\mathcal{M}) \cap T_M^*X \subset T_X^*X.$$

Let N be a real analytic submanifold of M of codimension $d \geq 1$ in M , and Y be a complexification of N in X . We assume that Y is non-characteristic for $\mathcal{M}$, *i.e.*

$$(2) \quad \text{char}(\mathcal{M}) \cap T_Y^*X \subset T_X^*X.$$

In this situation, we have the canonical morphisms

$$T_N^*M \xleftarrow[\rho]{} T_N^*X \xrightarrow[\varpi]{} T_X^*X.$$

Under the above notation we have

THEOREM 1. *The natural morphism*

$$\mathbf{R}\rho_* \mathbf{R}\underline{\text{Hom}}_{\mathcal{D}_X}(\mathcal{M}, \mathcal{C}_{N|X}^f) \leftarrow \mathbf{R}\underline{\text{Hom}}_{\mathcal{D}_X}(\mathcal{M}, T^{-\mu_N}(\mathcal{D}b_M)) \otimes \text{or}_{N/M}$$

is an isomorphism.

In the above theorem $\text{or}_{N/M}$ denotes the relative orientation sheaf of N in M . The sheaf $\mathcal{C}_{N|X}^f$ on T_N^*X is the tempered version of $\mathcal{C}_{N|X}$ and is given, with the tempered microlocalization due to E. Andronikof[An], by

$$\mathcal{C}_{N|X}^f := T^{-\mu_N}(\mathcal{O}_X) \otimes \text{or}_M[n].$$

We remark that the above object in the derived category is concentrated in degree 0. For a point $\overset{\circ}{x} \in T_N^*X$, the stalk of $\mathcal{C}_{N|X}^f$ at $\overset{\circ}{x}$ is given, with the aid of local cohomology with bounds, by

$$\mathcal{C}_{N|X, \overset{\circ}{x}}^f \simeq \varinjlim \mathbb{H}_{[Z]}^n(\mathcal{O}_X)_{\pi_X(\overset{\circ}{x})}.$$

Here π_X denotes the projection $\pi_X : T^*X \rightarrow X$ and the inductive limit is taken for all closed subanalytic sets Z in X satisfying the property

$$C_N(Z)_{\pi_X(\overset{\circ}{x})} \subset \{v \in T_N X; \langle \overset{\circ}{x}, v \rangle < 0\} \cup \{0\}.$$

Refer here to Kashiwara-Schapira[K-S2] for the notion of normal cones $C_N(\cdot)$. The sheaf $T-\mu_N(Db_M)$ on T_N^*M is also constructed by E. Andronikof[An]. We just explain that its stalk at $\overset{\circ}{x} \in T_N^*M$ is given by the isomorphism

$$T-\mu_N(Db_M)_{\overset{\circ}{x}} \simeq \varinjlim_Z \Gamma_Z(Db_M)_{\pi_M(\overset{\circ}{x})}.$$

Here the inductive limit is taken for any closed subanalytic set Z in M with the property

$$C_N(Z)_{\pi_M(\overset{\circ}{x})} \subset \{v \in T_N M; \langle \overset{\circ}{x}, v \rangle < 0\} \cup \{0\}$$

($\pi_M: T^*M \rightarrow M$).

Next we give another theorem, which is analogous to Theorem 6.3.1 of Kashiwara-Shapira [K-S1] (refer also to Kashiwara-Kawai[K-K] where we find the theorem of [K-S1] in its original form).

THEOREM 2. *Let $\tilde{\mathcal{M}} = \mathcal{E}_X \otimes_{\pi_X^{-1}\mathcal{D}_X} \pi_X^{-1}\mathcal{M}$. Then the natural morphism*

$$\mathbf{R}\underline{\mathrm{Hom}}_{\mathcal{E}_X}(\tilde{\mathcal{M}}, \mathcal{C}_{N|X}^f) \longleftarrow \mathbf{R}\underline{\mathrm{Hom}}_{\mathcal{E}_X}(\tilde{\mathcal{M}}, \mathcal{E}_{X \leftarrow Y}) \bigotimes_{\mathrm{End}(\mathcal{E}_{X \leftarrow Y})}^{\mathbf{L}} \mathbf{R}\underline{\mathrm{Hom}}_{\mathcal{E}_X}(\mathcal{E}_{X \leftarrow Y}, \mathcal{C}_{N|X}^f)$$

is an isomorphism outside of $T_N^*X \cap T_Y^*X$. This entails an isomorphism

$$\mathbf{R}\underline{\mathrm{Hom}}_{\mathcal{E}_X}(\mathcal{M}, \mathcal{C}_{N|X}^f) \simeq \mathbf{R}\underline{\mathrm{Hom}}_{\mathcal{E}_X}(\mathcal{M}, \mathcal{E}_{X \leftarrow Y}) \otimes_{p^{-1}\mathcal{E}_Y}^{\mathbf{L}} p^{-1}\mathcal{C}_N^f$$

on $T_N^*X \setminus T_Y^*X$ where p is the canonical morphism

$$p: T_N^*X \setminus T_Y^*X \rightarrow T_N^*Y.$$

In the above theorem, the object $\mathcal{C}_N^f$ on T_N^*Y is the sheaf of temperate microfunctions. This is a subsheaf of $\mathcal{C}_N$ and describes microlocal analytic singularities of distributions on N . By the notation of E. Andronikof[An], this sheaf is defined as

$$\mathcal{C}_N^f := T-\mu_N(\mathcal{O}_Y)[n-d] \otimes \mathrm{or}_{N/Y}.$$

The proof of this theorem is essentially the same as in Theorem 6.3.1 of [K-S1] and relies on the division theorem of temperate microfunctions with holomorphic parameters with respect to microdifferential operators. We also remark that only the non-charactericity of Y is utilized in its proof.

By combining the above theorems into one, we get the main theorem of this note. Let q denote the restriction of ρ to $\overset{\circ}{T}_N^*X \setminus T_M^*X$; $q: \overset{\circ}{T}_N^*X \setminus T_M^*X \rightarrow T_N^*M$ and p the projection $\overset{\circ}{T}_N^*X \setminus T_Y^*X \rightarrow \overset{\circ}{T}_N^*Y$. Then we have

THEOREM 3. *We have a canonical isomorphism on $\overset{\circ}{T}_N^* Y$*

$$\mathbf{R}q_* \left(\mathbf{R}\underline{\mathrm{Hom}}_{\mathcal{E}_X}(\tilde{\mathcal{M}}, \mathcal{E}_{X \leftarrow Y} |_{\overset{\circ}{T}_N^* X}) \overset{\mathbf{L}}{\otimes}_{p^{-1}\mathcal{E}_Y} p^{-1}\mathcal{C}_N^f \right) \simeq \mathbf{R}\underline{\mathrm{Hom}}_{\mathcal{D}_X}(\mathcal{M}, T-\mu_N(\mathcal{D}b_M)) \otimes or_{N/M}.$$

2. Idea of Proof

What is left to us is now to construct the morphism in Theorem 1 and to show it an isomorphism.

First we construct a commutative diagram

$$(A) \quad \begin{array}{ccc} \mathbf{R}\rho_! \mathcal{C}_{N|X}^f \otimes or_{N/X} & \longrightarrow & T-\mu_N(\mathcal{A}_M) \\ \downarrow & & \downarrow \\ \mathbf{R}\rho_* \mathcal{C}_{N|X}^f \otimes or_{N/X} & \longleftarrow & T-\mu_N(\mathcal{D}b_M) \end{array}$$

where $T-\mu_N(\mathcal{A}_M)$ is the tempered microlocalization of the sheaf $\mathcal{A}_M$ along N and is constructed by E. Andronikof[A]. This object is the Fourier transform of the tempered specialization $T-\nu_N(\mathcal{A}_M)$ whose stalk at $\overset{\circ}{v} \in T_N M$ is given by

$$T-\nu_N(\mathcal{A}_M)_{\overset{\circ}{v}} \simeq \varinjlim_U \{u \in \mathcal{A}(U); u \text{ is tempered on } M \text{ as a distribution}\}.$$

Here U in the inductive limit ranges through any open subanalytic set in M with the property

$$\overset{\circ}{v} \notin C_N(M \setminus U).$$

To construct (A), it is sufficient to construct its image by the inverse Fourier transformation

$$(A') \quad \begin{array}{ccc} \iota^{-1} T-\nu_N(\mathcal{O}_X) \otimes or_{N/X} & \longrightarrow & T-\nu_N(\mathcal{A}_M) \\ \downarrow & & \downarrow \\ \iota^! T-\nu_N(\mathcal{O}_X) \otimes or_{N/X} & \longleftarrow & T-\nu_N(\mathcal{D}b_M). \end{array}$$

Here ι is the canonical embedding

$$\iota : T_N M \longrightarrow T_N X,$$

and $T-\nu_N(\mathcal{O}_X)$ is the tempered specialization of the sheaf $\mathcal{O}_X$ along N , which is concentrated in degree 0. The stalk of $T-\nu_N(\mathcal{O}_X)$ at $\overset{\circ}{v} \in T_N X$ is given by

$$T-\nu_N(\mathcal{O}_X)_{\overset{\circ}{v}} \simeq \varinjlim_U \{u \in \mathcal{O}(U); u \text{ can be extended to } X \text{ as a distribution}\}$$

ELLIPTIC BOUNDARY VALUE PROBLEMS IN THE SPACE OF DISTRIBUTIONS

where U runs through all open subanalytic sets in X with $\overset{\circ}{v} \notin C_N(M \setminus U)$. The diagram (A') can be constructed easily if we scrutinize the construction by E. Andronikof[An].

Next we apply $\mathbf{R}\underline{\mathbf{Hom}}_{\mathcal{D}_X}(\mathcal{M}, \cdot)$ to the diagram (A') and obtain the commutative diagram

$$\begin{array}{ccc}
 \mathbf{R}\underline{\mathbf{Hom}}_{\mathcal{D}_X}(\mathcal{M}, \iota^{-1}T^{-\nu_N}(\mathcal{O}_X)) \otimes or_{N/X} & \xrightarrow{\Phi_1} & \mathbf{R}\underline{\mathbf{Hom}}_{\mathcal{D}_X}(\mathcal{M}, T^{-\nu_N}(\mathcal{A}_M)) \\
 \Phi_4 \downarrow & & \downarrow \Phi_2 \\
 \mathbf{R}\underline{\mathbf{Hom}}_{\mathcal{D}_X}(\mathcal{M}, \iota^!T^{-\nu_N}(\mathcal{O}_X)) \otimes or_{N/X} & \xleftarrow{\Phi_3} & \mathbf{R}\underline{\mathbf{Hom}}_{\mathcal{D}_X}(\mathcal{M}, T^{-\nu_N}(\mathcal{D}b_M)).
 \end{array}$$

It is easy to see from the ellipticity of $\mathcal{M}$ that Φ_4 and Φ_2 are isomorphisms. (To show Φ_4 is an isomorphism, it is easier to consider its image by Fourier transformation). Thus to prove that Φ_3 and thus its image by Fourier transformation are isomorphisms, it suffices to show that Φ_1 is an isomorphism. The problem for Φ_1 can be reduced to the case where $\mathcal{M}$ is a single equation; i.e. $\mathcal{M} = \mathcal{D}_X / \mathcal{D}_X P$. Moreover it is sufficient to show that

$$\underline{\mathbf{Hom}}_{\mathcal{D}_X}(\mathcal{D}_X / \mathcal{D}_X P, \iota^{-1}T^{-\nu_M}(\mathcal{O}_X)) \otimes or_{N/X} \longrightarrow \underline{\mathbf{Hom}}_{\mathcal{D}_X}(\mathcal{D}_X / \mathcal{D}_X P, T^{-\nu_N}(\mathcal{A}_M))$$

is surjective. This problem can be solved by using the construction of the elementary solution of P by means of Radon transformation and microdifferential operators.

REFERENCES

- [An] Andronikof, E., Microlocalisation tempérée des distributions et des fonctions holomorphes I, II. C.R. Acad. Sci. t.303, 347-350 (1986) et t.304, n°17, 511-514 (1987). see also Thèse d'Etat, Paris-Nord, (juin 1987) and paper to appear.
- [K-K] Kashiwara M. and T. Kawai, On the Boundary Value Problems for Elliptic Systems of Linear Differential Equations I, II, Proc. Japan Academy 48, pp. 712-715 (1972), 49, pp. 164-168 (1973).
- [K-S1] Kashiwara, M. and P. Schapira, Microhyperbolic Systems, Acta Math. 142, pp. 1-55 (1974).
- [K-S2] ———, MICROLOCAL STUDY of SHEAVES, Astérisque 128 (1985).
- [U] Uchida, M., *in these proceedings*.

E. Andronikof

Département de Mathématiques, Univ. Paris XIII
93430 Villetaneuse, France

N. Tose

Mathematics, General Education, Keio Univ.
4-1-1 Hiyoshi, Yokohama 223, Japan