TITLE:

The Cauchy problem fir hyperbolic operators of strong type

AUTHOR(S):

KAJITANI, KUNIHICO; NISHITANI, TATSUO;
WAKABAYASHI, SEIICHIRO

CITATION:

KAJITANI, KUNIHICO ...[et al]. The Cauchy problem fir hyperbolic operators of strong type.
数理解析研究所講究録 1995, 894: 43-66

ISSUE DATE:

1995-01

URL:

<http://hdl.handle.net/2433/84438>

RIGHT:

The Cauchy problem for hyperbolic operators of strong type

KUNIHICO KAJITANI (梶谷 邦彦)*

TATSUO NISHITANI (西谷 達雄)**

SEIICHIRO WAKABAYASHI (若林 誠一郎)*

1. Introduction

Let P be a differential operator of order m in an open set $\Omega \subset \mathbf{R}^n$, and write $P(x, D) = \sum_{j=0}^m P_j(x, D)$, where the $P_j(x, \xi)$ ($0 \leq j \leq m$) are homogeneous polynomials of ξ of degree j whose coefficients are C^∞ functions of x in Ω . Let $z^0 = (x^0, \xi^0) \in T^*(\Omega) \setminus 0$ be a characteristic point of order r of the principal part P_m of P . It is well known that for the Cauchy problem for P to be C^∞ -well posed the lower order term P_j must vanish at least of order $r - 2(m - j)$ at z^0 if $r - 2(m - j) > 0$, i.e., $j > [(2m - r)/2]$, where $[s]$ denotes the integral part of s for $s \in \mathbf{R}$ (see Ivrii-Petkov [5]). We call the above necessary condition for C^∞ -well posedness the Ivrii-Petkov condition here. On the other hand it is also known that the Cauchy problem is C^∞ -well posed for any lower order terms if and only if every characteristic point of P_m is at most of order 2 and P_m is effectively hyperbolic at every characteristic point of order 2 (see [5], Iwasaki [6] and the references in Nishitani [8]). Therefore, as a generalization of effective hyperbolicity it is natural to call P_m a hyperbolic operator of strong type if the Cauchy problem for P is C^∞ -well posed for every lower order term P_j satisfying the Ivrii-Petkov condition. So a question rises whether there exist hyperbolic operators of strong type. The answer is positive. Roughly speaking, we can prove that a hyperbolic operator is of strong type if its principal part has the real characteristic roots and the difference of any two roots is bounded from below by the absolute value of a time function, where the precise definition of "time function" will be given later.

In Kajitani-Wakabayashi [9] and [10] we introduced the microlocal *a priori* estimates to solve the Cauchy problem for linear partial differential operators in C^∞ , and as its application we investigated the Cauchy problem for such operators with time functions which are in involution. Nishitani studied the same problem for operators

*Institute of Mathematics, University of Tsukuba

**Department of Mathematics, College of General Education, Osaka University

with one time function in [14] and the propagation of singularities in [15]. In this article we shall consider hyperbolic operators of strong type. We shall give a large class of hyperbolic operators of strong type (see Theorem 1.1 below). Although we have no general results on the characterization of hyperbolic operators of strong type, we believe that our class is very close to the whole class of hyperbolic operators of strong type.

Let $P(x, D) = D_1^m + \sum_{|\alpha| \leq m, \alpha_1 < m} a_\alpha(x) D^\alpha$ be a partial differential operator with coefficients in $C^\infty(\mathbf{R}^n)$, where m is a positive integer, $x = (x_1, x') = (x_1, x_2, \dots, x_n) \in \mathbf{R}^n$ and $D = (D_1, D') = (D_1, D_2, \dots, D_n) = -i(\partial/\partial x_1, \dots, \partial/\partial x_n)$. We assume that [A.I] $p(x, \xi) = \xi_1^m + \sum_{|\alpha| = m, \alpha_1 < m} a_\alpha(x) \xi^\alpha$ is hyperbolic with respect to $\theta = (1, 0, \dots, 0)$, i.e.,

$$(1.1) \quad p(x, \xi - i\theta) \neq 0 \quad \text{for } x \in \mathbf{R}^n \text{ and } \xi \in \mathbf{R}^n.$$

It is well known that it follows from Lax-Mizohata's theorem [13] that (1.1) is necessary in order that the Cauchy problem for $P(x, D)$ is C^∞ -well posed.

Let $P(x, \xi)$ be the full symbol of $P(x, D)$ and denote by $P_j(x, \xi)$ the homogeneous part of degree j of $P(x, \xi)$. We define $h_j(x, \xi)$ ($j = 0, 1, \dots, m$) as $|p(x, \xi - is\theta)|^2 = \sum_{j=0}^m s^{2(m-j)} h_j(x, \xi)$ for $x \in \mathbf{R}^n$, $\xi \in \mathbf{R}^n$ and $s \in \mathbf{R}$. Writing $p(x, \xi) = \prod_{j=1}^m q_j(x, \xi)$, where $q_j(x, \xi) = \xi_1 - \lambda_j(x, \xi')$, we have

$$h_j(x, \xi) = \sum_{1 \leq \ell_1 < \ell_2 < \dots < \ell_j \leq m} |q_{\ell_1}(x, \xi)|^2 \cdots |q_{\ell_j}(x, \xi)|^2 \quad \text{for } 1 \leq j \leq m$$

and $h_0(x, \xi) = 1$. We assume that for $z^0 = (x^0, \xi^0) \in T^*(\mathbf{R}^n) \simeq \mathbf{R}^{2n}$ with $|\xi^0| = 1$

[A.II] $_{z^0}$ there are a conic neighborhood $\mathcal{U}$ of z^0 and $C > 0$ such that

$$|P_j(x, \xi)| \leq C |h_{2j-m}(x, \xi)|^{1/2}$$

for $(x, \xi) \in \mathcal{U}$ with $|\xi| = 1$ and $[m/2] + 1 \leq j \leq m - 1$.

It should be noted that $h_{2j-m}(x, \xi) \neq 0$ near z^0 if $j \leq (2m - r)/2$ and z^0 is a characteristic point of order r . We note that there are variants equivalent to the condition [A.II] $_{z^0}$ (see Lemma 2.5 in [10]). We remark that if the multiple characteristic set $\{(x, \xi) \in T^*(\mathbf{R}^n) \setminus 0; p(x, \xi) = dp(x, \xi) = 0\}$ of $p(x, \xi)$ is a C^∞ manifold, the

condition $[A.II]_{z^0}$ is necessary in order that the Cauchy problem for $P(x, D)$ is C^∞ -well posed (see [5]). We also note that if the coefficients of P belong to a Gevrey class $\mathcal{E}^{(s)}$ ($s \leq 2$) and the conditions $[A.I]$ and $[A.II]_{z^0}$ for every $z^0 \in T^*(\mathbf{R}^n)$ with $|\xi^0| = 1$ are valid, then the Cauchy problem is well posed in $\mathcal{E}^{(s)}$ (see Kajitani [8]).

We now introduce the notion of time functions. For a multiple characteristic point $z^0 \in T^*(\mathbf{R}^n) \setminus 0$ of p , we denote by $p_{z^0}(X)$ the localization of p at z^0 , i.e., $p(z^0 + sX) = s^\mu(p_{z^0}(X) + o(1))$ as $s \rightarrow 0$ and $p_{z^0}(X) \neq 0$ in $X \in T_{z^0}(T^*\mathbf{R}^n)$. Denote by $\Gamma(p_{z^0}, (0, \theta))$ the connected component of the set $\{X \in T_{z^0}(T^*\mathbf{R}^n); p_{z^0}(X) \neq 0\}$ containing $(0, \theta)$. We say that $t(x, \xi)$ is a time function of p at z^0 if $t(x, \xi)$ is real-valued, continuous in $T^*(\mathbf{R}^n) \setminus 0$ and positively homogeneous in ξ of degree 0 and if there are a conic neighborhood $\mathcal{U}$ of z^0 and a compact convex set $K \subset \Gamma(p_{z^0}, (0, \theta))$ such that $t(x, \xi)$ is Lipschitz continuous in $\mathcal{U} \cap \{1/2 \leq |\xi| \leq 2\}$ and $-(|\xi| \nabla_\xi t(x, \xi), -\nabla_x t(x, \xi)) \in K$ for a.e. $(x, \xi) \in \mathcal{U}$. In particular, $t(x, \xi)$ is a time function of p at z^0 , if $t(x, \xi)$ is a real valued function in $C^1(T^*(\mathbf{R}^n) \setminus 0)$ and positively homogeneous of degree 0 and $-H_t(z^0) \equiv -(\nabla_\xi t, -\nabla_x t)(z^0) \in \Gamma(p_{z^0}, (0, \theta))$. In fact, it follows from continuity in z of $-H_t(z)$ that there are a conic neighborhood $\mathcal{U}$ of z^0 and a compact convex set $K \subset \Gamma(p_{z^0}, (0, \theta))$ such that $-H_t(z) \in K$ for $z \in \mathcal{U}$.

For a multiple characteristic point $z^0 = (x^0, \xi^0) \in T^*(\mathbf{R}^n)$ with $|\xi^0| = 1$ we assume that

$[A.III]_{z^0}$ there are a conic neighborhood $\mathcal{U}$ of z^0 and time functions $t_\ell(x, \xi)$ ($1 \leq \ell \leq d$) of p at z^0 such that

$$(1.2) \quad h_{m-1}(x, \xi) \geq h_{m-2}(x, \xi)t(x, \xi)^2 \quad \text{for } (x, \xi) \in \mathcal{U} \text{ with } |\xi| = 1,$$

where $t(x, \xi) = \min_{1 \leq \ell \leq d} |t_\ell(x, \xi)|$.

In particular, $[A.III]_{z^0}$ implies that the multiple characteristic set of p is included in the union of the zeros of finitely many time functions. We note that the condition (1.2) is equivalent to

$$(1.3) \quad |q_j(x, \xi)| + |q_k(x, \xi)| \geq ct(x, \xi) \quad \text{for } (x, \xi) \in \mathcal{U} \text{ with } |\xi| = 1 \text{ and } j \neq k,$$

where $p(x, \xi) = \prod_{j=1}^m q_j(x, \xi)$ and $c > 0$ (see Lemma 2.4 in [10]). Since it follows from [1] that the q_j are Lipschitz continuous in $T^*(\mathbf{R}^n) \setminus 0$, (1.3) is valid if there are $c_{jk} \in \mathbf{R}$ such that $(c_{jk}q_j + c_{kj}q_k)|\xi|^{-1}$ are time functions of p at z^0 for $j \neq k$.

Now we can state our main result. Let Ω be a bounded domain in $\mathbf{R}^n$ containing the origin. Under the assumption [A.I] we can define the generalized flows issuing from x^0 in $\mathbf{R}^n$ by

$$K_{x^0}^\pm = \{x(t); \pm t \geq 0, \text{ and } \{x(t)\} \text{ is a Lipschitz continuous curve in } \mathbf{R}^n \\ \text{satisfying } dx(t)/dt \in \Gamma(p(x(t), \cdot), \theta)^* \text{ (a.e. } t) \text{ and } x(0) = x^0\},$$

where $\Gamma(p(x, \cdot), \theta)$ denotes the connected component of the set $\{\xi \in \mathbf{R}^n; p(x, \xi) \neq 0\}$ containing θ and $\Gamma^* = \{x \in \mathbf{R}^n; x \cdot \xi \geq 0 \text{ for any } \xi \in \Gamma\}$.

Theorem 1.1. Assume that the conditions [A.I] is satisfied and that [A.II] $_{z^0}$ and [A.III] $_{z^0}$ are satisfied for $z^0 = (x^0, \xi^0) \in T^*(\mathbf{R}^n)$ with $|\xi^0| = 1$, $x^0 \in \bar{\Omega}$, $x_1^0 \geq 0$ and $dp(z^0) = 0$. Then for any $f \in \mathcal{D}'$ with $\text{supp } f \subset \{x_1 \geq 0\}$ the Cauchy problem

$$(CP) \quad \begin{cases} P(x, D)u(x) = f & \text{in } \Omega, \\ \text{supp } u \subset \{x_1 \geq 0\} \end{cases}$$

has a solution $u \in \mathcal{D}'$. If $x^0 \in \Omega$, $K_{x^0}^- \cap \{x_1 \geq 0\} \subset \Omega$, and $u \in \mathcal{D}'$ satisfies (CP) with $f = 0$ near $K_{x^0}^-$, then $x^0 \notin \text{supp } u$. Moreover if $x^0 \in \Omega$, $K_{x^0}^- \cap \{x_1 \geq 0\} \subset \Omega$, $u \in \mathcal{D}'$ satisfies (CP) and $f \in C^\infty(\mathbf{R}^n)$, then $x^0 \notin \text{sing supp } u$, i.e., $u \in C^\infty$ near x^0 .

Here we give a simple example to elucidate the geometrical meanings of the conditions [A.II] $_{z^0}$ and [A.III] $_{z^0}$. Let $p(x, \xi)$ be factorized as

$$(1.4) \quad p(x, \xi) = e(x, \xi) \prod_{j=1}^r q_j(x, \xi)$$

in a conic neighborhood $\mathcal{U}$ of z^0 , where $e(x, \xi)$ and $q_j(x, \xi)$ are smooth near z^0 and positively homogeneous of degree $m - r$ and 1, respectively, and $e(z^0) \neq 0$, $dq_j(z^0) \neq 0$ and $q_j(x^0, \theta) > 0$. Assume, for simplicity, that dq_j are linearly independent at z^0 . Recall that the cone generated by the Hamilton vector fields $H_{q_j}(z^0)$ of q_j forms the propagation cone $\Gamma(p_{z^0}, (0, \theta))^\sigma$ of the localization $p_{z^0}(X)$, where $\Gamma^\sigma = \{X \in T_{z^0}(T^*(\mathbf{R}^n)); \sigma(Y, X) \geq 0 \text{ for any } Y \in \Gamma\}$ for $\Gamma \subset T_{z^0}(T^*(\mathbf{R}^n))$ and $\sigma = \sum d\xi_j \wedge dx_j$. Then the condition [A.III] $_{z^0}$ is fulfilled if and only if $\Gamma(p_{z^0}, (0, \theta))^\sigma$ is transversal to the tangent space at z^0 of each intersection of any two hypersurfaces $q_k = 0$, $q_j = 0$ (see [15]). On the other hand, the condition [A.II] $_{z^0}$ is satisfied if and only if $P_j(x, \xi)$ vanishes of order $r - 2(m - j)$ on each intersection of any two hypersurfaces $q_k = 0$,

$q_t = 0$ near z^0 whenever $r - 2(m - j) > 0$. Even if the $q_j(x, \xi)$ are not smooth in (1.4), we can show that (1.2) implies that

$$(1.5) \quad \Gamma(p_{z^0}, (0, \theta))^\sigma \cap \Sigma(p_{z^0}) = \{0\},$$

where $\Sigma(p_{z^0}) = \{X \in T_{z^0}(T^*(\mathbf{R}^n)); dp_{z^0}(X) = 0\}$. If z^0 is a characteristic point of order 2, or if $p(x, \xi)$ can be smoothly factorized as (1.4), then the converse is also true (see [8]). So one can conjecture that the condition (1.5) is necessary for (CP) to be C^∞ -well posed with arbitrary lower order terms P_j satisfying [A.II] $_{z^0}$.

The major part of the proof of Theorem 1.1 is devoted to the proof of Theorem 2.2 below. We now explain the basic ideas proving the microlocal *a priori* estimates (2.1) $_{z^0}$ in Theorem 2.2 for a fixed z^0 . To derive the microlocal *a priori* estimates we consider $P_\Lambda(x, D) = e^\Lambda(x, D)^{-1} P(x, D) e^\Lambda(x, D)$, where $e^\Lambda(x, D)$ is a pseudodifferential operator with symbol $e^{\Lambda(x, \xi)}$. Here the Hamilton vector field H_Λ of $\Lambda(x, \xi)$ satisfies

$$H_\Lambda \sim M \sum_{j=1}^d (t_j(x, \xi)^2 + N|\xi|^{-1})^{-1/2} H_{t_j},$$

where M and N are large parameters and the $t_j(x, \xi)$ are the time functions in [A.III] $_{z^0}$. The conjugated operator $P_\Lambda(x, D)$ admits the following expansion:

$$P_\Lambda(x, \xi) \sim \sum_{j=0}^m (iH_\Lambda)^j p(x, \xi)/j! + \dots$$

To estimate the norm of $P_\Lambda u$ from below, following the classical idea attributed to Leray [12] and Gårding [3] we separate P_Λ by $Q(x, D)$ and estimate

$$(1.6) \quad \text{Im} (P_\Lambda(x, D)u, Q(x, D)u)_{L^2} = (S(x, D)u, u)_{L^2},$$

where $S(x, D) = (2i)^{-1}(Q^* P_\Lambda - P_\Lambda^* Q)$ and $(f, g)_{L^2}$ denotes the inner product of $L^2(\mathbf{R}^n)$. A desired $Q(x, \xi)$ is given by

$$Q(x, \xi) = |\tilde{H}|^{-1} \tilde{H} \sum_{j=0}^m (iH_\Lambda)^j p(x, \xi)/j!,$$

where $\tilde{H} = (\langle \xi \rangle_h \nabla_\xi \Lambda(x, \xi), -\nabla_x \Lambda(x, \xi))$ and $\langle \xi \rangle_h = (h^2 + |\xi|^2)^{1/2}$. From the conditions [A.II] $_{z^0}$ and [A.III] $_{z^0}$ and microhyperbolicity for $p(x, \xi)$ with respect to $-H_{t_j}$ (see

[16]), $S(x, \xi)$ becomes a "weight" in the sense of Hörmander [4] and $S(x, \xi) - S_0(x, \xi) \in S(NM^{-2}S_0, g)$, where a suitable metric g associates with the $t_j(x, \xi)$ and $S_0(x, \xi)$ is the principal part of $S(x, \xi)$. Then, if we can choose $NM^{-2} \ll 1$, constructing an inverse of $S(x, D)$, we can show that (1.6) is bounded from below by $\|u\|_{H^{(m-1)/2}(\mathbf{R}^n)}^2$. Then P_Λ satisfies the microlocal *a priori* estimates and, therefore, P also does. From the microlocal *a priori* estimates we can prove Theorem 1.1, following the idea of the proof of Theorem 1.5 in [9].

In our argument, under the restriction that $NM^{-2} \ll 1$, we must construct an inverse of the pseudodifferential operator with symbol

$$e^{\Lambda(x, \xi)} \sim \prod_{j=1}^d ((t_j(x, \xi)^2 |\xi| + N)^{1/2} + t_j(x, \xi) |\xi|^{1/2})^M.$$

In [10] we consider the cases where the $t_j(x, \xi)$ are in involution. If the $t_j(x, \xi)$ are not in involution, we have serious difficulties in this construction, since $e^\Lambda(x, D)$ is a pseudodifferential operator of type 1/2, 1/2 with two large parameters. This construction is essential in proving Theorem 1.1, which will be given in §3.

2. Proof of Theorem 1.1

In this section we shall give an outline of the proof of Theorem 1.1 from the microlocal *a priori* estimates (see Theorem 2.2 below).

Let Ω be a bounded domain in $\mathbf{R}^n$. Denote $\Omega_+ = \Omega \cap \{x_1 > 0\}$ and $\Omega_{+, \delta} = \{x \in \mathbf{R}^n; \text{dis}(x, \Omega_+) < \delta\}$ for $\delta > 0$. Let $\delta > 0$ and choose $R > 0$ so that $\Omega_{+, 2\delta} \subset \{|x| < R\}$. Moreover, choose $\varphi_\delta(x) \in C^\infty(\mathbf{R}^n)$ and $\chi_{R, \delta}(x) \in C^\infty(\mathbf{R}^n; \mathbf{R}^n)$ such that $\varphi_\delta(x) = 1$ in $\Omega_{+, \delta}$ and $\varphi_\delta = 0$ in the complement of $\Omega_{+, 2\delta}$ and $\chi_{R, \delta}(x) = x$ in $\{|x| < R\}$ and $= Rx/|x|$ in $\{|x| \geq R + \delta\}$. For $P(x, \xi) = \sum_{|\alpha| \leq m} a_\alpha(x) \xi^\alpha$ we define

$$\begin{aligned} \tilde{P}(x, \xi) = & (1 - (1 - \varphi_\delta(x))^2 |\xi'|^2 (\partial/\partial \xi_1)^2)^{m-1} \sum_{|\alpha|=m} a_\alpha(\chi_{R, \delta}(x)) \xi^\alpha \\ & + \varphi_{\delta/2}(x) \sum_{|\alpha| < m} a_\alpha(x) \xi^\alpha. \end{aligned}$$

We denote by $\tilde{p}(x, \xi)$ the principal part of $\tilde{P}$. Then we have the following

Proposition 2.1. Assume that $P(x, \xi)$ satisfies [A.I], [A.II] $_{x^0}$ and [A.III] $_{x^0}$. Then, (i) $\tilde{P}(x, \xi) = P(x, \xi)$ for $x \in \Omega_{+, \delta/2}$, (ii) the coefficients of $\tilde{P}$ belong to $\mathcal{B}^\infty(\mathbf{R}^n)$

and $\tilde{p}(x, \xi)$ is strictly hyperbolic with respect to $\theta = (1, 0, \dots, 0)$ in the complement of $\Omega_{+, 2\delta}$, and (iii) $\tilde{P}(x, \xi)$ satisfies [A.I], [A.II] $_{z^0}$ and [A.III] $_{z^0}$.

Let $z^0 = (x^0, \xi^0) \in T^*(\mathbf{R}^n)$ be fixed so that $|\xi^0| = 1$ and $dp(z^0) = 0$. We use the following notations: (i) $P_\gamma(x, D) = P(x, D - i\gamma\theta)$ for $\gamma > 0$ and $\theta = (1, 0, \dots, 0) \in \mathbf{R}^n$, (ii) $t_\pm(x, \xi) \in \mathcal{B}^\infty(T^*(\mathbf{R}^n) \setminus 0)$, positively homogeneous in ξ of degree 0, satisfying $t_\pm(x, \xi) = \pm(x_1 - x_1^0) + |x - x^0|^2 + |\xi/|\xi| - \xi^0|^2$ in a conic neighborhood of z^0 , (iii) $\Theta(t) \in C_0^\infty(\mathbf{R})$ satisfying $0 \leq \Theta(t) \leq 1$ and $\Theta(t) = 1$ for $|t| \leq 1$ and $\Theta(t) = 0$ for $|t| \geq 2$, (iv) $\Theta_h(\xi) = \Theta(|\xi|/h)$ for $h > 0$, (v) $\varphi_i(x, \xi) \in C^\infty(T^*(\mathbf{R}^n) \setminus 0)$ ($i = 1, 2$) being positively homogeneous in ξ of degree 0, (vi) $\varphi_{i,h}(x, \xi) = (1 - \Theta_{h/2}(\xi))\varphi_i(x, \xi)$, (vii) $\zeta(x)$ satisfying $\nabla\zeta(x) \in \mathcal{B}^\infty(\mathbf{R}^n)$ and $\zeta(x) = \theta^0 \cdot (x - x^0) + k|x - x^0|^2$ near x^0 for fixed $x^0 \in \mathbf{R}^n$, $k > 0$ and $\theta^0 \in \Gamma(p(x^0, \cdot), \theta)$, (viii) $P(x, D; \gamma)u(x) = e^{-\gamma\zeta(x)}P(x, D)(e^{\gamma\zeta(x)}u(x))$, (ix) $\langle \xi \rangle_h = (h^2 + |\xi|^2)^{1/2}$ for $h > 0$, (x) $\Lambda_\pm(x, \xi) \equiv \Lambda_{\pm, a, b}(x, \xi) = \{(at_\pm(x, \xi) - b) \log \langle \xi \rangle_h\} \varphi_{2, h/2}(x, \xi)$ for $a \geq 1$ and $b \in \mathbf{R}$, and (xi) $P_{\gamma\Lambda_\pm}(x, D) = e^{-\Lambda_\pm(x, D)}P_\gamma(x, D)e^{\Lambda_\pm(x, D)}$, $P_{\Lambda_\pm}(x, D; \gamma) = e^{-\Lambda_\pm(x, D)}P(x, D; \gamma) \times e^{\Lambda_\pm(x, D)}$, where $e^{\Lambda_\pm(x, D)}$ are pseudodifferential operators with symbols $e^{\Lambda_\pm(x, \xi)}$. Similarly, we define $\tilde{P}(x, D; \gamma)$, $\dots$ and so on.

We begin with the microlocal *a priori* estimates for $\tilde{P}$ at z^0 .

Theorem 2.2. Let $z^0 = (x^0, \xi^0)$ with $|\xi^0| = 1$ be a multiple characteristic point of p . Assume that $P(x, D)$ satisfies [A.I], [A.II] $_{z^0}$ and [A.III] $_{z^0}$. Then there are conic neighborhoods $\mathcal{U}_1$ and $\mathcal{U}_2$ of z^0 , $\varphi_i(x, \xi) \in C^\infty(T^*(\mathbf{R}^n) \setminus 0)$ ($i = 1, 2$) and $\ell_i \in \mathbf{R}$ ($i = 1, 2, 3$) such that $\text{supp } \varphi_2 \subset \mathcal{U}_2$, $\varphi_i(x, \xi) = 1$ in $\mathcal{U}_1$ and for any $a \geq 1$ and $b \in \mathbf{R}$ there are $\gamma_0 \geq 1$ and $C > 0$ satisfying

$$(2.1)_{z^0} \quad \|\langle D \rangle_h^{\ell_1} v\|_{L^2} \leq C \{ \|\langle D \rangle_\gamma^{\ell_2} \tilde{P}_{\Lambda_+}(x, D; \gamma)v\|_{L^2} + \|\langle D \rangle_\gamma^{\ell_3} (1 - \varphi_{1, h}(x, D))v\|_{L^2} \}$$

if $v \in C_0^\infty(\mathbf{R}^n)$ and $h = \gamma \geq \gamma_0$.

An outline of the proof of this theorem will be given in §4. The inequality (2.1) $_{z^0}$ is called the microlocal *a priori* estimate for $\tilde{P}$. We can also obtain the microlocal *a priori* estimates with Λ_+ replaced by Λ_- for ${}^t\tilde{P}$. From Theorem 2.2 we have the following

Lemma 2.3 (Lemma 3.1 in [9]). There is $\varphi(x) \in C_0^\infty(\mathbf{R}^n)$ such that $\varphi(x) = 1$ near x^0 and for any $s \geq 0$ there are $\ell', \ell'' \in \mathbf{R}$ satisfying the following; for any $\ell \in \mathbf{R}$

there are $\gamma_0 \geq 1$ and $C > 0$ such that

$$\begin{aligned} \|\langle D \rangle_\gamma^\ell \varphi(x) u\|_{L^2} &\leq C \{ \|\langle D \rangle_\gamma^{\ell+\ell'} \chi(x_1) \tilde{P}(x, D; \gamma) u\|_{L^2} \\ &\quad + \|\langle D \rangle_\gamma^{\ell-s} u\|_{L^2} + \|\langle D \rangle_\gamma^{\ell+\ell''} \chi(x_1) \Theta_{2h}(D) u\|_{L^2} \} \end{aligned}$$

if $u \in C_0^\infty(\mathbf{R}^n)$, $\text{supp } u \subset \{x_1 > x_1^0 - \delta_0\}$ and $h = \gamma \geq \gamma_0$, where $\chi(x_1) = \Theta((x_1 - x_1^0)/(3\delta_0))$.

Let Ω_0 be a bounded domain in $\mathbf{R}^n$ such that $\nabla \zeta(x) \in \Gamma(\tilde{p}(x, \cdot), \theta)$ for $x \in \overline{\Omega}_0$. Using ellipticity of $\tilde{P}(x, \xi; \gamma)$ for $x \in \overline{\Omega}_0$ and $\xi \in \mathbf{R}^n$ with $|\xi| \leq C\gamma$, we have the following

Lemma 2.4. *For any $t \in \mathbf{R}$ and $\delta_0 > 0$ there is $\delta'_0 > 0$ such that for any $s \geq 0$ there is $\ell' \in \mathbf{R}$ satisfying the following; for any $\ell \in \mathbf{R}$ there are $\gamma_0 \geq 1$ and $C > 0$ such that*

$$\|\langle D \rangle_\gamma^\ell \tau_{\delta'_0}(x_1 - t) u\|_{L^2} \leq C \{ \|\langle D \rangle_\gamma^{\ell+\ell'} \tau_{\delta_0}(x_1 - t) \tilde{P}(x, D; \gamma) u\|_{L^2} + \|\langle D \rangle_\gamma^{\ell-s} u\|_{L^2} \}$$

if $u \in C_0^\infty(\Omega_0 \cap \{x_1 > t - \delta'_0\})$ and $\gamma \geq \gamma_0$, where $\tau_{\delta'}(x_1) = \Theta(2x_1/\delta')$.

In particular, Lemma 2.4 gives a Carleman estimate for $\tilde{P}(x, D)$. Therefore, we have the following

Theorem 2.5 (Theorem 1.2 in [9]). *Assume that P satisfies [A.I], [A.II] $_{z^0}$ and [A.III] $_{z^0}$ for any multiple characteristic point z^0 of p . Then for any $x^0 \in \mathbf{R}^n$ there is a neighborhood U_0 of x^0 such that $x^0 \notin \text{supp } u$ if $u \in \mathcal{D}'$, $\text{supp } \tilde{P}u \cap \overline{V} = \emptyset$ and $\text{supp } u \cap \{x \in \partial V; \zeta(x) \leq \zeta(x^0)\} = \emptyset$ for some neighborhood V of x^0 with $V \subset\subset U_0$.*

The method of sweeping out due to John [7] proves the following uniqueness theorem, noting that $\tilde{K}_x^- \cap \{x_1 \geq t\}$ is bounded for each $x \in \mathbf{R}^n$ and $t \in \mathbf{R}$, where $\tilde{K}_x^\pm$ denote the generalized flows issuing from x for $\tilde{P}$ (see, e.g., [12], [16] and [17]).

Theorem 2.6. *Assume that the hypotheses of Theorem 2.5 are fulfilled. Let $x^0 \in \mathbf{R}^n$ and $t \in \mathbf{R}$. Then $x^0 \notin \text{supp } u$ if $u \in \mathcal{D}'$, $\text{supp } u \subset \{x_1 \geq t\}$ and $\tilde{P}u = 0$ near $\tilde{K}_{x^0}^-$.*

Note that $-\theta \in \Gamma(\tilde{p}(x, \cdot), -\theta)$ for $x \in \mathbf{R}^n$. By Theorem 2.2 we obtain the microlocal *a priori* estimates with Λ_+ replaced by Λ_- for ${}^t\tilde{P}$. Applying the same argument as in the proof of Lemma 2.4, we can see that for any $t \in \mathbf{R}$ and $\delta_0 > 0$ there is $\delta'_0 > 0$

such that for any $s \geq 0$ there is $\ell' \in \mathbf{R}$ satisfying the following; for any $\ell \in \mathbf{R}$ there are $\gamma_0 \geq 1$ and $C > 0$ such that

$$\begin{aligned} & \| \langle D \rangle_\gamma^\ell \tau_{\delta'_0}(x_1 - t) u \|_{L^2} \\ & \leq C \{ \| \langle D \rangle_\gamma^{\ell+\ell'} \tau_{\delta_0}(x_1 - t)^t \tilde{P}(x, D + i\gamma\theta) u \|_{L^2} + \| \langle D \rangle_\gamma^{\ell-s} u \|_{L^2} \} \end{aligned}$$

if $u \in C_0^\infty(\Omega_0 \cap \{x_1 < t + \delta'_0\})$ and $\gamma \geq \gamma_0$. Define

$$\|u\|_{+, \gamma, (m, s)} = \inf \{ \| \langle D \rangle_\gamma^m \langle D' \rangle_\gamma^s U \|; U|_X = u \text{ and } U \in \mathcal{H}_{(m, s)}(\mathbf{R}^n) \},$$

where $X = \{x \in \mathbf{R}^n; x_1 > -6\delta\}$. By the same argument as in [9] we can see that there is $\ell' \in \mathbf{R}$ such that for any $\ell \in \mathbf{R}$ there are $\gamma_0 \geq 1$ and $C > 0$ satisfying

$$\|u\|_{+, \gamma, (\ell, 0)} \leq C \{ \| {}^t \tilde{P}(x, D + i\gamma\theta) u \|_{+, \gamma, (\ell+\ell', 0)} + \| {}^t \tilde{P}(x, D + i\gamma\theta) u \|_{+, \gamma, (0, \ell+\ell')} \}$$

if $u \in C_0^\infty(\Omega_0)$ and $\gamma \geq \gamma_0$. By the Hahn-Banach theorem and Theorem 2.6 we can prove Theorem 1.1. For the detail of the proof we refer to [11].

3. Construction of the inverse of $e^\Lambda(x, D)$

Let $t_k(x, \xi)$ ($1 \leq k \leq d$) be real valued bounded mesurable functions in $\mathbf{R}^{2n}$ satisfying

$$(3.1) \quad |t_k(x + y, \xi + \eta) - t_k(x, \xi)| \leq C \{ |y| + |\xi|^{-1} |\eta| \}$$

for $(x, \xi), (y, \eta) \in \mathbf{R}^{2n}$ and $|\eta| \leq |\xi|/2$. Put $t_{k,h}(x, \xi) = (1 - \Theta_h(\xi)) t_k(x, \xi)$ ($1 \leq k \leq d$), where $\Theta_h(\xi)$ was given in §2. Then $t_{k,h}$ satisfies

$$|t_{k,h}(x + y, \xi + \eta) - t_{k,h}(x, \xi)| \leq C \{ |y| + \langle \xi \rangle_h^{-1} |\eta| \}$$

for $(x, \xi), (y, \eta) \in \mathbf{R}^{2n}$, where $\langle \xi \rangle_h = (h^2 + |\xi|^2)^{1/2}$. Put, with parameters $h \geq 1$ and $N \geq d$,

$$n_k(x, \xi) = \{ t_{k,h}(x, \xi)^2 \langle \xi \rangle_h + N \}^{1/2}, \quad T_k(x, \xi) = n_k(x, \xi) + t_{k,h}(x, \xi) \langle \xi \rangle_h^{1/2},$$

$$\varphi_k(x, \xi) = \langle \xi \rangle_h^{-1/2} n_k(x, \xi), \quad \Psi_k(x, \xi) = \varphi_k(x, \xi) \langle \xi \rangle_h,$$

$$\varphi(x, \xi)^{-1} = \sum_{k=1}^d \varphi_k(x, \xi)^{-1}, \quad \Psi(x, \xi) = \varphi(x, \xi) \langle \xi \rangle_h,$$

$$g_{(x, \xi)}(y, \eta) = \varphi(x, \xi)^{-2} |y|^2 + \Psi(x, \xi)^{-2} |\eta|^2,$$

$$g_{(x, \xi)}^\sigma(y, \eta) = \Psi(x, \xi)^2 |y|^2 + \varphi(x, \xi)^2 |\eta|^2.$$

Following [4] we say that a positive real valued function $m(x, \xi)$ defined in $\mathbf{R}^{2n}$ is g continuous, if there are positive constants c_0 and C such that if $g_{(x, \xi)}(y, \eta) \leq c_0$,

$$(3.2) \quad C^{-1}m(x, \xi) \leq m(x + y, \xi + \eta) \leq Cm(x, \xi),$$

and that a g continuous function $m(x, \xi)$ is σ, g temperate, if there are $C > 0$ and $\ell \in \mathbf{R}$ such that

$$m(x + y, \xi + \eta) \leq Cm(x, \xi)(1 + g_{(x, \xi)}^\sigma(y, \eta))^\ell$$

for $(x, \xi), (y, \eta) \in \mathbf{R}^{2n}$. For simplicity we denote $A \sim B$ if A/B and B/A are bounded. For example $m(x + y, \xi + \eta) \sim m(x, \xi)$ means (3.2). By giving a series of lemmas without proof we want to show an outline of the construction of the inverse of $e^\Lambda(x, D)$. For the detail of the proof we refer to [11].

Lemma 3.1. Assume that (3.1) is valid and $h \geq N \geq d$. Then $\langle \xi \rangle_h$ and $n_k(x, \xi)$ are σ, g temperate.

It follows from Lemma 3.1 that φ_k , Ψ_k , φ and Ψ are σ, g temperate. Hence we have the following

Lemma 3.2. Assume that (3.1) is valid and $h \geq N \geq d$. Then the metric $g_{(x, \xi)}(y, \eta)$ is σ temperate.

Lemma 3.3. Assume that (3.1) is valid and $h \geq N \geq d$. Then, (i)

$$\varphi(x + y, \xi + \eta)^{-1} \leq C\varphi(x, \xi)^{-1}\{1 + N^{-2}g_{(x, \xi)}^\sigma(y, \eta)\}^{1/4}$$

for $(x, \xi), (y, \eta) \in \mathbf{R}^{2n}$. (ii)

$$\varphi(x + y, \xi + \eta) \leq C\varphi(x, \xi)\{1 + N^{-2}g_{(x, \xi)}^\sigma(y, \eta)\}^{1/2}$$

for $(x, \xi), (y, \eta) \in \mathbf{R}^{2n}$. (iii) Ψ also satisfies (i) and (ii), replacing φ by Ψ .

Lemma 3.4. Assume that (3.1) is valid and $h \geq N \geq d$. Then $T_k(x, \xi) = n_k(x, \xi) + t_{k, h}(x, \xi)\langle \xi \rangle_h^{1/2}$ ($1 \leq k \leq d$) are σ, g temperate and satisfy

$$T_k(x + y, \xi + \eta)^{\pm 1} \leq CT_k(x, \xi)^{\pm 1}(1 + N^{-2}g_{(x, \xi)}^\sigma(y, \eta))$$

for $(x, \xi), (y, \eta) \in \mathbf{R}^{2n}$.

We have to approximate φ and Ψ by functions in some Gevrey class. Let $\rho(z)$ be a function in $C_0^\infty(\mathbf{R}^{2n})$ satisfying $\rho(z) \geq 0$, $\rho(z) = 0$ for $|z| \geq c_0$, $\int \rho(z) dz = 1$ and

$$|D_z^\alpha \rho(z)| \leq CA^{|\alpha|} |\alpha|!^\kappa \quad \text{for } z \in \mathbf{R}^{2n},$$

where $\kappa > 1$ and $A > 0$. Define

$$\begin{aligned} & \tilde{\varphi}(x, \xi) \\ &= \int \int \rho((x-y)\varphi(y, \eta)^{-1}, (\xi-\eta)\Psi(y, \eta)^{-1}) (\varphi\Psi)(y, \eta)^{-n} \varphi(y, \eta) dy d\eta, \\ & \tilde{\Psi}(x, \xi) \\ &= \int \int \rho((x-y)\varphi(y, \eta)^{-1}, (\xi-\eta)\Psi(y, \eta)^{-1}) (\varphi\Psi)(y, \eta)^{-n} \Psi(y, \eta) dy d\eta, \\ (3.3) \quad & T(x, \xi) \\ &= \int \int \rho((x-y)\tilde{\varphi}(x, \xi)^{-1}, (\xi-\eta)\tilde{\Psi}(x, \xi)^{-1}) (\tilde{\varphi}\tilde{\Psi})(x, \xi)^{-n} \tilde{T}(y, \eta) dy d\eta, \end{aligned}$$

where $\tilde{T}(y, \eta) = \prod_{k=1}^d T_k(y, \eta)$.

Lemma 3.5. *There is a positive constant C such that (i) $C^{-1}\varphi(x, \xi) \leq \tilde{\varphi}(x, \xi) \leq C\varphi(x, \xi)$, $C^{-1}\Psi(x, \xi) \leq \tilde{\Psi}(x, \xi) \leq C\Psi(x, \xi)$, and $C^{-1}\tilde{T}(x, \xi) \leq T(x, \xi) \leq C\tilde{T}(x, \xi)$ for $(x, \xi) \in \mathbf{R}^{2n}$. (ii)*

$$\begin{aligned} & |\tilde{\varphi}_{(\beta)}^{(\alpha)}(x, \xi)| / \tilde{\varphi}(x, \xi) + |\tilde{\Psi}_{(\beta)}^{(\alpha)}(x, \xi)| / \tilde{\Psi}(x, \xi) + |T_{(\beta)}^{(\alpha)}(x, \xi)| / T(x, \xi) \\ & \leq C^{|\alpha+\beta|+1} |\alpha + \beta|!^\kappa \varphi(x, \xi)^{-|\beta|} \Psi(x, \xi)^{-|\alpha|} \end{aligned}$$

for $(x, \xi) \in \mathbf{R}^{2n}$. (iii) $\tilde{\varphi}$ and $\tilde{\Psi}$ satisfy (i), (ii) and (iii) in Lemma 3.3, replacing φ and Ψ by $\tilde{\varphi}$ and $\tilde{\Psi}$, respectively.

We note that it follows from Lemma 3.4 and (i) of Lemma 3.5 that T is σ, g temperate. For $M > 0$ we define

$$(3.4) \quad \Lambda(x, \xi) = \Lambda(x, \xi; M, N, h) = M \log T(x, \xi),$$

where T is given in (3.3). Denote

$$\omega_\beta^\alpha(\Lambda; x, \xi) = e^{-\Lambda(x, \xi)} D_x^\beta \partial_\xi^\alpha e^{\Lambda(x, \xi)}.$$

Then we have the following

Lemma 3.6. *There is a positive constant C , independent of M and N , such that*

(i)

$$|\Lambda_{(\beta)}^{(\alpha)}(x, \xi)| \leq MC^{|\alpha+\beta|+1} |\alpha + \beta|!^\kappa \varphi(x, \xi)^{-|\beta|} \Psi(x, \xi)^{-|\alpha|}$$

for $(x, \xi) \in \mathbf{R}^{2n}$ and $|\alpha + \beta| \geq 1$, and (ii)

$$|D_x^\delta \partial_\xi^\gamma \omega_\beta^\alpha(\Lambda; x, \xi)| \leq C(C\varphi^{-1})^{|\beta+\delta|} (C\Psi^{-1})^{|\alpha+\gamma|} \sum_{j=0}^{|\alpha+\beta|} (CM)^{|\alpha+\beta|-j} (|\gamma + \delta| + j)!^\kappa$$

for $(x, \xi) \in \mathbf{R}^{2n}$.

We obtain from (ii) of Lemma 3.6

$$(3.5) \quad |D_x^\delta \partial_\xi^\gamma \omega_\beta^\alpha(\Lambda; x, \xi)| \leq C^{|\alpha+\beta+\gamma+\delta|+1} |\gamma + \delta|!^\kappa (|\alpha + \beta|^\kappa + M)^{|\alpha+\beta|} \\ \times \varphi(x, \xi)^{-|\beta+\delta|} \Psi(x, \xi)^{-|\alpha+\gamma|} \quad \text{for } (x, \xi) \in \mathbf{R}^{2n},$$

noting that $M^{|\alpha+\beta|-j} (|\gamma + \delta| + j)!^\kappa \leq C_1^{|\alpha+\beta|+|\gamma+\delta|} |\gamma + \delta|!^\kappa (j^\kappa + M)^{|\alpha+\beta|}$ for $j \leq |\alpha + \beta|$. Following [4], for a positive function $m(x, \xi)$ and the metric $g = \varphi^{-2}|dx|^2 + \Psi^{-2}|d\xi|^2$ we define the symbol class $S(m, g)$ of pseudodifferential operators by the set of all $a(x, \xi) \in C^\infty(\mathbf{R}^{2n})$ satisfying

$$|a_{(\beta)}^{(\alpha)}(x, \xi)| \leq C_{\alpha\beta} m(x, \xi) \varphi(x, \xi)^{-|\beta|} \Psi(x, \xi)^{-|\alpha|} \quad \text{for } (x, \xi) \in \mathbf{R}^{2n}.$$

Denote by $e^\Lambda(x, D)$ the pseudodifferential operator whose symbol is $e^{\Lambda(x, \xi)}$ and by ${}^R e^{-\Lambda}(x, D)$ the reversed operator of $e^{-\Lambda}(x, D)$, i.e.,

$${}^R e^{-\Lambda}(x, D)u(x) = (2\pi)^{-n} \int \int e^{i(x-y) \cdot \xi - \Lambda(y, \xi)} u(y) dy d\xi$$

for $u \in C_0^\infty(\mathbf{R}^n)$. Put $q(x, D) = {}^R e^{-\Lambda}(x, D)e^\Lambda(x, D)$. Then we can express

$$q(x, \xi) = (2\pi)^{-n} os- \int \int e^{-iy \cdot \eta - \Lambda(x+y, \xi+\eta) + \Lambda(x+y, \xi)} dy d\eta,$$

where $os- \int \int$ means an oscillatory integral. By Taylor's expansion we have

$$(3.6) \quad D_x^\beta \partial_\xi^\gamma q(x, \xi) = \sum_{|\alpha| < \ell} D_x^{\alpha+\beta} \partial_\xi^\gamma \omega^\alpha(-\Lambda; x, \xi) / \alpha! + D_x^\beta \partial_\xi^\gamma q_\ell(x, \xi),$$

where $\omega^\alpha(-\Lambda; x, \xi) = e^{\Lambda(x, \xi)} \partial_\xi^\alpha e^{-\Lambda(x, \xi)}$ and

(3.7)

$$\begin{aligned} q_{\ell(\beta)}^{(\gamma)}(x, \xi) &\equiv D_x^\beta \partial_\xi^\gamma q_\ell(x, \xi) \\ &= (2\pi)^{-n} \text{os-} \int \int e^{-iy \cdot \eta} \sum_{|\alpha|=\ell} \alpha!^{-1} \\ &\quad \times \int_0^1 \ell(1-\theta)^{\ell-1} D_x^{\alpha+\beta} \partial_\xi^\gamma \{e^{\Lambda_1}(\partial_\xi^\alpha e^{-\Lambda_2})\} dy d\eta d\theta, \\ &= (2\pi)^{-n} \text{os-} \int \int e^{-iy \cdot \eta} \int_0^1 (1-\theta)^{\ell-1} e^{\Lambda_1-\Lambda_2} \omega_{\ell\beta}^\gamma(x+y, \xi, \theta\eta) dy d\eta d\theta, \end{aligned}$$

where $\Lambda_1 = \Lambda(x+y, \xi)$, $\Lambda_2 = \Lambda(x+y, \xi+\theta\eta)$ and

$$\begin{aligned} (3.8) \quad \omega_{\ell\beta}^\gamma(x+y, \xi, \theta\eta) &= \ell \sum_{|\alpha|=\ell} \alpha!^{-1} e^{\Lambda_2-\Lambda_1} D_x^{\alpha+\beta} \partial_\xi^\gamma \{e^{\Lambda_1}(\partial_\xi^\alpha e^{-\Lambda_2})\}. \\ &= \ell \sum_{|\alpha|=\ell} \sum_{\gamma' \leq \gamma, \beta' \leq \alpha+\beta} \alpha!^{-1} \binom{\alpha+\beta}{\alpha'} \binom{\gamma}{\gamma'} \\ &\quad \times \omega_{\beta'}^{\gamma'}(\Lambda; x+y, \xi) \omega_{\alpha+\beta-\beta'}^{\alpha+\gamma-\gamma'}(-\Lambda; x+y, \xi+\theta\eta). \end{aligned}$$

Denote

$$\begin{aligned} g_1 &= g_{(x+y, \xi)}(0, \eta), \quad \varphi_1 = \varphi(x+y, \xi), \quad \varphi_2 = \varphi(x+y, \xi+\theta\eta), \\ \Psi_1 &= \Psi(x+y, \xi), \quad \Psi_2 = \Psi(x+y, \xi+\theta\eta), \quad \omega_\beta^\alpha(\Lambda_1) = \omega_\beta^\alpha(\Lambda; x+y, \xi), \\ &\quad \omega_\beta^\alpha(-\Lambda_2) = \omega_\beta^\alpha(-\Lambda; x+y, \xi+\theta\eta). \end{aligned}$$

Then we have the following

Lemma 3.7. *There are $C > 0$ and $c_0 > 0$, independent of N and M , such that*

$$\begin{aligned} (3.9) \quad |D_{x,y}^\delta \partial_{\xi,\eta}^\rho \omega_{\ell\beta}^\gamma(x+y, \xi, \theta\eta)| \\ \leq C^{|\beta+\gamma|+|\delta+\rho|+1} (|\beta+\gamma|+|\delta+\rho|)!^\kappa \varphi_1^{-|\beta|-|\delta|} \Psi_1^{-|\gamma|-|\rho|} \{CM^{2\kappa-1}N^{-1}\}^\ell, \end{aligned}$$

for $\ell = M$ if $g_1 \leq c_0$.

By virtue of (3.5), noting that $\varphi\Psi \geq cN$, we can estimate $\omega^\alpha(-\Lambda)$ in (3.6) as follows:

$$\begin{aligned} (3.10) \quad |D_x^\beta \partial_\xi^\gamma D_x^\alpha \omega^\alpha(-\Lambda)|/\alpha! \\ \leq \alpha!^{-1} C^{|\gamma+\beta+\alpha|} |\gamma+\beta+\alpha|!^\kappa (|\alpha|^\kappa + M)^{|\alpha|} \varphi^{-|\beta+\alpha|} \Psi^{-|\alpha+\gamma|} \\ \leq C_1^{\beta+\gamma+\alpha} |\beta+\gamma|!^\kappa \{|\alpha|^{(2\kappa-1)} + |\alpha|^{\kappa-1} M\}^{|\alpha|} \varphi^{-|\beta|} \Psi^{-|\gamma|} N^{-|\alpha|} \\ \leq C_1^{\gamma+\beta} |\gamma+\beta|!^\kappa \{C_1 M^{2\kappa-1} N^{-1}\}^{|\alpha|} \varphi^{-|\beta|} \Psi^{-|\gamma|} \end{aligned}$$

for $|\alpha| \leq \ell = M$. Let us prove that

$$(3.11) \quad |q_{\ell(\beta)}^{(\alpha)}(x, \xi)| \leq C_{\alpha\beta} \{CM^{2\kappa-1}N^{-1}\}^\ell \varphi^{-|\beta|} \Psi^{-|\alpha|}$$

for $\ell = M$, where $C_{\alpha\beta}$ is independent of M and N , and C is independent of M , N , α and β . Let $\chi(t)$ be in $C_0^\infty(\mathbf{R})$ such that $\chi(t) = 1$ for $|t| \leq 1/2$ and $\chi(t) = 0$ for $|t| \geq 1$, and

$$|D_t^j \chi(t)| \leq CA^j j!^\kappa \quad \text{for any } j \text{ and } t \in \mathbf{R},$$

where $A > 0$. In brief we denote

$$\begin{aligned} z_1 &= (x + y, \xi), \quad z_2 = (x + y, \xi + \theta\eta), \\ \varphi_i &= \varphi(z_i), \quad \Psi_i = \Psi(z_i), \quad \tilde{\varphi}_i = \tilde{\varphi}(z_i), \quad \tilde{\Psi}_i = \tilde{\Psi}(z_i) \quad (i = 1, 2), \\ \chi_1 &= \chi(c_0^{-1} \tilde{\Psi}(z_1)^{-2} |\eta|^2). \end{aligned}$$

Then we can write from (3.7)

$$(3.12) \quad \begin{aligned} q_{\ell(\beta)}^{(\gamma)}(x, \xi) &= (2\pi)^{-n} \int \int \int_0^1 e^{-iy \cdot \eta} \chi_1 (1 - \theta)^{\ell-1} e^{\Lambda_1 - \Lambda_2} \omega_{\ell\beta}^\gamma dy d\eta d\theta \\ &+ (2\pi)^{-n} \int \int \int_0^1 e^{-iy \cdot \eta} (1 - \chi_1) (1 - \theta)^{\ell-1} e^{\Lambda_1 - \Lambda_2} \omega_{\ell\beta}^\gamma dy d\eta d\theta \equiv I_1 + I_2, \end{aligned}$$

where $\omega_{\ell\beta}^\gamma = \omega_{\ell\beta}^\gamma(x + y, \xi, \theta\eta)$ is given in (3.8).

Lemma 3.8. *There is $C > 0$ such that*

$$|D_y^\beta \partial_\eta^\alpha \chi_1| \leq C^{|\alpha+\beta|+1} |\alpha + \beta|!^\kappa \varphi_1^{-|\beta|} \Psi_1^{-|\alpha|}.$$

Now we shall estimate I_1 in (3.12). Noting that $\varphi_1 \sim \varphi_2$, $\Psi_1 \sim \Psi_2$, $e^{\Lambda_1 - \Lambda_2} \leq C^M$ and $|\eta| \leq \langle \xi \rangle_h / 2$ on $\text{supp } \chi_1$, we can estimate, by use of (3.5), (3.9) and (ii) of Lemma 3.3,

$$(3.13) \quad \begin{aligned} &|(\partial_\eta^\rho \{e^{\Lambda_1 - \Lambda_2} \chi_1 \omega_{\ell\beta}^\gamma(x + y, \xi, \theta\eta)\})| \\ &\leq \sum_{\rho'} \binom{\rho}{\rho'} |(\partial_\eta^{\rho'} e^{\Lambda_1 - \Lambda_2}) \partial_\eta^{\rho - \rho'} \{\chi_1 \omega_{\ell\beta}^\gamma(x + y, \xi, \theta\eta)\}| \\ &\leq \sum_{\rho'} \binom{\rho}{\rho'} C^M C^{|\rho'|} (|\rho'|^\kappa + M)^{|\rho'|} C^{|\rho - \rho' + \beta + \gamma|} |\rho - \rho' + \beta + \gamma|!^\kappa \\ &\quad \times (CM^{2\kappa-1}N^{-1})^\ell \varphi_1^{-|\beta|} \Psi_1^{-|\gamma + \rho|} \\ &\leq C_1^{|\rho + \beta + \gamma|} |\rho + \beta + \gamma|!^\kappa (C_1 M^{2\kappa-1} N^{-1})^\ell \varphi_1^{-|\beta|} \Psi_1^{-|\rho + \gamma|} \\ &\leq C_{\rho\beta\gamma} (C_1 M^{2\kappa-1} N^{-1})^\ell \Psi_1^{-|\rho|} \varphi^{-|\beta|} \Psi^{-|\gamma|} (1 + g_{(x+y, \xi)}^\sigma(y, 0))^{|\beta + \gamma|/2} \end{aligned}$$

for $\ell = M$. Since $g_{(x+y, \xi)}^\sigma(y, 0) = \Psi_1^2 |y|^2$, we can estimate from (3.13)

$$\begin{aligned}
 (3.14) \quad |I_1| &= |(2\pi)^{-n} \int \int \int_0^1 e^{-iy \cdot \eta} (1 - \theta)^{\ell-1} (1 - \Psi_1^2 \Delta_\eta)^p \\
 &\quad \times [(1 + \Psi_1^2 |y|^2)^{-p} \{\chi_1 e^{\Lambda_1 - \Lambda_2} \omega_{\ell\beta}^\gamma(x + y, \xi, \theta\eta)\}] dy d\eta d\theta| \\
 &\leq C_{p\beta\gamma} \{CM^{2\kappa-1} N^{-1}\}^\ell \varphi^{-|\beta|} \Psi^{-|\gamma|} \\
 &\quad \times \int \int \int_{\text{supp } \chi_1} (1 + \Psi_1^2 |y|^2)^{-p+|\beta+\gamma|/2} dy d\eta d\theta
 \end{aligned}$$

for $p \geq |\beta + \gamma|/2$. Since $|\eta| \leq C\Psi_1$ on $\text{supp } \chi_1$, taking account of Lemma 3.3 we get

$$\begin{aligned}
 &\int \int \int_{\text{supp } \chi_1} (1 + \Psi_1^2 |y|^2)^{-p+|\beta+\gamma|/2} dy d\eta d\theta \\
 &\leq C_n \int (1 + \Psi_1^2 |y|^2)^{-p+|\beta+\gamma|/2} \Psi_1^n dy \\
 &\leq C_{np} \int \{1 + \Psi^2 |y|^2 (1 + N^{-2} \Psi^2 |y|^2)^{-1}\}^{-p+|\beta+\gamma|/2} \\
 &\quad \times \Psi^n (1 + N^{-2} \Psi^2 |y|^2)^{n/2} dy \\
 &\leq C_1 \int (1 + \Psi |y|)^{-p+|\beta+\gamma|/2+n} \Psi^n dy \leq C_2,
 \end{aligned}$$

if we choose $p \geq |\beta + \gamma|/2 + 2n + 1$. Thus we obtain from (3.14)

$$|I_1| \leq C_{\beta\gamma} (CM^{2\kappa-1} N^{-1})^\ell \varphi^{-|\beta|} \Psi^{-|\gamma|} \quad \text{for } \ell = M.$$

Next we shall estimate I_2 in (3.12). From (3.7) and (3.8) we can write

$$\begin{aligned}
 (3.15) \quad (1 - \chi_1) e^{\Lambda_1 - \Lambda_2} \omega_{\ell\beta}^\gamma &= \ell \sum_{\alpha=\ell} \alpha!^{-1} (1 - \chi_1) \{e^{\Lambda_1} (e^{-\Lambda_2})^{(\alpha)}\}_{(\alpha+\beta)}^{(\gamma)} \\
 &= \ell \sum_{|\alpha|=\ell} \alpha!^{-1} \sum_{\beta' \leq \alpha+\beta, \gamma' \leq \gamma} \binom{\alpha+\beta}{\beta'} \binom{\gamma}{\gamma'} (1 - \chi_1) (e^{\Lambda_1})_{(\beta')}^{(\gamma')} \\
 &\quad \times (e^{-\Lambda_2})_{(\alpha+\beta-\beta')}^{(\alpha+\gamma-\gamma')},
 \end{aligned}$$

where $w_{(\beta)}^{(\alpha)}$ denotes $D_x^\beta \partial_\xi^\alpha w(x, \xi)$. For simplicity we put

$$r_1 = (1 - \chi_1) (e^{\Lambda_1})_{(\beta')}^{(\gamma')}, \quad r_2 = (e^{-\Lambda_2})_{(\alpha+\beta-\beta')}^{(\alpha+\gamma-\gamma')}.$$

Then it follows from (3.5) and Lemma 3.8 that

$$(3.16) \quad |D_y^\delta \partial_\eta^\rho r_1| \leq C_{\rho\beta'\gamma'} C^M \{C(|\delta|^\kappa + M)\}^{|\delta|} e^{\Lambda_1} \varphi_1^{-|\beta'+\delta|} \Psi_1^{-|\gamma'+\rho|}.$$

Similarly, we can estimate

$$(3.17) \quad |D_y^\delta \partial_\eta^\rho r_2| \leq C_{\rho\beta\gamma} C^M \{C(|2\alpha + \delta|^\kappa + M)\}^{2\alpha + \delta} e^{-\Lambda_2} \varphi_2^{-|\alpha + \beta - \beta' + \delta|} \Psi_2^{-|\alpha + \gamma - \gamma' + \rho|},$$

Therefore, we obtain from (3.15)-(3.17)

$$\begin{aligned} |I_2| &= |(2\pi)^{-n} \int \int \int_0^1 e^{-iy \cdot \eta} \{(1 - \theta)^{\ell-1} (1 - \chi_1)\} e^{\Lambda_1 - \Lambda_2} \omega_{\ell\beta}^\gamma dy d\eta d\theta| \\ &= |(2\pi)^{-n} \int \int \int_0^1 e^{-iy \cdot \eta} |\eta|^{-2s} \Delta_y^s [\{(1 - \theta)^{\ell-1} (1 - \chi_1) e^{\Lambda_1 - \Lambda_2}\} \\ &\quad \times \omega_{\ell\beta}^\gamma] dy d\eta d\theta| \\ &\leq (2\pi)^{-n} \ell \sum_{|\alpha|=\ell} \alpha!^{-1} \sum_{\beta' \leq \alpha + \beta, \gamma' \leq \gamma} \binom{\alpha + \beta}{\beta'} \binom{\gamma}{\gamma'} \sum_{|\delta|=s} s!^{-1} \\ &\quad \times \sum_{\delta' \leq 2\delta} \binom{2\delta}{\delta'} \left| \int \int \int_0^1 e^{-iy \cdot \eta} (1 - \Psi_1^2 \Delta_\eta)^p (1 + \Psi_1^2 |y|^2)^{-p} |\eta|^{-2s} \right. \\ &\quad \times \left. \{(1 - \chi_1) (e^{\Lambda_1})_{(\beta')}^{(\gamma')}\}_{(\delta')} \{(e^{-\Lambda_2})_{(\alpha + \beta - \beta')}^{(\alpha + \gamma - \gamma')}\}_{(2\delta - \delta')} (1 - \theta)^{\ell-1} dy d\eta d\theta \right|. \end{aligned}$$

By (3.16) and (3.17), a simple calculation yields

$$\begin{aligned} &|(1 - \Psi_1^2 \Delta_\eta)^p (1 + \Psi_1^2 |y|^2)^{-p} |\eta|^{-2s} D_y^{\delta'} r_1 D_y^{2\delta - \delta'} r_2| \\ &\leq C_{p\beta\gamma} C^M \{C(|2\alpha| + 2s)^\kappa + M\}^{2\alpha + 2s} (1 + \Psi_1^2 |y|^2)^{-p} (1 + \Psi_1/\Psi_2)^{2p} \\ &\quad \times e^{\Lambda_1 - \Lambda_2} |\eta|^{-2s} \varphi_1^{-|\beta' + \delta'|} \Psi_1^{-|\gamma'|} \varphi_2^{-|\alpha + \beta - \beta' + 2\delta - \delta'|} \Psi_2^{-|\alpha + \gamma - \gamma'|}, \end{aligned}$$

where $\beta' \leq \alpha + \beta$, $\gamma' \leq \gamma$, $\delta' \leq 2\delta$ and $|\delta| = s$, since $|\eta|^2 \geq c_0 \Psi_1^2/2$ on $\text{supp } (1 - \chi_1)$.

It follows from Lemma 3.4 that $e^{\Lambda_1 - \Lambda_2} \leq \{C(1 + N^{-2} g_1^\sigma)\}^{Md}$, where $g_1^\sigma = g_{z_1}^\sigma(0, \eta)$.

Therefore, by Lemma 3.3 we have

$$\begin{aligned} &e^{\Lambda_1 - \Lambda_2} \varphi_1^{-|\beta' + \delta'|} \Psi_1^{-|\gamma'|} \varphi_2^{-|\alpha + \beta - \beta' + 2\delta - \delta'|} \Psi_2^{-|\alpha + \gamma - \gamma'|} (1 + \Psi_1/\Psi_2)^{2p} \\ &\leq C^{2\alpha + \beta - \beta' + 2\delta - \delta' + \gamma - \gamma' + Md + p} \varphi_1^{-|\alpha + \beta| - 2s} \Psi_1^{-|\alpha + \gamma|} \\ &\quad \times (1 + N^{-2} g_1^\sigma)^{(|2\alpha + \beta - \beta' + \gamma - \gamma'| + 2s)/4 + Md + p/2} \\ &\leq C_{p\beta\gamma} C_1^{s+M} \{1 + N^{-2} g_1^\sigma\}^{(2\ell + 2s + |\beta + \gamma|)/4 + Md + p/2} (\varphi_1 \Psi_1)^{-\ell} \varphi_1^{-2s} \varphi_1^{-|\beta|} \Psi_1^{-|\gamma|} \\ &\quad \times \{1 + N^{-2} g_{(x+y, \xi)}^\sigma(y, 0)\}^{|\beta + \gamma|/2}, \end{aligned}$$

where $\beta' \leq \alpha + \beta$, $\gamma' \leq \gamma$, $\delta' \leq 2\delta$, $|\alpha| = \ell = M$ and $|\delta| = s$. Noting that

$$g_1^\sigma = \varphi_1^2 |\eta|^2 = (\varphi_1 \Psi_1)^2 g_1 \geq c_0 d^2 N^2/2 \quad \text{for } g_1 \geq c_0/2,$$

we have by virtue of Lemma 3.3

$$\begin{aligned}
|I_2| &\leq C_{p\beta\gamma} C_2^{s+M} M^{2\kappa(s+M)} \int \int \int_{\text{supp}(1-\chi_1)} (1 + \Psi_1^2 |y|^2)^{-p} \\
&\quad \times (\varphi_1 |\eta|)^{-2s} \{N^{-2} g_1^\sigma\}^{(2\ell+|\beta+\gamma|+2s)/4+Mdp/2} \\
&\quad \times \varphi^{-|\beta|} \Psi^{-|\gamma|} (\varphi_1 \Psi_1)^{-\ell} (1 + \Psi_1^2 |y|^2)^{|\beta+\gamma|/2} dy d\eta d\theta \\
&\leq C_{p\beta\gamma} C_3^\ell N^{-(2s-n-1)-\ell} M^{2\kappa(s+M)} \varphi^{-|\beta|} \Psi^{-|\gamma|},
\end{aligned}$$

if we choose $p \geq |\gamma+\beta|2+n+1$ and $2Md+p+|\gamma+\beta|/2+\ell+n+1 \leq s \leq CM+|\beta+\gamma|$.

In fact, noting that $(1 + \Psi_1 |y|)^{-1} \leq C(1 + \Psi_1 |y|)(1 + \Psi |y|)^{-1}$, Lemma 3.3 yields

$$\begin{aligned}
&\int \int \int_{g_1 \geq c_0/2} (1 + \Psi_1^2 |y|^2)^{-p} (\varphi_1 |\eta|)^{-(n+1)} (1 + \Psi_1 |y|)^{|\beta+\gamma|} dy d\eta d\theta \\
&\leq C_{\beta\gamma} \int \int \int (1 + \Psi |y|)^{-n-1} (1 + \varphi |\eta|)^{-n-1} (1 + \Psi_1 |y|)^{-2p+2n+2+|\beta+\gamma|} dy d\eta \\
&\leq C'_{\beta\gamma} \int \int (1 + \Psi |y|)^{-n-1} (1 + \varphi |\eta|)^{-(n+1)} dy d\eta \leq C''_{\beta\gamma}.
\end{aligned}$$

Choose s , N and ε so that

$$(3.18) \quad \begin{cases} N \geq M^{2-\varepsilon}, & 2 - \varepsilon > \kappa, \\ (2 - \varepsilon)(2s - n - 1) \geq 2\kappa s + M. \end{cases}$$

Then we have

$$|I_2| \leq C_{\beta\gamma} (CM^{2\kappa-1} N^{-1})^\ell \varphi^{-|\beta|} \Psi^{-|\gamma|}$$

if

$$(3.19) \quad 1 < \kappa < 2.$$

Note that we can choose $\varepsilon > 0$ so that $0 < \varepsilon < 2 - \kappa$ if (3.19) is valid. Therefore, we can choose $s > 0$ so that (3.18) is valid. Thus we have just proved (3.11). Moreover if $CM^{2\kappa-1} N^{-1} \leq 1$, i.e., $0 < \varepsilon < 3 - 2\kappa$, $1 < \kappa < 3/2$ and $M \gg 1$, then from (3.6), (3.10) and (3.11) we have the following

Proposition 3.9. Assume $1 < \kappa < 3/2$, $0 < \varepsilon < 3 - 2\kappa$ and $h \geq N \geq M^{2-\varepsilon} \gg 1$ are valid. Then $q(x, D) = {}^R e^{-\Lambda}(x, D) e^\Lambda(x, D)$ satisfies

$$|\{q(x, \xi) - \sum_{|\alpha| < k} D_x^\alpha \omega^\alpha(-\Lambda; x, \xi) \alpha!\}_{(\beta)}^{(\gamma)}| \leq C_{k\beta\gamma} (M^{2\kappa-1} N^{-1})^k \varphi^{-|\beta|} \Psi^{-|\gamma|},$$

for any $k \geq 0$ with $k \leq M$, where $\omega^\alpha(-\Lambda; x, \xi) = e^{\Lambda(x, \xi)} \partial_\xi^\alpha e^{-\Lambda(x, \xi)}$ and $C_{k\beta\gamma}$ is independent of M, N and h .

Repeating the same argument as in Proposition 3.9, we can prove the following

Proposition 3.10. Assume that the same conditions in Proposition 3.9 are satisfied. Let $m(x, \xi)$ be a positive function defined in $T^*(\mathbb{R}^n)$ which is σ, g temperate and $w(x, \xi) \in S(m, g)$ satisfying

$$|w_{(\beta)}^{(\alpha)}(x, \xi)| \leq m(x, \xi) A^{|\alpha+\beta|} |\alpha + \beta|!^\kappa \varphi^{-|\beta|} \Psi^{-|\alpha|} \quad \text{for } (x, \xi) \in \mathbb{R}^{2n}.$$

Denote $k(x, \xi) = e^{\Lambda(x, \xi)} w(x, \xi)$ and $q(x, D) = {}^R e^{-\Lambda}(x, D) k(x, D)$. Then we have

$$\begin{aligned} |\{q(x, \xi) - \sum_{|\alpha| < k} D^\alpha (\omega^\alpha(-\Lambda; x, \xi) w(x, \xi)) / \alpha!\}_{(\beta)}^{(\gamma)}| \\ \leq C_{k\beta\gamma} m(x, \xi) (M^{2\kappa-1} N^{-1})^k \varphi^{-|\beta|} \Psi^{-|\gamma|} \quad \text{for any } k \text{ with } k \leq M, \end{aligned}$$

where $\omega^\alpha(\Lambda; x, \xi) = e^{-\Lambda(x, \xi)} \partial_\xi^\alpha e^{\Lambda(x, \xi)}$ and $C_{k\beta\gamma}$ is independent of M, N and h .

Now we can construct the inverse of $e^\Lambda(x, D)$. Put

$$J(x, D) = {}^R e^{-\Lambda}(x, D) e^\Lambda(x, D) - I.$$

Then it follows from Proposition 3.9 that $J(x, \xi) \in S(1, g)$ satisfies

$$\|J(x, D)\|_{\mathcal{L}(L^2, L^2)} \leq C M^{2\kappa-1} N^{-1} \leq C M^{2\kappa-3+\varepsilon} < 1$$

if $\varepsilon < 3 - 2\kappa$, $1 < \kappa < 3/2$, $N \geq M^{2-\varepsilon}$ and $M \gg 1$. Therefore $\{I + J(x, D)\}^{-1} = \sum_{k=0}^{\infty} (-J(x, D))^k$ converges in $\mathcal{L}(L^2, L^2)$. Moreover it follows from Beals [2] that $(I + J(x, D))^{-1}$ is a pseudodifferential operator with symbol in $S(1, g)$. Thus we have obtained

$$(3.20) \quad (e^\Lambda(x, D))^{-1} = (I + J(x, D))^{-1} {}^R e^{-\Lambda}(x, D)$$

which is also a pseudodifferential operator with symbol in $S(C(M) e^{-\Lambda(x, \xi)}, g)$.

Let $g_0 = (B \log \langle \xi \rangle_h)^2 (|dx|^2 + \langle \xi \rangle_h^{-2} |d\xi|^2)$ be a metric, where $B \geq 1$ is a parameter, and $g = \varphi^{-2} |dx|^2 + \Psi^{-2} |d\xi|^2$ is the metric given in Lemma 3.3. In this case $H(x, \xi) = B \log \langle \xi \rangle_h \Psi(x, \xi)^{-1} \leq B d N^{-1} h^{-1/2} \log h \leq 1$ if $h^{1/2} (\log h)^{-1} \geq B$. Moreover $(g_0 + g)/2 \sim g$ and $g_0/g_0^g \leq 1$ are valid. Therefore it follows from [4] that

$\sigma(a(x, D)b(x, D))(x, \xi) \in S(m_1 m_2, g)$ for $a \in S(m_1, g)$ and $b \in S(m_2, g_0)$, where m_1 and m_2 are σ, g and σ, g_0 temperate, respectively.

For $p(x, \xi)$ in $S(\langle \xi \rangle_h^m, g_0)$ ($m \in \mathbf{R}$) we define

$$(3.21) \quad p_\Lambda(x, D) = (e^\Lambda(x, D))^{-1} p(x, D) e^\Lambda(x, D).$$

Then it follows from [4] that

$$(3.22) \quad \sigma(p(x, D)e^\Lambda(x, D))(x, \xi) - \sum_{|\alpha| < k} p^{(\alpha)}(x, \xi) (e^{\Lambda(x, \xi)})_{(\alpha)} \alpha!^{-1}$$

belongs to $S(C(M, k)e^{\Lambda(x, \xi)} \langle \xi \rangle_h^{m-k/2} (B \log \langle \xi \rangle_h)^k, g)$, where $C(M, k)$ is independent of h and B .

Lemma 3.11. *Let $a(x, \xi)$ and $b(x, \xi)$ be in $S(m_1, g)$ and $S(m_2, g_0)$ respectively.*

Then

$$a(x, \xi)b(x, \xi) - \sum_{|\alpha| < k} (-1)^{|\alpha|} \sigma(a^{(\alpha)}(x, D)b_{(\alpha)}(x, D))(x, \xi)/\alpha!$$

is in $S(m_1 m_2 (B \log \langle \xi \rangle_h)^k \langle \xi \rangle_h^{-k/2}, g)$ for any integer $k \geq 0$.

Applying Lemma 3.11 to $(e^{\Lambda(x, \xi)})_{(\alpha)} p^{(\alpha)}(x, \xi)$, we see that

$$(3.23) \quad \begin{aligned} & p^{(\alpha)}(x, \xi) (e^{\Lambda(x, \xi)})_{(\alpha)} \\ & - \sum_{|\beta| < \ell} (-1)^{|\beta|} \sigma(k_\alpha^{(\beta)}(x, D) p_{(\beta)}^{(\alpha)}(x, D)/\beta!)(x, \xi) \end{aligned}$$

is in $S(C(M)e^\Lambda \Psi^{-|\alpha|} \langle \xi \rangle_h^{m-\ell/2} (\log \langle \xi \rangle_h)^\ell, g)$, where $k_\alpha(x, \xi) = (e^{\Lambda(x, \xi)})_{(\alpha)}$. Since $k_\alpha^{(\beta)}(x, \xi) = \partial_\xi^\beta D_x^\alpha e^{\Lambda(x, \xi)} = e^{\Lambda(x, \xi)} \omega_\alpha^\beta(\Lambda; x, \xi)$, it follows from Proposition 3.10 and (3.5) that

$$(3.24) \quad \sigma(R e^{-\Lambda}(x, D) k_\alpha^{(\beta)}(x, D))(x, \xi) - \sum_{|\gamma| < \ell} D_x^\gamma \{ \omega^\gamma(-\Lambda; x, \xi) \omega_\alpha^\beta(\Lambda; x, \xi) / \gamma! \}$$

is in $S(C(\ell, |\alpha| + |\beta|)(|\alpha| + |\beta|^\kappa + M)^{|\alpha| + |\beta|} \varphi^{-|\alpha|} \Psi^{-|\beta|} (M^{2\kappa-1} N^{-1})^\ell, g)$, for any ℓ with $\ell \leq M$. Moreover, noting that

$$\sigma((I + J(x, D))^{-1} - I) = \sigma(-J(x, D)(I + J(x, D))^{-1}) \in S(M^{2\kappa-1} N^{-1}, g),$$

we get, from (3.20) and (3.24) with $\ell = 1$,

$$(3.25) \quad \sigma((e^\Lambda(x, D))^{-1} k_\alpha^{(\beta)}(x, D))(x, \xi) - \omega_\alpha^\beta(\Lambda; x, \xi)$$

in $S(C_{\alpha\beta}M^{|\alpha+\beta|}\varphi^{-|\alpha|}\Psi^{-|\beta|}M^{2\kappa-1}N^{-1}, g)$. From (3.22), (3.23) and (3.25) we have the following

Theorem 3.12. *Assume that the same conditions in Proposition 3.9 are satisfied and, moreover, that $h^{1/2}(\log h)^{-1} \geq B \geq 1$ and $N \leq M^{2\kappa}$. Let $p(x, \xi)$ be in $S(\langle \xi \rangle_h^m, g_0)$. Then for $p_\Lambda(x, D)$ given in (3.21) we can see that*

$$(3.26) \quad p_\Lambda(x, \xi) - \sum_{|\alpha+\beta| < k} p_{(\beta)}^{(\alpha)}(x, \xi) w_\alpha^\beta(x, \xi)$$

is in $S(C(M, k)(B \log \langle \xi \rangle_h)^k \langle \xi \rangle_h^{m-k/2}, g)$ for any $k \geq 0$, where $w_0^0(x, \xi) = 1$ and

$$(3.27) \quad w_\alpha^\beta(x, \xi) = (-\Lambda_\xi(x, \xi))^\beta (-i\Lambda_x(x, \xi))^\alpha / (\alpha! \beta!)$$

is in $S(C_{\alpha\beta}M^{|\alpha+\beta|}\varphi^{-|\alpha|}\Psi^{-|\beta|}M^{2\kappa-1}N^{-1}, g)$ for $|\alpha + \beta| > 0$.

4. Outline of the proof of Theorem 2.2

In this section we shall give an outline of the proof of Theorem 2.2. Let $z^0 = (x^0, \xi^0) \in T^*(\mathbf{R}^n)$ with $|\xi^0| = 1$ be a multiple characteristic point of $\tilde{p}$. We assume that $\tilde{P}(x, D)$ satisfies (i), (ii) and (iii) of Proposition 2.1. For simplicity we denote $\tilde{P}$ by P . Let r be the multiplicity of z^0 , i.e., $d^j p(z^0) = 0$ for $j \leq r-1$ and $d^r p(z^0) \neq 0$. Following the notation in §2, we have

$$\begin{aligned} P(x, D; \gamma)u(x) &= e^{-\gamma\zeta(x)} P(x, D)(e^{\gamma\zeta(x)} u(x)) \\ &= \sum_{|\alpha| \leq m} a_\alpha(x) \sum_{\alpha' \leq \alpha} \binom{\alpha}{\alpha'} \omega_{\alpha-\alpha'}(\gamma\zeta) D_x^{\alpha'} u(x), \end{aligned}$$

where we write $\omega_\alpha(\gamma\zeta) = e^{-\gamma\zeta} D_x^\alpha e^{\gamma\zeta}$. Note that $\omega_\alpha(\gamma\zeta)$ is a polynomial in γ and its principal part is $(-i\gamma\zeta_x)^\alpha$. Therefore, we can write

$$(4.1) \quad P(x, \xi; \gamma) = \sum_{|\alpha| \leq m} P^{(\alpha)}(x, \xi) \omega_\alpha(\gamma\zeta) / \alpha!,$$

where $\omega_\alpha(\gamma\zeta) - (-i\gamma\zeta_x)^\alpha$ is a polynomial in γ of degree $|\alpha| - 1$ whose coefficients are in $\mathcal{B}^\infty(\mathbf{R}^n)$. Since $\Lambda_+(x, \xi)$ given in §2 satisfies

$$|\Lambda_{+(\beta)}^{(\alpha)}(x, \xi)| \leq \begin{cases} C_{\alpha\beta}(a + |b|) \log \langle \xi \rangle_h \langle \xi \rangle_h^{-|\alpha|} & \text{for } (x, \xi) \in \mathbf{R}^{2n}, \\ C_{\alpha\beta}(a \log \langle \xi \rangle_h + |b|) \langle \xi \rangle_h^{-|\alpha|} & \text{if } \varphi_{2,h/2}(x, \xi) = 1 \text{ and } |\alpha + \beta| \geq 1, \end{cases}$$

we have $e^{\Lambda_+(x, \xi)} \in S(e^{\Lambda_+}, g_0)$, where $g_0 = \{(a + |b|) \log \langle \xi \rangle_h\}^2 \{|dx|^2 + \langle \xi \rangle_h^{-2} |d\xi|^2\}$. Moreover $\omega_\alpha^\beta(\Lambda_+; x, \xi) = e^{-\Lambda_+(x, \xi)} D_x^\beta \partial_\xi^\alpha e^{\Lambda_+(x, \xi)}$ belongs to $S(C_{\alpha\beta} \{(a + |b|) \log \langle \xi \rangle_h\}^{|\alpha+\beta|} \times \langle \xi \rangle_h^{-|\beta|}, g_0)$ and

$$\omega_\alpha^\beta(\Lambda_+; x, \xi) - (-\Lambda_{+\xi})^\alpha (-i\Lambda_{+x})^\beta$$

is in $S(C_{\alpha\beta} \{(a + |b|) \log \langle \xi \rangle_h\}^{|\alpha+\beta|-1} \langle \xi \rangle_h^{-|\beta|}, g_0)$ for $|\alpha + \beta| \geq 1$. Put

$$J_+(x, D) = e^{-\Lambda_+}(x, D) e^{\Lambda_+}(x, D) - I.$$

Then, $J_+(x, \xi)$ is in $S(\{(a + |b|) \log \langle \xi \rangle_h\}^2 \langle \xi \rangle_h^{-1}, g_0)$. So we have $\|J_+(x, D)\|_{\mathcal{L}(L^2, L^2)} < 1$ if $h \gg \{(a + |b|) \log h\}^2$ and the symbol of $(e^{\Lambda_+}(x, D))^{-1} = (I + J_+(x, D))^{-1} e^{-\Lambda_+}(x, D)$ belongs to $S(e^{-\Lambda_+}, g_0)$. On the other hand we have

$$(4.2) \quad P(\Lambda_+, x, \xi; \gamma) - \sum_{|\alpha+\beta| < \ell} P_{(\beta)}^{(\alpha)}(x, \xi; \gamma) w_{+\alpha}^\beta(x, \xi)$$

is in $S(C_\ell \langle \xi \rangle_h^m \{\langle \xi \rangle_h^{-1} (a + |b|) (\log \langle \xi \rangle_h)\}^\ell, g_0)$ for any $\ell \geq 0$ and $\gamma \leq h$, where $P(\Lambda_+, x, D; \gamma) = e^{\Lambda_+}(x, D)^{-1} P(x, D; \gamma) e^{\Lambda_+}(x, D)$ and

$$w_{+\alpha}^\beta(x, \xi) - (-i\Lambda_{+x})^\alpha (-\Lambda_{+\xi})^\beta / (\alpha! \beta!)$$

is in $S(C_{\alpha\beta} \{(a + |b|) \log \langle \xi \rangle_h\}^{|\alpha+\beta|-1} \langle \xi \rangle_h^{-|\beta|}, g_0)$ for $|\alpha + \beta| \geq 1$ and $w_{+0}^0(x, \xi) = 1$. Thus it follows from (4.1) and (4.2) that

$$(4.3) \quad P(\Lambda_+, x, \xi; \gamma) - \sum_{|\alpha+\beta| < \ell} P_{(\beta)}^{(\alpha)}(x, \xi) \tilde{w}_\alpha^\beta(x, \xi, \gamma)$$

is in $S(C_\ell \langle \xi \rangle_h^m \{\langle \xi \rangle_h^{-1} (a + |b|) \log \langle \xi \rangle_h\}^\ell, g_0)$ for any $\ell \geq 0$, where

$$(4.4) \quad \tilde{w}_\alpha^\beta(x, \xi, \gamma) - (-i\gamma\zeta_x - i\Lambda_{+x})^\alpha (-\Lambda_{+\xi})^\beta / (\alpha! \beta!)$$

is in $S(C_{\alpha\beta} \{\gamma + (a + |b|) \log \langle \xi \rangle_h\}^{|\alpha+\beta|-1} \langle \xi \rangle_h^{-|\beta|}, g_0)$ for $|\alpha + \beta| \geq 1$ ($C_{\alpha\beta}$ is independent of γ and h) and $\tilde{w}_0^0(x, \xi, \gamma) = 1$.

It suffices to show that $P(\Lambda_+, x, D; \gamma)$ satisfies (2.1)_{zo}. Then we can see that $P_{\Lambda_+}(x, D; \gamma) = e^{-\Lambda_+}(x, D) P(x, D; \gamma) e^{\Lambda_+}(x, D)$ also satisfies (2.1)_{zo}.

From now on we choose parameters h, M, N, γ, a and b such that $h = \gamma$, $N = M^{2-\varepsilon} \geq d$, $h^{1/2} / \log h \geq a + |b|$, $a \geq 1$ and $b \in \mathbf{R}$, where $0 < \varepsilon < 3 - 2\kappa$

and $1 < \kappa < 3/2$. Let $t_k(x, \xi)$ ($k = 1, \dots, d$) be the time functions of p at z^0 which appear in $[A.III]_{z^0}$, and let $\Lambda(x, \xi)$ be defined by (3.4). We conjugate again $P(\Lambda_+) = P(\Lambda_+, x, D; \gamma)$ by $e^\Lambda(x, D)$. Then it follows from (3.26), (3.27) of Theorem 3.12, (4.3) and (4.4) that

$$(4.5) \quad P(\Lambda; x, \xi) - \sum_{|\alpha+\beta|<\ell} P_{(\beta)}^{(\alpha)}(x, \xi) w_\alpha^\beta(x, \xi)$$

is in $S(C(M, a, b)\langle \xi \rangle_h^{m-\ell/2}(\log\langle \xi \rangle_h)^\ell, g)$ for any $\ell \geq 0$, where $P(\Lambda; x, D) = (e^\Lambda(x, D))^{-1} \times P(\Lambda_+, x, D; \gamma)e^\Lambda(x, D)$ and

$$(4.6) \quad \rho_\alpha^\beta(x, \xi) \equiv w_\alpha^\beta(x, \xi) - (-i\gamma\zeta_x - i\Lambda_{+x} - i\Lambda_x)^\alpha (-\Lambda_{+x} - \Lambda_x)^\beta / (\alpha! \beta!)$$

is in $S(C_{\alpha\beta} M^{2\kappa-1} N^{-1} [\{\gamma + (a + |b|) \log\langle \xi \rangle_h\} \varphi + M]^{|\alpha+\beta|} \varphi^{-|\alpha|} \Psi^{-|\beta|}, g)$.

We define the principal part of $P(\Lambda) = P(\Lambda; x, D)$ by

$$p(\Lambda; x, \xi) = \sum_{|\alpha+\beta|<m+2} p_{(\beta)}^{(\alpha)}(x, \xi) (-i\gamma\zeta_x - i\Lambda_{+x} - i\Lambda_x)^\alpha (-\Lambda_{+x} - \Lambda_x)^\beta / (\alpha! \beta!),$$

where $p(x, \xi)$ is the principal part of $P(x, \xi)$. Then we can write from (4.5) and (4.6)

$$\begin{aligned} P(\Lambda; x, \xi) = p(\Lambda; x, \xi) + \sum_{0<|\alpha+\beta|<m+2} p_{(\beta)}^{(\alpha)}(x, \xi) \rho_\alpha^\beta(x, \xi) \\ + \sum_{j=0}^{m-1} \sum_{|\alpha+\beta|<m+2} P_{j(\beta)}^{(\alpha)}(x, \xi) w_\alpha^\beta(x, \xi) + r(x, \xi), \end{aligned}$$

where $r(x, \xi) \in S(C(M, a, b)\langle \xi \rangle_h^{(m-2)/2}(\log\langle \xi \rangle_h)^{m+2}, g)$ and $w_\alpha^\beta(x, \xi)$ and $\rho_\alpha^\beta(x, \xi)$ are given in (4.6).

For $\zeta = (y, \eta) \in \mathbf{R}^{2n}$ we define

$$p(x, \xi; \zeta) = \sum_{|\alpha+\beta|<m+2} p_{(\beta)}^{(\alpha)}(x, \xi) (i\eta)^\alpha (-y)^\beta / (\alpha! \beta!).$$

Then we have

$$p(\Lambda; x, \xi) = p(x, \xi; H_{\Lambda_1}),$$

where $\Lambda_1 = \gamma\zeta(x) + \Lambda_+ + \Lambda$ and $H_{\Lambda_1} = (\Lambda_{1\xi}, -\Lambda_{1x})$. We put

$$\begin{aligned} Q(x, \xi) = & (\langle \xi \rangle_h^2 |\Lambda_{1\xi}|^2 + |\Lambda_{1x}|^2)^{-1/2} \\ & \times \sum_{j=1}^n (\Lambda_{1\xi_j} \partial / \partial x_j - \Lambda_{1x_j} \partial / \partial \xi_j) p(x, \xi; \zeta)|_{\zeta=H_{\Lambda_1}}, \end{aligned}$$

$$S(x, D) = (2i)^{-1} \{Q^*(x, D)P(\Lambda; x, D) - P^*(\Lambda; x, D)Q(x, D)\},$$

where we denote by Q^* the adjoint operator of Q as $(Qu, v)_{L^2} = (u, Q^*v)_{L^2}$ for u and $v \in C_0^\infty(\mathbf{R}^n)$. Taking $\psi_h(x, \xi) = (1 - \Theta_h(\xi))\psi(x, \xi)$, where $\psi \in C^\infty(T^*(\mathbf{R}^n))$ is homogeneous in ξ of degree 0 for $|\xi| \geq 1$ and $\psi = 1$ in a conic neighborhood of z^0 , we can prove that there are $c > 0$ and $C > 0$ such that

$$(4.7) \quad (S(x, D)\psi_h(x, D)v, \psi_h(x, D)v)_{L^2} \\ \geq c\|\langle D \rangle_h^{(m-1)/2}v\|_{L^2}^2 - C\|\langle D \rangle_h^{m-1/2}(1 - \psi_h)v\|_{L^2}^2$$

for $v \in \mathcal{S}$. Then (2.1)_{z⁰} easily follows from (4.7). We omit the proof of (4.7), since the proof is not short and the method is similar to one used in [10]. For the proof of (4.7) we refer to [11].

REFERENCES

- [1]. M.D. Bronshtein, *Smoothness of polynomials depending on parameters*, Sibirsk. Mat. Zh. **20** (1979), 493–509.
- [2]. R. Beals, *Weighted distribution spaces and pseudodifferential operators*, J. d'analyse Math. **39** (1981), 131–187.
- [3]. L. Gårding, *Solution directe du problème de Cauchy pour les equations hyperboliques*, Coll. Int. CNRS, Nancy, 1956, 71–90.
- [4]. L. Hörmander, “The Analysis of Linear Partial Differential Operators III,” Springer, Berlin-Heidelberg-New York-Tokyo, 1983.
- [5]. V. Ja. Ivrii and V. M. Petkov, *Necessary conditions for the Cauchy problem for non-strictly hyperbolic equations to be well posed*, Uspehi Mat. Nauk **29** (1974), 3–68.
- [6]. N. Iwasaki, *The Cauchy problem for effectively hyperbolic equations (general case)*, J. Math. Kyoto Univ. **25** (1985), 727–743.
- [7]. F. John, *On linear partial differential equations with analytic coefficients, unique continuation of data*, Comm. Pure Appl. Math. **2** (1949), 209–254.
- [8]. K. Kajitani and T. Nishitani, “The Hyperbolic Cauchy Problem,” Springer Lecture Note in Math. **1505**, 1991.
- [9]. K. Kajitani and S. Wakabayashi, *Microlocal a priori estimates and the Cauchy problem I*, to appear.

- [10]. K. Kajitani and S. Wakabayashi, *Microlocal a priori estimates and the Cauchy problem II*, to appear.
- [11]. K. Kajitani, S. Wakabayashi and T. Nishitani, *The Cauchy problem for hyperbolic operators of strong type*, to appear.
- [12]. J. Leray, "Hyperbolic Differential Equations," The Institute for Advanced Study, Princeton, 1953.
- [13]. S. Mizohata, *Some remarks on the Cauchy problem*, J. Math. Kyoto Univ. **1** (1961), 109–127.
- [14]. T. Nishitani, *Hyperbolic operators with symplectic multiple characteristics*, J. Math. Kyoto Univ. **29** (1989), 405–447.
- [15]. T. Nishitani, *Propagation of singularities for hyperbolic operators with transverse propagation cone*, Osaka J. Math. **27** (1990), 1–16.
- [16]. S. Wakabayashi, *Singularities of solutions of the Cauchy problem for symmetric hyperbolic systems*, Comm. in Partial Differential Equations **9** (1984), 1147–1177.
- [17]. S. Wakabayashi, *Generalized flows and their applications*, Proc. NATO ASI on Advances in Microlocal Analysis, Series C, D. Reidel (1986), 363–384.