TITLE:

Comparative analysis of DNA alkylation by conjugates between pyrrole-imidazole hairpin polyamides and chlorambucil or seco-CBI.

AUTHOR(S):

Minoshima, Masafumi; Bando, Toshikazu;
Shinohara, Ken-Ichi; Kashiwazaki, Gengo;
Nishijima, Shigeki; Sugiyama, Hiroshi

CITATION:

Minoshima, Masafumi ...[et al]. Comparative analysis of DNA alkylation by conjugates between pyrrole-imidazole hairpin polyamides and chlorambucil or seco-CBI.. Bioorganic & medicinal chemistry 2010, 18(3): 1236-1243

ISSUE DATE:

2010-02-01

URL:

<http://hdl.handle.net/2433/97888>

RIGHT:

c 2009 Elsevier Ltd. All rights reserved.; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。; This is not the published version. Please cite only the published version.

Comparative Analysis of DNA Alkylation by Conjugates between Pyrrole-Imidazole Hairpin Polyamides and Chlorambucil or *seco*-CBI

Masafumi Minoshima¹, Toshikazu Bando¹, Ken-ichi Shinohara¹, Gengo Kashiwazaki¹, Shigeki Nishijima¹, and Hiroshi Sugiyama^{1,2*}

¹Department of Chemistry, Graduate School of Science, Kyoto University, Kitashirakawa-Oiwakecho, Sakyo-Ku, Kyoto, 606-8502, Japan, ²Institute for Integrated Cell-Materials Science (iCeMS).

RECEIVED DATE ;E-mail: hs@kuchem.kyoto-u.ac.jp

Abstract: We investigated sequence-specific DNA alkylation using conjugates between the N-methylpyrrole (Py)-N-methylimidazole (Im) polyamide and the DNA alkylating agent, chlorambucil, or 1-(chloromethyl)-5-hydroxy-1,2-dihydro-3H-benz[e]indole (*seco*-CBI). Polyamide-chlorambucil conjugates **1–4** differed in the position at which the DNA alkylating chlorambucil moiety was bound to the Py-Im polyamide. High-resolution denaturing polyacrylamide gel electrophoresis (PAGE) revealed that chlorambucil conjugates **1–4** alkylated DNA at the sequences recognized by the Py-Im polyamide core moiety. Reactivity and sequence specificity were greatly affected by the conjugation position, which reflects the geometry of the alkylating agent in the DNA minor groove. Polyamide-*seco*-CBI conjugate **5** was synthesized to compare the efficacy of chlorambucil with that of *seco*-CBI as an alkylating moiety for Py-Im polyamides. Denaturing PAGE analysis revealed that DNA alkylation activity of polyamide-*seco*-CBI conjugate **5** was similar to that of polyamide-chlorambucil conjugates **1** and **2**. In contrast, the cytotoxicity of conjugate **5** was superior to that of conjugates **1–4**. These results suggest that the *seco*-CBI conjugate was distinctly active in cells compared to the chlorambucil conjugates. These results may contribute to the development of more specific and active DNA alkylating agents.

1. Introduction

DNA targeting with small molecules has been studied for several decades¹. DNA alkylating agents have been used as anticancer agents as they induce sufficient DNA damage to kill cancer cells². Conjugates between small molecules and alkylating agents have superior DNA binding affinity to that of alkylating agents alone. Conjugates between distamycin A, a DNA minor groove binder, and nitrogen mustard DNA alkylating agents have superior DNA alkylating activity and cytotoxicity against cancer cells³. Antibiotics such as CC-1065 and duocarmycin derivatives are strong alkylating agents that bind to DNA⁴. These compounds alkylate at the N-3 position of adenines flanked by AT-rich sequences. Despite the effort that has been devoted to the development of DNA alkylating agents, their severe side effects have not been overcome.

To address this problem, our group focused on the introduction of sequence specificities into DNA alkylating agents. Minor groove-binding hairpin polyamides that contain N-methylpyrrole (Py)-N-methylimidazole (Im) and uniquely recognize each of the four Watson-Crick base pairs were developed by Dervan et al.⁵. Pairing of Im opposite Py targets the G-C base pair, and Py-Im targets the C-G base pair. Py-Py targets the T-A and A-T base pairs. It has been demonstrated that Py-Im hairpin polyamides that bind competitively to transcription regulatory sequences inhibit target gene expression⁶. We demonstrated that hybrid molecules consisting of duocarmycin A and Py-Im hairpin polyamides alkylated sequences predetermined by the recognition rules of the Py-Im polyamides⁷. Furthermore, we showed that introduction of a vinyl linker between the polyamides and cyclopropapyrroloindole (CPI) resulted in efficient alkylation of double-stranded DNA and improved cytotoxicity against cancer cells⁸. A more chemically stable indole linker and the synthetic alkylating agent 1,2,9,9a-tetrahydrocyclopropa[1,2-c]benz[1,2-

e]indol-4-one (CBI)⁹, were recently introduced for the design of sequence-specific DNA alkylating agents. This type of alkylating agent has superior stability, and can be synthesized to contain longer recognition sequences than other types of alkylating agents¹⁰. We have demonstrated that alkylation at the coding region of a template strand produces truncated mRNA¹¹. We have also demonstrated that selective alkylation at the coding region of a template strand using Py-Im polyamides results in silencing of reporter genes (luciferase and GFP)¹².

Dervan et al. constructed a small library of conjugates between polyamides with different recognition sequences and chlorambucil¹³. One of these conjugates arrested the growth of several cancer cell lines. They demonstrated selective alkylation at the coding region of the histone *h4c* gene and inhibition of its expression, which indicates that sequence-specific DNA alkylating polyamides are promising gene-targeting drugs for cancer chemotherapy¹⁴. However, little is known about conjugates between the hairpin polyamide and chlorambucil. In this report, we present a comprehensive analysis of the DNA alkylating activities of hairpin polyamides conjugated with chlorambucil at various positions, their sequence specificities, and their effects on cell viability. We also compared the polyamide-chlorambucil conjugates to a polyamide-*seco*-CBI conjugate.

2. Molecular Design and synthesis

We designed agents in which the Py-Im polyamide was conjugated with chlorambucil at different positions: C-tail (**1**), N-tail (**2**), N-tail with a β -alanine linker (**3**), and turn (**4**) (Figure 1a). To investigate the effect of the position of the conjugate, all Py-Im polyamides were designed to recognize the same sequence (Figure 1b). The core moiety of the Py-Im polyamide recognizes the seven base pair sequence, 5'-WGCWGC-3' (W = A or T). All conjugates had one cationic tertiary amino group at the C-tail.

Conjugate **4** had an (R)- α -diaminobutyric acid as a turn moiety. This turn moiety has more specific DNA alkylation activity than (R)- γ -diaminobutyric acid and has more efficient DNA alkylation activity than the stereochemically inversed (S)- α -diaminobutyric acid turn¹⁵. We also designed a Py-Im polyamide-*seco*-CBI conjugate (**5**) with an indole linker (Figure 1a). Conjugate **5** has the identical recognition sequences as other polyamide

chlorambucil conjugates. The Py-Im polyamides were synthesized using Fmoc-based solid-phase synthesis¹⁶. Polyamides with amino group were coupled with chlorambucil to produce conjugates **1-4**. A polyamide with carboxylic acid was coupled with indole-*seco*-CBI alkylating unit to produce the conjugate **5** (schemes are shown in supporting information)¹⁷. All conjugates were purified by HPLC, and confirmed by ESI-TOF-MS or NMR.

Figure 1. (a) A chemical structure of Py-Im polyamide-chlorambucil conjugates **1-4**, Py-Im polyamide-*seco*-CBI conjugate **5**. (b) Schematic representation of binding by polyamide at consensus seven-base-pair recognition sequence. W indicates A or T base.

Figure 2. Illustration of the 209 base pair DNA fragment with the position of the sequence indicated. The recognition sites for polyamide core moiety are boxed.

Figure 3. (a) Thermally induced strand cleavage of 5'-Texas Red-labeled 209 bp DNA fragments inserted ODN1 by polyamide-chlorambucil conjugates 1-4. The concentrations of conjugates were 1000, 500, 100, 50, and 10 nM. Alkylation sites are shown in red arrows. (b) Schematic representation of the recognition of alkylation sites by 1-4. The arrows indicate the site of alkylation and the alkylating bases are shown in red.

Figure 4. (a) Thermally induced strand cleavage of 5'-Texas Red-labeled 209 bp DNA fragments inserted inverted ODN1 by polyamide-chlorambucil conjugates 1-4. The concentrations of conjugates were 1000, 500, 100, 50, and 10 nM. Alkylation sites are shown in red arrows. (b) Schematic representation of the recognition of alkylation sites by 1-4. The arrows indicate the site of alkylation and the alkylating bases are shown in red.

3. High-resolution denaturing PAGE analysis of labeled DNA fragment alkylated by conjugates 1–4

The sequence-selective DNA alkylation activity of chlorambucil conjugates 1–4 was investigated using a 5'-Texas Red-labeled DNA fragment and an automated DNA sequencer as described previously¹⁸. The 209 bp DNA fragment was prepared by inserting ODN1 (Figure 2), including two recognition sequences of the polyamides and several adenines at the 5'- and 3'-side of the recognition sequences. Nitrogen mustard DNA alkylating agents predominantly react at guanine bases such as the N7 positions; however, when located at the minor groove, they react mainly at adenine bases such as the N3 positions³. Alkylation was carried out at 23 °C for 18 h, followed by quenching with calf thymus DNA. The samples were heated at 95 °C under neutral conditions for 10 min. The alkylation sites were visualized by thermal cleavage of the DNA strand. Under these heating conditions, all the purine N3 alkylated sites in the DNA produced cleavage bands on the gel. The results of sequencing gel analysis of DNA fragments alkylated by conjugates 1–4 after heat treatment are shown in Figure 3. Figure 3 shows that DNA alkylation by conjugates 1, 2, and 3 occurred mainly at sites located on the 3'-side of adenines recognized by the polyamide, 5'-AGCAGTCAAAAA-3' (site 2). Conjugates 1, 2, and 3 showed high alkylating activities at nanomolar concentrations. At the highest concentration of conjugate 1, all the DNA bands were disappeared probably because of the non-specific alkylation at many sites. On the other hand, slight DNA alkylation by conjugate 4 occurred at two sites located on the 5'-side of the recognition sequence, 5'-AGCAGTC-3' (sites 1 and 3). This alkylation result for conjugate 4 correspond with those of a previous report on polyamide-chlorambucil conjugates at the hairpin turn position¹⁴. Conjugate 4 resulted in little alkylation, even at a concentration of 1 μ M.

To investigate alkylation of the complementary strand, we used conjugates 1–4 and a 209 bp DNA fragment inserted inverted ODN1 sequence (Figure 2). The results of sequence gel analysis are shown in Figure 4. Surprisingly, conjugate 1 showed efficient DNA alkylation at site 4 (5'-AAAAAGACTGCT-3'), whereas conjugates 2 and 3 only alkylated this site slightly. Conjugate 4 did not provide distinct gel bands for any site. This indicates that conjugate 1 alkylates adenines in both strands adjacent to the

recognition sequence of the Py-Im polyamide, whereas conjugates 2–4 selectively alkylate only one strand.

4. DNA alkylation activity and sites of polyamide-chlorambucil conjugates 1–4

The denaturing PAGE experiment revealed that the activity and site of DNA alkylation are dependent on the position of the conjugate between the DNA alkylating chlorambucil moiety and the polyamide. Conjugates at the C-tail and N-tail positions of the polyamide (1–3) had potent DNA alkylating activity, whereas the conjugate at the hairpin turn position (4) had slight DNA alkylating activity. In this experiment, the tail position of polyamides binding to the predetermined sequence was located at five adenine tracts. This sequence may allow efficient alkylation activity by the DNA alkylating agents. Moreover, the difference of DNA binding affinity of polyamides may also reflect the alkylation activity. Dervan and co-workers reported that polyamides with α -turn had lower affinity compared to that of polyamides with γ -turn¹⁹. The sequence specificity for the DNA alkylation site differed between conjugates. The geometry of chlorambucil conjugated with the hairpin polyamide determines whether the alkylation site will lie on the 3'-side (conjugates 1–3) or the 5'-side (conjugate 4). These results indicate that the microenvironment of the DNA minor groove to which polyamides bind is critical for activity and sequence specificity. Figure 5 shows schematic representations of the DNA recognition and alkylation sites of conjugates 1–4.

The conjugate at the C-tail of the polyamide (1) efficiently alkylated adenines on both strands two to four bases from the sequence recognized by the polyamide core moiety. In this conjugate, the chlorambucil moiety is flexibly conjugated with the cationic tertiary amine, which may have enabled DNA alkylation of adenines in both strands.

The conjugate at the N-tail of the polyamide (2) efficiently alkylated adenines on the 3'-side one to three bases from the sequence recognized by the polyamide core moiety. Moreover, conjugate 2 exclusively alkylated the adenines in the 5'-AAAAA-3' sequence, not the adenines in the 3'-AAAAA-5' sequence. This one-way direction was observed for several minor groove binder nitrogen mustard conjugates. For example, tallimustine (FCE-24517) alkylates the specific sequence 5'-TTTTTPuA-3', where Pu = G or A^{3c}.

Figure 5. Schematic representations of the alkylation sites of conjugates 1–4. Arrows indicate alkylation sites corresponding to the sites presented in Figure 3 and 4.

7. CONCLUSIONS

We investigated the DNA alkylation activity, sequence specificity, and cytotoxicity of Py-Im polyamides conjugated with the alkylating agents, chlorambucil or *seco*-CBI. Chlorambucil conjugates at the N-tail and C-tail positions had high DNA alkylating activities, which were similar to that of the *seco*-CBI conjugate. A chlorambucil conjugate at a turn position had low DNA alkylating activity, but specifically alkylated adenines located near the turn moiety. These results suggest that the geometry of alkylating agents in the DNA minor groove is important for alkylation activity and sequence specificity. Moreover, the *seco*-CBI conjugate is superior to chlorambucil conjugates with regard to the cytotoxicity in culture cells. The properties of an alkylating moiety are important determinants of the efficacy of conjugates in cells.

Previously, we demonstrated that the substitution of one Py with an Im changed cytotoxicity against 39 human cancer cell lines²⁴. We also demonstrated distinct differences in cytotoxicity between alkylating polyamides in our library²⁵. These reports suggest that screening of highly active alkylating polyamides in a library is an alternative approach to elucidating the potential of alkylating polyamides. The construction of small libraries of alkylating polyamides with different recognition sequences and evaluation of their effects against various cancer cell lines is now in progress.

8. Experiments

8.1 General.

Reagents and solvents were purchased from standard suppliers and used without further purification. Abbreviations for reagents are: Fmoc, fluorenylmethoxycarbonyl; Boc, tert-butoxycarbonyl; TFA, trifluoroacetic acid; DIEA, *N,N*-diisopropylethylamine; DMF, *N,N*-dimethylformamide; HCTU, 1-[Bis(dimethylamino)methylene]-5-chloro-1*H*-benzotriazolium-3-oxide hexafluorophosphate; PyBOP, benzotriazole-1-yl-oxy-trispyrrolidino-phosphonium hexafluorophosphate; TIS, triisopropylsilane. HPLC analysis was performed with a Jasco PU-2080 HPLC pump, a UV-2075 HPLC UV/VIS detector, a Chemcobond 5-ODS-H 4.6 x 150 mm column (Chemco Scientific). A Chemcobond 5-ODS-H 10 x 150 mm column (Chemco Scientific) and a YMC-Pack Pro C18 150 x 20 mm column (YMC) were used in purification of polyamides. NMR spectra were recorded using a JEOL JNM-FX 400 nuclear magnetic resonance spectrometer with tetramethylsilane as internal standard. Proton NMR spectra were recorded in parts per million (ppm) downfield relative to tetramethylsilane. The following abbreviations apply to spin multiplicity: s (singlet), d (doublet), t (triplet), q (quartet), qu (quintet), m (multiplet), and br (broad). Electrospray ionization mass spectrometry (ESI-MS) and electrospray ionization time-of-flight mass spectrometry (ESI-TOF-MS) were performed using API 150 (PE SCIEX) and BioTOF II (Bruker Daltonics) mass spectrometers. PCR amplification and DNA sequencing was carried out using the Dice mini thermal cycler (Takara), GoTaq Green Master Mix (Promega), and the Thermo Sequenase core sequencing kit (GE Healthcare). Loading dye was prepared with formamide and fuchsin red (Merck). The 5'-Texas Red-modified DNA oligomer (19 mer) was obtained from Sigma Genosys Co. Ltd. The 50% Long Ranger™ gel solution was obtained from Lonza Co. Ltd. Polyacrylamide gel electrophoresis was performed using a HITACHI SQ5500-E DNA sequencer.

8.2 Solid-phase synthesis of Py-Im polyamides.

AcImPyPy-β-ImPy-γ-ImPy-β-ImPyPy-(CH₂)₃-NCH₃-(CH₂)₃-NH₂ (6). All the Py-Im polyamides were synthesized in a stepwise reaction by Fmoc solid-phase protocol according to reported procedures¹⁶. Resulting AcImPyPy-β-ImPy-γ-ImPy-β-ImPyPy-CO₂-oxime resin was cleaved with 500 μL of 3,3'-diamino-*N*-methyl-dipropylamine for overnight at 55 °C, and purified by HPLC using a YMC-Pack Pro C18 150 x 20 mm column (H₂O with 0.1% AcOH containing 15-35% CH₃CN over a linear gradient for 30 min at a flow rate of 6 mL/min). The peak around 10 min was collected and lyophilized to produce **6** (6.3 mg, 3.8 μmol, 18%) as a yellow powder. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.29 (s, 2H; NH), 10.24 (s, 1H; NH), 10.22 (s, 1H; NH), 9.94 (s, 2H; NH), 9.89 (s, 4H; NH), 8.03 (br, 4H; NH), 7.45 (s, 2H; Im-H), 7.42 (s, 1H; Im-H), 7.41 (s, 1H; Im-H), 7.26 (s, 2H; Py-H), 7.23 (s, 1H; Py-H), 7.21 (s, 1H; Py-H), 7.19 (s, 1H; Py-H), 7.18 (s, 1H; Py-H), 7.17 (s, 1H; Py-H), 7.11 (s, 2H; Py-H), 6.95 (s, 1H; Py-H), 6.93 (s, 1H; Py-H), 6.85 (s, 1H; Py-H), 6.83 (s, 1H; Py-H), 3.95 (s, 3H; CH₃), 3.94 (s, 6H; CH₃), 3.92 (s, 3H; CH₃), 3.83 (s, 6H; CH₃), 3.80 (s, 6H; CH₃), 3.79 (s, 6H; CH₃), 3.44 (q, *J*=6.8 Hz, 6H; CH₂), 3.37 (q, *J*=6.8 Hz, 4H; CH₂), 3.18 (m, 4H; CH₂), 2.58 (t, *J*=6.8 Hz, 4H; CH₂), 2.29 (br, 4H; CH₂), 2.12 (s, 3H; CH₃), 2.01 (s, 3H; CH₃), 1.82 (m, 6H; CH₂), ESI-MS *m/z* calcd for C₇₅H₉₅N₃₀O₁₄ [M + H]⁺ 1639.8; found 1639.8.

NH₂-ImPyPy-β-ImPy-γ-ImPy-β-ImPyPy-(CH₂)₃-N(CH₃)₂ (7). NH₂-ImPyPy-β-ImPy-γ-ImPy-β-ImPyPy-CO₂-oxime resin was cleaved with 500 μL of *N,N*-dimethylaminopropaneamine for overnight at 55 °C, and purified by HPLC (the same condition as **6**) to produce **7** (11.1 mg, 7.1 μmol, 34%) as a yellow powder. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.29 (s, 2H; NH), 10.24 (s, 1H; NH), 9.98 (s, 1H; NH), 9.91 (s, 3H; NH), 9.89 (s, 2H; NH), 8.05 (br, 4H; NH), 7.45 (s, 2H; Im-H), 7.43 (s, 2H; Im-H), 7.26 (s, 1H; Py-H), 7.23 (s, 1H; Py-H), 7.21 (s, 1H; Py-H), 7.20 (s, 1H; Py-H), 7.17 (s, 2H; Py-H), 7.11 (s, 2H; Py-H), 6.95 (s, 1H; Py-H), 6.93 (s, 1H; Py-H), 6.84 (s, 1H; Py-H), 6.82 (s, 1H; Py-H), 3.94 (s, 3H; CH₃), 3.93 (s, 3H; CH₃), 3.92 (s, 3H; CH₃), 3.86 (s, 3H; CH₃), 3.83 (s, 3H; CH₃), 3.81 (s, 3H; CH₃), 3.80 (s, 3H; CH₃), 3.79 (s, 3H; CH₃), 3.78 (s, 6H; CH₃), 3.43 (q, *J*=6.4 Hz, 4H; CH₂), 3.17 (d, *J*=5.6 Hz, 4H; CH₂), 2.58 (t, *J*=6.8 Hz, 2H; CH₂), 2.34 (t, *J*=7.2 Hz, 2H; CH₂), 2.23 (t, *J*=7.2 Hz, 2H; CH₂), 2.12 (s, 6H; CH₃), 2.10 (br, 2H; CH), 1.77 (qu, *J*=6.8 Hz, 2H; CH₂), 1.59 (qu *J*=6.8 Hz, 2H; CH₂); ESI-MS *m/z* calcd for C₇₁H₈₉N₂₉O₁₃ [M + H]⁺ 1554.7; found 1554.8.

NH₂-β-ImPyPy-β-ImPy-γ-ImPy-β-ImPyPy-(CH₂)₃-N(CH₃)₂ (8). A synthetic procedure similar to that used for the preparation of compound **7** provided **8** (11.5 mg, 7.1 μmol, 32%) as a yellow powder. ESI-MS *m/z* calcd for C₇₄H₉₄N₃₀O₁₄ [M + H]⁺ 1625.8; found 1625.8.

AcImPyPy-β-ImPy-(R)^{NH₂}-γ-ImPy-β-ImPyPy-(CH₂)₃-N(CH₃)₂ (9). AcImPyPy-β-ImPy-(R)^{NHBoc}-γ-ImPy-β-ImPyPy-CO₂-oxime resin was cleaved with 500 μL of *N,N*-dimethylaminopropaneamine for overnight at 55 °C, and purified by HPLC (the same condition as **6**), to produce Boc-protected **9**. After deprotection Boc group with 20% TFA in dichloromethane,

9 was obtained (9.3 mg, 5.8 μ mol, 26%) as a yellow powder. ESI-MS m/z calcd for $C_{73}H_{91}N_{30}O_{14}$ $[M + H]^+$ 1611.7; found 1611.8.

AcImPyPy- β -ImPy- γ -ImPy- β -Im-CO₂H (10). AcImPyPy- β -ImPy- γ -ImPy- β -Im-CO₂-Wang resin was cleaved with 1 mL of TFA/water/TIS (95/2.5/2.5; v/v/v) for 1 h at room temperature. After ether precipitation, **10** was obtained (24.2 mg, 19 μ mol, 53%) as a yellow powder. **10** was used for the synthesis of polyamide-*seco*-CBI conjugate **5** without further purification. ESI-TOF-MS m/z calcd for $C_{56}H_{66}N_{23}O_{13}$ $[M + H]^+$ 1268.52; found 1268.52.

AcImPyPy- β -ImPy- γ -ImPy- β -ImPyPy-(CH₂)₃-NCH₃-(CH₂)₃-NH-Chlorambucil (1). To a solution of compound **6** (6.6 mg, 4.0 μ mol), PyBOP (6.1 mg, 12 μ mol), and chlorambucil (4.3 mg, 14 μ mol) in DMF (100 μ L), DIEA (2.4 μ L, 14 μ mol) was added. The reaction mixture was incubated overnight at room temperature. Evaporation of the solvent gave yellow crude, which was washed with ether (1 mL x 2) and dichloromethane (1 mL x 2). HPLC purification was performed using a Chemcobond 5-ODS-H 10 x 150 mm column (H₂O with 0.1% AcOH containing 0-66% CH₃CN over a linear gradient for 30 min at a flow rate of 3 mL/min). The peak around 17 min was collected and lyophilized to produce conjugate **1** was obtained (2.7 mg, 1.4 μ mol, 35%) as a white powder. **1** was used in the DNA alkylation reaction. Synthetic procedures were similar to that used for the preparation of compounds **1-4** from polyamides **6-9**. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.38 (s, 2H; NH), 10.31 (s, 1H; NH), 10.28 (s, 1H; NH), 9.95 (s, 3H; NH), 9.92 (s, 1H; NH), 9.88 (s, 2H; NH), 8.08 (br, 5H; NH), 7.44 (s, 2H; Im-H), 7.42 (s, 1H; Im-H), 7.41 (s, 1H; Im-H), 7.26 (s, 2H; Py-H), 7.20 (s, 1H; Py-H), 7.18 (s, 1H; Py-H), 7.12 (s, 2H; Py-H), 6.98 (s, 1H; Py-H), 6.96 (m, 2H; benzene-H), 6.94 (s, 1H; Py-H), 6.85 (s, 1H; Py-H), 6.83 (s, 1H; Py-H), 6.63 (m, 2H; benzene-H), 3.93 (s, 9H; CH₃), 3.92 (s, 3H; CH₃), 3.83 (s, 3H; CH₃), 3.80 (s, 9H; CH₃), 3.79 (s, 6H; CH₃), 3.67 (m, 8H; CH₂), 3.45 (m, 6H; CH₂), 3.19 (m, 4H; CH₂), 2.66 (m, 2H; CH₂), 2.58 (m, 4H; CH₂), 2.32 (m, 6H; CH₂), 2.11 (s, 3H; CH₃), 2.01 (s, 3H; CH₃), 1.77 (m, 8H; CH₂); ESI-TOF-MS m/z calcd for $C_{89}H_{112}Cl_2N_{31}O_{15}$ $[M + H]^+$ 1924.83; found 1924.96.

Chlorambucil-NH-ImPyPy- β -ImPy- γ -ImPy- β -ImPyPy-(CH₂)₃-N(CH₃)₂ (2). **2** was obtained as a white powder (1.3 mg, 0.71 μ mol, 24%) from polyamide **7**. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.28 (s, 2H; NH), 10.24 (s, 1H; NH), 10.20 (s, 1H; NH), 9.97 (s, 1H; NH), 9.95 (s, 1H; NH), 9.90 (s, 3H; NH), 9.89 (s, 2H; NH), 8.06 (br, 4H; NH), 7.45 (s, 3H; Im-H), 7.42 (s, 1H; Im-H), 7.25 (s, 2H; Py-H), 7.21 (s, 1H; Py-H), 7.19 (s, 1H; Py-H), 7.17 (s, 2H; Py-H), 7.11 (s, 1H; Py-H), 7.02 (d, J =8.4 Hz, 2H; benzene-H), 6.95 (s, 1H; Py-H), 6.93 (s, 1H; Py-H), 6.85 (s, 1H; Py-H), 6.82 (s, 1H; Py-H), 6.65 (d, J =8.0 Hz, 2H; benzene-H), 3.94 (s, 3H; CH₃), 3.93 (s, 6H; CH₃), 3.92 (s, 3H; CH₃), 3.83 (s, 6H; CH₃), 3.80 (s, 6H; CH₃), 3.79 (s, 6H; CH₃), 3.68 (s, 6H; CH₃), 3.43 (q, J =6.4 Hz, 4H; CH₂), 3.17 (d, J =5.6 Hz, 4H; CH₂), 3.00 (br, 4H; CH₂), 2.56 (t, J =7.2 Hz, 2H; CH₂), 2.32 (t, J =8.0 Hz, 2H; CH₂), 2.17 (br, 2H; CH₂), 1.78 (m, 2H; CH₂), 1.72 (m, 2H; CH₂), 1.61 (m, 2H; CH₂); ESI-TOF-MS m/z calcd for $C_{85}H_{105}Cl_2N_{30}O_{14}$ $[M + H]^+$ 1839.78; found 1839.85.

Chlorambucil-NH- β -ImPyPy- β -ImPy- γ -ImPy- β -ImPyPy-(CH₂)₃-N(CH₃)₂ (3). **3** was obtained as a white powder (1.3 mg, 0.71 μ mol, 24%) from polyamide **8**. ESI-TOF-MS m/z calcd for $C_{88}H_{110}Cl_2N_{31}O_{15}$ $[M + H]^+$ 1910.82; found 1910.81.

AcImPyPy- β -ImPy-(R)^{NH-chlorambucil}- γ -ImPy- β -ImPyPy-(CH₂)₃-N(CH₃)₂ (4). **4** was obtained as a white powder (3.9 mg, 2.1 μ mol, 71%) from polyamide **9**. ESI-TOF-MS m/z calcd for $C_{87}H_{108}Cl_2N_{31}O_{15}$ $[M + H]^+$ 1896.80; found 1896.81.

AcImPyPy- β -ImPy- γ -ImPy- β -Im-Indole-*seco*-CBI (5). **5** was synthesized according to reported procedure from polyamide **10**¹⁶. HPLC purification was performed using a Chemcobond 5-ODS-H 4.6 x 150 mm column (H₂O with 0.1% AcOH containing 35-75% CH₃CN over a linear gradient for 20 min at a flow rate of 1 mL/min). The peak around 12 min was collected and lyophilized to produce conjugate **5** was obtained (0.5 mg, 0.3 μ mol, 5%) as a brown powder. **5** was used in the DNA alkylation reaction. ESI-TOF-MS m/z calcd for $C_{78}H_{82}ClN_{26}O_{14}$ $[M + H]^+$ 1641.62; found 1641.64.

8.3 Cloning of 209 bp DNA fragments.

All DNA fragments and primers for cloning or DNA amplification were purchased from SIGMA genosys. The 32 bp DNA fragment (ODN1: 5'-AAGCAGTCAAAAACCGGAAAGCAGTCTTTT-A-3', and 5'-AAAAAGACTGCTTTCCGGTTTTTGACTGCT-TA-3') were ligated into pGEM-T Easy vectors (Promega), Escherichia coli DH5 α competent cells (Toyobo) were transformed and cultured on LB plates with 100 μ g/mL ampicillin and 32 μ g X-gal /400 μ g IPTG overnight at 37 °C. White colonies were identified by colony direct polymerase chain reaction (PCR) in 20 μ L the reaction mixtures containing 250 nM of primer set (T7: 5'-TAATACGACTCACTATAGGG-3', sp6: 5'-GATTTAG-GTGACACTATAG-3'), 200 μ M of deoxynucleotide triphosphates (Sigma Aldrich), 2 units Taq DNA polymerase and 1 X ThermoPol Reaction Buffer (New England Bio Labs). The reaction mix was incubated at 95 °C for 5 min then followed by 30 incubation cycles of 95 °C for 30 sec, 55 °C for 30 sec, and 72 °C for 30 sec with final extension step of 72 °C for 7 min. The appropriate colony was selected for transfer to 5 mL of LB medium with 100 μ g/mL ampicillin and cultured overnight at 37 °C. The plasmids with inserts were extracted using GenElute™ Plasmid Miniprep Kit (Sigma Aldrich) and identified by PCR. (program and reaction mixtures were the same as above)

8.4 Preparation of 5'-Texas Red-labeled DNA Fragments and High Resolution Gel Electrophoresis.

The 5'-Texas Red-modified 209 bp DNA fragment was prepared by PCR with 5'-TexasRed-modified sp6 and T7 primers from 1 ng/ μ L of the 32 bp fragments inserted pGEM-T Easy vector (program and reaction mixtures were the same as above). Fragments were purified by GenElute™ PCR Clean-up Kit (Sigma Aldrich), and their concentrations were determined by UV absorption.

8.5 High-Resolution Gel Electrophoresis.

The 5'-Texas Red-labeled DNA fragments (10 nM) were alkylated by various concentration of conjugates **1-5** in 10 μ L of 5 mM sodium phosphate buffer (pH 7.0) containing 10% DMF at 23 °C. After incubation for 18 h, the reaction was quenched by the

addition of 10 μ g of calf thymus DNA and heating for 10 min at 95 °C. The solution was concentrated by vacuum centrifugation. The pellet was dissolved in 7 μ L loading dye (formamide with fuschin red), heated at 95 °C for 25 min, and then immediately cooled to 0 °C. A 2 μ L aliquot was subjected to electrophoresis on a 6% denaturing polyacrylamide gel using a Hitachi DNA Sequencers.

8.6 Analysis of inhibition of the growth of the human K562 cell line.

Detailed assay procedures were reported previously²⁵. The human lymphocyte K562 cells were cultured in RPMI-1640 medium supplemented with 10% heat-inactivated fetal bovine serum, penicillin (100 IU/mL), and streptomycin (100 μ g/mL) at 37 °C in a humidified atmosphere of 95% air and 5% CO₂. Colorimetric assays with the WST-8 (Dojindo, Kumamoto, Japan) were carried out in 96-well plates. In the logarithmic growth phase, 5×10^3 cells were plated into each well in 50 μ L of culture medium. Then, medium (50 μ L) containing various concentrations of compounds 1–5 or 1% DMSO (50 μ L) as a control was added to the wells. After treatment with the compounds for 48 h, WST-8 reagent (10 μ L) was added to each well, and the cells were incubated for 3 h at 37 °C. Absorbance was measured at 450 and 600 nm using an MPR-A4I microplate reader (Tosoh). The absorbances of the control (C), treated wells (T) and treated wells at time 0 (t_0) were measured. The GI₅₀ was calculated as $100 \times [(T - t_0)/(C - t_0)] = 50$.

Acknowledgments

Mr. M. Minoshima was supported by a research fellowship of the Global COE (Center of Excellence) program, International Center for Integrated Research and Advanced Education in Material Science, Kyoto University, Japan. We would like to thank Ms. K. Hashiya for the purification of the polyamides. This work was supported by a Grant-in-Aid for Priority Research from the Ministry of Education, Culture, Sports, Science, and Technology, Japan and SORST of Japan Science and Technology (JST).

References

- Hurley, L. H. *Nat. Rev. Cancer* **2002**, 2, 188-200.
- Rajski, S. R.; Williams, R. M. *Chem. Rev.* **1998**, 98, 2723-2795.
- (a) Baker B. F.; Dervan, P. B. *J. Am. Chem. Soc.* **1985**, 107, 8266-8268. (b) Broggini, M.; Erba, E.; Ponti, M.; Ballinari, D.; Ceroni, C.; Spreafico, F.; D'Incaici, M. *Cancer Res.* **1991**, 51, 199-204. (c) Broggini, M.; Coley, H. M.; Mongelli, N.; Pesenti, E.; Wyatt, M. D.; Hartley, J. A.; D'Incaici, M. *Nucleic Acids Res.* **1995**, 23, 81-87.
- (a) Hurley, L. H.; Lee, C.-S.; McGovern, J. P.; Warpehoski, M. A.; Mitchell, M. A.; Kelly, R. C.; Aristoff, P. A. *Biochemistry* **1988**, 27, 3886-3892. (b) Yasuzawa, T.; Muroi, K.; Ichimura, M.; Takahashi, I.; Ogawa, T.; Takahashi, K.; Sano, H.; Saitoh, Y. *Chem. Pharm. Bull.* **1995**, 43, 378-391.
- For recent review, see: (a) Dervan, P. B.; Poulin-Kerstien, A. T.; Fechter, E. J.; Edelson, B. S. *Top. Curr. Chem.* **2005**, 253, 1-31. (b) Dervan, P. B.; Edelson, B. S. *Curr. Opin. Str. Biol.* **2003**, 13, 284-299.
- (a) Gottesfeld, J. M.; Neely, L.; Trauger, J. W.; Baird, E. E.; Dervan, P. B. *Nature* **1997**, 387, 202-205. (b) Olenyuk, B. Z.; Zhang, G.-J.; Klco, J. M.; Nickols, N. G.; Kaelin, Jr, W. G.; Dervan, P. B. *Proc. Natl. Acad. Sci. USA* **2004**, 101, 16768-16873. (c) Nickols, N. G.; Dervan, P. B. *Proc. Natl. Acad. Sci. USA* **2007**, 104, 10418-10423.
- For recent review, see: Bando, T.; Sugiyama, H. *Acc. Chem. Res.* **2006**, 39, 935-944.
- (a) Boger, D. L.; Johnson, D. S. *Angew. Chem. Int. Ed.* **1996**, 35, 1438-1474. (b) Boger, D. L.; Munk, S. A. *J. Am. Chem. Soc.* **1992**, 114, 5487-5496.
- Bando, T.; Iida, H.; Tao, Z.-F.; Narita, A.; Fukuda, N.; Yamori, T.; Sugiyama, H. *Chem. Biol.* **2003**, 10, 751-758.
- (a) Bando, T.; Sasaki, S.; Minoshima, M.; Shinohara, K.; Dohno, C.; Narita, A.; Sugiyama, H. *Bioconjugate Chem.* **2006**, 17, 715-720. (b) Minoshima, M.; Bando, T.; Sasaki, S.; Shinohara, K.; Shimizu, T.; Fujimoto, J.; Sugiyama, H. *J. Am. Chem. Soc.* **2007**, 129, 5384-5390. (c) Sasaki, S.; Bando, T.; Minoshima, M.; Shinohara, K.; Sugiyama, H. *Chem. Eur. J.* **2008**, 18, 864-870.
- Oyoshi, T.; Kawakami, W.; Narita, A.; Bando, T.; Sugiyama, H. *J. Am. Chem. Soc.* **2003**, 125, 4752-4754.
- (a) Shinohara, K.; Narita, A.; Oyoshi, T.; Bando, T.; Teraoka, H.; Sugiyama, H. *J. Am. Chem. Soc.* **2004**, 126, 5113-5118. (b) Shinohara, K.; Sasaki, S.; Minoshima, M.; Bando, T.; Sugiyama, H. *Nucleic Acids Res.* **2006**, 34, 1189-1195.
- Dickinson, L. A.; Burnett, R.; Melander, C.; Edelson, B. S.; Arora, P. S.; Dervan, P. B.; Gottesfeld, J. M. *Chem. Biol.* **2004**, 11, 1583-1594.
- (a) Alvarez, D.; Chou, C. J.; Latella, L.; Zeitlin, S. G.; Ku, S.; Puri, P. L.; Dervan, P. B.; Gottesfeld, J. M. *Cell Cycle* **2006**, 5, 1537-1548. (b) Chou, C. J.; Farkas, M. E.; Tsai, S. M.; Alvarez, D.; Dervan, P. B.; Gottesfeld, J. M. *Mol. Cancer Ther.* **2008**, 7, 769-778. (c) Chou, C. J.; O'Hare, T.; Lefebvre, S.; Alvarez, D.; Tyner, J. W.; Eide, C. A.; Druker, B. J.; Gottesfeld, J. M. *PLoS ONE* **2008**, 3, e3593-3600.
- Tsai, S. M.; Farkas, M. E.; Chou, C. J.; Gottesfeld, J. M.; Dervan, P. B. *Nucleic Acids Res.* **2007**, 35, 307-316.
- (a) Baird, E. E.; Dervan, P. B. *J. Am. Chem. Soc.* **1996**, 118, 6141-6146. (b) Wurtz, N. R.; Turner, J. M.; Baird, E. E.; Dervan, P. B. *Org. Lett.* **2001**, 3, 1201-1203. (c) Belitsky, J. M.; Nguyen, D. H.; Wurtz, N. R.; Dervan, P. B. *Bioorg. Med. Chem.* **2002**, 10, 2767-2774.
- Sasaki, S.; Bando, T.; Minoshima, M.; Shimizu, T.; Shinohara, K.; Takaoka, T.; Sugiyama, H. *J. Am. Chem. Soc.* **2006**, 128, 12162-12168.
- (a) Bando, T.; Narita, A.; Saito, I.; Sugiyama, H. *Chem. Eur. J.* **2002**, 8, 4781-4790. (b) Bando, T.; Narita, A.; Saito, I.; Sugiyama, H. *J. Am. Chem. Soc.* **2003**, 125, 3471-3485. (c) Bando, T.; Narita, A.; Asada, K.; Ayame, H.; Sugiyama, H. *J. Am. Chem. Soc.* **2004**, 126, 8948-8955. (d) Bando, T.; Narita, A.; Sasaki, S.; Sugiyama, H. *J. Am. Chem. Soc.* **2005**, 127, 13890-13895.
- Farkas, M. E.; Tsai, S. M.; Dervan, P. B. *Bioorg. Med. Chem.* **2007**, 15, 6927-6936.
- (a) Chang, A. Y.; Dervan, P. B. *J. Am. Chem. Soc.* **2000**, 122, 4856-4864. (b) Wurtz, N. R.; Dervan, P. B. *Chem. Biol.* **2000**, 7, 153-161. (c) Wang, Y.-D.; Dziegielewska, J.; Wurtz, N. R.; Dziegielewska, B.; Dervan, P. B.; Beerman, T. A. *Nucleic Acids Res.* **2003**, 31, 1208-1215.
- (a) Bosanquet, A. G. *Cancer Chemother. Pharmacol.* **1985**, 14, 83-95. (b) Giraud, I.; Rapp, M.; Maurizis, J.-C.; Madelmont, J.-C. *J. Med. Chem.* **2002**, 45, 2116-2119.
- Edelson, B. S.; Best, T. P.; Olenyuk, N. G.; Nickols, N. G.; Doss, R. M.; Foister, S.; Heckel, A.; Dervan, P. B. *Nucleic Acids Res.* **2004**, 32, 2802-2818.
- Brooks, N.; McHugh, P. J.; Lee, M.; Hartley, J. A. *Chem. Biol.* **2000**, 7, 659-668.
- Bando, T.; Narita, A.; Iwai, A.; Kihara, K.; Sugiyama, H. *J. Am. Chem. Soc.* **2004**, 126, 3406-3407.
- Shinohara, K.; Bando, T.; Sasaki, S.; Sakakibara, Y.; Minoshima, M.; Sugiyama, H. *Cancer Sci.* **2006**, 97, 219-225.