数理解析研究所講究録
664巻 1988年 74-78

NOTES ON P-VALENTLY BAZILEVIĆ FUNCTIONS

SHIGEYOSHI OWA (近畿大・理工　尾和重義)

ABSTRACT

The object of the present paper is to improve the former results for p-valently Bazilević functions which were recently proved by the author and others.

I. INTRODUCTION

Let A_p denote the class of functions of the form

$$f(z) = z^p + \sum_{n=p+1}^{\infty} a_n z^n \qquad (p \in N = \{1,2,3,\dots\}) \tag{1.1}$$

which are analytic in the unit disk $E = \{z: |z| < 1\}$. A function f(z) belonging to A_p is said to be p-valently starlike if and only if it satisfies the condition

$$\mathrm{Re}\left\{ \frac{zf'(z)}{f(z)} \right\} > 0 \qquad (z \in E). \tag{1.2}$$

We denote by S_p^* the subclass of A_p consisting of functions which are p-valently starlike in E.

A function f(z) belonging to the class A_p is said to be p-valently Bazilević of type β and order γ if there exists a function g(z) belonging to S_p^* such that

$$\mathrm{Re}\left\{ \frac{zf'(z)}{f(z)^{1-\beta} g(z)^{\beta}} \right\} > \gamma \qquad (z \in E) \tag{1.3}$$

for some β ($\beta > 0$) and γ ($0 \leq \gamma < p$).

1980 Mathematics Subject Classification (1985 Revision). Primary 30C45.

* This research of the author was carried out at Research Institute for Mathematical Sciences, Kyoto University while the author was visiting from Kinki University.

Also we denote by $B_p(\beta,\gamma)$ the subclass of A_p consisting of all p-valently Bazilevič functions of type β and order γ in E. The concept of Bazilevič functions was first introduced by Bazilevič [6]. Thomas [8] has called a function in the class $B_1(\beta,0)$ a Bazilevič function of type β. Further, Nunokawa [3] has proved that a function f(z) in the class $B_p(\beta,0)$ is p-valent in the unit disk E.

In particular, the class $B_p(\beta,\gamma)$ for g(z) = f(z) is called is the class of p-valently starlike functions of order γ. Further, we note that the class $B_p(\beta,0)$ for g(z) = f(z) is equivalent to S_p^*.

Let $A_p(\alpha,\beta)$ be the subclass of A_p consisting of functions which satisfy the condition

$$\text{(1.4)} \qquad \mathrm{Re}\left\{(1-\alpha)\frac{zf'(z)}{f(z)} + \alpha\left(1+\frac{zf''(z)}{f'(z)}\right)\right\} > \beta \qquad (z \in E)$$

for some real α and β.

The class $A_1(\alpha,0)$ when p = 1 and $\beta = 0$ was introduced by Mocanu [2], and was studied by Miller, Mocanu and Reade [1], and Sakaguchi and Fukui [7].

Let $C_p(\alpha,\beta)$ be the subclass of A_p consisting of functions satisfying the condition

$$\text{(1.5)} \qquad \mathrm{Re}\left\{(1-\alpha)\frac{zf'(z)}{f(z)} + \alpha\left(1+\frac{zf''(z)}{f'(z)}\right)\right\} < \beta \qquad (z \in E)$$

for some real α and β ($\beta > p$).

The class $C_p(\alpha,\beta)$ was recently introduced by Nunokawa and Owa [4].

2. SOME PROPERTIES

In order to derive our results, we need the following lemmas.

LEMMA I ([5]). If a function f(z) belongs to the class $A_p(\alpha,\beta)$ with $\alpha > 0$ and $0 \leq -\beta/\alpha \leq 1/2$, then $f(z) \in B_p(1/\alpha, 2^{2\beta/\alpha})$, and therefore f(z) is p-valent in the unit disk E.

LEMMA 2 ([4]). Let a function $f(z)$ belong to the class $C_p(\alpha,\beta)$ with $\alpha \neq 0$, $\beta > p$, and $|\beta/\alpha| \leqq 1/2$. Then $f(z)$ is p-valent in the unit disk E. Moreover, if $0 \leqq -\beta/\alpha \leqq 1/2$, then $f(z) \in B_p(1/\alpha, 2^{2\beta/\alpha})$.

Applying the above lemmas, we prove

THEOREM I. If a function $f(z)$ belongs to the class $A_p(\alpha,\beta)$ with $\alpha > 0$ and $0 \leqq -\beta/\alpha \leqq 1/2$, then $f(z) \in B_p(1/\alpha, p2^{2\beta/\alpha})$.

PROOF. For a function $f(z)$ in the class A_p, we define the function $g(z)$ by

$$(2.1) \qquad g(z) = f(z)^{1/p} = z + g_2 z^2 + g_3 z^3 + \cdots .$$

Then $g(z)$ is in the class A_1, and satisfies

$$(2.2) \qquad \frac{zf'(z)}{f(z)} = p\,\frac{zg'(z)}{g(z)}$$

and

$$(2.3) \qquad 1 + \frac{zf''(z)}{f'(z)} = 1 + \frac{zg''(z)}{g'(z)} + (p-1)\,\frac{zg'(z)}{g(z)} .$$

It follows from (2.2) and (2.3) that

$$(2.4) \qquad (1-\alpha)\,\frac{zf'(z)}{f(z)} + \alpha\left(1 + \frac{zf''(z)}{f'(z)}\right) = (p-\alpha)\,\frac{zg'(z)}{g(z)} + \alpha\left(1 + \frac{zg''(z)}{g'(z)}\right).$$

Therefore, we have

$$(2.5) \qquad f(z) \in A_p(\alpha,\beta) \iff \mathrm{Re}\left\{\left(1 - \frac{\alpha}{p}\right)\frac{zg'(z)}{g(z)} + \frac{\alpha}{p}\left(1 + \frac{zg''(z)}{g'(z)}\right)\right\} > \frac{\beta}{p}$$

$$\Longleftrightarrow g(z) \in A_1(\alpha/p, \beta/p).$$

Applying Lemma 1 for p = 1, we see that

$$g(z) \in A_1(\alpha/p, \beta/p) \Longrightarrow g(z) \in B_1(p/\alpha, 2^{2\beta/\alpha}).$$

It follows that

$$f(z) \in A_p(\alpha, \beta) \Longrightarrow g(z) \in B_1(p/\alpha, 2^{2\beta/\alpha})$$

$$\Longleftrightarrow \mathrm{Re}\left\{ \frac{zg'(z)}{g(z)^{1-p/\alpha} h(z)^{p/\alpha}} \right\} > 2^{2\beta/\alpha} \qquad (h(z) \in S_1^*)$$

$$\Longleftrightarrow \mathrm{Re}\left\{ \frac{zf'(z)}{f(z)^{1-1/\alpha} (h(z)^p)^{1/\alpha}} \right\} > p2^{2\beta/\alpha} \qquad (h(z)^p \in S_p^*)$$

$$\Longleftrightarrow f(z) \in B_p(1/\alpha, p2^{2\beta/\alpha}).$$

This completes the assertion of Theorem 1.

REMARK I. Noting $p2^{2\beta/\alpha} \geqq 2^{2\beta/\alpha}$, we see that

$$B_p(1/\alpha, p2^{2\beta/\alpha}) \subseteq B_p(1/\alpha, 2^{2\beta/\alpha}).$$

Thus Theorem 1 is the improvement of Lemma 1 by Nunokawa, Owa, Saitoh, Yaguchi and Lee [5].

Taking p = 1 in Theorem 1, we have

COROLLARY I. If $f(z) \in A_1(\alpha, \beta)$ with $\alpha > 0$ and $0 \leqq -\beta/\alpha \leqq 1/2$, then $f(z) \in B_1(1/\alpha, 2^{2\beta/\alpha})$.

Using the same manner as in the proof of Theorem 1, we have

THEOREM 2. If a function f(z) belongs to the class $C_p(\alpha, \beta)$ with

$\alpha \neq 0$, $\beta > p$, and $0 \leqq -\beta/\alpha \leqq 1/2$, then $f(z) \in B_p(1/\alpha, p2^{2\beta/\alpha})$.

Finally, letting $p = 1$ in Theorem 2, we have

COROLLARY 2. If $f(z) \in C_1(\alpha,\beta)$ with $\alpha \neq 0$, $\beta > 1$, and $0 \leqq -\beta/\alpha \leqq 1/2$, then $f(z) \in B_1(1/\alpha, 2^{2\beta/\alpha})$.

REMARK 2. We note that Theorem 2 is the improvement of Lemma 2 due to Nunukawa and Owa [4].

REFERENCES

[1] S. S. Miller, P. T. Mocanu and M. O. Reade, All α-convex functions are univalent and starlike, Proc. Amer. Math. Soc. 37(1973), 553 - 554.

[2] P. T. Mocanu, Une proprieté de convexté generalisée dans la théorie de la representation conforme, Mathematica (Cluj) 11(1969), 127 - 133.

[3] M. Nunokawa, On the Bazilevič analytic functions, Sci. Rep. Fac. Edc. Gunma Univ. 21(1972), 9 - 13.

[4] M. Nunokawa and S. Owa, A remark on certain multivalent functions, Chinese J. Math., to appear.

[5] M. Nunokawa, S. Owa, H. Saitoh, T. Yaguchi and S. K. Lee, On certain subclass of analytic functions. II, Sci. Rep. Fac. Edc. Gunma Univ. 36(1988), in press.

[6] E. Bazilevič, On a class of integrability in quadratures of the Loewner-Kufarev equation, Mat. Sb. 37(1955), 471 - 476.

[7] K. Sakaguchi and S. Fukui, On alpha-starlike functions and related functions, Bull. Nara Univ. Educ. 28(1979), 5 - 12.

[8] D. K. Thomas, On Bazilevič functions, Trans. Amer. Math. Soc. 132(1968), 353 - 361.

Department of Mathematics
Kinki University
Higashi-Osaka, Osaka 577
Japan