数理解析研究所講究録
700巻 1989年 64-84

CFT on $\mathbb{P}^1$ and Monodromy Representations of the Braid Groups

(Talk at Research Institute of Mathematical Sciences, Kyoto Univ.,July 19,1988)

Yukihiro Kanie

Mie University, Fac.of Education, Dept.of Math.

§0. From the differential equations of N-point functions of vertex operators in the conformal field theory on $\mathbb{P}^1$, arise the monodromy representations of the braid group B_N. In the meeting of last year, I reported that these monodromy representations give "all" irreducuble representations of the Hecke algebra $H_N(q)$ of type A_{N-1} (obtained by H.Wenzl[W]) associated with the affine Lie algebra of type $A_n^{(1)}$. In this meeting, I will report that associated with the affine Lie algebras of type $B_n^{(1)}$, $C_n^{(1)}$ and $D_n^{(1)}$, the monodromy representations of the group B_N give "all" irreducible representations of the Birman-Wenzl-Murakami algebra, the q-analogue of Brauer's centralizer algebras. Very important is Jimbo-Miwa-Okado's

calculations[JMO], and in the case of type $C_n^{(1)}$ the representations are equivalent to the ones obtained by J.Murakami[M].

§1. Let $\mathfrak{g}$ be the simple Lie algebra of type X_n, and $\hat{\mathfrak{g}} = \mathfrak{g}\otimes\mathbb{C}[t,t^{-1}] \oplus \mathbb{C}c$ be the affine Lie algebra of type $X_n^{(1)}$. Fix an integer $\ell\geq 1$ and introduce the number $\kappa=\ell+g$, where g is the dual Coxeter number of $\hat{\mathfrak{g}}$.

Denote by P_+ the set of dominant integral weights of $\mathfrak{g}$ and by P_ℓ the set of elements $\lambda\in P_+$ satisfying $(\theta,\lambda)\leq\ell$, where θ is the maximum root. For a weight $\lambda\in P_\ell$, we denote by V_λ the irreducible representation of $\mathfrak{g}$ of highest weight λ, by $\mathcal{H}_\lambda$ the integrable representation of $\hat{\mathfrak{g}}$ of highest weight $\ell\Lambda_0+\lambda$ and by $|\lambda\rangle$ the (fixed) highest weight cyclic vector of V_λ and $\mathcal{H}_\lambda$.

The Virasoro algebra also acts on $\mathcal{H}_\lambda$ by the Sugawara forms $L(m), m\in\mathbb{Z}$, and the space $\mathcal{H}_\lambda$ is graded by means of the eigenspace decomposition w.r.t. the operator $L(0)$:

$$\mathcal{H}_\lambda = \sum_{d\in\mathbb{Z}_{\geq 0}} \mathcal{H}_\lambda(d) \ , \ \mathcal{H}_\lambda(d) = \{v\in\mathcal{H}_\lambda;\ L(0)v = (\Delta_\lambda+d)v\} \ ,$$

where $\Delta_\lambda = \dfrac{(\lambda,\lambda+2\rho)}{2\kappa}$ and ρ is the half sum of positive roots of $\mathfrak{g}$. Note that $\dim\mathcal{H}_\lambda(d)<\infty$ and $\mathcal{H}_\lambda(0) \cong V_\lambda$.

There are dual right $\mathfrak{g}$ and $\hat{\mathfrak{g}}$-module V_λ^+ and $\mathcal{H}_\lambda^+$ of V_λ and $\mathcal{H}_\lambda$, and the nondegenerate invariant bilinear form $\langle\ |\ \rangle$ on $V_\lambda^+ \times V_\lambda$ and $\mathcal{H}_\lambda^+ \times \mathcal{H}_\lambda$ with the normalized condition $\langle\lambda|\lambda\rangle = 1$ where $\langle\lambda|$ is a fixed highest weight vector of $V_\lambda^+ = \mathcal{H}_\lambda^+(0)$ and $\mathcal{H}_\lambda^+$.

A triple $\mathbb{v} = \begin{pmatrix} \lambda \\ \mu_2 \mu_1 \end{pmatrix}$ of weights in P_ℓ is called a vertex and is drawn as

$\mathbb{v} =$ 

.

A multi-valued, holomorphic function

$$\Phi(z) : V_\lambda \otimes \mathcal{H}_{\mu_1} \longrightarrow \hat{\mathcal{H}}_{\mu_2} = \prod_{d \in \mathbb{Z}_{\geq 0}} \mathcal{H}_{\mu_2}(d)$$

on $\mathbb{P}^1 \setminus \{0, \infty\}$ is called a *vertex operator of type* $\begin{pmatrix} \lambda \\ \mu_2 \mu_1 \end{pmatrix}$ (sometimes called *of weight* λ), if it satisfies the following:

(Gauge Condition) $[X(m),\ \Phi(z)(u \otimes \cdot)] = z^m \Phi(z)(Xu \otimes \cdot)$

$(X \in \mathfrak{g}, m \in \mathbb{Z}, u \in V_\lambda)$;

(Eq.of Motion) $[L(m),\ \Phi(z)] = z^m \{ z\frac{d}{dz} + (m+1)\Delta_\lambda \} \Phi(z)$,

where $X(m)=X\otimes t^m$ and the number Δ_λ is called the *conformal dimension* of the vertex operator $\Phi(z)$.

Denote by $\mathcal{V}er(\mathbb{v})$ the space of all vertex operators of type $\mathbb{v}$, and introduce the space

$$\mathcal{V}er(\lambda) = \sum_{\mu_1,\mu_2\in P_\ell} \mathcal{V}er\left(\begin{bmatrix}\lambda\\ \mu_2,\mu_1\end{bmatrix}\right)$$

of all vertex operators of weight λ.

Introduce the subalgebra $\mathfrak{t}_\theta = \mathbb{C}\langle X_\theta, [X_\theta, X_{-\theta}], X_{-\theta}\rangle \cong \mathfrak{sl}(2;\mathbb{C})$ of $\mathfrak{g}$ and the subspace $\mathcal{V}(\mathbb{v})$ of $\mathrm{Hom}_{\mathfrak{g}}(V_\lambda\otimes V_{\mu_1}, V_{\mu_2})$ defined by

$$\mathcal{V}(\mathbb{v}) = \cap \operatorname{Ker} \pi_{\mathfrak{t}_\theta}(j,j_1,j_2)$$

where the intersection is taken over the set $\{j,j_1,j_2\in \frac{1}{2}\mathbb{Z}_{\geq 0};\ j+j_1+j_3>\ell\}$, and $\pi_{\mathfrak{t}_\theta}(j,j_1,j_2)(\varphi) \in \mathrm{Hom}_{\mathfrak{t}_\theta}(W_j\otimes W_{j_1}, W_{j_2})$ is defined as

$$\pi_{\mathfrak{t}_\theta}(j,j_1,j_2)(\varphi) = \mathrm{proj}_{W_{j_2}}\circ\varphi|_{W_j\otimes W_{j_1}} \quad (\varphi\in\mathrm{Hom}_{\mathfrak{g}}(V_\lambda\otimes V_{\mu_1}, V_{\mu_2}))$$

where W_j, W_{j_1}, W_{j_2} are $\mathfrak{t}_\theta$-simple submodules of $V_\lambda, V_{\mu_1}, V_{\mu_2}$ with spin j, j_1, j_2 respectively.

By Equation of Motion, Φ is expressed as a formal

Laurent series

$$\Phi(z) = \sum_{m\in\mathbb{Z}} \Phi(m)\ z^{-m-\hat{\Delta}(\mathbb{v})} ,$$

where $\hat{\Delta}(\mathbb{v}) = \Delta_\lambda+\Delta_{\mu_1}-\Delta_{\mu_2}$ and $\Phi(m)$ is homogeneous of degree m, i.e.

$$\Phi(m):\ V_\lambda\otimes\mathcal{H}_{\mu_1}(d) \longrightarrow \mathcal{H}_{\mu_2}(d-m) \qquad \text{for any } d.$$

The principal branch of $\Phi(z)$ is taken such as the value of $z^{-\hat{\Delta}(\mathbb{v})}$ is positive for $z\in\mathbb{R}_+=\{z\in\mathbb{R};\ z>0\}$ and uniquely continued to the region $\mathbb{C}_+=\{z\in\mathbb{C};\ \mathrm{Im}z>0\}$, and we refer this for the value of $\Phi(z)$ on $\mathbb{C}_+$.

For any vertex operator $\Phi\in\mathcal{V}er(\mathbb{v})$, its initial term $\varphi = \Phi(0)\big|_{V_\lambda\otimes\mathcal{H}_{\mu_1}(0)} = \mathrm{proj}_{V_{\mu_2}}\circ z^{\hat{\Delta}(\mathbb{v})}\Phi(z)\ \big|_{V_\lambda\otimes V_{\mu_1}}$ belongs to $\mathcal{V}(\mathbb{v})$. Under this correspondence,

Theorem 1. *The space $\mathcal{V}er(\mathbb{v})$ of N-point functions of type $\mathbb{v}$ is isomorphic with the space $\mathcal{V}(\mathbb{v})$ of initial terms of type $\mathbb{v}$.*

Call $\mathbb{v}$ ℓCG *(ℓ-constrained Clebsch-Gordan)* vertex, if $\mathcal{V}(\mathbb{v}) \neq 0$, and denote by ($\ell$CG) the set of all ℓCG vertices.

For each $\varphi\in\mathcal{V}(\mathbb{V})$, denote by Φ_φ the vertex operator with the initial term φ.

Notes. i) Even if we asuume that $\lambda\in P_+$ and $\mu_i\in P_\ell$, $\mathcal{V}\left(\begin{bmatrix}\lambda\\ \mu_2 & \mu_1\end{bmatrix}\right)\neq 0$ implies that $\lambda\in P_\ell$.

ii) Operator product expansions of *currents* $X(z) = \sum_{m\in\mathbb{Z}} X(m)z^{-m-1}$ $(X\in\mathfrak{g})$ and the *energy-momentum tensor* $T(z) = \sum_{m\in\mathbb{Z}} L(m)z^{-m-2}$ with vertex operators allow the extension of the vertex operators $\Phi(z)$ of type $\begin{bmatrix}\lambda\\ \mu_2\mu_1\end{bmatrix}$ to the operators $\Phi(z)\colon \mathcal{H}_\lambda\otimes\mathcal{H}_{\mu_1}\longrightarrow\hat{\mathcal{H}}_{\mu_2}$ by means of contour integrals. (*Nuclear Democracy*)

iii) By the same arguments as in §3, the analytic continuation of a vertex operator Φ of type $\begin{bmatrix}\lambda\\ \mu_2\mu_1\end{bmatrix}$ along the path γ_0 gives a vertex operator of type $\begin{bmatrix}\lambda\\ \mu_1\mu_2\end{bmatrix}$, where

$$\gamma_0(t)= z\ e^{\pi\sqrt{-1}t}\ ,\quad t\in[0,1]\ ,\ z\in\mathbb{R}_+\ .$$

This gives an isomorphism C_{γ_0} of $\mathcal{V}er\begin{bmatrix}\lambda\\ \mu_2\mu_1\end{bmatrix}$ to $\mathcal{V}er\begin{bmatrix}\lambda\\ \mu_1\mu_2\end{bmatrix}$ and the corresponding isomorphism

$$C_{\gamma_0}\colon \mathcal{V}\begin{bmatrix}\lambda\\ \mu_2\mu_1\end{bmatrix}\longrightarrow\mathcal{V}\begin{bmatrix}\lambda\\ \mu_1\mu_2\end{bmatrix}$$

is given by

$$C_{\gamma_0} = e^{\pi\sqrt{-1}\,\hat{\Delta}(\mathbb{v})}\,T\ ,$$

where T is the transposition:

$$T : \mathrm{Hom}(V_\lambda \otimes V_{\mu_1}, V_{\mu_2}) \longrightarrow \mathrm{Hom}(V_{\mu_1} \otimes V_\lambda, V_{\mu_2})$$

$$(T\varphi)(u\otimes v) = \varphi(v\otimes u).$$

A vertex operator $\Phi(z)$ of type $\mathbb{v}$ is also considered as an operator from $\mathcal{H}_{\mu_1}$ to $\hat{\mathcal{H}}_{\mu_2}$ parametrized by V_λ, i.e.

$$\Phi(u;z)(v) = \Phi(z)(u\otimes v) \qquad (u\in V_\lambda,\ v\in\mathcal{H}_{\mu_1}).$$

§2. It is convenient to introduce the spaces $\mathcal{H} = \sum_{\lambda\in P_\ell} \mathcal{H}_\lambda$ and $\hat{\mathcal{H}} = \sum_{\lambda\in P_\ell} \hat{\mathcal{H}}_\lambda$ and consider vertex operators as linear operators of $\mathcal{H}$ to $\hat{\mathcal{H}}$. The vacuum $|0\rangle$ of $\mathcal{H}_0$ is called a *Virasoro vacuum*, since $L(m)|0\rangle = 0$ for $m\geq -1$. Note that $V_0 = \mathbb{C}|0\rangle$.

For an N-ple $\Lambda=(\lambda_N,\cdots,\lambda_1)$ of weights in P_ℓ, denote

$$V_\Lambda = V_{\lambda_N}\otimes\cdots\otimes V_{\lambda_1} \quad \text{and} \quad V_g^\vee(\Lambda) = \mathrm{Hom}_g(V_\Lambda,\mathbb{C})\ .$$

For any vertex operators $\Phi^i(z_i)$ of weight $\lambda_i\,(1\leq i\leq N)$,

$$\langle 0|\Phi^N(z_N)\cdots\Phi^1(z_1)|0\rangle$$

is the coefficient of $|0\rangle$ the iterated application $\Phi^N(z_N)\cdots\Phi^1(z_1)|0\rangle$ to the vector $|0\rangle$, and this is a $V_g^\vee(\Lambda)$-valued formal Laurent series in $z_N,\cdots,z_1$ called the *N-point function of weight* Λ and is denoted by $\langle\Phi^N(z_N)\cdots\Phi^1(z_1)\rangle$. Denote by $\mathcal{V}er(\Lambda)$ the space of all N-point functions of weight Λ.

The space $V_g^\vee(\Lambda)$ is decomposed as

$$V_g^\vee(\Lambda) = \sum_{\mu} V_g^\vee(\Lambda)_\mu , \qquad \mu=(\mu_{N-1},\cdots,\mu_1) \in (P_+)^{N-1} ;$$

$$V_g^\vee(\Lambda)_\mu \xleftarrow[\cong]{C_\Lambda} \mathrm{Hom}_g(V_{\lambda_N}\otimes V_{\mu_{N-1}},V_0)\otimes\cdots\otimes\mathrm{Hom}_g(V_{\lambda_i}\otimes V_{\mu_{i-1}},V_{\mu_i})$$
$$\otimes\cdots\otimes\mathrm{Hom}_g(V_{\lambda_1}\otimes V_0,V_{\mu_1}),$$

where the identification C_Λ is given by

$$C_\Lambda(\varphi_N\otimes\cdots\otimes\varphi_1)(u_N\otimes\cdots\otimes u_1)$$
$$= \langle 0|\varphi_N(u_N\otimes\varphi_{N-1}(\cdots\otimes\varphi_2(u_2\otimes\varphi_1(u_1\otimes|0\rangle)\cdots)\rangle$$
$$= \langle 0|\varphi_N(u_N)\circ\cdots\circ\varphi_1(u_1)(|0\rangle) ,$$

for $\varphi_i\in \mathrm{Hom}_g(V_{\lambda_i}\otimes V_{\mu_{i-1}},V_{\mu_i}) \cong \mathrm{Hom}_g(V_{\lambda_i},\mathrm{Hom}(V_{\mu_{i-1}},V_{\mu_i}))$ $(1\le i\le N;\ \mu_N=\mu_0=0)$, and $u_N\otimes\cdots\otimes u_1\in V_\Lambda$.

Introduce the subspace $\mathcal{V}(\Lambda)$ of $V_{\mathfrak{g}}^{\vee}(\Lambda)$ defined, through C_{Λ}, by

$$\mathcal{V}(\Lambda) = \sum_{\mu} \mathcal{V}(\Lambda)_{\mu} , \qquad \mu=(\mu_{N-1},\cdots,\mu_1) \in (P_{\ell})^{N-1} ;$$

where

$$\mathcal{V}(\Lambda)_{\mu} = \mathcal{V}(v_N(\mu))\otimes\cdots\otimes\mathcal{V}(v_i(\mu)) \otimes\cdots\otimes\mathcal{V}(v_1(\mu)) \subset V_{\mathfrak{g}}^{\vee}(\Lambda)$$

and

$$v_N(\mu)=\begin{bmatrix}\lambda_N & \\ 0 & \mu_{N-1}\end{bmatrix},\cdots,\ v_i(\mu)=\begin{bmatrix}\lambda_i & \\ \mu_i & \mu_{i-1}\end{bmatrix},\cdots,\ v_1(\mu)=\begin{bmatrix}\lambda_1 & \\ \mu_1 & 0\end{bmatrix}.$$

Then the space $\mathcal{V}(\Lambda)$ is isomorphic to $\mathcal{V}er(\Lambda)$ of N-point functions of weight Λ as follows: to each $\varphi = C(\varphi_N\otimes\cdots\otimes\varphi_1)\in\mathcal{V}(\Lambda)$, assign the N-point function

$$\Phi_{\varphi_N\otimes\cdots\otimes\varphi_1}(z) = \langle\Phi_{\varphi_N}(z_N)\cdots\Phi_{\varphi_1}(z_1)\rangle \in \mathcal{V}er(\Lambda) .$$



Now introduce a system $KZ(\Lambda)$ of differential equations on $\mathrm{Hom}_{\mathfrak{g}}(V_{\Lambda},\mathbb{C})$-valued functions $\Phi(z)$ on $X_N = \{z=(z_N,\cdots,z_1) \in\mathbb{C}^N;\ z_i\neq z_j\ (i\neq j)\}$

$$\mathrm{KZ}(\Lambda) \qquad \left(\kappa\frac{\partial}{\partial z_i} - \sum_{\substack{k=1\\k\neq i}}^{N} \frac{\Omega_{ik}}{z_i - z_k}\right) \Phi(z) = 0 \qquad (1\leq i\leq N)$$

due to Knizhnik-Zamolodchikov[KZ], where

$$\Omega_{ik} = \sum_{a=1}^{\dim \mathfrak{g}} \rho_i(X^a)\ \rho_k(X_a)\ ,$$

ρ_i denotes the $\mathfrak{g}$-action on the i-th component of $\mathrm{Hom}(V_\Lambda, \mathbb{C})$ and $\{X^a\}$ and $\{X_a\}$ are dual bases of $\mathfrak{g}$.

Further introduce an additional ℓ-constraint condition, *i.e.* a system $\ell C(\Lambda)$ of algebraic equations

$$\ell C(\Lambda) \quad \sum_{|\mathbf{m}_i|=L_i} \binom{L_i}{\mathbf{m}_i} \prod_{k\neq i} (z_k - z_i)^{-m_k} \Phi(z)(X_\theta^{m_N} u_N, \cdots, |\lambda_i\rangle, \cdots, X_\theta^{m_1} u_1) = 0,$$

$$(1\leq i\leq N)$$

for any $u_k \in V_{\lambda_k}$ $(k\neq i)$, where $\mathbf{m}_i = (m_N, \cdots, \hat{m}_i, \cdots, m_1) \in (\mathbb{Z}_{\geq 0})^{N-1}$, $|\mathbf{m}_i| = \sum_{k\neq i} m_k$, $L_i = \ell - (\lambda_i, \theta) + 1$ and $\binom{L_i}{\mathbf{m}_i}$ is the multinomial coefficient.

Remark. The system $\mathrm{KZ}(\Lambda)$ of differential equations is completely integrable because of the infinitesimal pure braid relations among the operators Ω_{ik} (see [A]). The system $\ell C(\Lambda)$ is compatible with the system $\mathrm{KZ}(\Lambda)$.

Any N-point function of weight Λ satisfies the systems

of $KZ(\Lambda)$ and $\ell C(\Lambda)$. Hence

Theorem 2.

i) For any N-ple Λ of weights in P_ℓ, any N-point function $\langle\Phi^N(z_N)\cdots\Phi^1(z_1)\rangle$ of weight Λ is absolutely convergent in the region $\mathcal{R}_N$, and is analytically continued to a multivalued holomorphic function on X_N, where $\mathcal{R}_N$ is defined by

$$\mathcal{R}_N = \{z=(z_N,\cdots,z_1)\in\mathbb{C}_+^N;\ |z_N|>\cdots>|z_1|\} \subset X_N .$$

ii) The solution space of the joint system $KZ(\Lambda)$ and $\ell C(\Lambda)$ is isomorphic with $\mathcal{V}er(\Lambda)$, hence with $\mathcal{V}(\Lambda)$.

Note. If $v = \begin{bmatrix} \lambda & \\ \mu & 0 \end{bmatrix} \in (CG)$, then $\mu = \lambda$, $\hat{\Delta}(v) = 0$, and $\mathcal{V}(v) \cong \mathrm{Hom}_{\mathfrak{g}}(V_\lambda, V_\lambda) = \mathbb{C}\mathrm{id}$.

If $v = \begin{bmatrix} \lambda & \\ 0 & \mu \end{bmatrix} \in (CG)$, then $\mu = \lambda^+$, $\hat{\Delta}(v) = 2\Delta_\lambda$, and $\mathcal{V}(v) \cong \mathrm{Hom}_{\mathfrak{g}}(V_\lambda \otimes V_{\lambda^+}, \mathbb{C}) = \mathbb{C}\nu$, where the *anti-weight* λ^+ of λ is defined as $-\lambda^+ (=w_0\lambda)$ is the lowest weight of V_λ and ν is normalised as $\nu(|\lambda\rangle\otimes w_0|\lambda^+\rangle)=1$, where w_0 is the longest element of the Weyl group of $\mathfrak{g}$.

3-point functions are essentially nothing but vertex

operators. The assignation to $\varphi\in\mathcal{V}\left(\begin{bmatrix}\lambda\\ \mu_2 & \mu_1\end{bmatrix}\right)$ the element

$\nu\otimes\varphi\otimes\mathrm{id}\in\mathcal{V}(\mu_2^+,\lambda,\mu_1)$,

$$\nu\otimes\varphi\otimes\mathrm{id}(|u\rangle\otimes|v\rangle\otimes|w\rangle)$$
$$=\nu(|u\rangle\otimes\varphi(|v\rangle\otimes|w\rangle)\qquad\left[|u\rangle\otimes|v\rangle\otimes|w\rangle\in V_{\mu_2^+}\otimes V_\lambda\otimes V_{\mu_1}\right],$$

gives the isomorphism between them. Hence the space $\mathcal{V}er\left(\begin{bmatrix}\lambda\\ \mu_2 & \mu_1\end{bmatrix}\right)$ of vertex operators is isomorphic with the space $\mathcal{V}er(\mu_2^+,\lambda,\mu_1)$ of 3-point functions. More precisely, the classical sector $\mathrm{proj}_{V_{\mu_2}}\circ\Phi_\varphi(z)|_{V_\lambda\otimes V_{\mu_1}}$ of the vertex operator $\Phi_\varphi(z)$ is given by

$$\lim_{z_t\nearrow\infty}\lim_{z_s\searrow 0} z^{2\Delta_{\mu_2}}\langle\Phi_\nu(z_t)\Phi_\varphi(z)\Phi_{\mathrm{id}}(z_s)\rangle.$$

§3. Denote by $\mathcal{V}er(\mathbb{v}_2)\circ\mathcal{V}er(\mathbb{v}_1)$ the space of compositions $\Phi^2(z_2)\Phi^1(z_1)$ of vertex operators Φ^i of type $\mathbb{v}_i$. Then

$$\sum_{\mu\in P_\ell}\mathcal{V}er\left(\begin{bmatrix}\lambda_2\\ \mu_t & \mu\end{bmatrix}\right)\circ\mathcal{V}er\left(\begin{bmatrix}\lambda_1\\ \mu & \mu_s\end{bmatrix}\right)\cong\mathcal{V}er(\Lambda)\cong\mathcal{V}(\Lambda),$$

where $\Lambda=(\mu_t^+,\lambda_2,\lambda_1,\mu_s)$.

The composition $\Phi^2(z_2)\Phi^1(z_1)$ is determined by the classical sector $\mathrm{proj}_{V_{\mu_t}}\circ\Phi^2(z_2)\Phi^1(z_1)|_{V_{\mu_s}}\in$ $\mathrm{Hom}_{\mathfrak{g}}(V_{\lambda_2}\otimes V_{\lambda_1}\otimes V_{\mu_s},\ V_{\mu_t})$ and it is given by

$$\lim_{z_t \nearrow \infty} \lim_{z_s \searrow 0} z_t^{2\Delta_{\mu_t}} \langle \Phi_\nu(z_t)\Phi^2(z_2)\Phi^1(z_1)\Phi_{id}(z_s)\rangle .$$

Hence by Theorem 2, the composition $\Phi^2(z_2)\Phi^1(z_1)$ is absolutely convergent in the range $\mathcal{R}_2 = \{(z_2,z_1)\in\mathbb{C}_+^2;\ |z_2|>|z_1|>0\}$, so by the analytic continuation it defines the holomorphic (multivalued) function valued in $\mathrm{Hom}(V_{\lambda_2}\otimes V_{\lambda_1}, \mathrm{Hom}(\mathcal{H}_{\mu_s}, \hat{\mathcal{H}}_{\mu_t}))$ on the complex manifold $M_2 = \{(z_2,z_1)\in (\mathbb{C}\setminus\{0\})^2;\ z_1\neq z_2\}$.

Denote by $\Phi^2(u_2;z_1)\Phi^1(u_1,z_2) = C_\gamma\ (\Phi^2(u_2;z_2)\Phi^1(u_1,z_1))$ its analytic continuation along the path γ:

$$\gamma(t) = \left(\frac{z_2+z_1}{2} + e^{\pi\sqrt{-1}t}\,\frac{z_2-z_1}{2},\ \frac{z_2+z_1}{2} - e^{\pi\sqrt{-1}t}\,\frac{z_2-z_1}{2}\right), \quad t\in[0,1]$$

for $(w,z)\in\mathcal{R}$, then the corresponding analytic continuation

$$T\ \langle \Phi_\nu(z_t)\Phi^2(z_1)\Phi^1(z_2)\Phi_{id}(z_s)\rangle$$

satisfies the systems $KZ(T\Lambda)$ and $\ell C(T\Lambda)$ as a $\mathrm{Hom}(V_{\lambda_1}\otimes V_{\lambda_2}, \mathrm{Hom}(\mathcal{H}_{\mu_s}, \hat{\mathcal{H}}_{\mu_t}))$-valued function, where T is the transposition operator: $\mathrm{Hom}(V_{\lambda_2}\otimes V_{\lambda_1}, A) \longrightarrow \mathrm{Hom}(V_{\lambda_1}\otimes V_{\lambda_2}, A)$,

$$(T\varphi)(u_1\otimes u_2) = \varphi(u_2\otimes u_1) \qquad (u_2\otimes u_1\in V_{\lambda_1}\otimes V_{\lambda_2}),$$

and $T\Lambda = (\mu_t^+, \lambda_1, \lambda_2, \mu_s)$. Hence the analytic continuation

along γ gives an isomorphism between the spaces of compositions of vertex operators:

Theorem 3. (Commutation Relations)

For $\Lambda = (\mu_t^+, \lambda_2, \lambda_1, \mu_s)$, $C(\Lambda) = C_\gamma(\Lambda)$ is an isomorphism :

$$\begin{array}{ccc} C_\gamma(\Lambda):\ \mathcal{V}(\Lambda) & \longrightarrow & \mathcal{V}(T\Lambda) \\ \| \wr & & \| \wr \\ \sum_{\mu\in P_\ell} \mathcal{V}\left(\begin{bmatrix} & \lambda_2 & \\ \mu_t & & \mu \end{bmatrix}\right) \otimes \mathcal{V}\left(\begin{bmatrix} & \lambda_1 & \\ \mu & & \mu_s \end{bmatrix}\right) & \longrightarrow & \sum_{\mu\in P_\ell} \mathcal{V}\left(\begin{bmatrix} & \lambda_1 & \\ \mu_t & & \mu \end{bmatrix}\right) \otimes \mathcal{V}\left(\begin{bmatrix} & \lambda_2 & \\ \mu & & \mu_s \end{bmatrix}\right) . \end{array}$$

Remark. The isomorphisms $C_\gamma(\Lambda)$, $\Lambda\in P_\ell^4$ enjoy the braid relations: For any $N\geq 1$, $\mu_t, \mu_s \in P_\ell$, introduce the space

$$\mathcal{V}(N;\mu_t,\mu_s) = \sum_{\lambda_1,\cdots,\lambda_N\in P_\ell} \mathcal{V}(\mu_t^+, \lambda_N, \cdots, \lambda_1, \mu_s) .$$

Define the operators C_i $(1\leq i\leq N-1)$ on $\mathcal{V}(N;\mu_t,\mu_s)$ such that

$$C_i\ \mathcal{V}(\mu_t^+, \lambda_N, \cdots, \lambda_1, \mu_s) \subset \mathcal{V}(\mu_t^+, \lambda_N, \cdots, \lambda_i, \lambda_{i+1}, \cdots, \lambda_1, \mu_s)$$

and

$$\begin{aligned} & C_i(\varphi_N\otimes\cdots\otimes\varphi_1) \\ & = \varphi_N\otimes\cdots\otimes\varphi_{i+2}\otimes C(\mu_{i+1}, \lambda_{i+1}, \lambda_i, \mu_{i-1})(\varphi_{i+1}\otimes\varphi_i)\otimes\varphi_{i-1}\otimes\cdots\otimes\varphi_1 \end{aligned}$$

for $\varphi_N \otimes \cdots \otimes \varphi_1 \in \mathcal{V}(\mu_t^+, \lambda_N, \cdots, \lambda_1, \mu_s)_{(\mu_{N-1}, \cdots, \mu_1)}$

$$= \mathcal{V}\left(\begin{bmatrix} & \lambda_N & \\ \mu_t & & \mu_{N-1} \end{bmatrix}\right) \otimes \cdots \otimes \mathcal{V}\left(\begin{bmatrix} & \lambda_i & \\ \mu_i & & \mu_{i-1} \end{bmatrix}\right) \otimes \cdots \otimes \mathcal{V}\left(\begin{bmatrix} & \lambda_1 & \\ \mu_1 & & \mu_s \end{bmatrix}\right).$$

Then

$$C_i \; C_{i+1} C_i = C_{i+1} \; C_i \; C_{i+1}$$

as isomorphisms of $\mathcal{V}(N; \mu_t, \mu_s)$ to itself.

§4. The composition $\Phi^2(u_2; w)\Phi^1(u_1; z)$ is singular at $w=z$ and its behaviour near $w=z$ is described as follows.

For $\Lambda = (\mu_t^+, \lambda_2, \lambda_1, \mu_s)$, the space $V_g^\vee(\Lambda)$ has another decomposition

$$V_g^\vee(\Lambda) \xleftarrow[F]{\cong} \sum_{\nu \in P_+} \mathrm{Hom}_g(V_{\lambda_2} \otimes V_{\lambda_1}, V_\nu) \otimes \mathrm{Hom}_g(V_\nu \otimes V_{\mu_s}, V_{\mu_t}) \ ,$$

where the identification F is given by

$$F(\varphi_2 \otimes \varphi_1)(u_2 \otimes u_1 \otimes u_s) = \varphi_1(\varphi_2(u_2 \otimes u_1) \otimes u_s) \qquad (u_i \in V_{\lambda_i}, u_s \in V_\mu),$$

For $\varphi_2 \in \mathcal{V}\begin{bmatrix} & \nu & \\ \mu_t & & \mu_s \end{bmatrix}$ and $\varphi_1 \in \mathcal{V}\begin{bmatrix} & \lambda_2 & \\ \nu & & \lambda_1 \end{bmatrix}$, a "vertex operator" $\Phi^f_{\varphi_2 \otimes \varphi_1}(z)$ of $\mathcal{H}_{\mu_s}$ to $\hat{\mathcal{H}}_{\mu_t}$ parametrized by $V_{\lambda_2} \otimes V_{\lambda_1}$ defined by

$$\Phi^f_{\varphi_2 \otimes \varphi_1}(u_2 \otimes u_1; z) = \Phi_{\varphi_1}(\varphi_2(u_2 \otimes u_1); z) \qquad (u_i \in V_{\lambda_i}).$$

Theorem 4.(Short range expansion or Fusion rule)

i) Near $w=z$ $((w,z)\in\mathcal{R}_2))$,

$$\Phi^2(u_2;w)\Phi^1(u_1;z) = \sum_{\nu\in P_\ell} (w-z)^{-\hat{\Delta}(w_1)} \left[\Phi^f_{\psi_\nu}(u_2\otimes u_1;z) + O(w-z)\right]$$

$$\sim (w-z)^{-(\Delta_{\lambda_1}+\Delta_{\lambda_2})} \sum_{\nu\in P_\ell} (w-z)^{\Delta_\nu}\, \Phi^f_{\psi_\nu}(u_2\otimes u_1;z) \ ,$$

where $\psi_\nu\in\mathcal{V}\left(\begin{bmatrix} & \lambda_2 & \\ \nu & & \lambda_1\end{bmatrix}\right)\otimes\mathcal{V}\left(\begin{bmatrix} & \nu & \\ \mu_t & & \mu_s\end{bmatrix}\right)$, *and* $O(w-z)$ *is holomorhic near* $w=z$ *an vanishes at* $w=z$*:*



The value of $(w-z)^{-\hat{\Delta}(w_1)}$ *is chosen as it is positive for* $(w,z)\in\mathcal{R}_2\cap\mathbb{R}^2$.

ii) For $\Lambda = (\mu_t^+,\lambda_2,\lambda_1,\mu_s)$, *the fusion gives an isomorphism*

$$F(\Lambda):\mathcal{V}(\Lambda)\cong\sum_{\mu\in P_\ell}\mathcal{V}\left(\begin{bmatrix} & \lambda_2 & \\ \mu_t & & \mu\end{bmatrix}\right)\otimes\mathcal{V}\left(\begin{bmatrix} & \lambda_1 & \\ \mu & & \mu_s\end{bmatrix}\right) \longrightarrow \sum_{\nu\in P_\ell}\mathcal{V}\left(\begin{bmatrix} & \lambda_2 & \\ \nu & & \lambda_1\end{bmatrix}\right)\otimes\mathcal{V}\left(\begin{bmatrix} & \nu & \\ \mu_t & & \mu_s\end{bmatrix}\right)$$

defined by

$$F(\Lambda)(\varphi^2\otimes\varphi^1) = \sum_{\nu\in P_\ell}\psi_\nu \qquad \left(\varphi^2\otimes\varphi^1\in\mathcal{V}\left(\begin{bmatrix} & \lambda_2 & \\ \mu_t & & \mu\end{bmatrix}\right)\otimes\mathcal{V}\left(\begin{bmatrix} & \lambda_1 & \\ \mu & & \mu_s\end{bmatrix}\right)\right),$$

where ψ_ν *are the ones obtained in i) for* $\Phi^i = \Phi_\varphi i$.

Theorem 5.

For $\Lambda = (\mu_t^*, \lambda_2, \lambda_1, \mu_s)$, *the following diagram commutes:*

$$\begin{array}{ccc}
\mathcal{V}(\Lambda) & \xrightarrow{F(\Lambda)} & \sum_{\nu\in P_\ell} \mathcal{V}\left(\begin{bmatrix} & \nu & \\ \mu_t & & \mu_s \end{bmatrix}\right)\otimes\mathcal{V}\left(\begin{bmatrix} & \lambda_2 & \\ \nu & & \lambda_1 \end{bmatrix}\right) \\
\Big\downarrow C_\gamma & & \Big\downarrow \mathrm{id}\otimes C_{\gamma_0} \\
\mathcal{V}(T\Lambda) & \xrightarrow{F(T\Lambda)} & \sum_{\nu\in P_\ell} \mathcal{V}\left(\begin{bmatrix} & \nu & \\ \mu_t & & \mu_s \end{bmatrix}\right)\otimes\mathcal{V}\left(\begin{bmatrix} & \lambda_1 & \\ \nu & & \lambda_2 \end{bmatrix}\right)
\end{array}$$

Remark. The equation KZ(Λ) in the limit $z_4 \nearrow \infty$, $z_1 \searrow 0$ is reduced to a differential equation (reduced KZ-system) RKZ(Λ) on $V_g(\Lambda)$- functions of one variable $\zeta = z_3/z_2$. The equation RKZ(Λ) has only regular singularities at $\zeta = 0, 1, \infty$. The isomorphisms $C_\gamma(\Lambda)$ and $F(\Lambda)$ are essentially nothing but the connection matrices from the space of its solutions regularized at $\zeta=0$ to the spaces of solutions regularized at $\zeta=\infty$ and $\zeta=1$ respectively.

§5. Naturally arises a problem to determine the

isomorphisms $C_\gamma(\Lambda)$ and $F(\Lambda)$, but it is very difficult to carry out for all cases. We succeeded (last year) in the case where $\hat{g}$ is an affine Lie algebra of type $A_n^{(1)}$ and $\Lambda = (\mu_t, \square, \square, \mu_s)$, where $\square$ means a Young diagram consisting of one node and represent the vector representation of $g = sl(n+1, \mathbb{C})$.

Now let $\hat{g}$ be an affine Lie algebra of type $A_n^{(1)}$, $B_n^{(1)}$, $C_n^{(1)}$, $D_n^{(1)}$, and P_+^0 be the set of weights $\lambda \in P_+$ such that the simple g-module V_λ can appear in some tensor products of the vector representations $V_\square$ of $g = sl(n+1;\mathbb{C})$, $o(2n;\mathbb{C})$, $sp(2n;\mathbb{C})$, $o(2n+1;\mathbb{C})$ respectively.

For each $\tau \in P_\ell$, introduce the space

$$\mathcal{V}_N(\tau) = \sum_{\mu} \mathcal{V}_N(\tau)_\mu ,$$

$$\mathcal{V}_N(\tau)_\mu = \mathcal{V}\left(\begin{bmatrix} & \square & \\ \tau & & \mu_{N-1} \end{bmatrix}\right) \otimes \cdots \otimes \mathcal{V}\left(\begin{bmatrix} & \square & \\ \mu_i & & \mu_{i-1} \end{bmatrix}\right) \otimes \cdots \otimes \mathcal{V}\left(\begin{bmatrix} & \square & \\ \mu_1 & & 0 \end{bmatrix}\right),$$

where the summation is taken over the set $P_\ell^{N-1} \ni \mu = (\mu_1, \cdots, \mu_{N-1})$. Then $\mathcal{V}_N(\tau)$ is the subspace of $\mathcal{V}(N;\tau,0)$ which is invariant under the operators $C_i (1 \le i \le N-1)$.

The braid group B_N with N-strings of $\mathbb{C}$ has a system $\{b_i; 1 \le i \le N-1\}$ of generators with the fundamental relations:

$$\text{(BR)} \quad \begin{cases} b_i b_j = b_j b_i \quad (|i-j| \ge 2) \\ b_i b_{i+1} b_i = b_{i+1} b_i b_{i+1} \quad (1 \le i \le N-2). \end{cases}$$

These generators b_i are represented by the curves on $\mathbb{C}$ defined by

$$b_i(t)=\left[N,N-1,\cdots,i+\frac{1}{2}(1+e^{\pi\sqrt{-1}t}),i+\frac{1}{2}(1-e^{\pi\sqrt{-1}t}),\cdots 2,1\right]$$

$$t\in[0,1],$$

We now define a monodromy representation π_N^τ of B_N on the space $\mathcal{V}_N(\tau)$ as $\pi_N^\tau(b_i) = C_i$ $(1\leq i\leq N-1)$. The we get the main theorems.

Theorem 6.

If $\mathfrak{g}$ is of type A_n, then the monodromy representation $r^{1/(n+1)}\pi_N^\lambda$ in $\mathcal{V}_N(\tau)$ factors through the Iwahori-Hecke algebra $H_N(r)$, where $r=\exp(\frac{\pi\sqrt{-1}}{\ell+n+1})$.

Note. The algebra $H_N(r)$ is defined by generators $\{\tau_i,\tau_i^{-1}(1\leq i\leq N-1)\}$ with the defining relations:

$$\tau_i\tau_i^{-1}=\tau_i^{-1}\tau_i=1\ ,\quad \tau_i-\tau_i^{-1}=(r-r^{-1})\quad (1\leq i\leq N-1)\ ,$$

$$\tau_i\tau_{i+1}\tau_i=\tau_{i+1}\tau_i\tau_{i+1}\quad\text{and}\quad \tau_i\tau_j=\tau_j\tau_i\quad (|i-j|\geq 2).$$

Theorem 7. If $\mathfrak{g}$ is the simple Lie algebra of type B_n, C_n or D_n. Then the monodromy representation π_N^λ in $\mathcal{V}_N(\tau)$ factors through the Birmann-Wenzl-Murakami algebra $C_N(\mathfrak{g};r)$

where $r=\exp(\frac{\pi\sqrt{-1}}{\ell+g})$, $g(B_n) = 2n-1$, $g(C_n) = n+1$, $g(D_n) = 2n-2$; $C_N(B_n;r) = C_N(r^{-n-1/2},r)$, $C_N(C_n;r) = C_N(r^n,r)$ and $C_N(D_n;r)= C_N(r^{-n},r)$.

Note. The algebra $C_N(a,r)$ is defined by generators $\{\tau_i,\tau_i^{-1},\varepsilon_i(1\le i\le N-1)\}$ with the defining relations:

$$\tau_i\tau_i^{-1}=\tau_i^{-1}\tau_i=1 , \quad \tau_i\varepsilon_i = \varepsilon_i\tau_i = -(a^2r)^{-1}\varepsilon_i ,$$

$$\tau_i - \tau_i^{-1} = (r - r^{-1})(1-\varepsilon_i) \qquad (1\le i\le N-1),$$

$$\tau_i\tau_{i+1}\tau_i=\tau_{i+1}\tau_i\tau_{i+1}, \ \varepsilon_i\varepsilon_{i+1}\varepsilon_i=\varepsilon_i, \ \varepsilon_{i+1}\varepsilon_i\varepsilon_{i+1} = \varepsilon_{i+1},$$

$$\tau_i^{\pm1}\varepsilon_{i+1}\varepsilon_i = \tau_{i+1}^{\mp1}\varepsilon_i , \ \tau_{i+1}^{\pm1}\varepsilon_i\varepsilon_{i+1} = \tau_i^{\mp1}\varepsilon_{i+1} ,$$

$$\varepsilon_i\varepsilon_{i+1}\tau_i^{\pm1} = \varepsilon_i\tau_{i+1}^{\mp1} , \ \varepsilon_{i+1}\varepsilon_i\tau_{i+1}^{\pm1} = \varepsilon_{i+1}\tau_i^{\mp1} \qquad (1\le i\le N-2),$$

$$\tau_i\tau_j = \tau_j\tau_i, \ \varepsilon_i\tau_j = \tau_j\varepsilon_i, \ \varepsilon_i\varepsilon_j = \varepsilon_j\varepsilon_i \qquad (|i-j|\ge2),$$

The proof is carried out by the explicit calculation of a differential equation of 4-point function in a very special case and the algebraic arguments for the algebras $H_N(r)$ and $C_N(a,r)$.

References

[A] K.Aomoto, *Gauss-Manin connection of integral of difference products*, J.Math.Soc.of Japan,**39-2**(1987),

191-208.

[BW] J.S.Birman and H.Wenzl, *Braids, link polynomials and a new algebra*, preprint.

[JMO] M.Jimbo, T.Miwa and M.Okado, *Solvable Lattice Models related to the Vector Representation of Classical Simple Algebas*, CMP,**116**(1988),507-525.

[KZ] K.G.Knizhnik and A.B.Zamolodchikov, *Current Algebra and Wess-Zumino models in two dimensions*, Nuclear Physics, **B247** (1984),83-103.

[M] J.Murakami, *The representations of the* q-*analogue of Brauer's centralizer algebras and the Kauffman polynomial of links*, preprint Osaka Univ(1988).

[TK] A.Tsuchiya and Y.Kanie, *Vertex operators in conformal field theory on* $\mathbb{P}^1$ *and monodromy representations of braid group*, Advanced Studies in pure Math.,**16** (1988), 297-372.

[W1] H.Wenzl, *Hecke algebras of type* A_n *and subfactors*, Invent. math.,**92**(1988),349-383.

[W2] H.Wenzl, *On the structure of Brauer's centralizer algebras*, Ann.of Math.,**125**(1988),173-193.