

Abelian varieties attached to Hilbert modular surfaces

Takayuki ODA

§ Introductory speculations.

In this note, I would like to consider the Hodge realization of the following speculations via motives (cf. Deligne [1]).

Let  $F$  be a real quadratic field over the rational number field  $Q$ . Suppose that a Hilbert modular cusp form  $f$  is given, which is a common eigenfunction of all Hecke operators. Let us call such a form  $f$  ( $\neq 0$ ) primitive. And let  $K_f$  be the subspace of the complex number field  $C$ , generated by the eigenvalues of all Hecke operators. Then we have the following conjecture.

Tensor Product Conjecture. (0.1).

(i) We expect that there exists a motif  $M(f)$  defined over  $Q$  attached to  $f$ , on which  $K_f$  acts as endomorphisms on  $M(f) \times_{\text{Spec}(Q)} \text{Spec}(F)$ . Moreover the rank of  $M(f)$  over  $K_f$  should be  $2^2=4$ .

(ii) For any  $f$  which is given above, we expect that there exist two motives  $M_{f,1}$  and  $M_{f,2}$  defined over  $F$  with  $K_f$  actions on them, such that  $M(f) \times_{\text{Spec}(Q)} \text{Spec}(F) \cong M_{f,1} \otimes_{K_f} M_{f,2}$  (an isomorphism as  $K_f$ -motives). Moreover these two motives  $M_{f,1}$  and  $M_{f,2}$ , which will be of rank two over  $K_f$ , should be conjugate with respect to the extension  $F/Q$ .

(iii) Let  $f^\sigma$  be a primitive Hilbert cusp form obtained from  $f$ , applying an automorphism  $\sigma$  of  $C$  over  $Q$  to the Fourier coefficients of  $f$ . Then  $M_{f^\sigma,1} \cong M_{f,1}$  and  $M_{f^\sigma,2} \cong M_{f,2}$ , or  $M_{f^\sigma,1} \cong M_{f,2}$  and  $M_{f^\sigma,2} \cong M_{f,1}$ .

Let us apply the above conjecture to the problem of counting the number of algebraic 2-cycles on Hilbert modular surfaces. Let  $M^2(S)$  be the 2-motif defined over  $\mathbb{Q}$  attached to a Hilbert modular surface  $S$ , and let  $W_2 M^2(S)$  be the pure part of  $M^2(S)$ . Let  $\Phi$  be a representative system of equivalence classes of primitive Hilbert modular cusp forms of weight 2 with respect to the following equivalence relation:

Two primitive forms  $f$  and  $g$  are equivalent, if and only if  $f^c = cg$  for some  $c \in \text{Aut}(\mathbb{C}/\mathbb{Q})$  and  $c \in \mathbb{C}$ .

Then we should have a decomposition of the motif  $W_2 M^2(S)$

$$W_2 M^2(S) \times_{\text{Spec}(\mathbb{Q})} \text{Spec}(\mathbb{F}) \simeq \mathbb{F}[-1] \oplus \mathbb{F}[-1] \oplus \left( \bigoplus_{f \in \Phi} M(f) \right),$$

where  $\mathbb{F}[-1]$  the Tate motif of "poids" 2 over  $\mathbb{F}$ .

By Tensor Product Conjecture (0.1), there exist two abelian varieties  $A_{f,1}$  and  $A_{f,2}$  defined over  $\mathbb{F}$  of dimension  $d=[K_f:\mathbb{Q}]$  with endomorphism ring  $K_f$ , such that

$$M(f) \simeq A_{f,1} \otimes_{K_f} A_{f,2},$$

where we regard these  $\check{A}$  abelian varieties as 1-motives,   
 i.e. abelian varieties up to  $K_f$ -isogeny.

Therefore,

$$W_2 M^2(S) \times_{\text{Spec}(\mathbb{Q})} \text{Spec}(\mathbb{F}) \simeq \mathbb{F}[-1]^{\oplus 2} \oplus \left\{ \bigoplus_{f \in \Phi} (A_{f,1} \otimes_{K_f} A_{f,2}) \right\}.$$

Now let us consider the Tate twist of  $W_2 M^2(S) \times_{\text{Spec}(\mathbb{Q})} \text{Spec}(\mathbb{F})$ :

$$(W_2 M^2(S) \times_{\text{Spec}(\mathbb{Q})} \text{Spec}(\mathbb{F})) [1] \simeq \mathbb{F}[0] \oplus \mathbb{F}[0] \oplus \left( \bigoplus_{f \in \Phi} A_{f,1} [1] \otimes_{K_f} A_{f,2} \right).$$

Since  $A_{f,1}$  is an abelian variety, there exists a polarization

$$A_{f,1} \otimes_{K_f} A_{f,1} \longrightarrow K_f[-1].$$

Hence  $A_{f,1} [1]$  is isomorphic to the dual motif  $\check{A}_{f,1}$  of  $A_{f,1}$ .

Therefore, we have

$$(W_2 M^2(S) \times_{\text{Spec}(Q)} \text{Spec}(F)) [1] \cong F[0] \oplus F[0] \oplus \left\{ \bigoplus_{f \in \Phi} \text{Hom}_{K_f} (A_{f,1}, A_{f,2}) \right\},$$

where  $\text{Hom}_{K_f} (A_{f,1}, A_{f,2})$  is the Hom-object in the category of  $K_f$ -motives.

Let  $(W_2 M^2(S) \times_{\text{Spec}(Q)} \text{Spec}(F))_{\text{alg}}$  be the submotif of  $(W_2 M^2(S) \times_{\text{Spec}(Q)} \text{Spec}(F))$

$\text{Spec}(F)$ , generated by algebraic 2-cycles.

Then Hodge-Tate conjecture for motives implies that

$$(\#) \quad (W_2 M^2(S) \times_{\text{Spec}(Q)} \text{Spec}(F))_{\text{alg}} [1] \cong F[0] \oplus F[0] \oplus \left\{ \bigoplus_{f \in \Phi} \text{Hom}_{K_f} (A_{f,1}, A_{f,2}) \right\}.$$

Here  $\text{Hom}_{K_f} (A_{f,1}, A_{f,2})$  is the  $K_f$ -module of  $K_f$ -linear homomorphism of

$K_f$ -isogeny class of abelian varieties defined by  $A_{f,1}$ , to  $K_f$ -isogeny class of abelian varieties defined by  $A_{f,2}$ .

So, if (#) is true, we can reduce the problem of determination of algebraic cycles, to determination of endomorphisms of abelian varieties.

Here we consider the Hodge realization of the above argument. In this case, the validity of (#) is guaranteed by the Lefschetz criterion on algebraic cycles on surfaces:

A 2-cycle on an algebraic surface is algebraic, if and only if it is rational and of (1.1)-type.

§ 0. Main results.

Let  $F$  be a real quadratic field with discriminant  $D$  and with class number 1. And suppose that  $F$  has a unit with negative norm. Let  $f$  be a primitive form of weight 2 with respect to  $\Gamma = \text{SL}_2(\mathcal{O}_F)$ . For any primitive form  $f$  of weight 2, we can attach, in general, two abelian varieties  $A_{f,1}$  and  $A_{f,2}$  of dimension  $d = [K_f : \mathbb{Q}]$ , which have  $K_f$  as  $\mathbb{A}$  subring of  $\text{End}(A_{f,i}) \otimes_{\mathbb{Z}} \mathbb{Q}$  ( $i=1,2$ ).  $\left. \vphantom{[K_f : \mathbb{Q}]}\right\}$  defined over  $\mathbb{C}$

Definition. (0.1). A primitive form  $f(z_1, z_2)$  is called self-conjugate, iff  $f(z_1, z_2) = f(z_2, z_1)$ .

Remark. We avoid the term "symmetric", because the associated 2-form  $\omega_f = (2\pi i)^2 f(z_1, z_2) dz_1 \wedge dz_2$  is not symmetric  $\left. \vphantom{f(z_1, z_2)}\right\}$  with respect to but anti-symmetric the involution of Hilbert modular surfaces obtained from the mapping ~~on passing to~~ ~~XXXXXX~~  $(z_1, z_2) \mapsto (z_2, z_1)$  on passing to the quotients. There is also a theoretical reason to call these forms self-conjugate.

For  $\mathbb{A}$  self-conjugate form  $f$ , it is known that there exists a reellen Neben Typus cuspidal form  $\psi$  of weight 2 with respect to (elliptic)  $\Gamma_0(D)$  with multiplier defined by Jacobi symbol  $\left(\frac{D}{x}\right)$ , such that ~~XXXXXX~~ ~~XXXXXXXX~~ ~~XXXXXXXX~~ ~~XXXXXX~~ ~~XXXXXX~~ ~~XXXXXXXX~~  $f$  is the lifting of  $\psi$  (cf. Doi-Naganuma [11], Naganuma [12], Zagier [13], etc).

Let us recall the results of Shimura [20]. Let  $K_\psi$  be the field generated by the eigenvalues of Hecke operator over  $\mathbb{Q}$ . And let  $k_\psi$  be the totally real subfield contained in  $K_\psi$  with  $[K_\psi : k_\psi] = 2$ . Then Shimura [20] naturally attached abelian variety



Remark. The equality  $\lambda \leq 3p_{sc} + 4p_{nsc}$  is proved by Hirzebruch [6].

Let us explain the outline of our proofs. In the first four sections, we construct abelian variety  $A(f)$  of dimension  $4d$  defined over  $\mathbb{C}$ , attached to a primitive form  $f$ , where  $d = [K_f : \mathbb{Q}]$ . Here we use an idea of Satake [9], Kuga-Satake [8], and Deligne [2]. More precisely speaking, we attach a polarized Hodge structure  $H(f)$  of "poids" 2 with  $K_f$  action to each primitive form ( § 2 ). And next, following Kuga-Satake [8], we construct abelian varieties, by using the Clifford algebras attached to polarization forms on  $H(f)$ .

In section 5, we consider two topological involutions on Hilbert modular surfaces. And by means of these involutions, we show that our Clifford algebras are isomorphic to two  $2 \times 2$  matrix algebra over  $K_f$  (a direct product of). Accordingly, we have a decomposition up to isogenie  $A(f) \sim A_{f,1} \times A_{f,1} \times A_{f,2} \times A_{f,2}$ .

In section 6, we calculate explicitly the period lattices of these two abelian varieties. In this calculation, the period relation of Riemann-Hodge plays a key role.

The rest of this note <sup>es</sup> discusses the application of these result to the problem of counting the number of transcendental cycles on Hilbert modular surfaces. The sources are following: Manin's idea [1] to represent the period integrals of modular forms by the twisted L-functions of the modular forms; liftings of Doi-Naganuma [11], and Naganuma [12]; Shimura's results on period integrals of elliptic modular forms; the determination of endomorphisms rings of abelian varieties attached to elliptic cusp forms <sup>up to</sup> by Ribet [18] and Momose [19].

§ 1. Polarized Hodge structures attached to Hilbert modular surfaces.

(1.0) Definitions.

We fix a real quadratic field  $F = \mathbb{Q}(\sqrt{d})$  with discriminant  $d > 0$ , and assume that  $F$  has a unit  $\epsilon$  with negative norm  $N\epsilon = -1$ . Let  $H$  be the complex upper half plane  $\{z \in \mathbb{C} \mid \text{Im } z > 0\}$ , and let  $SL_2(\mathbb{R})$  be the special linear group of degree 2 with entries in the real number field  $\mathbb{R}$ . As usual  $SL_2(\mathbb{R})$  acts on  $H$  via

$$g(z) = (az+b)/(cz+d) \quad \text{for } g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbb{R}), \text{ and } z \in H.$$

The product group  $SL_2(\mathbb{R}) \times SL_2(\mathbb{R})$  acts on the product  $H \times H$  factorwise.

Let  $\alpha_1$  and  $\alpha_2$  be two embeddings of an element  $\alpha$  of  $F$  into  $\mathbb{R}$ .

Then the mapping

$$\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \longrightarrow \left( \begin{pmatrix} \alpha_1 & \beta_1 \\ \gamma_1 & \delta_1 \end{pmatrix}, \begin{pmatrix} \alpha_2 & \beta_2 \\ \gamma_2 & \delta_2 \end{pmatrix} \right)$$

defines an injective homomorphism of groups  $SL_2(F) \longrightarrow SL_2(\mathbb{R}) \times SL_2(\mathbb{R})$ .

By means of this injection, the group  $SL_2(F)$  acts on  $H \times H$ , and

accordingly its subgroup  $\Gamma = SL_2(\mathcal{O}_F)$  also acts on  $H \times H$ . Here  $\mathcal{O}_F$  is

the ring of integers of  $F$ .  $\Gamma$  acts properly discontinuously on  $H \times H$ ,

and the quotient analytic space  $S = \Gamma \backslash H \times H$  has a natural structure of quasi-projective algebraic surface (cf. Baily-Borel [30] for example),

which is smooth except finite number of quotient singularities

corresponding to elliptic fixed points on  $H \times H$  with respect to  $\Gamma$ .

The surface  $S$  is called the Hilbert modular surface attached to  $\Gamma$ ,

or to  $F$ .

Let  $\bar{S}$  be the standard compactification, which is a union of  $S$  and  $SL_2(\mathcal{O}_F) \backslash P^1(F)$  as a set. Here  $P^1(F) = F \cup \{\infty\}$  is the 1-dimensional projective line over  $F$ . As is well known, the cardinality of the finite set  $SL_2(\mathcal{O}_F) \backslash P^1(F)$  is equal to the class number  $h_F$  of

F. An element of  $SL_2(O_F) \backslash P^1(F)$  is called an equivalence class of cusps, or by an abuse of languages, simply called a cusp.  $\bar{S}$  has singularities at these cusps, whose resolution is precisely studied by Hirzebruch [5].

(1.1) Cohomology groups of Hilbert modular surfaces.

Let us recall some basic facts on the homology and cohomology groups of  $S$  and  $\bar{S}$ . Let  $\pi: \tilde{S} \rightarrow \bar{S}$  be a resolution of the quotient and cusp singularities of  $\bar{S}$ . Then the cohomology group  $H^2(\tilde{S}, Q)$  is a direct sum of the image of  $\pi^*: H^2(\bar{S}, Q) \rightarrow H^2(\tilde{S}, Q)$ , and the subspace of  $H^2(\tilde{S}, Q)$  generated by algebraic cycles which are obtained as irreducible components of the inverse image loci of singularities by  $\pi$ . Restricting the intersection form on  $H^2(\tilde{S}, Q)$  to  $\pi^*H^2(\bar{S}, Q)$ , we can define a nondegenerate pairing on  $\pi^*H^2(\bar{S}, Q)$ .

Lemma. Let  $(\tilde{S}', \pi')$  be another desingularization of  $\bar{S}$ . Then  $\pi^*H^2(\bar{S}, Q)$  and  $\pi'^*H^2(\bar{S}, Q)$  are canonically isomorphic as vector spaces with inner products defined by intersection forms.

Proof. Routine. Q.E.D.

By this lemma we can define a unique natural intersection form on the coimage of  $\pi^*: H^2(\tilde{S}, Q) \rightarrow H^2(\bar{S}, Q)$ . Now let us investigate the kernel of  $\pi^*$ . Let  $r_1, r_2, \dots, r_k$  (resp.  $c_1, c_2, \dots, c_h$ ) be the set of quotient (resp. cusp) singularities on  $\bar{S}$ . Put  $R_i = \pi^{-1}(r_i)$  and  $C_i = \pi^{-1}(c_i)$ . Then, considering the exact sequences of the relative cohomology groups, we have

$$\begin{array}{ccccccc}
 \longrightarrow & 0 & \longrightarrow & H^2(S \text{ mod } \{r_1, \dots, r_k\} \cup \{c_1, \dots, c_h\}, Q) & \xrightarrow{j} & H^2(\bar{S}, Q) & \longrightarrow 0 \\
 & \downarrow & & \cup & & \downarrow & \downarrow \\
 - & H^1(\bigcup_{i=1}^k R_i \cup \bigcup_{i=1}^h C_i, Q) & \xrightarrow{\cong} & H^2(\tilde{S} \text{ mod } \bigcup_{i=1}^k R_i \cup \bigcup_{i=1}^h C_i, Q) & \longrightarrow & H^2(\tilde{S}, Q) & \longrightarrow *
 \end{array}$$



Here the isomorphism  $\#$  is obtained by the excision theorem.

By means of this diagram, we have  $\text{Image } \partial = \text{Ker } \pi^*$ . Since  $r_i$  ( $1 \leq i \leq k$ ) are quotient singularities,  $H^1(R_i, \mathbb{Q}) = 0$ . Therefore, we have the following proposition.

Proposition.     The sequence

$$\bigoplus_{i=1}^h H^1(C_i, \mathbb{Q}) \xrightarrow{j\#^{-1}\partial} H^2(\bar{S}, \mathbb{Q}) \longrightarrow H^2(\tilde{S}, \mathbb{Q})$$

is exact.

We apply this exact sequence for calculation of the mixed Hodge structure of  $H^2(\bar{S}, \mathbb{Q})$ . From now on we consult with Deligne [4].

(1.2) The mixed Hodge structures of Hilbert modular surfaces.

Let us consider the mixed Hodge structure of  $H^2(S, \mathbb{Q})$ ,  $H_c^2(S, \mathbb{Q})$  and  $H^2(\bar{S}, \mathbb{Q})$ . By means of the previous proposition, we have an exact sequence of cohomological descent (cf. Saint-Donat [1]), we can calculate the mixed Hodge structure of  $H^2(\bar{S}, \mathbb{Q})$  by this (Deligne [4]). By results of Hirzebruch [5], we can take as  $C_i$  a stable curve of genus 1, which is not an elliptic curve (the so-called Neron's  $N$ -polygon). Therefore,  $H^1(C_i, \mathbb{Q}) \cong \mathbb{Q}$ , and the weight filtration of the mixed Hodge structure of  $H^1(C_i, \mathbb{Q})$  is given by

$$W_k H^1(C_i, \mathbb{Q}) = 0 \quad (k < 0), \quad W_0 H^1(C_i, \mathbb{Q}) \cong \mathbb{Q} \quad (k = 0), \quad W_k H^1(C_i, \mathbb{Q}) = H^1(C_i, \mathbb{Q}) \quad (k \geq 0).$$

Therefore the weight filtration of  $H^2(\bar{S}, \mathbb{Q})$  is given by

$$\begin{aligned} W_k H^2(\bar{S}, \mathbb{Q}) &= 0 \quad (k < 0), \\ W_0 H^2(\bar{S}, \mathbb{Q}) &\cong \bigoplus \mathbb{Q} \quad (k = 0), \\ W_1 H^2(\bar{S}, \mathbb{Q}) &= W_0 H^2(\bar{S}, \mathbb{Q}) \quad (k = 1), \\ W_2 H^2(\bar{S}, \mathbb{Q}) &= H^2(\bar{S}, \mathbb{Q}) \quad (k \geq 2). \end{aligned}$$

Moreover

$$\text{Gr}_2^W H^2(\bar{S}, \mathbb{Q}) = W_2 H^2(\bar{S}, \mathbb{Q}) / W_1 H^2(\bar{S}, \mathbb{Q})$$

is a polarized homogeneous sub-Hodge structure of the polarized Hodge structure of  $H^2(S, \mathbb{Q})$ .

By means of the trivial isomorphism  $H_c^2(S, \mathbb{Q}) \cong H^2(\bar{S}, \mathbb{Q})$ , we can transcribe the mixed Hodge structure on  $H^2(\bar{S}, \mathbb{Q})$  to  $H_c^2(S, \mathbb{Q})$ .

Since  $S$  is rationally smooth, we have the Poincaré duality

$$H^2(S, \mathbb{Q}) \times H_c^2(S, \mathbb{Q}) \longrightarrow \mathbb{Q}[-2].$$

Therefore the weight filtration of the mixed Hodge structure on  $H^2(S, \mathbb{Q})$  is given by

$$\begin{aligned} W_k H^2(S, \mathbb{Q}) &= 0 \quad (k \leq 1), \\ W_2 H^2(S, \mathbb{Q}) &= W_3 H^2(S, \mathbb{Q}), \\ W_k H^2(S, \mathbb{Q}) &= H^2(S, \mathbb{Q}) \quad (k \geq 4), \end{aligned}$$

and  $W_2 H^2(S, \mathbb{Q})$  is a homogeneous polarized Hodge structure of "poids" 2, and  $\text{Gr}_4 W H^2(S, \mathbb{Q}) \cong \mathbb{Q}[-2]$ .

(1.3) The Hodge decomposition of the pure part  $W_2 H^2(S, \mathbb{Q})$  of  $H^2(S, \mathbb{Q})$ .

From now on, for simplicity, we assume that the class number  $h_F$  of  $F$  is 1. Let  $\mathcal{K} = \mathcal{K}(\Gamma, \Delta)$  be the Hecke algebra  $\Gamma = \text{SL}_2(\mathcal{O}_F)$  with respect to the commensurator  $\Delta = \{\alpha \in M_2(\mathcal{O}_F) \mid \det \alpha \in \mathcal{O}_F - \{0\}\}$ . Since we can naturally regard any element of  $\mathcal{K}$  as an algebraic correspondence of  $S$ , the Hecke algebra  $\mathcal{K}$  acts on  $H^2(S, \mathbb{Q})$ ,  $H_c^2(S, \mathbb{Q})$ , and  $H^2(S, \mathbb{Q})$ .

Remark. We can write the cohomology groups  $H^2(S, \mathbb{Q})$ , and  $H_c^2(S, \mathbb{Q}) \cong H^2(\bar{S}, \mathbb{Q}) \cong H^2(\bar{S} \text{ mod cusps}, \mathbb{Q})$  as relative cohomology groups of the discrete group  $\Gamma = \text{SL}_2(\mathcal{O}_F)$ . Therefore these cohomology groups are given as cohomology groups of certain complex of  $\Gamma$ -equivariant cochains on  $H \times H$ . Therefore, the commensurator  $\Delta$  acts naturally on these cohomology groups.

The Hecke algebra  $\mathcal{H}$  acts also on the space  $S_2(\Gamma)$  of holomorphic cusp forms on  $H \times H$  with respect to

By means of the mapping

$$f(z_1, z_2) \in S_2(\Gamma) \longmapsto (2\pi i)^2 f(z_1, z_2) dz_1 \wedge dz_2,$$

we can regard any element of  $S_2(\Gamma)$  as a holomorphic 2-form on  $S$ .

And it is known that these 2-forms are prolonged to holomorphic

2-forms on a smooth model  $\tilde{S}$ , and moreover by this mapping  $S_2(\Gamma)$

is canonically isomorphic to  $\Gamma(\tilde{S}, \Omega_{\tilde{S}}^2)$  (cf. Freitag [31]).

Especially, we have

$$\dim_{\mathbb{C}} S_2(\Gamma) = p_g(\tilde{S}).$$

Now the Hodge decomposition of  $W_2 H^2(S, \mathbb{Q})$  is given by the following theorem.

Theorem (1.1). (Hirzebruch [6]).

Let  $F$  be a real quadratic field with has units with negative norm. Let  $\varepsilon$  be such a unit, whose two embeddings into  $\mathbb{R}$  satisfies  $\varepsilon_1 > 0, \varepsilon_2 < 0, \varepsilon_1 \varepsilon_2 = -1$ . Then as  $\Gamma$ -modules, we have a natural isomorphism

$$\begin{aligned} W_2 H^2(S, \mathbb{Q}) \otimes_{\mathbb{Q}} \mathbb{C} &\simeq \left\{ (2\pi i)^2 f(z_1, z_2) dz_1 dz_2 \mid f \in S_2(\Gamma) \right\} \\ &\oplus \left\{ (2\pi i)^2 f(\varepsilon_1 z_1, \varepsilon_2 \bar{z}_2) dz_1 \wedge d\bar{z}_2 \mid f \in S_2(\Gamma) \right\} \\ &\oplus \left\{ (2\pi i)^2 f(\varepsilon_2 \bar{z}_2, \varepsilon_1 z_1) d\bar{z}_1 \wedge dz_2 \mid f \in S_2(\Gamma) \right\} \\ &\oplus \left\{ (2\pi i)^2 f(-\bar{z}_1, -\bar{z}_2) d\bar{z}_1 \wedge d\bar{z}_2 \mid f \in S_2(\Gamma) \right\} \\ &\oplus \mathbb{C} \frac{dz_1 \wedge d\bar{z}_1}{y_1^2} \oplus \mathbb{C} \frac{dz_2 \wedge d\bar{z}_2}{y_2^2}. \end{aligned}$$

Here  $z_i = x_i + \sqrt{-1}y_i$  ( $i=1, 2$ ).

(1.4) The primitive part  $W_2H^2(S, \mathbb{Q})_{\text{pr}}$  of  $W_2H^2(S, \mathbb{Q})$ .

Let  $L_1$  and  $L_2$  be the invertible sheaves corresponding to the automorphic factors  $j_i(z, g) = (c_i z_i + d_i)$  ( $i=1, 2$ ), respectively.

Here

$$z = (z_1, z_2) \in \mathbb{H} \times \mathbb{H}, \text{ and } g = \left( \begin{pmatrix} a_1 & b_1 \\ c_1 & d_1 \end{pmatrix}, \begin{pmatrix} a_2 & b_2 \\ c_2 & d_2 \end{pmatrix} \right) \in \text{SL}_2(\mathbb{R}) \times \text{SL}_2(\mathbb{R}).$$

We can prolong these two sheaves to two invertible sheaves  $L_1$  and  $L_2$  on a smooth proper model of  $S$  by the results of Ueno-Van der Geer [24]. The Chern classes of these invertible sheaves define two elements of  $H^2(S, \mathbb{Q})$ , and accordingly two elements of  $W_2H^2(S, \mathbb{Q})$ .

Let  $W_2H^2(S, \mathbb{Q})_{\text{pr}}$  be the orthogonal complement of the space spanned by these two elements, with respect to the intersection form. Then the Hodge decomposition of  $W_2H^2(S, \mathbb{Q})_{\text{pr}}$  is given by the right hand side of the isomorphism of Theorem (1.1), without the last two direct factors

$$\mathbb{C} \frac{dz_1 \wedge \bar{d}z_1}{y_1^2} \oplus \mathbb{C} \frac{dz_2 \wedge \bar{d}z_2}{y_2^2}.$$

Because the Chern forms of  $L_1$  and  $L_2$  are given the above two (1,1)-type 2-forms.

§ 2. Polarized Hodge structures attached to primitive cusp forms of weight 2.

Definition. A Hilbert modular cusp form  $f$  ( $\neq 0$ ) of weight 2 with respect to  $\Gamma$  is called primitive, if it is a common eigenfunction of all Hecke operators:  $T_\alpha f = a_\alpha f$  ( $\alpha \in \mathcal{O}_F$ ).

We denote by  $K_f$  a subfield of  $\mathbb{C}$ , generated by eigenvalues  $a$  over  $\mathbb{Q}$ .

Lemma.  $K_f$  is contained in  $\mathbb{R}$ .

Proof. Well-known. Recall that in our case all Hecke operators  $T_\alpha$  are self-adjoint with respect to the Petersson metric  $(,)$  on  $S_2(\Gamma)$ . Q.E.D.

Let  $\phi_f$  be the homomorphism of  $\mathcal{H}$  into  $K_f$ , defined by the mapping  $\phi_f: T_\alpha \longrightarrow a_\alpha$ .

Put  $\tilde{H}_2(S, \mathbb{Q})_{\text{pr}} = \text{Coimage}(j_*: H_2(S, \mathbb{Q}) \longrightarrow H_2(\tilde{S}, \mathbb{Q})) / \{ \text{the space generated by the homology classes corresponding to the line bundles } L_1 \text{ and } L_2 \}$ ,

and define  $H_2(f)$  by

$$H_2(f) = \tilde{H}_2(S, \mathbb{Q})_{\text{pr}} \otimes_{(\mathcal{K}, \phi_f)} K_f.$$

Then the dual space of  $H_2(f)$  over  $K_f$  is canonically isomorphic to

$$H^2(f) = \{ \eta \in W_2 H^2(S, \mathbb{Q})_{\text{pr}} \otimes_{\mathbb{Q}} K_f \mid T_\alpha \eta = a_\alpha \eta \}.$$

Theorem (2.1). Let  $f$  be a primitive form of weight 2.

(i) The restriction  $\psi_f: H^2(f) \times H^2(f) \longrightarrow K_f$  of the extension of scalars

$$\psi \otimes_{K_f} : W_2 H^2(S, \mathbb{Q})_{\text{pr}} \otimes_{\mathbb{Q}} K_f \times W_2 H^2(S, \mathbb{Q})_{\text{pr}} \otimes_{\mathbb{Q}} K_f \longrightarrow K_f$$

of the intersection form  $\psi$ , is a nondegenerate symmetric bilinear form.

(ii) The extension of scalars of  $\psi_f$  with respect to the injection  $K_f \hookrightarrow \mathbb{R}$

$$\psi_f \otimes_{K_f} \mathbb{R} : H^2(f) \otimes_{K_f} \mathbb{R} \times H^2(f) \otimes_{K_f} \mathbb{R} \longrightarrow \mathbb{R}$$

is of signature  $(2+, 2-)$ , and a polarization of a homogeneous Hodge structure of "poids" 2 given by (iii)

(iii)  $H^2(f) \otimes_{K_f} \mathbb{C}$  has a Hodge decomposition:

$$\begin{aligned} H^2(f) \otimes_{K_f} \mathbb{C} \simeq & \mathbb{C}(2\pi i)^2 f(z_1, z_2) dz_1 \wedge dz_2 \oplus \mathbb{C}(2\pi i)^2 f(\xi_1 z_1, \xi_2 \bar{z}_2) dz_1 \wedge d\bar{z}_2 \\ & \oplus \mathbb{C}(2\pi i)^2 f(\xi_2 \bar{z}_1, \xi_1 z_2) d\bar{z}_1 \wedge dz_2 \oplus \mathbb{C}(2\pi i)^2 f(-\bar{z}_1, -\bar{z}_2) d\bar{z}_1 \wedge d\bar{z}_2. \end{aligned}$$

Epecially  $\dim_{K_f} H^2(f) = 4$ .

Proof. We first recall the following basic fact.

Multiplicity One Theorem. Let  $f$  and  $g$  be two primitive forms of weight 2. If  $T_n f = a_n f$  and  $T_n g = a_n g$  for all integral ideal of  $O_F$ . Then  $f$  is a constant multiple of  $g$ .

Evidently, we can canonically identify  $H^2(f) \otimes_{K_f} \mathbb{C}$  with

$$\{ \eta \in W_2 H^2(S, \mathbb{Q})_{\text{pr}} \otimes_{\mathbb{Q}} \mathbb{C} \mid T_n \eta = a_n \eta \}.$$

By Multiplicity One Theorem ~~XXXXXXXXXXXX~~ and Theorem (1.1), this space is isomorphic to the right hand side of the statement (iii). So (iii) is proved.

Let  $f_1, f_2, \dots, f_{p_g}$  be basis of  $S_2(\Gamma)$  consisting of primitive forms. Then we have

$$W_2 H^2(S, \mathbb{Q})_{\text{pr}} \otimes_{\mathbb{Q}} \mathbb{C} = \bigoplus_{i=1}^{p_g} H^2(f_i) \otimes_{K_{F_i}} \mathbb{C}.$$

$$\text{Hence } W_2 H^2(S, \mathbb{Q}) \otimes_{\text{pr } \mathbb{Q}} \mathbb{R} = \bigoplus_{i=1}^{p_g} H^2(f_i) \otimes_{K_{f_i}} \mathbb{R}.$$

Let us consider  $\psi \otimes \mathbb{R}$ . Let  $f_1$  and  $f_2$  be two primitive form such that  $f_1 \neq cf_2$  for any  $c \in \mathbb{C}$ , and denote by  $a_{\sigma,1}$  and  $a_{\sigma,2}$  the eigenvalues of  $T_{\sigma}$  for  $f_1$  and  $f_2$ , respectively. Then, for  $\eta_1 \in H^2(f_1) \otimes_{K_{f_1}} \mathbb{R}$  and  $\eta_2 \in H^2(f_2) \otimes_{K_{f_2}} \mathbb{R}$ , we have

$$a_{\sigma} \langle \eta_1, \eta_2 \rangle = \langle a_{\sigma,1} \eta_1, \eta_2 \rangle = \langle T_{\sigma} \eta_1, \eta_2 \rangle = \langle \eta_1, T_{\sigma} \eta_2 \rangle = \langle \eta_1, a_{\sigma,2} \eta_2 \rangle = a_{\sigma,2} \langle \eta_1, \eta_2 \rangle$$

where  $\langle , \rangle$  is the intersection form  $\psi \otimes \mathbb{R}$ .

Since  $f_1 \neq cf_2$  for any  $c \in \mathbb{C}$ , by Multiplicity One Theorem, there exist some ideals  $\sigma$  such that  $a_{\sigma,1} \neq a_{\sigma,2}$ . Hence  $\langle \eta_1, \eta_2 \rangle = 0$

Therefore the bilinear form  $\psi \otimes_{\mathbb{Q}} \mathbb{R}$  is a direct sum of  $\psi_{f_i} \otimes \mathbb{R}$

( $i=1, 2, \dots, p_g$ ), and consequently each  $\psi_{f_i}$  is nondegenerate.

Thus (i) is proved.

Because of the index theorem of Hodge, the bilinear form  $\psi \otimes_{\mathbb{Q}} \mathbb{R}$  has signature  $(2p_g^+, 2p_g^-)$ , where  $p_g = \dim_{\mathbb{C}} S_2(\Gamma)$ . For we took the "primitive part" of  $W_2 H^2(S, \mathbb{Q})$ . The Hodge decomposition of  $H^2(f) \otimes_{K_f} \mathbb{C}$  and the fact that Hecke operator commute with the  $C$  operator of Weil imply (ii). Q.E.D.

3.  $\mathbb{Q}$ -basis theorem.

Let  $S_2(\Gamma; \mathbb{Q})$  be the space of Hilbert cusp forms, whose Fourier coefficients are rational numbers. Then the following theorem is known.

$\mathbb{Q}$ -Basis Theorem (3.1). (Shimura [14], or Rapoport [10]).

(i)  $S_2(\Gamma; \mathbb{Q})$  spans  $S_2(\Gamma)$ . In other words, there is a natural isomorphism  $S_2(\Gamma; \mathbb{Q}) \otimes_{\mathbb{Q}} \mathbb{C} \cong S_2(\Gamma)$ .

(ii) Let  $f(z_1, z_2)$  be an element of  $S_2(\Gamma)$  with Fourier expansion

$$f(z_1, z_2) = \sum_{\nu \in \delta_+^{-1}} a(\nu) \exp[2\pi i(\nu_1 z_1 + \nu_2 z_2)],$$

where  $\delta_+^{-1}$  is the intersection of the codifferent and the set of totally positive elements.

For an element  $\sigma \in \text{Aut}(\mathbb{C}/\mathbb{Q})$ , we define  $f^\sigma(z_1, z_2)$  by a formal Fourier expansion

$$f^\sigma(z_1, z_2) = \sum_{\nu \in \delta_+^{-1}} a(\nu)^\sigma \exp[2\pi i(\nu_1 z_1 + \nu_2 z_2)].$$

Then  $f^\sigma(z_1, z_2)$  belongs to  $S_2(\Gamma)$ .

Corollary (3.2). Let  $f$  be a primitive form of weight 2 such that  $T_{\mathfrak{a}} f = a_{\mathfrak{a}} f$ . And let  $\{\sigma_1 = \text{id}_{K_f}, \sigma_2, \dots, \sigma_d\}$  be the set of all embeddings of  $K_f$  into  $\mathbb{R}$ , where  $d = [K_f : \mathbb{Q}]$ . Then there exist primitive forms in  $S_2(\Gamma)$ , such that

$$T_{\mathfrak{a}} f_i = \sigma_i(a_{\mathfrak{a}}) f_i \quad (i=2, 3, \dots, d)$$

for all integral ideal  $\mathfrak{a}$ . Especially we have  $K_{f_i} = \sigma_i(K_f)$

( $1 \leq i \leq d$ ), and  $K_f$  is a totally real field.



§ 4. Clifford algebras and abelian varieties.

Let us recall the data given in the previous section.

Data (4.1)

$K_f$ : a totally real algebraic number field of degree  $d$  over  $\mathbb{Q}$ .

$\{\epsilon_1, \epsilon_2, \dots, \epsilon_d\}$  be the set of all embeddings of  $K_f$  to  $\mathbb{R}$ .

$H(f)$ : a  $K_f$ -module of rank 4 on which a symmetric non-degenerate bilinear form  $\psi_f$  with values in  $K_f$  is defined, which satisfies the following condition:

(i) For any  $\epsilon_i$  ( $i=1, \dots, d$ ),  $H(f) \otimes_{K_f, \epsilon_i} \mathbb{R}$  is equipped with a homogeneous Hodge structure of "poids" 2, and  $\psi \otimes_{K_f, \epsilon_i} \mathbb{R}$  gives a polarization with respect to this Hodge structure.

Let start from this data, and construct ~~an~~ abelian variety attached to this data. First we choose an integral structure. From now on, until the last part of this section, we omit the subscript  $f$  to simplify the notation. Now let  $O_K$  be an order of  $K$ , and let  $H_Z$  be an  $O_K$ -module of rank 4 in  $H$ . And moreover we can choose  $H_Z$  such that the values of  $\psi$  on  $H_Z \times H_Z$  is contained in  $O_K$ .

Let  $C^+(H_Z)$  be the even Clifford algebra attached to over  $O_K$ . Thus this algebra is of rank  $2^3=8$  over  $O_K$ . There is a sophisticated description to attach abelian variety by Deligne [2]. We recall here more naive definition of [8].

First, we consider a real torus of dimension  $8d$

$$C^+(H_Z) \otimes_{\mathbb{Z}} \mathbb{R} / C^+(H_Z).$$

Clearly on this real torus, the Clifford algebra  $C^+(H_Z)$  acts by left multiplication. Next, we define a complex structure on this real torus.

Let us note the natural isomorphism

$$C^+(H_Z) \otimes_{\mathbb{Z}} \mathbb{R} \cong \bigoplus_{i=1}^d C^+(H_Z) \otimes_{O_{K, \epsilon_i}} \mathbb{R} \cong \bigoplus_i C^+(H_Z \otimes_{O_{K, \epsilon_i}} \mathbb{R}).$$

Let us consider the intersection

$$H_Z \otimes_{O_{K, \epsilon_i}} \mathbb{R} \cap (H^{2,0} \oplus H^{0,2}),$$

for each  $i$  ( $i=1,2,\dots,d$ ). Choose an orthogonal basis  $e_i^+, e_i^-$  of this intersection subspace with respect to  $\psi_{\epsilon_i} \otimes \mathbb{R}$ . Then,

$J_i = e_i^+ e_i^-$  ( $i=1,2,\dots,d$ ) defines a complex structure on each factor  $C^+(H_Z \otimes_{O_{K, \epsilon_i}} \mathbb{R})$  by means of right multiplication. Therefore the direct sum  $J = \bigoplus_i J_i$  of  $J_i$  defines a complex structure on our real torus by right multiplication.

**Theorem (4.2)** This complex torus has a structure of abelian variety.

**Proof.** It is well-known that a complex torus with sufficient many endomorphisms becomes automatically an abelian variety, as in our case. ~~And the homogeneous~~ See also Kuga-Satake, or Deligne for  $K=\mathbb{Q}$ . [8] [2]  
Q.E.D.

**Remark.** The dimension of this abelian variety is equal to  $4d$ .

The isogeny class of this abelian variety does not depend upon the choice of the integral structure  $H_Z$  in  $H$ .

§ 5. Involutions of Hilbert modular surfaces and decomposition of abelian varieties.

Let  $\xi$  be a unit such that  $\xi_1 > 0$  and  $\xi_2 < 0$ . Let  $G_\infty, H_\infty,$  and  $F_\infty$  be involutions on  $H \times H$  defined by

$$G_\infty: (z_1, z_2) \in H \times H \longmapsto (\xi_1 z_1, \xi_2 \bar{z}_2) \in H \times H,$$

$$H_\infty: (z_1, z_2) \in H \times H \longmapsto (\xi_2 \bar{z}_1, \xi_1 z_2) \in H \times H,$$

$$F_\infty: (z_1, z_2) \in H \times H \longmapsto (-z_1, -z_2) \in H \times H.$$

Clearly, we have  $G_\infty H_\infty = H_\infty G_\infty = F_\infty$ .

On passing to the quotient  $S$ , we obtain involutions on  $H \times H$ , which we denote by the same symbol

$$G_\infty: S \longrightarrow S, H_\infty: S \longrightarrow S, F_\infty = H_\infty G_\infty = G_\infty H_\infty: S \longrightarrow S.$$

Evidently these involutions act on the homology groups and cohomology groups  $H_2(S, \mathbb{Q}), H^2(S, \mathbb{Q}), W_2 H^2(S, \mathbb{Q})$  etc..

Note here that  $G_\infty$  and  $H_\infty$  changes the orientation. To check this apply  $G_\infty$  and  $H_\infty$  to  $dz_1 \wedge \bar{dz}_1 \wedge dz_2 \wedge \bar{dz}_2$ .

Remark.  $F_\infty$  is the Frobenius at infinity. Namely let  $S_{\mathbb{R}}$  be a canonical model of  $S$  defined over the real number field  $\mathbb{R}$ . Then  $F_\infty$  coincides with the action of the nontrivial element of  $\text{Gal}(\mathbb{C}/\mathbb{R})$  on  $\mathbb{C}$ -valued points  $S$  of  $S_{\mathbb{R}}$ .

Definition. - Proposition (5.1) Put

$$H_{++}(f) = \{ \alpha \in H(f) \mid G_\infty \alpha = \alpha, H_\infty \alpha = \alpha \},$$

$$H_{+-}(f) = \{ \alpha \in H(f) \mid G_\infty \alpha = \alpha, H_\infty \alpha = -\alpha \},$$

$$H_{-+}(f) = \{ \alpha \in H(f) \mid G_\infty \alpha = -\alpha, H_\infty \alpha = \alpha \},$$

$$H_{--}(f) = \{ \alpha \in H(f) \mid G_\infty \alpha = -\alpha, H_\infty \alpha = -\alpha \}.$$

Then we have a direct sum decomposition

$$H(f) = H_{++}(f) \oplus H_{+-}(f) \oplus H_{-+}(f) \oplus H_{--}(f).$$

Moreover  $H_{++}(f), H_{+-}(f), \dots, H_{--}(f)$  are all of rank one over  $K_f$ .

Proof. Note that the Hecke operators commutes with  $G_\infty, H_\infty,$  and  $F_\infty$ .  
And use the Hodge decomposition (Theorem (1.1)).

Noting that  $G_\infty$  and  $H_\infty$  change the orientation of  $S$ , we have the following proposition.

Proposition (5.2). For the intersection form  $\psi_f$ , we have

$$\begin{aligned} \psi_f(H_{++}(f), H_{++}(f)) &= \psi_f(H_{++}(f), H_{+-}(f)) = \psi_f(H_{++}(f), H_{-+}(f)) \\ &= \psi_f(H_{+-}(f), H_{--}(f)) = \psi_f(H_{-+}(f), H_{--}(f)) = \psi_f(H_{--}(f), H_{--}(f)) = 0. \end{aligned}$$

And  $H_{++}(f)$  and  $H_{--}(f)$ , and  $H_{+-}(f)$  and  $H_{-+}(f)$  are mutually dual with respect to  $\psi_f$  each other.

Corollary (5.3).  $\psi_f$  is a kernel form.

Corollary (5.4)  $C^+(H(f))$  the even Clifford algebra over  $H(f)$  with respect to  $\psi_f$  is a direct product of two copies of  $2 \times 2$  matrices with entries in  $K_f$ ;  $C^+(H(f)) \cong M_2(K_f) \oplus M_2(K_f)$ .

Corollary (5.4) Let  $A(f)$  be an abelian variety constructed in the previous section. Then we have a decomposition upto isogenie:

$$A(f) \underset{\text{isogenie}}{\sim} A_{f,1} \times A_{f,1} \times A_{f,2} \times A_{f,2}.$$

Here  $A_{f,i}$  ( $i=1,2$ ) are both of dimension  $d=[K_f:Q]$ , with endomorphism rings  $K_f \hookrightarrow \text{End}_{\mathbb{C}}(A_{f,i}) \otimes_{\mathbb{Z}} \mathbb{Q}$ .

6. Explicite calculation of period lattices of  $A_{f,1}$  and  $A_{f,2}$ .

In this section, we give the period lattices of the isogenie classes of abelian varieties represented by  $A_{f,1}$  and  $A_{f,2}$ .

Let  $f$  be a primitive form of weight 2, and let  $K_f$  be the field of the eigenvalues. Let  $\{\sigma_1 = \text{id}_{K_f}, \sigma_2, \dots, \sigma_d\}$  be the set of embeddings of  $K_f$  into  $\mathbb{R}$ , where  $d = [K_f : \mathbb{Q}]$ . Let  $f^{\sigma_i}$  be the primitive forms obtained from  $f$  by applying  $\sigma_i$  to the Fourier coefficients of  $f$  (We normalize  $f$ ). Let  $H_2(f^{\sigma_i})$  be the homology group attached to  $f^{\sigma_i}$  ( $i=1,2,\dots,d$ ). Let consider the action of  $G_\infty$  and  $H_\infty$  on  $H_2(f^{\sigma_i})$ . And define  $H_2^{++}(f)$  etc. similarly as the case of cohomology groups. We normalize  $f$  such that the "first" Fourier coefficient of  $f$  is a rational number.

Put

$$\omega_f = (2\pi i)^2 f(z_1, z_2) dz_1 \wedge dz_2$$

for such form  $f$ . Then we have the following lemma.

Lemma (6.1)

If  $\gamma^{++} \in H_2^{++}(f)$ , and  $\gamma^{--} \in H_2^{--}(f)$ . Then the period integrals

$$\int_{\gamma^{++}} \omega_f \quad \text{and} \quad \int_{\gamma^{--}} \omega_f \quad \text{are real numbers,}$$

and if  $\gamma^{+-} \in H_2^{+-}(f)$ , and  $\gamma^{-+} \in H_2^{-+}(f)$ , then ~~we have~~ the period integrals

$$\int_{\gamma^{+-}} \omega_f \quad \text{and} \quad \int_{\gamma^{-+}} \omega_f \quad \text{are purely imaginary numbers.}$$

Definition. Fix the above four cycles  $\gamma^{++}, \gamma^{+-}, \gamma^{-+}, \gamma^{--}$  in  $H_2(f) = H_2(S, \mathbb{Q}) \otimes_{\mathcal{K}(f)} K_f$ . And let  $\gamma_i^{++}$ , etc be the element of  $H_2(f^{\sigma_i})$ ,

obtained from  $\gamma^{++}$  etc. with respect to conjugation of  $K_f$  over  $Q$ .

Then we put

$$W_{++}(f^{\epsilon_i}) = \int_{\gamma_i^{++}} \omega_f^{\epsilon_i}, \quad W_{+-}(f^{\epsilon_i}) = \dots\dots\dots$$

Theorem (6.2) The period lattices of  $A_{f,1}$  and  $A_{f,2}$  are given by

$$L_1 = \left\{ v = (v_1, v_2, v_3, \dots, v_d) \in \mathbb{C}^d \mid \right. \\ \left. v_i = \epsilon_i(\alpha) + \epsilon_i(\beta) W_{+-}(f^{\epsilon_i}) / W_{++}(f^{\epsilon_i}) \quad (i=1, 2, \dots, d), \right. \\ \left. \text{for some } \alpha, \beta \text{ in } K_f. \right\},$$

and

$$L_2 = \left\{ v = (v_1, v_2, \dots, v_d) \in \mathbb{C}^d \mid \right. \\ \left. v_i = \epsilon_i(\alpha) + \epsilon_i(\beta) W_{-+}(f^{\epsilon_i}) / W_{++}(f^{\epsilon_i}) \quad (i=1, 2, \dots, d), \right. \\ \left. \text{for some } \alpha, \beta \text{ in } K_f. \right\}.$$

In the calculation of these periods, we need the following theorem.

Theorem (6.3) (The period relation of Riemann-Hodge).

If  $\hat{\psi}_f(\gamma^{++}, \gamma^{--}) = \hat{\psi}_f(\gamma^{+-}, \gamma^{-+})$  for the intersection form  $\hat{\psi}_f$  of  $H_2(f)$ , we have

$$W_{++}(f)W_{--}(f) = W_{+-}(f)W_{-+}(f).$$

Proof of Theorem (6.2). A very easy computation of Clifford algebras. We omit it here, because of a short of time.

以下, 締切時期が過ぎ、過ぎていくので日本語で.

前節までで, Abel多様体の構成に関する, *general nonsense* はおしまいである.

注. さて京都での研究集会のときに質問された, Shimura [23], Mountjoy [22], Hida [21] 等によって定義されている, アーベル多様体と, 上に構成したものとの関係についてであるが, これについてはどのように考えている: 上の3論文は, 本質的には, <sup>ある条件をみたす</sup> rational polarized Hodge structure が, あるいは, それよりアーベル多様体が構成できるというものである (この条件は Mountjoy [22] に書かれている). 最初にとり, rational polarized Hodge structure の "poids"  $n$  とするとき, これらの構成法はある意味で, これを, rational polarized Hodge structure of "poids" 1 と見做すことになる. 従って  $n \neq 1$  のとき, このアーベル多様体は, "自然な" ものとは思われない. 少なくとも, Grothendieck-Deligne の motives の考え方では説明不可能である.

さて, 特に, やはり Hilbert modular cusp forms にアーベル多様体を attach した Hida [21] との関係があるが, この場合 "poids"  $n = [F:\mathbb{Q}]$  (但し  $F$  は考えている総実数体) で,  $n \neq 1$  である. これを納得できない人は, 前節, 定理 (6.2) の周期と [21] で"の周期を調べてみれば, この二つは全くちが, 互いであることが容易に check できるはずである.

§7. Self-conjugate forms に付随する  $\Gamma$ - $\Gamma$  多様体.

序文に書かれた定理 A を証明するのはこの節の目標である。この証明は容易であるが長い。概略のみ書く。

定理 (7.1) (= 定理 A)

$\varphi$  を realen Nebentypus (elliptic modular) cusp form of weight 2,  $\varphi$  は primitive とする。  $f$  を  $\varphi$  の lifting to Hilbert modular forms とする。

$A_\varphi$  を Shimura [20] の  $\Gamma$ - $\Gamma$  多様体,  $A_\varphi \sim B_\varphi \times B'_\varphi$  とする分解とする。  $B_\varphi \sim B'_\varphi$  (isogeny) は知られている。

さてこのとき,  $\mathbb{C}$  上の  $B_\varphi$  と  $A_{f,1}, A_{f,2}$  とは isogenie で異なる。  $\blacktriangle$

(証明の概略) 定理 (6.2) による period lattices の計算による。

$W_{+-}(f)/W_{-+}(f) \in K_f$  はすでに知られる。よって  $A_{f,1}$  と  $A_{f,2}$  とは  $K_f$ -isogenous。問題は  $W_{++}(f), W_{+-}(f), W_{-+}(f), W_{--}(f)$  の計算である。  $K_\varphi = K_f$  はよく知られている。まずこのことがわかる。

命題 (7.2)  $W_{++}(f)/W_{++}(f)^2 \in K_f, \quad W_{--}(f)/W_{--}(f)^2 \in K_f. \quad \blacktriangle$

(証明の概略)  $W_{++}(f), W_{--}(f)$  は Shimura [15] [16] で定義されたものである。この命題は lifting の本来の定義 (Dai-Nagayama [11] Nagayama [12]) から容易に出る。  $\blacktriangle$



問題は  $w_{+-}(\phi), w_{-+}(\phi)$  の計算である.  $w_{+-}(\phi) = w_{-+}(\phi)$  として

よ. Riemann-Hodge の period relation (定理 (6.3)) より

$$\{w_{+-}(\phi)\}^2 / \{w_{++}(\phi)w_{--}(\phi)\} = 1$$

よて

$$\left\{ \frac{w_{+-}(\phi)}{w_{++}(\phi)w_{--}(\phi)} \right\}^2 \in K_f$$

であることは示さねばならない. 仮定からいけば  $w_{+-}(\phi) / w_{++}(\phi)w_{--}(\phi) \in K_f$

はわかる.  $w_{+-}(\phi) / w_{++}(\phi)w_{--}(\phi) \in K_f$  は別の方向で証明する.

基本的な考え方は, Oda [ ] の定理 1 を使って, Poincaré map の adjoint map を考える. すると同期積分と Fourier 係数とが, realen Hecke typus elliptic group forms が表わされる.

(Hirzebruch-Zagier は, これらの同期積分と交点数とを考慮する.)

Typical な Fourier 係数は  $\mathbb{C}$  とする.  $w_{+-}(\phi) / c_{f=1} \in K_f$  は自明にわかる. これは,  $f^0 / w_{++}(\phi)w_{--}(\phi)$  を表わすことである.

Petersson metric  $\langle \cdot, \cdot \rangle_{\text{ellipt.}}$   $\langle \cdot, \cdot \rangle_{\text{Hil.}}$  で表わすことである. (適当な normalization が必要である)

定理 (7.3) (Oda [25] 4  $\alpha$ )

$$c \frac{(\varphi, \varphi)_{\text{ellipt.}}}{(f, f)_{\text{Hil.}}} \in K_f$$

命題 (7.4)

$$(\varphi, \varphi) / w_{++}(\phi)w_{--}(\phi)\sqrt{-1} \in \mathbb{R}_\varphi = K_f,$$

$$(f, f) / w_{++}(\phi)w_{--}(\phi) \in K_f$$

$$\text{系 (7.5)} \quad c \frac{W_+(\varphi)W_-(\varphi)\sqrt{-1}}{W_{++}(f)W_{--}(f)} \in K_f \quad \blacktriangleleft$$

±で  $W_{++}(f)W_{--}(f) / W_+(\varphi)^2 W_-(\varphi)^2 \in K_f$  である。よって  
 $c\sqrt{-1} / W_+(\varphi)W_-(\varphi) \in K_f$ 。 ±で  $W_{+-}(f) / c\sqrt{-1} \in K_f$  である。  
 から、これをより

$$W_{+-}(f) / W_+(\varphi)W_-(\varphi) \in K_f,$$

よ、定理 (7.1) は証明された。  $\blacktriangleleft$

### §8. 定理 B の証明.

これも全く初等的な同期の計算である。但し定理 B の  
 系の証明では、Shimura [15] [16] の結果を少し一般化する必要  
 がある (Neben type elliptic modular の場合に)。ある種の同期の成  
 立ないことをいう必要がある。ここでも再び [25] の Theorem 1 を使  
 う。よってこの命題を得る。

命題 (8.1)  $f$  は self-conjugate primitive form of weight 2 である。

Neben type elliptic cusp form  $\varphi$  より lifting を得るものとす。

$$\text{このとき} \quad W_{++}(f) = W_+(\varphi)^2, \quad W_{+-}(f) = W_{-+}(f) = W_+(\varphi)W_-(\varphi),$$

$W_{--}(f) = W_-(\varphi)^2$  は  $K_f$  に linear independent な数である。

(証明) これは Ribet [18], Murase [19] の結果に帰着される。

定理 (8.2). ([18], [19])  $\varphi$  Nebentypus elliptic modular cusp form of weight 2. Suppose that  $\varphi$  primitive. Let  $B_\varphi$  be the abelian variety attached to  $\varphi$  by Shimura [20]. Then  $B_\varphi$  has no complex multiplication.

命題 (8.1) の否定  $\Rightarrow B_\varphi$  は CM type abelian variety となる。▲

§9.  $l$ -adic realization of the tensor product conjecture と問題.

Tensor product conjecture の  $l$ -adic realization は  $S$  が  $K3$  曲面と双有理同値なときは, Deligne [2] の結果によつて, 非常に多量の形であることが知られている。  $S$  自身でなく symmetric Hilbert modular surfaces が  $K3$  のときと同様であるとされている。

問題.  $A_{f,1}, A_{f,2}$  の定義体  $F$  を代数体  $F$  まで下げることに予想は  $F$  まで下がることを期待している。 Level が 1 ではない Hilbert modular surface については, これは容易に解ける場合が多いとありそうである。 Level が 1 の場合は難しく見える。 Deligne [2] の証明の論法を  $F$  まで一般化しようとするのは,

Hilbert modular surfaces の変形と考える必要がある。

$F$  が  $\mathbb{Q}(\sqrt{53}), \mathbb{Q}(\sqrt{61}), \mathbb{Q}(\sqrt{73})$  のとき  $S$  は elliptic fibre space

over  $\mathbb{P}^1$  と双有理同値で、これについては AKas<sub>1</sub> の結果がある。

$F$  が  $\mathbb{Q}(\sqrt{89}), \mathbb{Q}(\sqrt{97})$  のときは  $S$  は  $P_g = 3, c_1^2 = 2$ , して

$c_1^2 = 2P_g - 4$  の general type の曲面と双有理同値である。これ

の変形についても Horikawa [27] の結果がある。これらの結果か

ら出発して少なくとも特別な場合に、定義体を下げることで、

Tensor product conjecture の  $l$ -adic realization の弱い形での証明は

可能かも知れない。しかし証明に至るいくつかの step はまだ

霧の中にある。いすれにせよ、定理 A によって lifting ができる

Hilbert modular forms  $f$  に対しては  $A_{f,1} \sim A_{f,2}$  の定義は、これ

が  $B_p$  と isogenous なことから自然に下がる。

$A_{f,1}, A_{f,2}$  の定義は一般化された Birch-Swinnerton-Dyer Conjecture

とも compatible であることは容易にわかる。とにかく  $A_{f,1}, A_{f,2}$

の定義が自然なものであることは明らかである。

Tensor product conjecture の  $l$ -adic realization と考えること

と関連の問題である。

## References.

1. Deligne, P.: Valeurs de fonctions L et periodes d'integrales.  
Automorphic forms, representations, and L-functions.  
Proc. of sympo. in pure math., vol XXXIII, part 2,  
313-346 (1979).
2. ———.: La conjecture de Weil pour les surfaces K3. Invent. math.  
15, 206-226 (1972).
3. ———.: Theorie de Hodge. II.
4. ———.: Theorie de Hodge. III. Inst. Hautes Etudes Sci. Pub. Math.  
44, 5-77 (1974).
5. Hirzebruch, F.: Hilbert modular surfaces. L'Ens. Math. 19, 183-281  
(1973).
6. ———.: Kurven auf den Hilbertschen Modulflachen und Klassen-  
zahlrelationen. Classification of algebraic varieties  
and compact complex manifold. Lecture Notes in Math.  
412, pp.75-93. Berlin-Heidelberg-New York : Springer  
1974.
7. ———., Van de Ven, A.: Hilbert modular surfaces and the classi-  
fication of algebraic surfaces. Inventiones math. 23,  
1-29 (1974).
8. Kuga, M., Satake, I.: Abelian varieties attached to polarized K3  
surfaces. Math. Annalen 169, 239-242 (1967).
9. Satake, I.: Clifford algebras and families of abelian varieties.  
Nagoya J. Math. 27-2, 435-446 (1966).
10. Rapoport, M.: Compactifications de l'espace de modules de Hilbert-  
Blumenthal. Compositio math. 36, 255-335 (1978).

11. Doi, K., Naganuma, H.: On the functional equation of certain Dirichlet series. *Inventiones math.* 9, 1-14 (1969).
12. Naganuma, H.: On the coincidence of two Dirichlet series associated with cusp forms of Hecke's "Neben"-type and Hilbert modular forms over real quadratic field. *J. Math. Soc. Japan* 25, 547-555 (1973).
13. Zagier, D.: Modular forms associated to real quadratic fields. *Inventiones math.* 30, 1-46 (1975).
14. Shimura, G.: On some arithmetic properties of modular forms of one and several variables, *Ann. of Math.* (2) 102, 491-515, (1975).
15. ———.: On the periods of modular forms. *Math. Ann.* 229, 211-221 (1977).
16. ———.: The special values of zeta functions associated with cusp forms. *Communications on pure and applied math.*, XXIX. 783-804 (1976).
17. Saint-Donant.: Descent cohomologique. S.G.A.7.   
Lecture Notes in Math.
18. Ribet, K.A.: ——— *to appear Math. Ann.*
19. Momose, F.: ——— *Preprint.*
20. Shimura, G.: Introduction to the arithmetical theory of automorphic function.

21. Hida, H.: On abelian varieties with complex multiplication as factors of the abelian variety attached to Hilbert modular forms. Japan J. Math., 5, 157-208(1979).
22. Mountjoy, R.H.: Abelian varieties attached to representations of discontinuous groups. Amer. J. of Math. 89, 149-224(1967).
23. Shimura, G.: Sur les integrales attachees aux formes automorphes. Journal of Math. Soc. of Japan, vol.11, 291-311(1959).
24. Ueno, K., van der Geer, : ————— A preprint
25. Oda, T. : ————— Math. Ann (1977)
26. Kas, A.: On the deformation types of regular elliptic surfaces. Complex Analysis and Algebraic Geometry, Iwanami-shoten (1977)
27. Horikawa, E.: Algebraic surfaces of general type with small  $c_2$ , I
28. Manin, Yu.: Parabolic forms and —————
30. Paily - Poul. : ————— 31. Freitag. —————

注意 文献表で題名が正しいは, preprint 等 2, 4, 9, 12 にも注意,  
 印刷事故下とい。