数理解析研究所講究録
438巻 1981年 1-5

Topics on Kac-Moody Lie algebras

Yōhei Tanaka (Nagoya Univ.)

In this note we introduce Kac-Moody Lie algebras, their representation theory, and relation with theory of differential equations.

I. Kac-Moody Lie algebras ([6],[I4],[I5])

Let $C=(c_{ij})$ be a $n\times n$-matix satisfying following conditions;

$$c_{ij}\in\mathbb{Z}\ ,\ c_{ii}=2\ ,\ c_{ij}\leq 0\ (i\neq j)\ ,\ c_{ij}=0 \Leftrightarrow c_{ji}=0\ ,$$

and define Lie algebra $\mathcal{L}(C)$ over $\mathbb{C}$ with generators $\{h_i, e_i, f_i\ \ i=I,\cdots,n\}$ and relations;

$$\begin{cases} [h_i,h_j]=0\ ,\ [h_i,e_j]=c_{ij}e_j\ ,\ [h_i,f_j]=-c_{ij}f_j\ ,\ [e_i,f_j]=\delta_{ij}h_i\ , \\ (\mathrm{ad}\ e_i)^{-c_{ij}+I}e_j=0\ ,\ (\mathrm{ad}\ f_i)^{-c_{ij}+I}f_j=0\ . \end{cases}$$

$\mathcal{L}=\mathcal{L}(C)$ is called Kac-Moody Lie algebra.

$\mathcal{L}$ has $\Gamma=\mathbb{Z}^n$ grading with $\deg h_i=(0,\cdots,0)$, $\deg e_i=(0,\cdots,\overset{i}{I},\cdots,0)=a_i$, and $\deg f_i=-a_i$.

Put $\mathcal{L}_a=\{x\in\mathcal{L}\mid \deg x=a, (a\in\Gamma)\}$, $m_a=\dim\mathcal{L}_a$, $\Delta=\{a\in\Gamma\setminus\{0\}\mid m_a\neq 0\}$.

From defining relations $\mathcal{L}$ has vector space decomposition

$$\mathcal{L}=\mathcal{H}\oplus\mathcal{L}_+\oplus\mathcal{L}_-\ \text{ where } \mathcal{H}=\sum_i\mathbb{C}h_i\ ,\ \mathcal{L}_+=\langle e_I,\cdots,e_n\rangle\ ,\ \mathcal{L}_-=\langle f_I,\cdots,f_n\rangle\ .$$

So Δ is disjoint union of $\Delta_+=\Gamma_+\cap\Delta$ and $\Delta_-=(-\Gamma_+)\cap\Delta$

$(\Gamma_+=\mathbb{Z}_+a_I+\cdots+\mathbb{Z}_+a_n)$

For each i , define s_i $GL(\Gamma)$ by

$$s_i(a_j)=a_j-c_{ij}a_i$$

and $W=\langle s_i \mid i=I,\cdots,n\rangle$.

If C is decomposable, i.e. has a permutation matrix P such that

$PCP^{-I}=\left(\begin{array}{c|c} C_I & 0 \\ \hline 0 & C_2 \end{array}\right)$, then $\mathcal{L}(C)\cong\mathcal{L}(C_I)+\mathcal{L}(C_2)$, $W(C)\cong W(C_I)\times W(C_2)$.

So we assume C is indecomposable.

A finite-dimensional complex simple Lie algebra is isomorphic to a $\mathcal{L}(C)$ whose W(C) is finite group. Equivalent condition on C to finiteness of W(C) is that there exist positive numbers $d_I,\cdots,d_n$ such that $\begin{pmatrix} d_I & & \\ & \ddots & \\ & & d_n \end{pmatrix}C$ is a positive definite symmetric matrix.

If we replace "positive definite" by "positive semi-definite" in this condition, we get a new class of infinite-dimensional Lie algebras. They are called Euclidean (or affine) Lie algebras. We deal with Euclidean case hereafter.

2. Character formula and denominator formula

Put $V=\mathbb{C}\otimes_{\mathbb{Z}}\Gamma$, and for $\lambda\in\mathfrak{h}^*$ let $c_\lambda:W\to V$ be the I-cocycle, i.e. satisfies

$c_\lambda(w_Iw_2)=w_Ic_\lambda(w_2)+c_\lambda(w_I)$, such that $c_\lambda(s_i)=\lambda(h_i)a_i$ $(i=I,\cdots,n)$.

If $\lambda(h_i)\in\mathbb{Z}$ for all i, λ is said integral, and moreover if $\lambda(h_i)\geq 0$, λ is said dominant integral. It is easy to see

$$\begin{cases} \lambda:\text{integral} \Rightarrow c_\lambda(W)\subset\Gamma \\ \lambda:\text{dominat integral} \Rightarrow c_\lambda(W)\subset\Gamma_+ \end{cases}$$

To each dominant integral, there exists $\mathcal{L}$-module V^λ, unique up to isomorphism, with the following property;

$$0\neq{}^{\exists}v\in V \text{ s.t. } \begin{cases} e_iv=0 \ , \ h_iv=\lambda(h_i)v \ , \ f_i^{\lambda(h_i)}v=0 \quad (i=I,\cdots,n) \\ V=\sum\mathbb{C}f_{i_I}f_{i_2}\cdots f_{i_k}v \end{cases}$$

V^λ is called standard module with highest weight λ.

For $a\in\Gamma_+$, let $V_a^\lambda=\sum_{a_{i_1}+\cdots a_{i_k}=a}\mathbb{C}f_{i_I}\cdots f_{i_k}v$, and put

$\mathrm{ch}\,V^\lambda=\sum_{a\in\Gamma_+}\dim V_a^\lambda e(-a)$ ($\mathrm{ch}V^\lambda$ is in $\mathbb{Z}[[e(-a_I),\cdots,e(-a_n)]]$.)

Let $\rho \in \mathfrak{h}^*$ be defined by $\rho(h_i)=1$ for all i. We have

character formula ([5],[8],[10])

$$\mathrm{ch}\, V^{\lambda} = \frac{\sum_{w\in W} \det(w) e(-c_{\lambda+\rho}(w))}{\sum_{w\in W} \det(w) e(-c_{\rho}(w))}$$

and

denominator formula

$$\sum_{w\in W} \det(w) e(-c\ \ (w)) = \prod_{a\in\Delta_+} (1-e(-a))^{m_a}$$

If we make specialization $e(-a_i) \mapsto q^{m_i}$ $(m_1,\cdots,m_n \geq 0)$ to both sides of denominator (or character) formula, we have various combinatorial identities. ([7],[8],[12],[16])

3. K-dV equation and construction of representation

Let $u(x), V_{nl}(x)$ $(n=1,2,3,\cdots, 1\leq l\leq 2n-2)$ be functions in infinite many variables $x=(x_1,x_3,x_5,\cdots)$. From the compatibility conditions of linear partial differential equations

$$\left.\begin{aligned} &\left\{\left(\frac{\partial}{\partial x_1}\right)^2 + u(x)\right\}\psi(x) = \lambda\psi(x) \\ &\frac{\partial\psi}{\partial x_{2n-1}} = \left\{\left(\frac{\partial}{\partial x_1}\right)^{2n-1} + V_{n1}\left(\frac{\partial}{\partial x_1}\right)^{2n-3} + \cdots + V_{n\,2n-2}\right\}\psi \quad (n=1,2,\cdots) \end{aligned}\right\}$$

we obtain non-linear differential equations on u, and V_{nl} are expressed by differential polynomials of u. These differential equations are called K-dV equations. A function $\tau(x)$ is called τ-function, if $u=\left(\frac{\partial}{\partial x_1}\right)^2 \log\tau$ is a solution of K-dV equations. The space of infinitesimal trasformations of τ-functios forms a Lie algebra $\mathcal{L}$ of linear trans-formations on $C[x_1,x_3,\cdots]$. Date, Jimbo, Kashiwara and Miwa find ([1]) that $\mathcal{L} \cong \mathcal{L}\begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix}$ and $\mathcal{L}$-module $C[x_1,x_3,\cdots]$ coincides with the standard module constructed by Lepowsky and Wilson. ([11]). Kac, Kazhdan, Lepowsky and Wilson construct standard modules for other Euclidean Lie algebras in analogous way to L-W, and D-J-K-M show

they correspond to some non-linear equations like K-dV.([2],[3] ,[9])

4. Remarks

(I) About classification and realizations of Euclidean Lie algebras,c.f. [6],[I5].

(2) Let $\mathfrak{R} \subset \mathcal{L}(C)$ be the maximam homogenious ideal with $\mathfrak{R} \cap \mathfrak{h} = 0$ .Strictly speaking, $\mathfrak{g}(C) = \mathcal{L}(C)/\mathfrak{R}$ is Kac-Moody Lie algebra.But finite or Euclidean case $\mathfrak{R} = 0$,and we conjecture that $\mathfrak{R} = 0$ for any case.

(3) Character formula and denominator formula are analogy of finite case, and proved for $\mathfrak{g}(C)$ with symmetrizable C,i.e. there exist positive numbers $d_I, \cdots, d_n$ such that $\begin{pmatrix} d_I & & \\ & \ddots & \\ & & d_n \end{pmatrix} C$ is symmetric matrix.

(4) Original K-dV equation is obtained by to put $x_5, x_7, \cdots$ constant.

(5) Frenkel and Kac ([4]) construct standard modules in different way from K-K-L-W.It will be interesting to describe isomorphism explicitly between K-K-L-W's module and F-K's.

References

[I] E.Date,M.Kashiwara and T.Miwa,Transformation groups for soliton equations II, RIMS preprint 357.

[2] E.Date,M.Jimbo,M.Kashiwara and T.Miwa, " IV,RIMS preprint 359 .

[3] " " " ", RIMS preprint 362.

[4] I.B.Frenkel and V.G.Kac,Basic representations of affine Lie algebras and dual resonance models,Inv.Math.,62(I980),23-66.

[5] H.Garland and J.Lepowsky,Lie algebra homology and the Macdonald-Kac formulas,Inv.Math.,34(I976),37-76.

[6] V.G.Kac,Simple irreducible graded Lie algebras of finite growth,Math.USSR Izvestija,2(I968),I27I-I3II.

[7] V.G.Kac,Infinite-dimensional Lie algebras and Dedekind's η-function,

Func.Analy.and its App.,8(I974),68-70.

[8] V.G.Kac,Infinite-dimensional algebras,Dedekind's η-function,classical Mobius function and the very strange formula,Adv.in Math.,30(I978),85-I36.

[9] V.G.Kac,D.A.Kazhdan,J.Lepowsky and R.L.Wilson,Realization of the basic representations of the Euclidean Lie algebra,Adv.in Math.,to appear.

[I0] J.Lepowsky,Lectures on Kac-Moody Lie algebras,Université Paris VI,I978.

[II] J.Lepowsky and R.L.Wilson,Construction of the affine Lie algebra $A_1^{(1)}$, Comm.Math.Physics,62(I978),43-53.

[I2] J.Lepowsky and R.L.Wilson,A Lie theoretic interpretation and proof of Rogers-Ramanujan identities, preprint.

[I3] I.G.Macdonald,Affine root systems and Dedekind's η-function,Inv.Math., I5(I972),9I-I43.

[I4] R.V.Moody,A new class of Lie algebra,J.of Alg.,I0(I968),2II-230.

[I5] ",Euclidean Lie algebra,Canad.J.Math.,2I(I969),I432-I454

[I6] " ,Macdonald identities and Euclidean Lie algebras,Proc.A.M.S. 48(I975),43-52.