

204

On the Uniformization of Analytic Sets with the Countable Sections
and Related Results

By Yutaka YASUDA*

Institute of Educational Technology, Tokai University, Hiratuka

Introduction. In a letter of Hadamard to Borel [1], Hadamard discussed an effective choice of an element from a given Borel set. Then Luzin [8] introduced the general problem of uniformization, and announced several results. One of these is that every analytic set is uniformized by the difference of two analytic sets. In 1978, Steel and Martin gave an example of an analytic set which can not be uniformized by the difference of two analytic sets (Cf. Mochovakis [10]).

In this paper, we will state some results concerning uniformization of analytic and Borel sets with special properties and enumerability of Borel, analytic and co-analytic sets. Our main aim is to give the positions in the σ -algebra generated by the analytic sets. We use the recursion theoretic methods, or Effective Descriptive Set Theory. We assume familiarity with [12,14], and use the notations of them. In our proofs we shall often use the following uniformization theorem which was first proved by Kondô in the classical case:

Number Uniformization Theorem (Kondô [3], Kreisel [7]). Every
 \prod_1^1 set in $\omega^\omega \times \omega^\omega$ can be made uniform by a set of the same class.

*The author is partly supported by Grant-in-Aid for Scientific Research, Proj, No.434007.

On the Uniformization of Analytic Sets

We would express our thanks to Professor Hisao Tanaka for his valuable aid in the preparation of the paper.

§ 1. Our main tools in this paper are Harrison's Effective Perfect Set Theorem [2] and

Theorem 1. There is an \prod_1^1 relation $C(\alpha, n)$ and a mapping $(\alpha, n) \mapsto \delta(\alpha, n) \in \omega$ defined on the set C such that

$$\beta \in \Delta_1^1(\alpha) \Leftrightarrow \exists n [C(\alpha, n) \& \beta = \delta(\alpha, n)]$$

and the relation " $\beta = \delta(\alpha, n)$ " is Δ_1^1 uniformly on $(\alpha, n) \in C$. Conversely there is a mapping $(\alpha, \beta) \mapsto \delta^{-1}(\alpha, \beta) \in \omega$ defined on the set $\{(\alpha, \beta) \mid \beta \in \Delta_1^1(\alpha)\}$ such that

$$\beta \in \Delta_1^1(\alpha) \Rightarrow \delta(\alpha, \delta^{-1}(\alpha, \beta)) = \beta$$

and the relation " $n = \delta^{-1}(\alpha, \beta)$ " is Δ_1^1 uniformly on $\beta \in \Delta_1^1(\alpha)$.

The first part of Theorem 1 is due to Kechris [3].

§ 2. We can extend Steel-Martin result (Cf. Introduction) as follows

Theorem 2. There is an \sum_1^1 set in $\omega \times \omega$ which can not uniform by an $(\sum_1^1)_\rho$ set.

This shows that Watanabe's result [17], which is an improvement of Luzin [10], (Cf. Tanaka [15]) is the best possible.

For the analytic sets with the countable sections, we have

Theorem 3. Every \sum_1^1 set in $\omega \times \omega$ with the countable sections can be uniform by an $(\sum_1^1)_\rho$ set.

For the negative side as Theorem 1 we have

Theorem 4. There is an \sum_1^1 set in $\omega \times \omega$ with the countable sections which can not be uniform by an analytic set.

Corollary 5. There is an \sum_1^1 set in $\omega \times \omega$ with the countable sections which can not be uniform by a co-analytic set.

§3. Luzin [10] showed that every Borel(analytic) set with the countable sections is the union of countably many Borel(analytic) curves(Cf.Kondô [7] and Tanaka [16]). We can prove

Theorem 6. Let B be an Δ_1^1 set in $\omega^{\omega} \times \omega^{\omega}$ such that if $B^{\langle \alpha \rangle}$ is non-empty then $B^{\langle \alpha \rangle}$ is denumerable. Then there is an Δ_1^1 set B^* in $\omega^{\omega} \times \omega^{\omega}$ such that

- (i) $B^{*\langle n \rangle}$ s are Δ_1^1 pairwise disjoint uniformizers of B,
- (ii) $B(\alpha, \beta) \Leftrightarrow \exists n B^*(n, \alpha, \beta)$.

Corollary 7. Let A be an Σ_1^1 set in $\omega^{\omega} \times \omega^{\omega}$ such that if $A^{\langle \alpha \rangle}$ is non-empty then $A^{\langle \alpha \rangle}$ is denumerable. Then there is an Σ_1^1 set A^* in $\omega^{\omega} \times \omega^{\omega}$ such that

- (i) $A^{*\langle n \rangle}$ s are Σ_1^1 pairwise disjoint curves,
- (ii) $A(\alpha, \beta) \Leftrightarrow \exists n A^*(n, \alpha, \beta)$.

§4. Sierpiński [14] showed that every denumerable G_δ set is effectively denumerable. We have

Theorem 8. Let E be a denumerable Δ_1^1 set of reals. Then the members of E can be enumerated by an Δ_1^1 mapping.

Kondô [5] showed that every denumerable analytic set is effectively denumerable. Sampei [12] and Tanaka [15] showed that every denumerable Σ_1^1 set of reals can be enumerated by an Δ_2^1 mapping. We can prove

Theorem 9. Let E be a denumerable Σ_1^1 set of reals. Then the members of E can be enumerated by an $(\Sigma_1^1)_{p\sigma} \cap (\Sigma_1^1)_{p\sigma c}$ mapping.

Theorem 10. There is a denummable Σ_1^1 set of reals whose members can not be enumerated by an Σ_1^1 mapping.

Corollary 11. There is a denumerable Σ_1^1 set of reals whose members can not be enumerated by an \prod_1^1 mapping.

For the denumerable \prod_1^1 set of reals we have

Theorem 12. Let E be a denumerable \prod_1^1 set of reals. Then the members of E can be enumerated by an $(\Delta_2^1)_{\rho\sigma}$ mapping.

This is an answer of a problem of Kondô [6].

In the neare future, we will publish the detaild proofs of the above theorems elsewhere.

References

- [1] Cinq lettre sur la théorie des fonctions, Bulletin de la Société mathématique de France, 1904.
- [2] Harrison, J., Doctoral Dissertation, Stanford Univ., 1967.
- [3] Kechris, A. S., Lecture Notes on Descriptive Set Theory, M. I. T., 1973.
- [4] Kondô, M., Sur l'uniformisation des complémentaires analytiques et les ensembles projectifs de la seconde classe, Japanese J. Math., 15(1938), 197-230.
- [5] Kondô, M., On denumerable analytic sets(in Japanese), Tokyo Buturi-Gakkô Zasi, No.567(1939), 1-6.
- [6] Kondô, M., On denumerable co-analytic sets(in Japanese), Isô-Sûgaku, Vol.2, No.1(1939), 69.
- [7] Kondô, M., Les Problèmes Fondamentaux parus dans Cinq Letters sur la Théorie des Ensembles, Proc. Faculty of Sci., Tokai Univ., 9(1973), 21-35.
- [8] Kreisel, G., The axiom of choice and the class of hyperarithmetic functions, Indag. Math., 24(1962), 307-319.

- [9] Luzin, N., Sur le problème de M. Jacques Hadamard d'uniformisation des ensembles, Mathematica, 9(1930), 54-66.
- [10] Luzin, N., Leçon sur les ensembles analytiques et leurs applications, Gauthier-Villars, 1930.
- [11] Moschovakis, Y. N., Descriptive Set Theory, North-Holland, 1980.
- [12] Sampei, Y., On the principle of effective choice and its applications, Comment. Math. Univ. St. Paul., 15(1966), 29-42.
- [13] Shoenfield, J. R., Mathematical logic, Addison-Wesley, 1967.
- [14] Sierpiński, W., Une démonstration du théorème sur la structure des ensembles de points, Fund. Math., 1(1920), 1-6.
- [15] Tanaka, H., Some results in the effective descriptive set theory, RIMS, Kyoto Univ. Ser. A, 3(1967), 11-52.
- [16] Tanaka, H., Recursion theory in analytical hierarchy, Comment. Math. Univ. St. Paul., 27(1978), 113-132.
- [17] Watanabe, H., Une remarque sur l'uniformisation des ensembles analytiques plans, Tôhoku Math. J., 5(1954), 79-82.