

The Enumeration Theorems under (Permutation) Group Action
and Their Applications to Combinatorial Problems

by

Hiroshi Narushima

Department of Mathematical Sciences, Faculty of Science, Tokai University,
Hiratsuka, Kanagawa, Japan

We present a short survey for enumerative combinatorial theory and then describe some topics on enumeration of mapping systems. Roughly speaking, I think, the theory is divided into the following three branches, (1) the enumeration theorems under (permutation) group action and their applications, (2) the principle of inclusion-exclusion and the Möbius function on partially ordered sets, and (3) a trial for uniting (1) and (2), in which (1) and (2) are rather established but (3) is not still done. We now have a general discussion on "enumeration" and "characterization" of objects. Let Ω be a nonempty set of objects. Let A be a set of attributes, mathematically speaking, an abstract set with a given structure. Let \mathfrak{C} be a relation between Ω and A . Then a Galois correspondence

$$p(\Omega) \begin{array}{c} \xleftarrow{\tilde{\mathfrak{C}}} \\ \xrightarrow{\mathfrak{E}} \end{array} p(A)$$

is induced by the relation \mathfrak{C} and the inverse relation \mathfrak{E} (see Ore [55] for the Galois correspondence induced by a relation). The notation \mathfrak{C} and \mathfrak{E} have a suggestive look of "characterization" and "enumeration". The map $\tilde{\mathfrak{C}}$ is closely related to the structure theory and so the map $\tilde{\mathfrak{E}}$ to the enumeration theory. More generally speaking, the two maps have a close connection with "analysis" and "synthesis". Also, this correspondence seems to be a set theoretic representation of Hamilton's diagram in classical logics, which shows the relationships between "extension" and "intension" of a concept. Speaking in respect of "enumeration", if $\tilde{\mathfrak{E}}(A) = \phi$ then it means that any object with the abstract property A does not exist, if $\tilde{\mathfrak{E}}(A) \neq \phi$ then it means that some objects with the property A exist, resulting in a problem of

enumerating the objects. Thus, the Galois correspondence induced by a relation is very useful as a framework in enumeration. Let's consider more concretely in connection with our subject.

1. The Enumeration Theorems Under Group Action Let Ω be a nonempty finite set and A be a permutation group G on Ω . We define a relation \mathcal{Q} between Ω and G in the following : for each x in Ω and α in G , $x\mathcal{Q}\alpha$ if and only if $\alpha(x) = x$. Then, a Galois correspondence

$$\rho(\Omega) \xrightleftharpoons[\tilde{N}]{\mathcal{Q}} \rho(G)$$

is induced by the relation \mathcal{Q} and the inverse relation \mathcal{H} . We see easily that for each X in $\rho(\Omega)$ $\tilde{\mathcal{Q}}(X) (= \bigcap_{x \in X} \tilde{\mathcal{Q}}(x))$ is the invariant group of X and that for each α in G $|\tilde{N}(\alpha)|$ is the character of α , that is, $\tilde{N}(\alpha) = \{x \in \Omega \mid \alpha(x) = x\}$. Note that $|\tilde{N}(\alpha)|$ is equal to the number of cycles of length 1 in α . The following theorem considered one of the fundamental theorems in enumerative combinatorial theory seems to be the origin of a series of enumeration theorems under group action by Frobenius, Redfield, Pólya, De Bruijn, Harary and Palmer [3,17,15,1,7,14].

Theorem (Frobenius). Let Ω/G denote the set of orbits in Ω relative to G and O_x be the orbit containing an element x in Ω . Then the following identities hold,

$$(1) |O_x| = |G| / |\tilde{\mathcal{Q}}(x)|$$

$$(2) |\Omega/G| = \frac{1}{|G|} \sum_{\alpha \in G} |\tilde{N}(\alpha)|.$$

This theorem shows that the computability of $|\tilde{\mathcal{Q}}(x)|$ and $|\tilde{N}(\alpha)|$ is essential in enumeration. The author has vaguely known that Burnside took up this theorem in his textbook[3]. On the other hand, at this symposium he is precisely taught by Professor Peter M. Neumann that the theorem was formulated by Frobenius (from Cauchy through Netto) and Burnside took up the theorem in his textbook. Since Frobenius's theorem, Redfield [17], Pólya[15], De Bruijn[1,2], Harary and Palmer[6-10,14] fruitfully used the cycle index of a permutation group G as a generating function, and they and others (Davis[4], Read[16], Harrison[11,12], Robinson[14],...) applied their methods to

enumeration of mapping patterns, chemical structures, graphs and machines. It is worth noticing that Redfield's work was referred by Littlewood[13], Read[16], Foulkes[5] and appreciated by Harary and Palmer[8]. The cycle index (group reduction function by Redfield) of a permutation group is defined as follows.

$$P(G; x_1, \dots, x_n) \text{ (or } Z(G)) = \frac{1}{|G|} \sum_{\alpha \in G} x_1^{p_1} x_2^{p_2} \dots x_n^{p_n},$$

where n is the degree of G and the cycle structure $t(\alpha)$ (or cycle type) of α is (p_1, \dots, p_n) , that is, p_i is the number of cycles of length i in α .

We now show some examples in illustration of the theorems. Example 1 is due to De Bruijn[1,2], Harary and Palmer[7] (the special case is due to Davis[4]) and it is elementary but essential in enumeration of finite mapping systems such as mapping patterns (schemata), chemical structures, graphs and machines. Example 2 is one which Harary and Palmer[10] showed to illustrate one of Redfield's theorems[17] (later Read[16]). Example 3 is one which the author[44,48] introduced by the motive of simplifying figure-works in enumerating the reducible or irreducible types of finite systems such as mappings, finite automata and sequential machines.

Example 1. Let $\mathcal{F}(S,T)$ denote the set of mappings from a finite set S to a finite set T . Let G_S be a permutation group on S and G_T be a permutation group on T . Then we define $f \approx g$ for each f and g in $\mathcal{F}(S,T)$ if and only if there are α in G_S and β in G_T such that $\beta(f(s)) = g(\alpha(s))$ for all s in S . The product group $G_S \times G_T$ is easily considered to be a permutation group on $\mathcal{F}(S,T)$. Therefore, for each (α, β) in $G_S \times G_T$ we obtain the following identity.

$$|\tilde{N}(\alpha, \beta)| = \prod_{1 \leq i} \left(\sum_{j|i} j q_j \right)^{p_i},$$

where $t(\alpha) = (p_1, \dots, p_i, \dots)$ and $t(\beta) = (q_1, \dots, q_i, \dots)$. Thus, the number of equivalence classes under the relation \approx (orbits in $\mathcal{F}(S,T)$ relative to $G_S \times G_T$, called patterns or schemata) is obtained from Frobenius'es theorem.

Example 2. Let D_n denote the dihedral group of degree n generated by the cycle $(1\ 2\ \dots\ n)$ and the relation $(1\ n)(2\ n-1)\dots$. Then the cycle index of D_n is as follows.

$$Z(D_n) = \frac{1}{2}Z(C_n) + \begin{cases} \frac{1}{2}x_1x_2^{(n-1)/2} & n:\text{odd} \\ \frac{1}{4}(x_2^{n/2} + x_1^2x_2^{(n-2)/2}) & n:\text{even} \end{cases}$$

where C_n is the cyclic group of degree n generated by the cycle $(1\ 2\ \dots\ n)$. The cycle index of C_n also is as follows.

$$Z(C_n) = \frac{1}{n} \sum_{k|n} \varphi(k) x_k^{n/k},$$


where $\varphi(k)$ is the Euler φ -function. The cycle indices of D_n and C_n were shown by Redfield[17]. On the other hand, since the automorphism group of a cycle graph of order n is D_n , by the Redfield's theorem (later by Read[16]) the number of different superpositions of 2 cycle graphs of order n with the same set of unlabeled points is equal to $Z(D_n) \cap Z(D_n)$, where \cap is the cap operation (originally denoted Ω by Redfield). For $m (\geq 2)$ monomials $x_1^{i_1}x_2^{i_2}\dots x_r^{i_r}, x_1^{j_1}x_2^{j_2}\dots x_r^{j_r}, \dots$, the cap \cap is defined by

$$\begin{aligned} & x_1^{i_1}x_2^{i_2}\dots x_r^{i_r} \cap x_1^{j_1}x_2^{j_2}\dots x_r^{j_r} \cap \dots \\ &= \begin{cases} \left(\prod_{k=1}^r k^{i_k} i_k!\right)^{m-1} & i_k = j_k = \dots \text{ for all } k \\ 0 & \text{otherwise (also } 0^0=1) \end{cases} \end{aligned}$$

By linearity, the cap operation may be extended to arbitrary polynomials in the variables. For example, for $n = 5$

$$\begin{aligned} Z(D_5) &= \frac{1}{2} \times \frac{1}{5} (x_1^5 + 4x_5) + \frac{1}{2} x_1 x_2^2 = \frac{1}{10} (x_1^5 + 4x_5 + 5x_1 x_2^2), \\ Z(D_5) \cap Z(D_5) &= \frac{1}{100} (x_1^5 \cap x_1^5 + 16x_5 \cap x_5 + 25x_1 x_2^2 \cap x_1 x_2^2) \\ &= \frac{1}{100} (120 + 80 + 200) = 4. \end{aligned}$$

The four superpositions are as follows :



Example 3. Let $PL(S)$ denote the partition lattice of a n -set S . Let $\mathcal{G}(S)$ be the symmetric group on S . Then a permutation group $\{\tilde{\alpha} | \alpha \text{ in } \mathcal{G}(S)\}$ on $PL(S)$, written $G(PL(S))$, is induced by $\tilde{\alpha}(\pi) = \{\alpha(B) | B \in \pi\}$ for each π in $PL(S)$. Furthermore, a permutation group $\{\tilde{\tilde{\alpha}} | \tilde{\alpha} \in G(PL(S))\}$ on the set $C(PL(S))$ of chains in $PL(S)$, written $G(C(PL(S)))$, is induced by $\tilde{\tilde{\alpha}}(\mathbf{c}) = \{\tilde{\alpha}(\pi) | \pi \in \mathbf{c}\}$ for each \mathbf{c} in $C(PL(S))$. In the sequel, we have

$$\mathcal{G}(S) \cong G(PL(S)) \cong G(C(PL(S)))$$

where \cong denotes "group isomorphic". We now define a cardinal congruence relation $\equiv_{\mathbf{c}}$ on $C(PL(S))$ in the following: for each \mathbf{c} and \mathbf{c}' in $C(PL(S))$ $\mathbf{c} \equiv_{\mathbf{c}} \mathbf{c}'$ if and only if there is α in $\mathcal{G}(S)$ such that $\tilde{\tilde{\alpha}}(\mathbf{c}) = \mathbf{c}'$. Since the cardinal congruence classes under the equivalence relation $\equiv_{\mathbf{c}}$ are the orbits in $C(PL(S))$ relative to $G(C(PL(S)))$, the following identity is obtained.

$$|O_{\mathbf{c}}| = n! / |G(C(PL(S)))_{\mathbf{c}}|.$$

When $\ell(\mathbf{c})$ (length of $\mathbf{c} = |\mathbf{c}| - 1$) = 0, it results in the well known formula, that is, for π in $PL(S)$

$$|O_{\pi}| = n! / \prod_{i=1}^n (i!)^{p_i} (p_i!),$$

where $\mathbf{t}(\pi)$ (the type of π) = $[(1)^{p_1} (2)^{p_2} \dots (n)^{p_n}]$ in the set of $P(n)$ of partitions of n . For π in $PL(S)$, $G(PL(S))_{\pi}$ is equal to the automorphism group of π defined by Ore[53]. We next consider the case of $\ell(\mathbf{c}) = 1$. Let $\gamma(\mathbf{x}) = (p_1, p_2, \dots, p_k)$ for $\mathbf{x} = [(1)^{p_1} (2)^{p_2} \dots (k)^{p_k}]$ in $P(k)$. Let π and τ be any elements in $PL(S)$ and $X(\pi) = \{B \text{ in } \pi | B \subseteq X\}$ for X in τ . Let $[\pi, \mathbf{l}]$ be an interval in $PL(S)$ such that $\gamma(\mathbf{t}(\pi)) = \mathbf{v}$ and $P(\mathbf{v})$ be the set of partitions of a vector \mathbf{v} , where \mathbf{l} is a unique maximal element in $PL(S)$. Then an extended type function $\mathbf{t}_{\pi}: [\pi, \mathbf{l}] \rightarrow P(\mathbf{v})$ is well defined by

$$\mathbf{t}_{\pi}(\tau) = [\gamma(\mathbf{t}(X(\pi))) | X \text{ in } \tau].$$

It is shown that $\pi < \tau \equiv_{\mathbf{c}} \pi' < \tau'$ if and only if $\mathbf{t}_{\pi}(\tau) = \mathbf{t}_{\pi'}(\tau')$, and the following formula is obtained.

$$|O_{\pi < \tau}| = n! / \left(\prod_{i=1}^n (i!)^{p_i} \prod_{j=1}^r (q_j! \left(\prod_{i=1}^n v_{ij}! \right)^{q_j} \right),$$

where $\mathbf{t}(\pi) = [(1)^{p_1} \dots (n)^{p_n}]$, $\mathbf{t}_{\pi}(\tau) = [v_1^{q_1} \dots v_r^{q_r}]$ and $v_j = (v_{1j}, \dots, v_{nj})$ for $1 \leq j \leq r$. For example, for

$\pi = \{\overline{1}, \overline{2}, \overline{3}, \overline{4}, \overline{5}, \overline{6}, \overline{7}, \overline{8}\}$ and $\tau = \{\overline{1}, \overline{3}, \overline{4}, \overline{2}, \overline{5}, \overline{6}, \overline{7}, \overline{8}\}$,
 $t(\pi) = [(1)^2(2)^3]$ and $t_\pi(\tau) = [(1,1)^2(0,1)]$. Then

$$|O_\pi| = 8!/((1!)^2(2!)^32!3!) = 420$$

$$|O_{\pi < \tau}| = 8!/((1!)^2(2!)^32!(1!1!)^21!(0!1!)) = 2520.$$

Furthermore, let d be a chain in $PL(S)$ and $C[d]$ denote the set of each chain in $PL(S)$ containing d as a subchain. Then, it is shown that $G(C(PL(S)))_d$ is a permutation group on $C[d]$ and that $c \equiv c'$ for each c and c' in $C[d]$ if and only if there is α in $G(C(PL(S)))_d$ such that $\tilde{\alpha}(c) = c'$. Let O_c^d denote the cardinal congruence class (orbit) on $C[d]$ containing c . Then, since for each subchain d of a chain c $G(C(PL(S)))_c$ is a subgroup of $G(C(PL(S)))_d$, the following identity is obtained,

$$|O_c^d| = |G(C(PL(S)))_d| / |G(C(PL(S)))_c| = |O_c| / |O_d|.$$

For the previous example π and τ , we have

$$|O_{\pi < \tau}^\pi| = |O_{\pi < \tau}| / |O_\pi| = 6.$$

It is noted that the formulas for $l(c) \geq 3$ are open.

2. The Principles of Inclusion-Exclusion on Partially Ordered Sets The enumeration theorems under group action are very useful in counting non-isomorphic types of finite systems, but nevertheless they can not answer for the problems of enumerating reducible or irreducible types of finite systems. So, in order to deal with the problems, we have been developing, so called "the enumeration theorems under lattice action", with the Harrison's problem "determine the number of minimal(irreducible) machines with n states" for a background. We now present the fringe of the theory. The principle of inclusion-exclusion on semilattices[45, Theorem 1] is as follows.

Theorem (Inclusion-Exclusion on Semilattices). Let Ω be a nonempty set and (L, \vee) be a finite join-semilattice. Let $f: L \rightarrow \wp(\Omega)$ be a map satisfying $f(x) \cap f(y) \subseteq f(x \vee y)$ for each x and y in L . Then for any measure m on $\wp(\Omega)$ the following identity holds.

$$m\left(\bigcup_{x \in L} f(x)\right) = \sum_{c \in C} (-1)^{l(c)} m\left(\bigcap_{x \in c} f(x)\right),$$

where C is the set of chains in L and $\ell(c)$ denotes the length of a chain c . The theorem can be dualized.

The theorem was applied to a Boolean lattice and a product partition lattice [45, Proposition 1, Theorem 2]. Also, the theorem has been restated in terms of valuations on distributive lattices instead of measures on $\wp(\Omega)$ [46]. The three different proofs have been given, one in which the Rota's theorem from [60, Theorem 1] plays an important role, that is, the closure relation is used, and the others are elemental. Furthermore, recently, the theorem has been extended on partially ordered sets (posets) as follows [49].

Theorem (Inclusion-Exclusion on Posets). Let Ω be a nonempty set and P be a finite partially ordered set with a unique maximal element. Let $f: P \rightarrow \wp(\Omega)$ be a map satisfying $f(x) \cap f(y) \subseteq f(z)$ for each x and y in P and for some minimal element z in the subposet (of P) of all upper bounds of $\{x, y\}$. Then for any measure m on $\wp(\Omega)$ the following identity holds.

$$m\left(\bigcup_{x \in P} f(x)\right) = \sum_{c \in C} (-1)^{\ell(c)} m\left(\bigcap_{x \in c} f(x)\right),$$

where C is the set of all chains in P and $\ell(c)$ denotes the length of a chain c . Also the theorem can be dualized, which results in other three cases. Furthermore, the theorem can be restated in terms of valuations on distributive lattices instead of measures on $\wp(\Omega)$.

We next describe the principle of inclusion-exclusion on partition semilattices and an application of the principle to enumeration of the reducible or irreducible mapping systems. The most simple case is explained (see [48] for the full information). Let $\mathcal{F}(S)$ be the set of mappings from S into S . Let's recall the Galois correspondence $(\tilde{\mathcal{E}}, \tilde{\mathcal{E}})$ in Introduction. Then, regarding $\mathcal{F}(S)$ as Ω and $PL(S)$ as \mathcal{A} , we define a relation \mathcal{L} between $\mathcal{F}(S)$ and $PL(S)$ regarded the relation \mathcal{g} in the following way: for each f in $\mathcal{F}(S)$ and π in $PL(S)$ $f \mathcal{L} \pi$ if and only if for each s and t in S $s \pi t$ implies $f(s) \pi f(t)$, where for any x and y in S $x \pi y$ denotes that x and y are contained in a same block of

π . Therefore, a Galois correspondence

$$\wp(\mathcal{F}(S)) \xrightleftharpoons[\tilde{\mathcal{J}}]{\tilde{\mathcal{L}}} \wp(\text{PL}(S))$$

is induced by the relation \mathcal{L} and the inverse relation \mathcal{J} . We see easily that for each F in $\wp(\mathcal{F}(S))$ $\tilde{\mathcal{L}}(F)$ is a sublattice of $\text{PL}(S)$ called a reduction diagram of F and that for each π in $\wp(\text{PL}(S))$ $\tilde{\mathcal{J}}(\pi)$ is a subsemigroup of the semigroup $\mathcal{F}(S)$ under map composition. Note that the map $\tilde{\mathcal{L}}$ is introduced by making an abstract of a sublattice of partitions with substitution property on a sequential machine studied by Hartmanis and Stearns[37]. Now, the important subset ${}_{\tau}\mathcal{F}_{\pi}$ of $\mathcal{F}(S)$ is defined by

$${}_{\tau}\mathcal{F}_{\pi} = \{f \in \mathcal{F}(S) \mid \max(\tilde{\mathcal{L}}(f) \cap [0, \tau]) = \pi\},$$

where τ and π are any elements in $\text{PL}(S)$ and 0 is a unique minimal element in $\text{PL}(S)$. Here, $[0, \tau]$ is called a reduction domain of f and $\tilde{\mathcal{L}}(f) \cap [0, \tau]$ is called a reduction diagram of f relative to $[0, \tau]$. In other words, ${}_{\tau}\mathcal{F}_{\pi}$ is the set of f in $\mathcal{F}(S)$ which is at most reduced to $f: \pi \rightarrow \pi$ relative to $[0, \tau]$. Therefore, ${}_{\tau}\mathcal{F}_0$ is the set of all irreducible mappings relative to $[0, \tau]$. In the computation of $|{}_{\tau}\mathcal{F}_{\pi}|$, the following theorem[45, Corollary] is essentially used.

Theorem (Inclusion-Exclusion on Partition Semilattices).

Let $\tilde{\mathcal{J}}$ be the map $\text{PL}(S) \rightarrow \wp(\mathcal{F}(S))$ induced by the relation \mathcal{J} . Let L be any subsemilattice in $\text{PL}(S)$. Then for any measure m on $\wp(\mathcal{F}(S))$ the following identity holds.

$$m\left(\bigcup_{x \in L} \tilde{\mathcal{J}}(x)\right) = \sum_{c \in C} (-1)^{\ell(c)} m(\tilde{\mathcal{J}}(c)),$$

where C is the set of all chains in L .

The map $f: P \rightarrow \wp(\Omega)$ in the principle of inclusion-exclusion on posets is called a weak morphism on P . In an application of the principle, for a given poset P and map $f: P \rightarrow \wp(\Omega)$, it is of interest whether f is a weak morphism or not. It is shown that the map $\tilde{\mathcal{J}}$ is a weak morphism on L , and then the theorem follows from the principle. It is also shown that for each chain c in $\text{PL}(S)$ $|\tilde{\mathcal{J}}(c)|$ is characterized by the arithmetic operations. On the other hand, the set ${}_{\tau}\mathcal{F}_{\pi}$ is characterized by the map $\tilde{\mathcal{J}}$, and the theorem is applied to compute $|{}_{\tau}\mathcal{F}_{\pi}|$. Let's recall the

permutation group $G(PL(S))$ in Example 3. Let α be any element in $\mathcal{C}(S)$ and $\hat{\alpha}$ be a permutation on $\mathcal{F}(S)$ induced by

$$\hat{\alpha}(f) = \begin{pmatrix} \alpha(1) & \cdots & \alpha(n) \\ \alpha(f(1)) & \cdots & \alpha(f(n)) \end{pmatrix}$$

for each f in $\mathcal{F}(S)$. Then, it is shown that for each α in $\mathcal{C}(S)$

$$\tilde{\alpha}(\tilde{\mathcal{L}}(f)) = \tilde{\mathcal{L}}(\hat{\alpha}(f)) \text{ and } \hat{\alpha}(\tau\tilde{\mathcal{F}}\pi) = \tilde{\alpha}(\pi)\tilde{\mathcal{F}}\tilde{\alpha}(\tau).$$

The present formulation is naturally extended to a relation between the set $(\mathcal{F}(S,T))^P$ of mapping systems and the product partition lattice $PL(S) \times PL(T)$, and then the method of counting the number of the reducible or irreducible mapping systems is established by the author[48]. Furthermore, "the relation between $(\mathcal{F}(S))^P$ and $PL(S)$ " and "a relation between machines and $PL(S)$ " are identified by a bijection from the set of transition functions to $(\mathcal{F}(S))^P$. Therefore, the enumeration of reducible or irreducible machines is transformed into the enumeration of reducible or irreducible mapping systems, resulting in the solution for the Harrison's problem. In this enumeration, the cardinal congruence relation described in Example 3 is used to simplify figure-works in it. Furthermore, by considering a permutation group on the set of machines induced by the symmetric group on the set of states, the identity on the number of non-state-isomorphic irreducible machines with n states is also obtained. Finally, it is noted that for each τ in $PL(S)$ $G(PL(S))_\tau$ is a permutation group on ${}_\tau\tilde{\mathcal{F}}_0$ and that the general computation method for $|{}_\tau\tilde{\mathcal{F}}_0/G(PL(S))_\tau|$ is not still established.⁽¹⁾ The author thinks that this note is a very simple entrance problem to the third branch described in Introduction.

References

1. N. G. De Bruijn, Generalization of Pólya's fundamental theorem in enumerative combinatorial analysis, *Nederl. Akad. Wetensch. Proc. Ser. A* 62(1959), 59-69.
2. N. G. De Bruijn, Enumeration of mapping patterns, *J. Combinatorial Theory, Ser.A* 12(1972), 14-20.
3. W. Burnside, "Theory of Groups of Finite Order", 2nd ed., Theorem VII, p.191, Cambridge, MA, 1911.
4. R. L. Davis, The number of structures of finite relations, *Proc. Amer. Math. Soc.*, 4(1953), 486-495.

5. H. O. Foulkes, On Redfield's group reduction functions, *Canad. J. Math.*, 15(1963), 272-284. On the Redfield's range-correspondences, *Canad. J. Math.* 18(1966), 1060-1071.
6. F. Harary, Enumeration under group action: Unsolved graphical enumeration problems, IV, *J. Combinatorial Theory*, 8(1970), 1-11.
7. F. Harary and E. M. Palmer, The power group enumeration theorem, *J. Combinatorial Theory*, 1(1966), 157-173.
8. F. Harary and E. M. Palmer, The enumeration methods of Redfield, *Amer. J. Math.* 89(1967), 373-384.
9. F. Harary and E. M. Palmer, Enumeration of finite automata, *Information and Control*, 10(1967), 499-508.
10. F. Harary and E. M. Palmer, "Graphical Enumeration", Academic Press, 1973.
11. M. A. Harrison, A census of finite automata, *Canad. J. Math.* 17(1965), 100-113.
12. M. A. Harrison, Note on the number of finite algebras, *J. Combinatorial Theory*, 1(1966), 395-397.
13. D. E. Littlewood, "The Theory of Group Characters and Matrix representations of Groups", 2nd ed., Oxford, 1950.
14. E. M. Palmer and R. W. Robinson, The matrix group of two permutation groups, *Bull. Amer. Math. Soc.* 73(1967), 204-207.
15. G. Pólya, Kombinatorische Anzahlbestimmungen für Gruppen, Graphen und Chemische Verbindungen, *Acta Math.* 68(1937), 145-253.
16. R. C. Read, The enumeration of locally restricted graphs, I and II, *J. London Math. Soc.* 34(1959), 417-436; 35(1960), 344-351.
17. J. H. Redfield, The theory of group reduced distributions, *Amer. J. Math.* 49(1927), 433-455.
18. M. Aigner, "Kombinatorik I, Grundlagen und Zähltheorie", Springer-Verlag, Berlin, 1975.
19. M. Aigner (ed.), "Higher Combinatorics", Proc. of the NATO Advanced Study Institute held in Berlin (West Germany), Sept. 1-10, 1976, D. Reidel Pub., 1977.
20. G. E. Andrews, "The Theory of Partitions", *Encyclopedia of Mathematics and Its Applications* (ed. G.-C. Rota), Vol.2, Addison-Wesley, 1976.
21. E. A. Bender and J. R. Goldman, On the applications of Möbius inversion in combinatorial analysis, *The Amer. Math. Monthly*, Vol.82, No.8(1975), 789-803.
22. C. Berge, "Principles of Combinatorics", Academic Press, New York, 1971.
23. G. Berman and D. M. Jackson, "Forward Citations in Enumeration", Dept. of Combinatorics and Optimization, Univ. of Waterloo, Ontario, 1976.
24. G. Birkhoff, "Lattice Theory", Vol. XXV, *Amer. Math. Soc. Coll. Pub.*, 1967.
25. P. J. Cameron (ed.), "Combinatorial Surveys", Proc. of the Sixth British Combinatorial Conference, Academic Press, 1977.
26. L. Comtet, "Advanced Combinatorics", D. Reidel, Pub. Dordrecht, 1974.

27. H. H. Crapo, Möbius inversion in lattices, *Archiv der Math.* 19(1968), 595-607.
28. H. H. Crapo and G.-C. Rota, On the foundations of combinatorial theory II, "Combinatorial Geometries", M.I.T. Press, 1970.
29. P. Erdős, A. Rényi and Vera T. Sós (ed.), "Combinatorial Theory and Its Applications I, II, III", *Coll. Math. Soc. János Bolyai*, 4, 1970.
30. P. Erdős (ed.: J. Spencer), "The Art of Counting", M.I.T. Press, 1973.
31. S. Even, "Algorithmic Combinatorics", The Macmillan C., 1973.
32. L. Geissinger, Valuations on distributive lattices I, II, III, *Archiv der Math.*, 24(1973), 230-239, 337-345, 475-481.
33. C. Greene, On the Möbius algebra of a partially ordered set, *Advances in Math.*, 10(1973), 177-187.
34. C. Greene and D. J. Kleitman, The structure of Sperner k-families, *J. of Combinatorial Theory (A)*, 20(1976), 41-68.
35. D. Foata (ed.), "Combinatoire et Représentation du Groupe Symétrique", *Lecture Notes in Math.*, 579, Springer-Verlag, 1977.
36. M. Hall, "Combinatorial Theory", Chap.2, Blaisdell, Waltham, MA, 1967.
37. J. Hartmanis and R. E. Stearns, "Algebraic Structure Theory of Sequential Machines", Chap.2,3, Prentice Hall, Englewood Cliffs, NJ, 1966.
38. D. A. Holton (ed.), "Combinatorial Mathematics", *Proc. of the 2nd Australian Conference*, *Lecture Notes in Math.*, 403, Springer-Verlag, 1974.
39. D. E. Knuth, "The Art of Computer Programming I, II, III", Addison-Wesley, 1968, 1969, 1973.
40. C. L. Liu, "Introduction to Combinatorial Mathematics", Chap. 4,5, McGraw-Hill, New York, 1968.
41. C. L. Liu, Lattice functions, pair algebras, and finite state machines, *JACM*, Vol.16, No.3 (1969), 442-454.
42. C. L. Liu, "Elements of Discrete Mathematics", McGraw-Hill, 1977.
43. H. Narushima, Order preserving maps and classification of partially ordered sets, *Proc. Fac. Sci. Tokai Univ.* VI(1971), 27-45. (*Math. Reviews*, 45, #8586).
44. H. Narushima, Order maps and cardinal congruence classes on partitions, *Proc. Fac. Sci. Tokai Univ.* IX(1974), 1-13, (*Math. Reviews*, 50, #4325).
45. H. Narushima, Principle of inclusion-exclusion on semilattices, *J. of Combinatorial Theory (A)*, 17(1974), 196-203.
46. H. Narushima and H. Era, A variant of inclusion-exclusion on semilattices, *Discrete Math.*, 21(1978).
47. M. Kamii and H. Narushima, Further results on order maps, *Proc. Fac. Sci. Tokai Univ.* VIII(1973), 1-7. (*Math. Reviews*, 47, #8363).
48. H. Narushima, "Principle of Inclusion-Exclusion on Semilattices and Its Applications", Doctor's Thesis, Waseda Univ., Dec. 1977.
49. H. Narushima, Principle of inclusion-exclusion on partially ordered sets, to appear.
50. H. Narushima, On the number of chains in a Hasse diagram, to appear.

51. A. Nijenhuis and H. S. Wilf, "Combinatorial Algorithms", Academic Press, 1975.
52. A. Nozaki, Another proof of principle of inclusion-exclusion on semilattices, private paper.
53. O. Ore, Theory of equivalence relations, Duke Math. J., 9(1942), 573-627.
54. O. Ore, Galois connections, Trans. Amer. Math. Soc., 55(1944), 493-513.
55. O. Ore, "Theory of Graphs", Vol. XXXVIII, Amer. Math. Soc. Coll. Pub. Chap. 11, Providence, RI, 1962.
56. Randall Rustin (ed.), "Combinatorial Algorithms", Courant Computer Science Symposium 9, Algorithmics Press, INC., New York, 1973.
57. R. C. Read (ed.), "Graph Theory and Computing", Academic Press, 1972.
58. E. M. Reingold, J. Nievergelt and N. Deo, "Combinatorial Algorithms", Theory and Practice, Prentice-Hall, INC., 1977.
59. J. Riordan, "An Introduction to Combinatorial Analysis", Wiley, 1958.
60. G.-C. Rota, On the foundations of combinatorial theory I, Theory of Möbius functions, Z. Wahrscheinlichkeitstheorie und Verw. Gebiete 2(1964), 340-368.
61. G.-C. Rota, On the combinatorics of Euler characteristics, in: Studies in Pure Mathematics; (to R. Rado), Ed. Mirsky, (Academic Press, 1971), 221-233.
62. H. J. Ryser, "Combinatorial Mathematics", published by the Mathematical Association of America, distributed by Wiley, 1963.
63. D. Smith, Incidence functions as generalized arithmetics functions I, II, Duke Math. J., 34(1967), 617-634; 36(1969), 15-30.
64. L. Solomon, The Burnside algebra of a finite group, J. of Combinatorial Theory 2(1967), 603-615.
65. J. N. Srivastava (ed. with the cooperation of F. Harary, C. R. Rao, G.-C. Rota and S. S. Shrikhande), "A Survey of Combinatorial Theory", (to R. C. Bose), North-Holland Pub., 1973.
66. R. P. Stanley, Combinatorial reciprocity theorems, Advances in Math., 14(1974), 194-253.
67. W. T. Tutte (ed.), "Recent Progress in Combinatorics", Proc. of the Third Waterloo Conference on Combinatorics (May 1968), Academic Press, 1969.
68. H. S. Wilf, Hadamard determinants, Möbius functions, and the chromatic number of a graph, Bull. Amer. Math. Soc., 74, No.5 (1968), 960-964.
69. R. T. Yeh, Generalized pair algebra with applications to automata theory JACM, 15, No.2 (1968), 304-316.

(1)

Note that $\{\alpha \mid \alpha \text{ in } \mathcal{C}(S)\}$ is a permutation group on $\mathcal{F}(S)$, written $G(\mathcal{F}(S))$, and that we have $G(\text{PL}(S)) \cong \mathcal{C}(S) \cong G(\mathcal{F}(S))$.