

Comparison of Tests with
Same Bahadur Efficiency

By

Tapas K. Chandra
Presidency College

J. K. Ghosh
Indian Statistical Institute
and
University of Missouri

Abstract

This paper discusses the problem of discrimination between two test procedures whose Bahadur-efficiencies are equal. Since it is usually believed that equal Bahadur-efficiency is equivalent to equal Cochran-efficiency, we have discussed the problem from the point of view of Cochran as well; it is shown that at the level of deficiency, this equivalence does not hold true.

Comparison of Tests with
Same Bahadur-Efficiency

0. Introduction.

In [9], Hodges and Lehman studied the problem of discrimination between two statistical procedures which are, according to some criterion, equally "efficient"; deficiency is essentially a quantitative measure of this discrimination. In the same spirit, we have discussed here the problem of discrimination between two test procedures which have equal Bahadur-efficiency.

It is suggested by Bahadur ([1],[2]) that in many cases alternative test procedures might be compared on the basis of the associated limiting "attained levels." Following his suggestion we have introduced the notion of Bahadur-deficiency for two test procedures which are equally efficient from Bahadur's view-point. It appears that this approach of discrimination involves some difficulties; for example, the quantities involved are, in general, unlikely to be constants almost surely.

On the other hand, Cochran ([5]) measured the efficiency of a test procedure by the rate of convergence (to zero) of its size, when the power is held fixed against a specified alternative. It is well-known that the Cochran's approach to efficiency usually leads precisely to the same conclusions as Bahadur's approach does. Motivated by this fact we have introduced

in section 2 the notion of Cochran-deficiency (to be referred to as BCD for reasons explained in the next paragraph) and have shown by means of an example that at the level of deficiency, the above equivalence between Cochran's and Bahadur's view-points is no longer true. A necessary and sufficient condition for the existence of Cochran-deficiency is proved. In most cases this condition does not hold and so Cochran-deficiency will rarely exist. When appropriate asymptotic expansions of the significance levels are available, an "approximate" Cochran-deficiency is calculated as a compensation. Conditions under which the said expansions are valid are also investigated.

As Bahadur-deficiency will, in general, be random, one may like to go to the considerations of taking some sort of average of Bahadur-deficiency. But since the computations involved in such considerations appear to be quite difficult, we proceed along a somewhat different route in section 4, leading to a new interpretation of Cochran deficiency more in line with Bahadur's approach. In view of this interpretation we shall refer to Cochran deficiency as Bahadur-Cochran-deficiency (BCD).

1. Notations and Preliminaries.

Let (X, B) be a measurable space; let $\{P_\theta: \theta \in \mathbb{H}\}$ be a family of probability distributions on X .

Let $s = (x_1, x_2, \dots)$ be an infinite sequence of independent observations on x . Let $T \equiv \{T_n(s): n \geq 1\}$ be a real-valued statistic such that, for each n , $T_n(s)$ depends on s only through (x_1, \dots, x_n) . In the next paragraph, a brief synopsis of Cochran's efficiency is given; for details consult [5], [1] and [2].

Let \mathbb{H}_0 be a proper subset of \mathbb{H} . We are interested in testing $H_0: \theta \in \mathbb{H}_0$ against $H_1: \theta \in \mathbb{H} - \mathbb{H}_0$. For this purpose, we consider a test procedure which is based on a test statistic T and which regards the large values of $T_n(s)$ to be significant; i.e., the critical region W_n of the test procedure is of the form

$$(1.1) \quad W_n = \{s: T_n(s) \geq k_n\}.$$

Fix a θ in $\mathbb{H} - \mathbb{H}_0$ and a β such that $0 < \beta < 1$. Choose $\{k_n: n \geq 1\}$ such that

$$(1.2) \quad P_\theta(W_n) \rightarrow \beta$$

as $n \rightarrow \infty$. Note that k_n will depend on β as well as on θ . Let $\alpha_n(\beta) \equiv \alpha_n(\theta, \beta)$ be the resulting size of the test, i.e., let

$$(1.3) \quad \alpha_n(\theta, \beta) = \sup\{P_{\theta_0}(W_n): \theta_0 \in \mathbb{H}_0\}.$$

Cochran argued that the rate at which $\alpha_n(\theta, \beta)$ converges (to zero) is an indication of asymptotic efficiency of T against θ . Equivalently, one may proceed in the following way which is more suitable for our purpose: for each $\delta, 0 < \delta < 1$, let $M(\delta) \equiv M(\theta, \beta, \delta)$ be the least integer $m \geq 1$ such that $\alpha_n(\beta) < \delta$ for all $n \geq m$; otherwise let $M(\delta)$ be infinity. Henceforth, we shall assume that $\alpha_n(\beta) \rightarrow 0$ as $n \rightarrow \infty$, which ensures that $M(\delta)$ is finite for all δ . The Cochran-efficiency of the test procedure, when it exists, is equal to the limit of $[2 \log(1/\delta)/M(\delta)]$ as $\delta \rightarrow 0$.

Suppose now that Θ_1, Θ_2 are two distinct subsets of $\Theta - \Theta_0$ with $\Theta_2 \subset \Theta_1$; consider the testing problem $H_0: \theta \in \Theta_0$ vs. $H_1: \theta \in \Theta_1$ as well as $H_0: \theta \in \Theta_0$ vs. $H_2: \theta \in \Theta_2$. The first testing procedure is based on the test statistic $T_1(s) = \{T_{1n}(s): n \geq 1\}$, while the second is based on the test statistic $T_2(s) = \{T_{2n}(s): n \geq 1\}$. Fix a $\theta \in \Theta_2$ and also a β such that $0 < \beta < 1$. We want to discriminate between these two procedures when θ obtains. Define $M_i(\delta)$, $i = 1, 2$, in the usual way. Clearly, the limit of $[M_2(\delta)/M_1(\delta)]$ as $\delta \rightarrow 0$ gives the Cochran-efficiency of T_1 relative to T_2 when θ obtains. When this efficiency is 1, $[M_1(\delta) - M_2(\delta)]/M_1(\delta) \rightarrow 0$. In typical cases, however, $[M_1(\delta) - M_2(\delta)]$ remains bounded as $\delta \rightarrow 0$, and so for the purpose of a more subtle distinction, one may use the limit of $[M_1(\delta) - M_2(\delta)]$ as $\delta \rightarrow 0$, whenever this limit exists.

Definition 1.1. The lower (upper) Bahadur-Cochran-deficiency (BCD) at θ of the first testing procedure w.r.t the second is

$$\underline{D}_C(\theta, \beta) = \liminf_{\delta \rightarrow 0} [M_1(\delta) - M_2(\delta)]$$

$$(\overline{D}_C(\theta, \beta) = \limsup_{\delta \rightarrow 0} [M_1(\delta) - M_2(\delta)]) .$$

In case these two deficiencies are equal, we say that the BCD at θ exists and is equal to the common value.

Of course, $\underline{D}_C = \overline{D}_C = +\infty$ or $-\infty$ if $\lim[M_1(\delta)/M_2(\delta)]$ exists and $\neq 1$. The main use of deficiency is to discriminate tests for which $\lim[M_1(\delta)/M_2(\delta)]$ is 1. Note that although the relative Cochran-efficiency of two test procedures is usually free from β (see proposition 11, []), their relative BCD need not be so.

Let $\Phi(x)$ stand for the distribution function of the standard normal distribution and $\phi(x)$ stand for its density function. For $0 < \beta < 1$, we define z_β by requiring that $\Phi(z_\beta) = 1 - \beta$. The following results will be needed in the sequel.

Lemma 1.1. (See Chapter VII, [6]).

If x is positive,

$$1 - \Phi(x) = \frac{\phi(x)}{x} \left(1 - \frac{1}{x^2} + O(x^{-4})\right) .$$

Lemma 1.2. (See Chapter XV, [7]).

Let $\{X_i\}$ be i.i.d. random variables with the common

distribution $F(x)$ and with $E(X_1) = 0$, $E(X_1^2) = \sigma^2$. Let $F_n(x)$ be the normalized n -fold convolution of $F(x)$. If $F(x)$ is not a lattice distribution and if

$$m_3 \equiv E(X_1^3) \text{ is finite}$$

then one has

$$F_n(x) = \phi(x) + \frac{m_3}{6\sigma^3\sqrt{n}}(1-x^2)\phi(x) + o(n^{-1/2})$$

uniformly in x .

For the next result, let $\{Y_n\}$ be a sequence of i.i.d. random vectors with values in R^m ($m \geq 1$). Let f_1, \dots, f_k be real-valued Borel measurable functions on R^m . In the below, j stands for a positive integer ≥ 2 . Assume

$$(A_{1j}): \quad E|f_i(Y_1)|^j < +\infty \text{ for } 1 \leq i \leq k.$$

Write

$$Z_n = (f_1(Y_n), \dots, f_k(Y_n))$$

$$\mu = EZ_1, \quad V = \text{Cov } Z_1.$$

Assume

$$(A_2): \quad V \text{ is nonsingular.}$$

Let H be a real-valued function defined on some neighborhood N of μ . Assume

(A_{3,j}): H has bounded continuous derivatives on N of all orders up to and including j.

Let

$$\ell = (D_1 H, \dots, D_k H)(\mu)$$

where D_i denotes differentiation w.r.t. the i^{th} coordinate.

Assume

(A₄): $\ell \neq 0$.

Define H arbitrarily (but measurably) on all of R^k . We are interested in the asymptotic expansion of the distribution function of the statistic

$$W_n = \sqrt{n}(H(\bar{Z}) - H(\mu)),$$

where

$$\bar{Z} = \frac{1}{n} \sum_{i=1}^m Z_i.$$

Lemma 1.3. (See theorem 2, Bhattacharya and Ghosh, [4].)

Assume (A_{1j}), (A_{3j}) (for some integer $j \geq 2$), (A₂) and (A₄) hold. If in addition the distribution function of $(Z_1 - \mu)$ satisfies the Cramer's condition, namely,

$$\limsup_{\|t\| \rightarrow \infty} \left\| \int_{R^k} \exp\{i \langle t, x \rangle\} Q(dx) \right\| < 1,$$

then there exists polynomials q_r , $1 \leq r \leq j-2$, whose coefficients depend only on the cumulants of $(Z_1 - \mu)$ (of order j and less) and

the derivatives of H at μ (of order $j-1$ and less) such that

$$\sup_{u \in \mathbb{R}^1} \left| \text{Prob}(W_n \leq u) - \int_{-\infty}^{u/\sigma} (v) \left[1 + \sum_{r=1}^{j-2} n^{-r/2} q_r(v) \right] dv \right|$$

$$= o(n^{-(j-2)/2}),$$

where σ^2 is the variance of $\langle \ell, z_1 - \mu \rangle$.

In [4], the methods of computing the polynomials q_r are also given. For details, the reader is referred to this paper.

2. Cochran's Approach.

In this section, we discuss Cochran's approach to our problem. It is proved that for the existence of a finite BCD, the size functions of the test-procedures must be related in a very special way. Under appropriate asymptotic expansions of these size functions, the bounds of the upper and the lower BCD's are found out; also the methods of evaluating these deficiencies are discussed. All proofs are deferred to the appendix.

2.1. Existence of a finite BCD:

Recall that θ is a fixed element in $\mathbb{H}_2 \subset \mathbb{H} - \mathbb{H}_0$ and β is a fixed real number s.t. $0 < \beta < 1$.

Obviously, $\lim_{\delta \rightarrow 0} (M_1(\delta) - M_2(\delta))$, if it exists, will be either an integer or, one of the two infinities; also

$$(2.1.1) \quad \lim_{\delta \rightarrow 0} (M_1(\delta) - M_2(\delta)) \text{ exists and is equal to} \\ \text{an integer } d \equiv d(\theta, \beta),$$

iff for all sufficiently small δ , $M_1(\delta) - M_2(\delta)$ is identically equal to d . Thus at least in one case, e.g., when

$$(2.1.2) \quad \text{there exists an integer } d \text{ such that} \\ \alpha_{2,n} = \alpha_{1,n+d} \text{ for all sufficiently large } n,$$

(2.1.1) holds true and hence the BCD is d . Our main theorem 2.1.1 of this section states that the converse implication is

also true. For the proof of this theorem, we need the part (c) of the following proposition, which may be of independent interest.

Let $\{\alpha_n\}$ be a sequence of real numbers in $[0,1]$, and put $\alpha_n^* = \sup_{m \geq n} \alpha_m$ ($n \geq 1$). For each δ , $0 < \delta \leq 1$, let $M(\delta)$ be the smallest integer m for which $\alpha_m^* < \delta$. Considering the graph of the function $\delta \rightarrow M(\delta)$, one can easily convince oneself of the following proposition.

Proposition 2.1.1.

Let $0 \leq \alpha_n \leq 1$, $n \leq 1$ and $\alpha_n \rightarrow 0$. Then

(a) The function $\delta \rightarrow M(\delta)$ from $(0,1] \rightarrow I_+$ is a left continuous, decreasing step function. (I_+ is the set of natural numbers.)

(b) δ is a point of continuity of $M(\delta)$ iff

$$\alpha_{M(\delta)-1}^* > \delta \text{ or } M(\delta) = 1.$$

(c) The function $\delta \rightarrow M(\delta)$ determines α_m^* uniquely for all $m \geq 1$. More precisely, let $S = \{M(\delta) > 1: 0 < \delta \leq 1\}$ and let the elements of S arranged in ascending order be $1 < m_1 < m_2 < \dots$; let δ_i be such that $M(\delta_i) = m_i$ and δ_i is a point of discontinuity of $M(\delta)$. Then,

$$\alpha_n^* = \begin{cases} \delta_1 & \text{if } n < m_1 \\ \delta_i & \text{if } m_{i-1} < n < m_i \\ 0 & \text{if } S \text{ has a maximum element } m_k \\ & \text{and } n \geq m_k. \end{cases}$$

We now state Theorem 2.1.1.

Theorem 2.1.1.

Suppose that for each $i=1,2$, α_{in} is a decreasing function of n for all sufficiently large n . Then (2.1.1) and (2.1.2) are equivalent.

Remark 2.1.1. It should be noted that the main reason why the existence of a finite Cochran's deficiency imposes a strong condition like (2.1.2) on the functions α_{in} is the discrete nature of $M_i(\delta)$. Unfortunately, any attempt to make the sizes continuous by taking resort to mixtures, as done by Hodges and Lehman in [9] doesn't seem to work here.

2.2. The bounds for the upper and the lower Cochran's deficiencies; the notion of approximate Cochran-deficiency.

As we know from the analysis of the previous section that the BCD will exist rarely, we now turn to the problem of finding the bounds for the upper and the lower BCD's --- of course, under suitable assumptions on the significance levels of the two test procedures.

We assume that

(2.2.1) Each of $\{\alpha_{1,n}\}$ and $\{\alpha_{2,n}\}$ is a decreasing function of n for all sufficiently large n ;

(2.2.2) For each $i=1,2$, there exists an extension $\{\alpha_{i,x}\}$ of $\{\alpha_{i,n} : n \in I_+\}$ for non-integral values of x , $1 \leq x < \infty$, which is also a decreasing function of x ;

for the next assumption, define for each $n \geq 1$, a real number $m(n)$ such that $\alpha_{2,n} = \alpha_{1,m(n)}$. Assume that

(2.2.3) The limit of $(m(n)-n)$ exists (the limit may be infinite).

We let $d(\beta) \equiv d(\theta, \beta)$ stand for this limit; in the rest of this paper, $d(\beta)$ will have this meaning only (unless otherwise is stated). Note that $d(\beta)$ need not be an integer.

Definition 2.2.1. The approximate BCD of the first testing procedure w.r.t. the second one is the real number $d(\theta, \beta)$.

Note that the approximate BCD depends on the particular extensions $\{\alpha_{i,x}\}$, $i=1,2$, we are using. We shall, however, suppress this dependence.

The above assumptions (2.2.1)-(2.2.3) are valid under appropriate asymptotic expansions of $\{\alpha_{i,n}\}$. (See lemma 2.3.1.)

Note that, under (2.2.1), $M_1(\delta)$ is simply the first integer $m \geq 1$ such that $\alpha_{im} < \delta$. Consequently, if we let, for each δ ($0 < \delta \leq 1$), $M_{11}(\delta)$ and $M_{12}(\delta)$ satisfy

$$\alpha_{1, M_{11}(\delta)} = \alpha_{2, M_2(\delta) - 1} \quad , \quad \alpha_{1, M_{12}(\delta)} = \alpha_{2, M_2(\delta)} \quad ,$$

then

$$(2.2.4) \quad M_{11}(\delta) \leq M_1(\delta) \leq M_{12}(\delta) + 1 .$$

Because of the assumption (2.2.3), one has

$$\underline{D}_C(\theta, \beta) \geq d(\theta, \beta) - 1, \quad \bar{D}_C(\theta, \beta) \geq d(\theta, \beta) + 1;$$

or equivalently,

$$(2.2.5) \quad -[-d(\theta, \beta)] - 1 \leq \underline{D}_C(\theta, \beta) \leq \bar{D}_C(\theta, \beta) \leq [d(\theta, \beta)] + 1,$$

where $[t]$ stands for the greatest integer $\leq t$.

Remark 2.2.1. It follows that the BCD is $+\infty$ or $-\infty$ according as the approximate BCD is $+\infty$ or $-\infty$.

Remark 2.2.2. If the BCD exists (it may be infinite) then it must be equal to d and so d will be an integer; d may, however, be an integer even if the BCD doesn't exist. (See examples 2.4.1 and 2.4.2.)

If d is non-integral (and finite), say, $d = m + t$ ($0 < t < 1$, m is an integer), then the bounds given by (2.2.5) are sharp enough to conclude that

$$\underline{D}_C(\theta, \beta) = m \quad \text{and} \quad \bar{D}_C(\theta, \beta) = m + 1.$$

In this case, it is possible to give the following interpretation of \underline{D}_C and \bar{D}_C : For each δ ($0 < \delta \leq 1$), let $M_{13}(\delta)$ be the smallest integer k such that $\alpha_{2, M_2(\delta)} \geq \alpha_{1, k}$. Assumptions (2.2.2) and (2.2.3) will then imply that

$$\lim_{\delta \rightarrow 0} (M_{13}(\delta) - M_2(\delta)) = m+1 = \bar{D}_C$$

Similarly for \underline{D}_C .

Remark 2.2.3. If d is an integer, the bounds given by (2.2.5) cannot immediately be used to find out the values of \underline{D}_C and \bar{D}_C . However, when $m(n) - n$ always lies strictly on one side of d , one of these bounds can be improved upon as described in the next paragraph.

An improvement of the upper bound of $M_1(\delta)$ given by (2.2.4) is $M_1(\delta) \leq [M_{12}(\delta)] + 1$; also from the assumption (2.2.3), $[M_{12}(\delta)] = M_2(\delta) + [d]$ or $M_2(\delta) + [d] - 1$ for all sufficiently small δ .

Suppose now that d is an integer and that $m(n) - n - d > 0$ for all sufficiently large n . Then $[M_{12}(\delta)] \leq M_2(\delta) + d - 1$ and hence $\bar{D}_C \leq d$. Clearly the BCD cannot exist. Thus one has from (2.2.5)

$$\underline{D}_C = d-1, \quad \bar{D}_C = d.$$

Similarly when d is an integer and $m(n) - n - d < 0$ for all sufficiently large n , one has $\underline{D}_C = d$, $\bar{D}_C = d+1$.

When asymptotic expansions of the size functions $\{\alpha_{in}\}$, $i=1,2$, are available, it is possible to determine the exact rate of convergence (to zero) of $m(n) - n - d$ and hence to verify whether for all sufficiently large n , $m(n) - n - d$ is positive or negative. (See the section 2.4(A) for details.)

2.3. Determination of the approximate BCD

We assume throughout this section that the significance levels $\{\alpha_{in}\}$, $i=1,2$, of the test procedures admit of the following asymptotic expansions:

$$(2.3.1) \quad \log \alpha_{in}(\theta, \beta) = -na_i(\theta, \beta) + \sqrt{n} b_i(\theta, \beta) + c_i(\theta, \beta) \log n + d_i(\theta, \beta) + o_i(1) \quad (i=1,2)$$

where $a_i(\theta, \beta) > 0$, $i=1,2$.

In typical cases, $a_i(\theta, \beta)$ will be free from β --- this will be the case if the Bahadur-slopes of T_1 and T_2 exist; for a precise result, see Theorem 2 of Ragharachari [11]. Note that $M_1(\delta) | M_2(\delta) \rightarrow 1$ iff $a_1(\theta, \beta) = a_2(\theta, \beta) = a(\theta, \beta)$, say. Henceforth we shall assume that this is the case. For convenience, we shall suppress the dependence on θ, β of the quantities a, b_i, c_i, d_i .

The following lemma connects the two sets of assumptions made in the present and the previous sections.

Lemma 2.3.1. Assume that (2.3.1) holds. Then (2.2.1), (2.2.2) and (2.2.3) are valid. In fact, (2.2.2) holds in the following strong sense: There exist extensions $\{\alpha_{ix}\}$, $i=1,2$, which satisfy (2.3.1) for non-integral values of x as well.

We shall work with such extensions $\{\alpha_{ix}\}$ only. The next theorem gives the possible values of the approximate BCD $d(\theta, \beta)$.

Theorem 2.3.1.

Let $\{\alpha_{i,n}\}$, $i=1,2$, satisfy (2.3.1). Then one has

- (a) if $b_1 = b_2$ and $c_1 = c_2$, then $d = (d_1 - d_2)/a$;
- (b) if $b_1 \neq b_2$, then d is $+\infty$ or $-\infty$, according as $b_1 > b_2$ or $b_1 < b_2$;
- (c) if $b_1 = b_2$ and $c_1 \neq c_2$, then d is $+\infty$ or $-\infty$, according as $c_1 > c_2$ or $c_1 < c_2$.

Remark 2.3.1. In general, the value of the approximate BCD will depend both on θ and on β . However, in the examples discussed here, it will be free from β . In these examples, the situation (a) occurs and $d_1(\theta, \beta) - d_2(\theta, \beta)$ is free from both θ and β ; as observed earlier, $a(\theta, \beta)$ will usually be free from β .

2.4. Examples.

In this section, we discuss two examples. As these examples will clarify different points of the next section 2.5, we prefer to go through the details. For simplicity, we determine the constants $k_n(\theta, \beta)$ (cf. (1.2)) such that

$$P_\theta(T_n > k_n(\theta, \beta)) \equiv \beta.$$

(A) In many examples, the size functions $\{\alpha_{in} : n \in I_+\}$, $i=1,2$ admit of extensions $\{\alpha_{ix} : x \in [1, \infty)\}$, $i=1,2$ which are decreasing functions of x and moreover, for large values of x , the following asymptotic expansions of these extensions are valid:

$$(2.4.1) \quad \log \alpha_{ix}(\theta, \beta) = -xa(\theta) + \sqrt{x} b(\theta, \beta) - \frac{1}{2} \log x + d_i(\theta, \beta) \\ + e(\theta, \beta)x^{-\frac{1}{2}} + o_i(x^{-\frac{1}{2}}) \quad (i=1,2)$$

where $a(\theta) > 0$, $b(\theta, \beta) = 0$ iff $\beta = \frac{1}{2}$ and finally $d_1(\theta, \beta) - d_2(\theta, \beta)$ is always nonzero. Moreover, when $\beta = \frac{1}{2}$.

$$(2.4.2) \quad \log \alpha_{ix}(\theta, \frac{1}{2}) = -xa(\theta) - \frac{1}{2} \log x + d_i(\theta, \frac{1}{2}) + e(\theta, \frac{1}{2})x^{-\frac{1}{2}} \\ + f(\theta)x^{-1} + o_i(x^{-1}) \quad (i=1,2)$$

From Theorem 2.3.1, the approximate BCD is $d(\theta, \beta) = (d_1(\theta, \beta) - d_2(\theta, \beta))/a(\theta)$. We want to compute $\underline{D}_C(\theta, \beta)$ and $\overline{D}_C(\theta, \beta)$, making use of the analysis made in the remark 2.2.3. For this let us first observe the following result about the rate of convergence to zero of $t_n \equiv m(n) - n - d(\theta, \beta)$.

Lemma 2.4.1. Assume that the size functions $\{\alpha_{i,n}\}$, $i=1,2$ satisfy (2.4.1) and (2.4.2). Then $\sqrt{n} t_n \rightarrow (d(\theta, \beta) \cdot b(\theta, \beta))/2a(\theta)$. If $\beta = \frac{1}{2}$, $nt_n \rightarrow d(\theta, \frac{1}{2})/2a(\theta)$.

We assume below, without loss of generality, that $d(\theta, \beta)$ is positive for all β . Three cases may arise.

Case I. Let β be such that $b(\theta, \beta) > 0$. In this case, $\sqrt{n} t_n$ converges to some positive number. Consequently, $m(n) - n - d(\theta, \beta)$ is positive for all sufficiently large n . It then follows from the remark 2.2.3 that the BCD doesn't exist and $\overline{D}_C(\theta, \beta) = d(\theta, \beta)$ while $\underline{D}_C(\theta, \beta) = d(\theta, \beta) - 1$.

Case II. Let β be such that $b(\theta, \beta) < 0$. In this case, $m(n) - n - d(\theta, \beta)$ is negative for all sufficiently large n . So the BCD doesn't exist and $\bar{D}_C(\theta, \beta) = d(\theta, \beta) + 1$ while $\underline{D}_C(\theta, \beta) = d(\theta, \beta)$.

Case III. Let β be such that $b(\theta, \beta) = 0$, i.e., let $\beta = \frac{1}{2}$. In this case nt_n converges to some positive number. So the conclusions of the Case I are valid.

(B) Example 1. (The Normal Distribution)

Let \mathbb{H} be the real line $(-\infty, +\infty)$, $\mathbb{H}_0 = \{0\}$, and $\mathbb{H}_1 = \mathbb{H} - \{0\}$, $\mathbb{H}_2 = (0, \infty)$. For $\theta \in \mathbb{H}$, let P_θ stand for the normal distribution with mean θ and variance 1. Fix a θ in \mathbb{H}_2 .

For the testing problem H_0 : the population mean is zero against the alternative that it is nonzero, the critical region of the most powerful unbiased invariant test is given by

$\{|\sqrt{n} \bar{X}_n| > k_{1,n}\}$ where $\bar{X}_n = n^{-1} \sum_{i=1}^n X_i$ and $k_{1,n}$ is such that

$\beta = 1 - \Phi(k_{1n} - \sqrt{n} \theta) + \Phi(-k_{1n} - \sqrt{n} \theta)$. Its power at θ is β

and its size is $\alpha_{1n}(\theta, \beta) = 2(1 - \Phi(k_{1n}))$. Bahadur's (as

well as the Cochran's) efficiency of $T_1 = \{|\sqrt{n} \bar{X}_n| : n \geq 1\}$ at θ is θ^2 .

For the testing problem H_0 : the population mean is zero against the alternative that it is positive, the critical region of the most powerful test is given by $\{\sqrt{n} \bar{X}_n > k_{2,n}\}$ where k_{2n}

$$(2.4.5) \quad \exp(x\theta^2) (\log \alpha_{1x} - \log \alpha_{2x} - \log 2) \rightarrow 0 .$$

To this end, first observe that k_{2x} serves as a good asymptotic approximation to k_{1x} ; (2.4.4) implies that $(k_{1x} - \sqrt{x}\theta) \rightarrow z_\beta$, i.e., $(k_{1x} - k_{2x}) \rightarrow 0$. In fact, we have

$$(2.4.6) \quad \lim_{x \rightarrow \infty} \exp(2x\theta^2) (k_{1x} - k_{2x}) = 0 ;$$

for one has, from the definition of k_{2x} and (2.4.4), $1 - \Phi(k_{1x} + \sqrt{x}\theta) = (k_{1x} - k_{2x})\phi(\xi(x))$ for some $\xi(x)$ satisfying $z_\beta < \xi(x) < k_{1x} - \sqrt{x}\theta$. By what has been proved, $\xi(x) \rightarrow z_\beta$ as $x \rightarrow \infty$. Thus

$$\begin{aligned} \exp(2x\theta^2) (k_{1x} - k_{2x}) &\leq \frac{\exp(2x\theta^2)}{\phi(\xi(x))} \cdot (1 - \Phi(k_{2x} + \sqrt{x}\theta)) \\ &= o(x^{-\frac{1}{2}}) , \text{ by Lemma 1.1.} \end{aligned}$$

This completes the proof of (2.4.6). Now the l.h.s. of (2.4.5) is $\exp(x\theta^2) (k_{2x} - k_{1x})\phi(\xi_1(x)) / (1 - \Phi(\xi_1(x)))$ for some $\xi_1(x)$ satisfying $k_{2x} < \xi_1(x) < k_{1x}$; in particular, $\xi_1(x) = \sqrt{x}\theta(1 + o(1))$. So (2.4.6) and Lemma 1.1 now complete the proof of (2.4.5).

Example 2. (The Student's Distribution)

Here $\mathbb{H} = \{(\mu, \sigma) : -\infty < \mu < \infty, 0 < \sigma < \infty\}$, $\mathbb{H}_0 = \{0\} \times (0, \infty)$, $\mathbb{H}_1 = \mathbb{H} - \mathbb{H}_0$, $\mathbb{H}_2 = (0, \infty) \times (0, \infty)$. For $\theta = (\mu, \sigma)$ in \mathbb{H} , let P_θ stand for the normal distribution with mean μ and variance σ^2 .

is such that $\beta = 1 - \Phi(k_{2n} - \sqrt{n} \theta)$, i.e., $k_{2n} = \sqrt{n} \theta + z_\beta$. Its power at θ is β and its size is $\alpha_{2n}(\theta, \beta) = 1 - \Phi(k_{2n})$. The Bahadur (as well as the Cochran) efficiency at θ of $T_2 = \{\sqrt{n} \bar{X}_n : n \geq 1\}$ is θ^2 .

Thus the two test procedures are equally efficient. At the level of deficiency, however, their performances are different. In this example, the situation described in (A) above holds. In fact, for non-integral values of x , one defines.

$$(2.4.3) \quad \alpha_{1x} = 2(1 - \Phi(k_{1x})) , \quad \alpha_{2x} = 1 - \Phi(k_{2x})$$

where k_{1x} is determined from

$$(2.4.4) \quad \beta = 1 - \Phi(k_{1x} - \sqrt{x} \theta) + \Phi(-k_{1x} - \sqrt{x} \theta)$$

and $k_{2x} = \sqrt{x} \theta + z_\beta$. We claim that (2.4.1) and (2.4.2) hold with $a(\theta) = \theta^2/2$, $b(\theta, \beta) = -z_\beta \theta$, $d_1(\theta, \beta) = -(z_\beta^2 + \log(2\pi\theta^2))/2 + \log 2$, $d_2(\theta, \beta) = -(z_\beta^2 + \log(2\pi\theta^2))/2$, $e(\theta, \beta) = -z_\beta \theta^{-1}$ and $f(\theta) = \theta^{-2}$. From (A), we can therefore conclude that the BCD doesn't exist and that the approximate BCD = $2 \log 2/\theta^2$. Also, $\bar{D}_C(\theta, \beta) = 2 \log 2/\theta^2$ if $\beta \geq \frac{1}{2}$; $= 2 \log 2/\theta^2 + 1$ if $\beta < \frac{1}{2}$ while $\underline{D}_C(\theta, \beta) = 2 \log 2/\theta^2 - 1$ if $\beta \geq \frac{1}{2}$; $= 2 \log 2/\theta^2$ if $\beta < \frac{1}{2}$. Thus the approximate BCD doesn't depend on β . The upper and the lower BCD do depend on β , but in a very weak sense.

We now proceed to the proof of our claim. That $\{\alpha_{2x}\}$ satisfies (2.4.1) and (2.4.2) is easy --- one has to use Lemma 1.1. To show the same for $\{\alpha_{1x}\}$, it is sufficient to prove that as $x \rightarrow \infty$

Fix a $\theta_1 = (\mu_1, \sigma_1)$ in Θ_2 ; put $\mu = \mu_1 \sigma_1^{-1}$, $\theta = (\mu, 1)$ and $\theta_0 = (0, 1)$. Put $T_{2n} = \sqrt{n} \bar{X}_n / s_n$ and $T_{1n} = |T_{2n}|$ where $n\sigma_1^2 = \sum_1^n (X_i - \bar{X}_n)^2$. Note that under θ_0 , $\sqrt{n/(n-1)}$, T_{2n} is distributed as a Student's t-variable with $(n-1)$ d.f.

For the testing problem H_0 : the population mean is zero against the alternative that it is non-zero (the s.d. being unknown), the "best" test is based on the critical region $\{T_{1n} > k_{1n}\}$ where k_{1n} is such that $\beta = P_{\theta_1}(T_{1n} > k_{1n})$. Its power at θ_1 then is β and its size is $\alpha_{1n}(\theta_1, \beta) = 2P_{\theta_0}(T_{2n} > k_{1n})$. The Bahadur (as well as the Cochran) efficiency at θ_1 of T_1 is $\log(1 + \mu^2)$.

For the testing problem H_0 : the population mean is zero against the alternative that it is positive (the s.d. being unknown), the "best" test is based on the critical region $\{T_{2n} > k_{2n}\}$ where k_{2n} is such that $\beta = P_{\theta_1}(T_{2n} > k_{2n})$. Its power at θ_1 is β and its size is $\alpha_{2n}(\theta_1, \beta) = P_{\theta_0}(T_{2n} > k_{2n})$. The Bahadur (as well as the Cochran) efficiency at θ_1 of T_2 is $\log(1 + \mu^2)$.

Here too the two test procedures are equally efficient, though their deficiency is not zero. We shall show that $\{\alpha_{in} : n \in I_+\}$, $i = 1, 2$, satisfy (2.4.1) and (2.4.2) with $a(\theta_1) = \frac{1}{2} \log(1 + \mu^2)$, $b(\theta_1, \beta) = z_\beta \cdot \mu(1 + \frac{1}{2} \mu^2)^{\frac{1}{2}}(1 + \mu^2)^{-1}$ and $d(\theta_1, \beta) - d(\theta_1, \beta) = \log 2$. In this example, we cannot define

the extensions $\{\alpha_{ix} : x \in [1, \infty)\}$, $i=1,2$, as we did in the example 1. One has to take linear averages of $\log \alpha_{in}$ and $\log \alpha_{i(n+1)}$; for details see the proof of Lemma 2.3.1.

From (A) we therefore conclude that the BCD doesn't exist and that the approximate BCD is $d(\theta_1, \beta) = 2 \log 2 / \log(1+\mu^2)$. Also, $\bar{D}_C(\theta_1, \beta) = d(\theta_1, \beta)$ if $\beta \geq \frac{1}{2}$; $= d(\theta_1, \beta) + 1$ if $\beta < \frac{1}{2}$ while $\underline{D}_C(\theta_1, \beta) = d(\theta_1, \beta) - 1$ if $\beta \geq \frac{1}{2}$; $= d(\theta_1, \beta)$ if $\beta < \frac{1}{2}$. Thus in this example too, the approximate BCD is free from β . The upper and the lower BCD do depend on β , but in a very weak sense.

The proofs of the different facts mentioned above run essentially along the same line as that of the example 1; however, some of the steps have to be justified in different ways. Lemma 1.1, e.g., is to be replaced by the following one, the proof of which depends on integration by parts.

Lemma 2.4.2. Let $n \geq 5$ and $\alpha > 0$. Put

$$I_n(\alpha) = \int_{\alpha}^{\infty} (1+x^2)^{-n/2} dx \quad \text{and} \quad \gamma_n(\alpha) = (n-2)^{-1} \alpha^{-1} (1+\alpha^2)^{-n-2/2}.$$

Then one has

$$\gamma_n(\alpha) [1 - (n-4)^{-1} (\alpha^{-2} + 1)] \leq I_n(\alpha) \leq \gamma_n(\alpha) [1 - 3(n-4)^{-1} (n-6)^{-1} (\alpha^{-2} + 1)].$$

Using this lemma, one can verify

Lemma 2.4.3. Let p_n stand for $P_{\theta_0}(T_{2n} > a\sqrt{n} + b + cn^{-\frac{1}{2}} + dn^{-1})$,

$a > 0$. Then

$$\log p_n = -\frac{n}{2} \log(1+a^2) - \sqrt{n} ab(1+a^2)^{-1} - \frac{1}{2} \log n \\ + K_1(a,b,c) + K_2(a,b,c,d)n^{-\frac{1}{2}} + o(n^{-\frac{1}{2}}),$$

where K_1 and K_2 are constants (free from n) which depend on a, b, c , etc. as indicated above.

We also need to use the following result, which is a direct consequence of Lemma 1.3.

Lemma 2.4.4. Let $W_n = T_{2n}^* - \sqrt{n}\mu$. There exist two polynomials p_1 and p_2 whose coefficients are free from n such that

$$\sup_{u \in \mathbb{R}^1} \sqrt{n} \left| P_{\theta_1}(W_n > u\tilde{\sigma}) - \int_u^{\infty} \phi(t) dt - \left(\frac{p_1(u)}{\sqrt{n}} + \frac{p_2(u)}{n} \right) \phi(u) \right| \rightarrow 0$$

$$\text{where } \tilde{\sigma} = (1 + \frac{1}{2}\mu^2)^{\frac{1}{2}}.$$

Define $G_n(u) = 1 - \phi(u) + (p_1(u)n^{-\frac{1}{2}} + p_2(u)n^{-1})\phi(u)$;
determine the constants (free from n) d, d_1, d_2 such that
 $G_n(d + d_1n^{-\frac{1}{2}} + d_2n^{-1}) = \beta + o(n^{-1})$; one may verify that $d = z_\beta$
(we do not need the exact values of d_1 and d_2). Put
 $k'_{2n} = \sqrt{n}\mu + (d + d_1n^{-\frac{1}{2}} + d_2n^{-1})\tilde{\sigma}$. Then $k_{2n} - k'_{2n} = o(n^{-1})$ ---
for a justification imitate the proof of the Lemma 2.5.1. What
we have achieved so far is simply an approximation k'_{2n} of k_{2n}

which guarantees that $\log P_{\theta_0}(T_{2n} > k_{2n}) - \log P_{\theta_0}(T_{2n} > k'_{2n}) = o(n^{-\frac{1}{2}})$;

to verify this, one uses lemma 2.4.2. An application of lemma 2.4.3 then shows that $\{\alpha_{2n}\}$ satisfies (2.4.1) with the said values of $a(\theta)$ and $b(\theta, \beta)$. Consider now the case of $\{\alpha_{1n}\}$. Here $\beta = P_{\theta_1}(W_n > k_{1n} - \sqrt{n} \mu) + P_{\theta_1}(W_n < -k_{1n} - \sqrt{n} \mu)$. Because of Lemma 2.4.4, $P_{\theta_1}(W_n < -k_{1n} - \sqrt{n} \mu)$ is $o(n^{-1})$. (In fact it can be shown that it is $o(n^{-j})$ for each positive integer j .) This implies that $(k_{1n} - k'_{2n}) = o(n^{-1})$; and so

$\log P_{\theta_0}(T_{2n} > k_{1n}) - \log P_{\theta_0}(T_{2n} > k'_{2n}) = o(n^{-\frac{1}{2}})$. Thus

$\log \alpha_{1n} = \log 2 + \log \alpha_{2n} + o(n^{-\frac{1}{2}})$. This completes the proof of the fact that $\{\alpha_{1n}\}$ satisfies (2.4.1) and that

$d_1(\theta, \beta) - d_2(\theta, \beta) = \log 2$. The proof of the fact that $\{\alpha_{1n}\}$ and $\{\alpha_{2n}\}$ satisfy (2.4.2) should now be clear.

Remark 2.4.2. Suppose we replace (1.2) by $\lim P_{\theta}(W_{in}) = \beta$ and put $R_{in}(\theta, \beta) = P_{\theta}(W_{in}) - \beta$, ($i=1,2$); then in the above examples it can be shown that (a) if $R_{in}(\theta, \beta) = o(n^{-\frac{1}{2}})$ then the value of the approximate BCD remains unchanged; (b) if $R_{in}(\theta, \beta) = o(n^{-1})$ and $R_{in}(\theta, \frac{1}{2}) = o(n^{-3/2})$, the BCD doesn't exist and the values of the upper and the lower BCD remain unchanged.

Remark 2.4.3. In example 2, it follows from Lemma 2.4.3. that the Bahadur slope at (μ, σ) of T_2 is $\log(1 + \mu^2 \sigma^{-2})$. Our derivation is simpler than any other available in the literature; compare, e.g., with the section 5 of [2] and [10].

2.5. On the validity of (2.3.1).

Here we shall find the conditions under which the asymptotic expansion of the form (2.3.1) of the significance level of a test procedure is valid. We motivate ourselves by considering a test procedure in which the critical region consists of the large values of the sum of some sequence of i.i.d. random variables. We have the following general result in this direction; compare with [3].

Let $\{Y_n: n \geq 1\}$ be a sequence of i.i.d. r.v.s. with the m.g.f. $M(t)$. Put $T_n = n^{-\frac{1}{2}} \sum_1^n Y_i$. Let μ be a constant ($\neq 0$) and $\{q_n\}$ be a bounded sequence of real numbers. Define $p_n = \text{Prob}(T_n > \sqrt{n} \mu + q_n)$. Assume that the distribution of X_1 and μ satisfy the following conditions: the distribution of X_1 is non-lattice; if T stands for $\{t: M(t) \text{ is finite}\}$, then T is a nondegenerate interval; there exists a positive τ in the interior of T such that $\exp(-\mu\tau)M(\tau) = \inf\{\exp(-\mu t)M(t): t \text{ in } T\} = \rho$ say.

Proposition 2.5.1. Assume the above set-up. Then

$$(2.5.1) \quad \log p_n = n \log \rho - \sqrt{n} \alpha q_n - \frac{1}{2} \log n - (a + q_n^2/2) + o(1)$$

where a, α, ρ are constants (free from n); $a > 0$, $\alpha > 0$, $0 < \rho < 1$.

Remark 2.5.1. If we assume that the distribution of X_1 satisfies Cramer's condition, we can get an asymptotic expansion of $\log p_n$ similar to the one given in the theorem 2 of [3].

Consider now the set-up of section 1. Our main interest is to find an asymptotic expansion of $P_\theta(T_n > k_n)$ where k_n is to be determined from the condition (1.2);

Assume that the distribution of $\{T_n\}$ under P_θ and $\{P_{\theta_0} : \theta_0 \in \mathbb{H}_0\}$ satisfies the following conditions:

(2.5.2) There exist constants (free from n) $\mu \equiv \mu(\theta)$ and $\tilde{\sigma} \equiv \tilde{\sigma}(\theta) > 0$ and a polynomial q_θ such that if we let $F_n(x) = P_\theta(T_n - \sqrt{n}\mu \leq \tilde{\sigma}x)$, then $F_n(x) = \Phi(x) + n^{-1/2}q_\theta(x)\phi(x) + o(n^{-1/2})$, uniformly in x .

(2.5.3) Whenever μ is a real number and $\{q_n\}$ is a bounded sequence of real numbers, (2.5.1) holds good with $P_n = \sup\{P_{\theta_0}(T_n > \sqrt{n}\mu + q_n) : \theta_0 \text{ in } \mathbb{H}_0\}$.

Lemma 2.5.1. Assume (2.5.2). Then there exists a constant $d \equiv d(\theta, \beta)$ such that $k_n = \sqrt{n}\mu + z_\beta \tilde{\sigma} + n^{-1/2}d\tilde{\sigma} + o(n^{-1/2})$.

Theorem 2.5.1. Assume (2.5.2) and (2.5.3). Then

$$\log \alpha_n = n \log \rho - \sqrt{n} \alpha z_\beta \tilde{\sigma} - \frac{1}{2} \log n + (a - \alpha \tilde{\sigma} d - \frac{1}{2} z_\beta^2 \tilde{\sigma}^2) + o(1) .$$

Remark 2.5.2. It is well known that $\{n^{-1/2} \sum_{i=1}^n Y_i: n \geq 1\}$ satisfies (2.5.2) where $\{Y_i: i \geq 1\}$ is a sequence of i.i.d.r.v.s. with a finite third moment. The main result of Bhattacharya and Ghosh (1976) indicates that (2.5.2) is satisfied for a large collection of statistics.

3. Bahadur's and Other Approaches.

In this section, we shall consider other approaches to deficiency. It is shown by means of an example that Bahadur-deficiency need not exist even though the Bahadur-Cochran-deficiency exists. A new interpretation of the latter is suggested in section 3.2.

3.1. Bahadur's Approach.

Assume the set-up of section 1. For each real t , let $F_{in}(t) = \sup\{P_{\theta_0}(T_{1n} > t) : \theta_0 \in \Theta_0\}$ and define $L_{in}(s) = F_{in}(T_{in}(s))$. For each δ ($0 < \delta \leq 1$) and for each s , let $N_i(\delta, s) \equiv N_i$ be the least integer $m \geq 1$ such that $L_{in}(s) < \delta$ for all $n \geq m$; otherwise let $N_i(\delta, s)$ be infinity.

Definition 3.1.1. The random lower (upper) Bahadur-deficiency at of the first testing procedure w.r.t. the second is

$$\begin{aligned} \underline{D}_B(\theta; \beta) &= (\text{a.s. } P_\theta) \liminf (N_1(\delta, s) - N_2(\delta, s)) \\ \left(\overline{D}_B(\theta; \beta) &= (\text{a.s. } P_s) \limsup (N_1(\delta, s) - N_2(\delta, s)) \right) . \end{aligned}$$

In case the above two deficiencies are equal, we say that the Bahadur-deficiency exists and is equal to the common value.

As in the case of Cochran-deficiency, the main use of studying these random deficiencies is to discriminate tests with the same Bahadur-efficiency, i.e., tests for which the (a.s. P_θ) limit of $(N_2(\delta, s)/N_1(\delta, s))$ is 1.

In this approach, the main source of difficulty is that the quantities $\sup\{L_{in}(s) : n \geq m\}$, $m \geq 1$ is difficult to expand --- any possible expansion would seem to depend on the particular sample sequence considered.

Example 3 (The Uniform Distribution):

Let θ be such that $0 < \theta < 1$; let $f_1(x)$ and $f_2(x)$ be respectively the densities of the uniform distributions over $[0, \theta]$ and $[0, 1]$. Consider the problem of testing

$H_0: f = f_2$ vs. $H_1: f = f_1$ on the basis of the following two statistics, $T_{1n} = 1 - x_{(n)}$, $T_{2n} = 1 - y_{(n)}$ where

$x_{(n)} = \max(x_4, x_6, \dots, x_{2n})$ and $y_{(n)} = \max(x_1, x_3, \dots, x_{2n-1})$, $(n \geq 2)$. We choose the constants $k_n(\theta, \beta)$ such that

$P_{H_1}(T_n > k_n(\theta, \beta)) \equiv \beta$ (cf. (1.2)). Then $\alpha_{1n} = \beta\theta^{n-1}$ while $\alpha_{2n} = \beta\theta^n$ so that the BCD exists and is 1 for all β .

We are going to show that the (a.s. or stochastic) limit of $(N_1(\delta, s) - N_2(\delta, s))$, if it exists, cannot be degenerate. Note that $L_{1n}(s) = (x_{(n)})^{n-1}$ and $L_{2n}(s) = (y_{(n)})^n$. Clearly, $P_\theta(N_1(s, \delta) = m) = P_\theta(N_2(s, \delta) = m+1)$, for all $m \geq 2$. The lemma below gives the exact distribution of $N_2(s, \delta)$ under P_θ .

Lemma 3.1.1. Let $p \equiv p(\theta, \delta)$ be the integer such that

$$(3.1.1) \quad \frac{\log \delta}{\log \theta} \leq p < \frac{\log \delta}{\log \theta} + 1.$$

Then the distribution function of $N_2(\delta, s)$ has the following expression:

$$(3.1.2) \quad P_\theta(N_2(s, \delta) \leq m) = \begin{cases} (\delta/\theta^{p-1}) \delta^{\sum_{j=1}^{p-1} j - m - 1} & \text{if } m \leq p-2 \\ \delta/\theta^{p-1} & \text{if } m = p-1 \\ 1 & \text{if } m = p \end{cases}$$

The proof is straightforward. The next lemma studies the weak convergence under P_θ of $p(\delta) - N_2(s, \delta)$ as $\delta \rightarrow 0$. Then $p(\delta) = p(\theta, e(\delta))$.

Lemma 3.1.2. For each c , $0 \leq c \leq 1$, let X_c be a r.v. such that $P_\theta(X_c = 0) = 1 - \theta^c$ and $P_\theta(X_c = i) = (1 - \theta) \cdot \theta^{c+i-1}$, $i \geq 1$. Let $e(\delta)$ be the excess over $(p(\delta)-1)$ of $\log \delta / \log \theta$, ($0 < e(\delta) \leq 1$). Then

(a) if $e(\delta_n) \rightarrow c$ and $\delta_n \rightarrow 0$, $p(\delta_n) - N_2(\delta_n; s)$ converges weakly to X_c under P_θ .

(b) if $p(\delta_n) - N_2(\delta_n; s)$ converges weakly to X under P_θ and $\delta_n \rightarrow 0$, $\{e(\delta_n)\}$ is a convergent sequence; moreover, $X = X_c$ where c is the limit of $e(\delta_n)$.

Proof: (a) By the definition of $e(\delta)$, $\delta = \theta^{(p-1)+e(\delta)}$ so that $\delta_n / \theta^{p(\delta_n)-1} = \theta^{e(\delta_n)} \rightarrow \theta^c$. (3.1.1) implies that

$$\theta^{-1} \cdot \delta^{(p(\delta)-1)^{-1}} < 1 \leq \theta^{-1} \delta^{(p(\delta))^{-1}}.$$

Thus for each $k \geq 1$, $\delta_n^{(p(\delta_n) - k)^{-1}} \rightarrow \theta$. It then follows from the lemma 3.1.1 that $P_\theta(p(\delta_n) - N_2(\delta_n, s) = m) \rightarrow P_\theta(X_C = m)$ $m \geq 0$, which completes the proof of (a).

(b) Because of (a), every convergent sub-sequence of $\{e(\delta_n)\}$, which is a bounded sequence, converges to the same real number.

It follows from the above lemma that the (a.s. P_θ or the stochastic) limit of $(N_1(\delta, s) - N_2(\delta, s))$ cannot be degenerate; to see this, one need only note that $N_1(\delta, s)$ and $N_e(\delta, s)$ are independent and then use Theorem 3.2, Chapter VIII of [7]. So in this case we cannot hope to get a single numerical value of deficiency from Bahadur's point of view.

3.2. Another interpretation of BCD.

The following result is due to Ragharachari [9] (see his Theorem 2). The set-up is the same as that given in section 1.

Lemma 3.2.1. For all β ($0 < \beta < 1$), $\lim_{n \rightarrow \infty} (\log \alpha_n(\theta, \beta)/n) = -c(\theta)$ iff $\log L_n(s)/n \rightarrow -c(\theta)$ (a.s. P_θ).

This fact leads to the following definition.

Definition 3.2.1. Fix a $\theta \in \mathbb{H}_0$, an ϵ with $0 < \epsilon < 1$ and a δ with $0 < \delta < 1$. Then $V(\epsilon, \delta) \equiv V(\theta, \epsilon, \delta)$ is the smallest integer $m \geq 1$ such that whenever $n \geq m$, $P_\theta(L_n(s) < \delta) > 1 - \epsilon$.

The next lemma gives the asymptotic behavior of $V(\epsilon, \delta)$ as $\delta \rightarrow 0$.

Lemma 3.2.2. Assume that $\log L_n(s)/n \rightarrow -c(\theta)$ (a.s. P_θ). Then for each ε , $0 < \varepsilon < \frac{1}{2}$, $\log \delta/V(\varepsilon, \delta) \rightarrow -c(\theta)$ as $\delta \rightarrow 0$.

Proof. The proof depends on routine calculations and hence is omitted.

The lemma 3.2.2 suggests the following measure of deficiency; consider the set-up of the section 1 and define $V_1(\varepsilon, \delta)$ and $V_2(\varepsilon, \delta)$ similarly using $L_{1n}(s)$, $L_{2n}(s)$ for $L_n(s)$.

Definition 3.2.2. Fix a $\theta \notin \mathbb{H}_0$, an ε with $0 < \varepsilon < \frac{1}{2}$. Then the lower (the upper) deficiency of the first testing procedure w.r.t. the second at θ is

$$\liminf_{\delta \rightarrow 0} (V_1(\varepsilon, \delta) - V_2(\varepsilon, \delta))$$

$$\limsup_{\delta \rightarrow 0} (V_1(\varepsilon, \delta) - V_2(\varepsilon, \delta)) .$$

Let $F_{in}(t)$ be a strictly decreasing continuous function of t , $i=1,2$. For each $\theta \notin \mathbb{H}_0$, we make the same assumption about $P_\theta\{T_{in} > t\}$.

For each $0 < \delta < 1$, let $t_{in}(\delta) = F_{in}^{-1}(\delta)$. Consider the sequence of tests $\phi_{in}(\delta)$:

$$\text{Reject } H_0 \text{ iff } T_{in} > t_{in}(\delta) .$$

Then the error of first kind for this test is δ . We denote its power by $\bar{\beta}_{in}$.

Fix $\theta \notin H_0$. For each $0 < \beta < 1$, define the test $\psi_{in}^{(\beta)}$:
 Reject H_0 iff $T_{in} > c_{in}(\beta)$ where $c_{in}(\beta)$ is such that
 $P_{\theta}\{T_{in} > c_{in}(\beta)\} = \beta$. Let its error of first kind be denoted
 by α_{in} .

Using the tests $\psi_{in}(\beta)$ define $M_i(\beta, \delta) \equiv M_i(\theta, \beta, \delta)$ as
 in section 1. Then

$$(3.2.1) \quad M_i(\beta, \delta) = V_i(1-\beta, \delta) .$$

To see this, note that if $n \geq V_i(1-\beta, \delta)$ then by definition
 of V_i , the tests ϕ_{in} have error of first kind = δ and
 power (at θ) $> \beta$. Hence for $n \geq V_i(1-\beta, \delta)$ the tests ψ_{in} which
 have power = β , must have error of first kind $\alpha_{in} < \delta$. By
 definition of $M_i(\beta, \delta)$ this means $M_i(\beta, \delta) \leq V_i(1-\beta, \delta)$.
 Similarly the reverse inequality can be proved.

Thus BCD, upper, lower or approximate, agrees with the
 corresponding notion as defined here.

4. Appendix: Proofs of the Results Given in Section 2.

Proof of Proposition 2.1.1: Easy.

Proof of Theorem 2.1.1: Direct consequence of Proposition 2.1.1.

Proof of Lemma 2.3.1: Assume (2.3.1). (2.2.1) is then immediate.

For (2.2.2) we define $\{\alpha_{ix}\}$ for non-integral x as follows:

let $n < x < n+1$; then $x = n\lambda + (n+1)\mu$ for some λ and μ

such that $0 < \lambda < 1$, $\lambda + \mu = 1$. We define $\log \alpha_{ix} = \lambda \log \alpha_{in} +$

$\mu \log \alpha_{i(n+1)}$. We claim that $\log \alpha_{ix} = -x a_i + \sqrt{x} b_i + c_i \log x + d_i + o(1)$

for this observe that $n = x - \mu$ and $n+1 = x + \lambda$ and so $\lambda \sqrt{n} + \mu \sqrt{n+1} =$

$\sqrt{x} + o(x^{-1})$. Also $\lambda \log n + \mu \log (n+1) = \log x + o(x^{-1})$. The

definition of α_{ix} shows that it is a decreasing function of x .

The proof of (2.2.3) is included in that of the theorem 2.3.1.

Proof of Theorem 2.3.1: First note that since $\log \alpha_{in}/n \rightarrow -a$

for each $i=1,2$, $m(n)/n \rightarrow 1$.

(a) From the definition of $m(n)$, one has

$$(m(n)-n) \left[a - \frac{b_1}{\sqrt{n} + \sqrt{m}} \right] - c_1 \log \frac{m(n)}{n} \rightarrow (d_1 - d_2)$$

i.e., $m(n) - n \rightarrow (d_1 - d_2)/a$.

(b) Here we have,

$$\frac{m(n)-n}{\sqrt{n}} \left[a - \frac{b_1}{\sqrt{n} + \sqrt{m}} \right] - (b_1 - b_2) + c_2 \frac{\log m(n)}{\sqrt{n}} - c_1 \frac{\log n}{\sqrt{n}} \rightarrow 0.$$

So $(m(n)-n)/\sqrt{n} \rightarrow (b_1 - b_2)/a$. From this, (b) is immediate.

(c) follows similarly.

Proof of Lemma 2.4.1: Similar to that of Theorem 2.3.1.

Proof of Proposition 2.5.1: We shall follow Bahadur and Ranga Rao [3]. Let H_n be the distribution function of the standardized n -fold convolution, and v the s.d., of the conjugate distribution of $x_1 - \mu$. Put $\alpha = v\tau$. Proceeding exactly in the same way of the Lemma 2 of [3], we have $p_n = \int_{q_n}^{\infty} e^{-\sqrt{n}\alpha x} dH_n(x)$ where

$$I_n = \int_{q_n}^{\infty} e^{-\sqrt{n}\alpha x} dH_n(x).$$

Because of the Lemma 1.2, $H_n(x) = \Phi(x) + \frac{K(1-x^2)}{\sqrt{n}} \phi(x) + \gamma_n(x)n^{-1/2}$

where $\gamma_n(x) \rightarrow 0$ uniformly in x and K is a constant.

(I) The contribution of $\Phi(x)$ to I_n is

$$\int_{q_n}^{\infty} e^{-\sqrt{n}\alpha x} d\Phi(x) = \frac{\exp(-\sqrt{n}\alpha q_n - q_n^2/2)}{\sqrt{2\pi} \sqrt{n} \alpha} \left\{ 1 - \frac{q_n}{\sqrt{n} \alpha} + \frac{q_n^2 - 1}{n\alpha^2} + o(n^{-1}) \right\}$$

(use Lemma 1.1).

(II) The contribution of $K(1-x^2)\phi(x)/\sqrt{n}$ to I_n is

$$\begin{aligned}
& \frac{\kappa}{\sqrt{n}} \int_{q_n}^{\infty} e^{-\sqrt{n} \alpha x} (x^3 - 3x) \phi(x) dx = \\
& = \frac{\kappa \exp(\frac{1}{2} n \alpha^2)}{\sqrt{n}} \int_{q_n + \sqrt{n} \alpha}^{\infty} \{(y - \sqrt{n} \alpha)^3 - 3(y - \sqrt{n} \alpha)\} \phi(y) \\
& = \frac{\exp(-\sqrt{n} \alpha q_n - q_n^2/2)}{\sqrt{2\kappa} \sqrt{n} \alpha} \cdot o(1) .
\end{aligned}$$

(III) Let $\varepsilon > 0$. As $\sup_x |\gamma_n(x)| < \varepsilon/2$ for all sufficiently large n ,

$$\begin{aligned}
n^{-\frac{1}{2}} \left| \int_{q_n}^{\infty} \exp(-\sqrt{n} \alpha x) d\gamma_n(x) \right| & \leq \alpha \int_{q_n}^{\infty} \exp(-\sqrt{n} \alpha x) |\gamma_n(x) - \gamma_n(q_n)| dx \\
& \leq \varepsilon n^{-\frac{1}{2}} \exp(-\sqrt{n} \alpha q_n) .
\end{aligned}$$

Thus the contribution of $n^{-\frac{1}{2}} \gamma_n(x)$ to I_n is

$$n^{-\frac{1}{2}} \exp(-\sqrt{n} \alpha q_n - q_n^2/2) \cdot o(1) .$$

From (I), (II) and (III), (2.5.1) follows.

Proof of Lemma 2.5.1. Because of the uniformity condition involved in (2.5.2), $F_n(z_\beta + dn^{-\frac{1}{2}}) = \Phi(z_\beta) + n^{-\frac{1}{2}}(d + q_\theta(z_\beta))\phi(z_\beta) + o(n^{-\frac{1}{2}})$.

Thus if we let $d = -q_\theta(z_\beta)$, then $F_n(z_\beta + dn^{-\frac{1}{2}}) = (1-\beta) + o(n^{-\frac{1}{2}})$.

Put $G_n(x) = \Phi(x) + n^{-\frac{1}{2}}q_\theta(x)\phi(x)$, $\varepsilon_{1n} = \sup_x |F_n(x) - G_n(x)|$

and $\varepsilon_{2n} = F_n(z_\beta + dn^{-\frac{1}{2}}) - (1-\beta)$. Choose $\eta_n \rightarrow 0$ ($\eta_n \neq 0$) such that $n^{\frac{1}{2}}\varepsilon_{in} = o(\eta_n)$ for each $i=1,2$. Then

$$\begin{aligned} & \frac{\sqrt{n}}{|\eta_n|} \{F_n(z_\beta + (d+\eta_n)n^{-\frac{1}{2}}) - (1-\beta)\} \\ &= \frac{\sqrt{n}}{|\eta_n|} \{G_n(z_\beta + (d+\eta_n)n^{-\frac{1}{2}}) - G_n(z_\beta + dn^{-\frac{1}{2}})\} + o(1) \\ &= \frac{\sqrt{n}}{|\eta_n|} \cdot \frac{n}{\sqrt{n}} \cdot G'_n(\xi_n) + o(1) \end{aligned}$$

for some ξ_n lying between $z_\beta + \frac{d}{\sqrt{n}}$ and $z_\beta + \frac{d}{\sqrt{n}} + \frac{\eta_n}{\sqrt{n}}$; in particular, $\xi_n \rightarrow z_\beta$. As $\{G'_n(u)\}$ is bounded away from zero in a neighborhood of z_β , one gets

$$\frac{\sqrt{n}}{|\eta_n|} \{F_n(z_\beta + (d + |\eta_n|)n^{-\frac{1}{2}}) - F_n(\frac{k_n - \sqrt{1}\mu}{\tilde{\sigma}})\} > 0$$

for all sufficiently large n .

This implies that $(k_n - \sqrt{n}\mu)/\tilde{\sigma} < z_\beta + dn^{-\frac{1}{2}} + |\eta_n|n^{-\frac{1}{2}}$ for all sufficiently large n . Similarly, $(k_n - \sqrt{n}\mu)/\tilde{\sigma} > z_\beta + dn^{-\frac{1}{2}} - |\eta_n|n^{-\frac{1}{2}}$ for all sufficiently large n . As $\eta_n \rightarrow 0$, the proof of the lemma 2.5.1 is complete.

Proof of Theorem 2.5.1:

Let $k_n^* = \sqrt{n} \mu + z_\beta \tilde{\sigma} + d \tilde{\sigma} n^{-1/2}$ and $\alpha_n^* = \sup\{P_{\theta_0}(T_n > k_n^*) : \theta_0 \in \Theta_0\}$.

Because of (2.5.3),

$$\log \alpha_n^* = n \log \rho - \sqrt{n} \alpha z_\beta \tilde{\sigma} - \frac{1}{2} \log n + (a - \alpha \tilde{\sigma} d - \frac{1}{2} z_\beta^2 \tilde{\sigma}^2) + o(1).$$

It suffices to show that

$$(*) \quad \log \alpha_n = \log \alpha_n^* = o(1).$$

Put $q_n = (k_n - \sqrt{n} \mu) / \tilde{\sigma}$ and $q_n^* = (k_n^* - \sqrt{n} \mu) / \tilde{\sigma}$. Then both the sequences $\{q_n\}$ and $\{q_n^*\}$ are bounded. By (2.5.3), one gets

$$\begin{aligned} & \log \alpha_n - \log \alpha_n^* \\ &= \sqrt{n} \alpha (q_n^* - q_n) + \frac{1}{2} (q_n^{*2} - q_n^2) + o(1) \\ &= \sqrt{n} \alpha \frac{(k_n^* - k_n)}{\tilde{\sigma}} + \frac{1}{2} \cdot (q_n + q_n^*) \frac{(k_n^* - k_n)}{\tilde{\sigma}} + o(1) \\ &= o(1) \text{ by lemma 2.5.1.} \end{aligned}$$

This completes the proof of (*).

References

- [1] Bahadur, R. R.: Rates of convergence of estimates and test statistics, Ann. Math. Stat., vol. 38, 303-304, 1967.
- [2] _____: Some limit theorems in statistics, SIAM, Philadelphia, 1971.
- [3] Bahadur, R. R. and Rao, R. Ranga: On deviations of the sample mean, Ann. Math. Stat., vol. 31, 1015-1027, 1960.
- [4] Bhattacharya, R. N. and Ghosh, J. K.: On the validity of the formal Edgeworth expansions, (To appear), 1976.
- [5] Cochran, W. G.: The χ^2 -goodness of fit test, Ann. Math. Stat., vol. 23, 507, 1952.
- [6] Feller, W.: An introduction to probability theory and its applications, vol. I.
- [7] _____: An introduction to probability theory and its applications, vol. II.
- [8] Gnedenko, B. V. and Kolmogorov, A. N.: Limit theorems for sums of independent random variables. English translation by K. L. Chung, Addison-Wesley, Reading, Mass. (1954).
- [9] Hodges, J. L., Jr. and Lehman, E. L.: Deficiency, Ann. Math. Stat., vol. 41, 783-801, 1970.
- [10] Killen, H. and Sievers, G. L.: An elementary theorem on the probability of large deviations, Ann. Math. Stat., vol. 43, 181-192, 1972.
- [11] Raghavachari, M.: On a theorem on the rate of convergence of test statistics, Ann. Math. Stat., vol. 41, 1695-1699, 1970.

List of Changes

Page	Line	Change	To
14	12 from top	" $m(n)-n-d > 0$ "	" $m(n)-n-d < 0$ "
14	7 from bottom	" $m(n)-n-d < 0$ "	" $m(n)-n-d > 0$ "
18	3 from top	" $\bar{D}_c(\theta, \beta) = d(\theta, \beta) + 1$ "	" $\bar{D}_c(\theta, \beta) = d(\theta, \beta)$ "
18	3 from top	" $\underline{D}_c(\theta, \beta) = d(\theta, \beta)$ "	" $\underline{D}_c(\theta, \beta) = d(\theta, \beta) - 1$ "
17	last two lines	" $\bar{D}_c(\theta, \beta) = d(\theta, \beta)$ " " $\underline{D}_c(\theta, \beta) = d(\theta, \beta) - 1$ "	" $\bar{D}_c(\theta, \beta) = d(\theta, \beta) + 1$ " " $\underline{D}_c(\theta, \beta) = d(\theta, \beta)$ "
19	8,9 from bottom	"Also, $\bar{D}_c(\theta, \beta) =$ $2 \log 2/\theta^2$ "	"Also, if $2 \log 2/\theta^2$ is an integer (see remark 2.2.2 for the contrary case) $\bar{D}_c(\theta, \beta) = 2 \log 2/\theta^2$ "
22	6 from top	"Also, $\bar{D}_c(\theta_1, \beta) =$ $d(\theta_1, \beta)$ "	"Also if $2 \log 2/(1+\mu^2)$ is an integer (see remark 2.2.2 for the contrary case) $\bar{D}_c(\theta_1, \beta) = d(\theta_1, \beta)$ "
22	2 from bottom	"upper bound for $I_n(\alpha)$ "	" $r_n(\alpha)[1 - (n-4)^{-1}(\alpha^{-2} + 11)$ $+ 3(n-4)^{-1}(n-6)^{-1}(1+\alpha^{-2})]$ "
25	11, 13, 21 from top	" X_1 "	" Y_1 "
31	7 from bottom	" $\log L_n(s) \rightarrow$ $-c(\theta)(a.s. P_\theta)$ "	" $\log L_n(s) \xrightarrow{P_\theta} -c(\theta)$ "
32	1 from top	" $\log L_n(s)$ $-c(\theta)(a.s. P_\theta)$ "	" $\log L_n(s) \xrightarrow{P_\theta} -c(\theta)$ "
32	2 from top	" $\epsilon < \frac{1}{2}$ "	" $\epsilon < 1$ "
32	8 from top	" $\epsilon < \frac{1}{2}$ "	" $\epsilon < 1$ "