数理解析研究所講究録
340巻 1978年 4-11



# Non-existence of bounded functions on the universal covering of $\{\mathbb{P}^n - n+2$ hyperplanes in general position$\}$ $(n\geqslant 2)$.

by Isao Wakabayashi

Let X be the complement of n+2 hyperplanes in general position in $\mathbb{P}^n$, and let $\pi:\tilde{X}\longrightarrow X$ be the universal covering of X. Then we have

THEOREM 1. If $n\geqslant 2$, there are no bounded, holomorphic and non-constant functions on $\tilde{X}$. ([3])

We derive this theorem from the following theorem.

THEOREM 2. There are no bounded, holomorphic and non-constant functions on the "homological covering" of $\{\mathbb{P}^1 - 3$ points$\}$. ([4])

## 1°) RELATION BETWEEN THE UNIVERSAL COVERING $\tilde{X}$ AND THE HOMOLOGICAL COVERING OF $\{\mathbb{P}^1 - 3$ POINTS$\}$.

DEFINITION. The *homological covering* of a variety is the covering determined by the commutator group of the fundamental group of the variety.

Notation. We denote by R the homological covering of $D=\mathbb{P}^1-\{0,1,\infty\}$.

Let $z_1,\cdots,z_n$ be the inhomogeneous coordinates of $\mathbb{P}^n$. We may suppose that the n+2 hyperplanes in general position which we take away from $\mathbb{P}^n$ are $\{z_1=0\},\cdots,\{z_n=0\}$, $\{z_1+\cdots+z_n-1=0\}$ and the infinite hyperplane. Let $L(z_1,\cdots\overset{i}{\vee}\cdots,z_n)$ be the complex line consisting of the points whose j-th coordinate is equal to $z_j$, $j=1,\cdots,i-1,i+1,\cdots,n$.

Let us suppose that $z_1+\cdots+z_{i-1}+z_{i+1}+\cdots+z_n-1\neq 0$, then by a topological obsevation we have

PROPOSITION. Every connected component of the inverse image of $L(z_1,\cdots\overset{i}{\vee}\cdots,z_n)$ by $\pi$ is holomorphically isomorphic to the homological covering R of D.

From this proposition and Theorem 2, the proof of Theorem 1 is immediate, because there are hense in $\tilde{X}$ sufficiently many analytic subsets admitting no bounded functions.

2°) TRIANGULATION OF R.

Let us pose $\Delta_1=\{z\in \mathbf{C} \mid \operatorname{Im} z\geqslant 0,\ z\neq 0,1\}$ and $\Delta_2=\{z\in \mathbf{C} \mid \operatorname{Im} z\leqslant 0,\ z\neq 0,1\}$. Let A be the segment (0, 1), B be the segment $(1,\infty)$ and let C be the segment $(\infty, 0)$. We consider $\Delta_1$ and $\Delta_2$ as triangles with 3 sides A, B and C. The inverse image $\pi^{-1}(\Delta_1)$ of $\Delta_1$ (resp. $\pi^{-1}(\Delta_2)$ of $\Delta_2$) by the projection $\pi:R\longrightarrow D$ consists of an infinite number of triangles which are holomorphically isomorphic to $\Delta_1$ (resp. $\Delta_2$). We number the triangles of $\pi^{-1}(\Delta_1)$ and those of $\pi^{-1}(\Delta_2)$ as follows: Let p be a point in $\Delta_1$. Let $\alpha$ (resp. $\beta$) be a closed curve of origin p and of end point p winding the point 0 (resp. the point 1) in the positive sense. Then every element of $\pi_1(D, p)/[\pi_1(D, p), \pi_1(D, p)]$ is represented uniquely by a curve of the form $\alpha^i\beta^j$ $(i,j\in \mathbf{Z})$. This element determines a point $p^{i,j}$ situated over p, so we denote by $\Delta_1^{i,j}$ the triangle of $\pi^{-1}(\Delta_1)$ which contains $p^{i,j}$. We further denote by $\Delta_2^{i,j}$ the triangle of $\pi^{-1}(\Delta_2)$ whose side B is identical with the side B of $\Delta_1^{i,j}$.

3°) TOPOLOGY OF R.

We prepare a countable number of triangles $\underline{\Delta}_1^{i,j}$ and $\underline{\Delta}_2^{i,j}$ $(i,j \in \mathbf{Z})$. We place them as Fig. 1.

Fig. 1.



We consider each pair of sides which are situated face to face in Fig. 1. We identify them oppositely so that the given orientations coinside. Let us denote by R' the surface constructed in this way. Then we can prove that the triangles $\Delta_1^{i,j}$ and $\Delta_2^{i,j}$ are patched in R exactly in the same way as the triangles $\underline{\Delta}_1^{i,j}$ and $\underline{\Delta}_2^{i,j}$ in R'. Hense

PROPOSITION. R is homeomorphic to R'.

4°) CRITERION OF A. PFLUGER.

To show Theorem 2 we use a criterion of A. Pfluger which gives a sufficient condition for a Riemann surface to belong to the class $O_{AB}$, i.e. the class of surfaces with no bounded holomorphic functions.

Let W be an open Riemann surface and let $\{A_n^k\}$, $k=1,2,\cdots$, $k(n)<\infty$, $n=1,2,\cdots$, be a family of doubly connected domains in W

satisfying the following conditions:

(1) $A_n^k$ is bounded by two closed curves $\alpha_n^k$ and $\alpha'^k_n$,

(2) $A_n^k \cap A_{n'}^{k'} = \phi$ if $(k,n) \neq (k',n')$,

(3) the complement of $\bigcup_{k=1}^{k(n)} A_n^k$ in W has a unique compact connected component $B_n$,

(4) $B_n$ is bounded by k(n) curves $\{\alpha_n^k\}_{k=1}^{k(n)}$ and contains all $A_{n'}^{k'}$ such that $n' < n$.

Let us denote by $\mu_n^k$ the harmonic module of $A_n^k$, where the *harmonic module* of a domain conformally equivalent to $\{z \in \mathbf{C} \mid r < |z| < r'\}$ is by definition $\log r'/r$. We pose

$$\mu_n = \operatorname*{Min}_k \mu_n^k, \text{ and } K(N) = \operatorname*{Max}_{n \leq N} k(n).$$

CRITERION (A. Pfluger). ([1], [2]) If

$$\overline{\lim_{N\to\infty}} \left\{ \sum_{n=1}^{N} \mu_n - \frac{1}{2}\log K(N) \right\} = \infty,$$

then $W \in O_{AB}$.

5°) CONSTRUCTION OF POLYGONES $\{P_n\}$, AND ITS BOUNDARY.

We construct on our surface R a family of polygones $\{P_n\}$ such that

(i) each $P_n$ consists of a finite number of triangles in 2°,

(ii) $P_n \subset P_{n+1}$,

(iii) $\bigcup_{n=1}^{\infty} P_n = R$.

Actually $P_n$ is constructed as Fig.2.

We find that the boundary of $P_n$ consists of 2n+1 connected

Figure 2.


$\gamma_n^{n-1}$
$\Delta_1^{n,-n}$
$\gamma_n^{k}$
$\Delta_2^{n+1,-k}$
$\Delta_1^{k+1,-n}$
$\Delta_2^{k+1,-n}$
$\Delta_1^{n,-k-1}$
$\Delta_2^{n,-k}$
$P_n$
$\Delta_2^{n+1,n+1}$
$\Delta_2^{-n,-n}$
$\Delta_1^{0,0}$
$\Delta_1^{-n,-n}$
$\Delta_1^{n,n+1}$
$\Delta_1^{-n-1,-n}$
$\Delta_1^{k+1,n+1}$
$\Delta_1^{-n-1,-k-1}$
$\Delta_2^{-n,-k}$
$\gamma_n^{-n-1}$
$\Delta_2^{-n,n+1}$

components. For each connected component of the boundary of $P_n$, we describe a closed curve $\gamma_n^k$ $(k=-n-1,\cdots,n-1)$ which runs sufficiently near to it.
$\{\gamma_n^k\}$ are given in Fig. 2.

6°) CONSTRUCTION OF DOUBLY CONNECTED DOMAINS $\{A_n^k\}$.

Let r, r' be two real numbers such that $r < r'$. Let us denote by $\Delta_\nu^{i,j}(0,r,r')$ the part of $\Delta_\nu^{i,j}$ $(\nu=1,2)$ lying over $\{z \in \mathbf{C} \mid r < |z| < r'\}$. Let $\Delta_\nu^{i,j}(1,r,r')$ be the part of $\Delta_\nu^{i,j}$ lying over $\{z \in \mathbf{C} \mid r < |z-1| < r'\}$. We pose $\Delta_\nu^{i,j}(\infty,r',r)=\Delta_\nu^{i,j}(0,r,r')$. Let $\Delta_\nu^{i,j}(0,r;1,r';\infty,r'')$ be the complement in $\Delta_\nu^{i,j}$ of the part lying over the set $\{|z|<r\} \cup \{|z-1|<r'\} \cup \{|z|>r''\}$.

We construct $A_n^k$ as the union of the following sets. We take the set of the form $\Delta_\nu^{i,j}(0,r,r')$ (resp. $\Delta_\nu^{i,j}(1,r,r')$ and $\Delta_\nu^{i,j}(\infty,r',r)$) if $\Delta_\nu^{i,j}$ is situated in the interior of $P_n$ and if $\gamma_n^k$ passes near 0 (resp. 1 and $\infty$) in $\Delta_\nu^{i,j}$. And we take the set of the form $\Delta_\nu^{i,j}(0,r;1,r';\infty,r'')$ if $\gamma_n^k$ passes the triangle $\Delta_\nu^{i,j}$ which is situated on the boundary of $P_n$. For example, in $\Delta_1^{i,-k-1}$ (resp. $\Delta_2^{i+1,-k}$) for $-n \leq i \leq n-1$, we take the part $\Delta_1^{i,-k-1}(0,r_{n+1},r_n)$ (resp. $\Delta_2^{i+1,-k}(0,r_{n+1},r_n)$), where $r_n=Ce^{-2\pi n}$ $(C>0)$. And in $\Delta_2^{n+1,-k}$ (resp. $\Delta_1^{n,-k-1}$), we take the part $\Delta_2^{n+1,-k}(0,r_{n+1};1,\varepsilon;\infty,1/r_{\frac{n-k+1}{2}})$ (resp. $\Delta_1^{n,-k-1}(0,r_{n+1};1,\varepsilon;\infty,1/r_{\frac{n-k-1}{2}})$) with $1 \gg \varepsilon > 0$. We put $A_n^k$ the union of all these sets.

7°) HARMONIC MODULE OF $A_n^k$.

We divide $A_n^k$ into some suitable parts, so that we have rectangles as their images by the map $w=\log z$ or $w=\log(z-1)$. We see that the width of each rectangle is $2\pi$ and the total length of all these rectangles is almost equal to $24\pi n$. Here $24\pi n$ is obtained as follows: $A_n^k$ passes $12n+1$ triangles among those which are in the interior of $P_n$. The intersection of $A_n^k$ and each triangle of this kind is mapped by the above map to a rectangle of width $2\pi$ and of length $\pi$. So the total length of the intersection of $A_n^k$ and all triangles in the interior of $P_n$ is almost equal to $12\pi n$. On the other hand, we see that the total length of the intersection of $A_n^k$ and all the triangles which are situated on the boundary of $P_n$ is almost equal to $12\pi n$. $24\pi n$ is the sum of these quantities.

We can verify that these rectangles are patched together only slightly twisted and so $A_n^k$ is almost like an annulus of width $2\pi$ and of length $24\pi n$ when we consider its harmonic module. Consequently the harmonic module $\mu_n^k$ of $A_n^k$ is almost equal to $\log\frac{12n+2\pi}{12n}$, hense $\mu_n=\operatorname{Min}_k \mu_n^k$ is almost equal to $\log\frac{12n+2\pi}{12n}$.

Since $K(N)=\operatorname{Max}_{n\leq N} k(n)=2N+1$, and $\pi/6=0.52\cdots>1/2$, we obtain

$$\overline{\lim_{N\to\infty}}\left\{\sum_{n=1}^{N}\mu_n-\frac{1}{2}\log K(N)\right\}\geqslant\overline{\lim_{N\to\infty}}\left\{\sum_{n=1}^{N}\frac{\pi}{6n}-\frac{1}{2}\sum_{n=1}^{N}\left(\frac{\pi}{6n}\right)^2-\frac{1}{2}\log(2N+1)\right\}$$

$$\geqslant\overline{\lim_{N\to\infty}}\left\{\frac{\pi}{6}\log N-\frac{1}{2}\log(2N+1)-\frac{1}{2}\sum_{n=1}^{N}\left(\frac{\pi}{6n}\right)^2\right\}=\infty.$$

Hense by virtue of the Pfluger's criterion, our Riemann surface R belongs to the class $O_{AB}$ as required.

BIBLIOGRAPHIE

[1] Mori (A.). A note on unramified abelian covering surfaces of a closed Riemann surface, J. Math. Soc. Japan, Vol. 6 (1954) p. 162-176.

[2] Pfluger (A.). Sur l'existence de fonctions non constantes, analytiques, uniformes et bornées sur une surface de Riemann ouverte, C.R. Acad. Sci. Paris, tom 230 (1950) p. 166-168.

[3] Wakabayashi (I.). Non-existence de fonctions holomorphes bornées non constantes sur le revêtement universel de $\{\mathbf{P}^n-(n+2)$ hyperplans$\}$, dans le cas $n \geqslant 2$, C.R. Acad. Sci. Paris, tom 285 (1977) p. 373-374.

[4] Wakabayashi (I.). Non-existence de fonctions bornées sur le revêtment homologique de $\{\mathbf{P}^1-$ 3 points$\}$, to appear.

Department of Mathematics,
Tokyo University
of
Agriculture and Technology,