数理解析研究所講究録
342巻 1979年 33-38

On certain non-continuous functions and shape

Hiroo Fukaishi

In the shape category of topological spaces a shape morphism is constructed by a system of maps (= continuous functions) ; it is, in general, not generated by a single map.
Hence we have the following questions :

Question 1. Is it possible that a kind of non-continuous function induces a shape morphism ?

Question 2. Can a shape equivalence be generated by a certain non-continuous function ?

Definition 1. Let X and Y be topological spaces. A function $f : X \to Y$ is a connectivity function if for any connected $C \subset X$, the graph $G(f|C)$ of $f|C$ is connected.

Definition 2. A function $f : X \to Y$ is almost continuous if for any open set $N \subset X \times Y$ containing $G(f)$ there is a continuous function $g : X \to Y$ such that $G(g) \subset N$.
These notions have been considered to generalize Brouwer's fixed point theorem (cf. Stallings[10]).

Each of the following is intermediate to answer the questions.

Proposition 1. Let $f : X \to Y$ be an almost continuous function between compact metric spaces. Then there are ANR-sequences $\underline{X} = \{X_i, p_{ij}\}$ and $\underline{Y} = \{Y_i, q_{ij}\}$ with limits X and Y, respectively, and a system $\underline{f} : \underline{X} \to \underline{Y}$ of almost continuous functions $f_i : X_i \to Y_i$ such that $f_i p_{ij} = q_{ij} f_j$ for $i \leq j$.

Proof. There are ANR-sequences $\underline{X}$ and $\underline{Y}$ with limits X and Y, respectively, and each projection p_i, q_i surjective, since both X and Y are compact metric. Define a function $f_i : X_i \to Y_i$ for each i, by the formula $f_i p_i = q_i f$. The almost continuity of f implies that of f_i.

Proposition 2. A bijective connectivity function with connectivity inverse function does not induce a shape equivalence.

Proof. By the example of Stallings [10,p.262]. Let X be the circle represented as the real numbers mod 1. Define a function $f : X \to X \times X$ by the formula

$$f(x \bmod 1) = 1/x \bmod 1, \text{ where } 0 < x \leq 1.$$

Let Y be the graph of f and $f^* : X \to Y$ such that

$$f^*(x) = (x, f(x)).$$

Then f^* is a bijection, and both f^* and f^{*-1} are connectivity functions, but $Sh(X) \neq Sh(Y)$, because their 1-dimensional Čech cohomology groups are different.

REFERENCES ON NON-CONTINUOUS FUNCTIONS

[1] Noguchi,H., A generalization of absolute neighborhood retracts, Kôdai Math. Sem. Rep.,5(1953),20-22.

[2] Fan,K. & Struble,R.A., Continuity in terms of connectedness, Indag. Math.,16(1954),161-164.

[3] Tanaka,T., On the family of connected subsets and the topology of spaces, J. of Math. Soc. Japan,7(1955),389-393.

[4] Nash,J., Generalized Brouwer theorem, Bull. Amer. Math. Soc., 62(1956),76, (Research problem).

[5] Hamilton,O.H., Fixed points for certain noncontinuous transformations, Proc. Amer. Math. Soc.,8(1957),750-756.

[6] O'Neill,B. & Straus,E.G., A fixed point theorem, Proc. Amer. Math. Soc.,8(1957),1148-1151.

[7] O'Neill,B., Induced homology homomorphisms for set-valued maps, Pacific J. Math.,7(1957),1179-1184.

[8] Pervin,W.J. & Levine,N., Connected mappings of Hausdorff spaces, Proc. Amer. Math. Soc.,9(1958),488-496.

[9] Ponomarev,V.I., A new space of closed sets and many-valued mappings, D. A. N. SSSR,118(1958),1081-1084.

[10] Stallings,J., Fixed point theorems for connectivity maps, Fund. Math.,47(1959),249-263.

[11] Long,P.E., Properties of certain non-continuous transformations, Duke Math. J.,28(1961),639-645.

[12] Hildebrand,S.K. & Sanderson,D.E., Connectivity functions and retracts, Fund. Math.,57(1965),237-245.

[13] Roberts,J.H., Zero-dimensional sets blocking connectivity functions, Fund. Math.,57(1965),173-179.

[14] Cornette,J.L., Connectivity functions and images on Peano continua, Fund. Math.,58(1966),183-192.

[15] Hagan,M.R., Equivalence of connectivity maps and peripherally continuous transformations,Proc. Amer. Math. Soc.,17(1966), 175-177.

[16] Naimpally,S.A., Graph topology for function spaces, Trans. Amer. Math. Soc.,123(1966),267-272.

[17] Whyburn,G.T., Connectivity of peripherally continuous functions, Proc. Nat. Acad. Sci. U. S. A.,55(1966),1040-1041.

[18] Cornette,J.L. & Girolo,J.E., Connectivity retracts of finitely coherent Peano continua, Fund. Math.,61(1967),177-182.

[19] Poppe,H., Über Graphentopologien für Abbildungsräume, I, Bull. Acad. Polon. Sci.,40(1967),71-80.

[20] Fuller,R.V., Relations among continuous and various non-continuous functions, Pacific J. Math.,25(1968),495-509.

[21] Gmurczyk,A., On approximative retracts, Bull. Acad. Polon. Sci.,16(1968),9-14.

[22] Granas,A., Fixed point theorems for the approximative ANR-s, Bull. Acad. Polon. Sci.,16(1968),15-19.

[23] Hyman,D.M., ANR divisors and absolute neighborhood contractibility, Fund. Math.,62(1968),61-73.

[24] Lin,S.T., Fixed point properties and inverse limit spaces, Pacific J. Math.,25(1968),117-122.

[25] Hyman,D.M., On decreasing sequences of compact absolute retracts, Fund. Math.,64(1969),91-17.

[26] Hoyle,III,H.B., Connectivity maps and almost continuous functions, Duke Math. J.,37(1970),671-680.

[27] Naimpally,S.A. & Pareek,C.M.,Graph topologies for function spaces,II, Annales Soc. Math. Polon. Ser.I.,13(1970),221-231.

[28] Brown,J.B., Totally discontinuous connectivity functions, Colloq. Math.,23(1971),53-60.

[29] Clapp,M.H., On a generalization of absolute neighborhood retracts, Fund. Math.,70(1971),117-130.

[30] Gmurczyk,A., Approximative retracts and fundamental retracts, Colloq. Math.,23(1971),61-63.

[31] Chewning,W.C., Connectivity retracts of unicoherent Peano continua in R^n, Fund. Math.,75(1972),25-27.

[32] Kellum,K.R. & Garrett,B.D., Almost continuous real functions, Proc. Amer. Math. Soc.,33(1972),181-184.

[33] Powers,M.J., Fixed point theorems for non-compact approximative ANR's, Fund. Math.,75(1972),62-68.

[34] Kellum,K.R., Almost continuous functions on I^n, Fund. Math., 79(1973),213-215.

[35] Bogatyi,S.A., Approximative and fundamental retracts, Math. USSR Sbornik,22(1974),91-103.

[36] Felt,J.E., ε-continuity and shape, Proc, Amer. Math. Soc.,46 (1974),426-430.

[37] Kellum,K.R., Sums and limits of almost continuous functions, Colloq. Math.,31(1974),125-128.

[38] Cannon, J. W., Taming cell-like embedding relations, Lecture Notes in Math., 439, Springer, (1975), 66-118.

[39] Kellum, K. R., On a question of Borsuk concerning non-continuous retracts I, Fund. Math., 87(1975), 90-92.

[40] Kellum, K. R., Non-continuous retracts, Stavrakas & Allen ed., Studies in topology, Academic Press, (1975), 255-261.

[41] Miller, Jr., M. H., Discontinuous G_δ graphs, ibid., 383-392.

[42] Kellum, K. R., On a question of Borsuk concerning non-continuous retracts II, Fund. Math., 92(1976), 135-140.

[43] Kellum, K. R., The equivalence of absolute almost continuous retracts and ε-absolute retracts, Fund. Math., 96(1977), 229-235.

[44] Ranchin, D. V., On modification of topologies in connection with the given discontinuous mappings of the space into itself, Bull. Acad. Polon. Sci., 26(1978), 149-156.