

ON REDUCIBLE PLANE CURVES

Makoto Yamamoto

Waseda university

§1. Introduction.

Let $f: \mathbb{C}^2 \rightarrow \mathbb{C}$ be a polynomial function such that $f(0)=0$ and f has an isolated critical point at the origin, and let L be the intersection of $V=f^{-1}(0)$ with a small sphere S_ϵ^3 . If f is analytically irreducible, then L is an iterated torus knot whose type is determined by the Puiseux characteristic pairs associated for f . If f is reducible and factored by irreducible components f_i as $f=f_1 \cdots f_r$, then L is a link of r components whose components are associated knots for f_i . Let K_i be the intersection of $V_i=f_i^{-1}(0)$ with a small sphere S_ϵ^3 for $i=1, \dots, r$. For the complement X of L (in S^3), $H_1(X; \mathbb{Z})$ is a free abelian multiplicative group G on the symbols $\{t_i\}_{i=1}^r$, where each t_i is geometrically carried by a meridian circle with homological linking number δ_{ij} with the component K_j of L . Let Λ_r be the integral group ring $\mathbb{Z}G$, and let \tilde{X} be the universal abelian covering of X . Then $H_1(\tilde{X}; \mathbb{Z})$ is a Λ_r -module. The Alexander polynomial of L is defined to be the characteristic polynomial of $H_1(X; \mathbb{Z})$ and it is topological invariant of L (up to units in Λ_r).

Lê [1] showed that algebraic knots having the same Alexander Polynomial (mod. unit) are of the same knot type. In this paper we shall show the following; let L be an algebraic link of $r(\geq 2)$ components whose components have distinct tangent cones each

other.

THEOREM A. Let L and L' be as above. If L and L' have the same Alexander polynomial (mod. units), then They are of the same link type.

§2. Algebraic links and Alexander polynomials.

First of all, we review algebraic knots and associated Puiseux expansions. We mainly refer to F. Pham [3].

Let $f: \mathbb{C}^2 \rightarrow \mathbb{C}$ be an analytically irreducible polynomial such that $f(0)=0$ and f has an isolated critical points at the origin. Then f can be written as

$$f(x,y) = \prod_{\zeta^m=1} (y-y(\zeta x))$$

in a small neighborhood of the origin, where m is the order of y and ζ 's are all m -th roots of unity, and

$$y(\zeta x) = \sum_{n=1}^{+\infty} a_n \zeta^n x^{n/m} \quad a_n \in \mathbb{C}.$$

Let $\mu = n_0/m$ be the exponent of the first term of $y(\zeta x)$ which has a non-zero coefficient. By $\text{in}(y-y(\zeta x))$, we denote the terms of $y-y(\zeta x)$ which have the smallest degree; that is,

$$\text{in}(y-y(\zeta x)) = \begin{cases} y & \mu > 1, \\ y - a_0 x & \mu = 1, \\ -a_{n_0} \zeta^{n_0} x^{n_0/m} & \mu < 1. \end{cases}$$

DEFINITION. The tangent cone $\mathbb{C}(V,0)$ of V at the origin is

the limit of the tangents at v_j , where v_j are the points of V tending to the origin.

Since $\text{in}(f) = \text{in}(\prod_{\zeta=1}^m (y - y(\zeta x))) = \prod_{\zeta=1}^m \text{in}(y - y(\zeta x))$, then

$$\text{in}(f) = \begin{cases} y^m & \mu > 1, \\ (y - a_{n_0} x)^m & \mu = 1, \\ \eta (-a_{n_0})^m x^{n_0} & \mu < 1, \text{ where } \eta = \prod_{\zeta=1}^m \zeta^{n_0}. \end{cases}$$

Then the tangent cone $\mathcal{C}(V, 0)$ is the line defined by $\text{in}(f) = 0$.

By changing coordinates linearly, we take the x -axis as $\mathcal{C}(V, 0)$. Then, assuming that $\mathcal{C}(V, 0)$ is the x -axis and by taking $\zeta = 1$, we have the Puiseux expansion of f

$$y = p(x) + a_1 x^{n_1/m_1} + \sum_{j=1}^{\ell_1} a_{1,j} x^{n_1+j/m_1} + a_2 x^{n_2/m_1 m_2} + \sum_{j=1}^{\ell_2} a_{2,j} x^{n_2+j/m_1 m_2} \\ + \dots + a_q x^{n_q/m_1 \dots m_q} + \sum_{j=1}^{+\infty} a_{q,j} x^{n_q+j/m_1 \dots m_q},$$

where $p(x)$ is a polynomial of x . We call the sequence of pairs of relatively prime positive numbers $\{(n_j, m_j)\}_{j=1}^q$ the Puiseux characteristic pairs of f .

Let K be the associated knot $V \cap S_{\varepsilon}^3$ for f . Then K is the iterated torus knot of type $\{(m_j, \lambda_j)\}_{j=1}^q$ inductively constructed as follows; let K^0 be the unknotted circle $\mathcal{C}(V, 0) \cap S_{\varepsilon'}^3$, where ε' is the positive number such that $\varepsilon' < \varepsilon$ and ε' is sufficiently close to ε . We call K^0 the primitive core of K . Let K^1 be the torus knot of type (m_1, n_1) in a small tubular neighborhood of K^0 , where the first coordinate m_1 and the second coordinate n_1 are

the longitudinal winding number and the meridional winding number respectively. This notation is opposite to that of Lê [1] and Sumners and Woods [4]. We suppose that the $(q-1)$ -st iteration K^{q-1} has been constructed. Let T and T_{q-1} be an unknotted torus and a small tubular neighborhood of K^{q-1} respectively. Let $\varphi: T \rightarrow T_{q-1}$ be the orientation preserving diffeomorphism from T to T_{q-1} which carries the longitude to the longitude. Then K is defined to be the image $K^q = \varphi(k)$ of a torus knot k of type (m_q, λ_q) in T , where

$$(2.1) \quad \begin{aligned} \lambda_1 &= n_1 \\ \lambda_j &= n_j - n_{j-1} m_j + \lambda_{j-1} m_{j-1} m_j \quad j=2, \dots, q. \end{aligned}$$

Next, we consider the case that f is reducible. Let $f: \mathbb{C}^2 \rightarrow \mathbb{C}$ be a polynomial such that $f(0)=0$ and f has an isolated critical point at the origin. We suppose that f is factored by irreducible components f_i as $f=f_1 \cdots f_r$ ($r \geq 2$). Let the Puiseux characteristic pairs of f_i be $\{(n_{ij}, m_{ij})\}_{j=1}^{q_i}$, and let K_i be the associated iterated torus knot of type $\{(m_{ij}, \lambda_{ij})\}_{j=1}^{q_i}$ for f_i . In this paper, we consider the case that all tangent cones $(V, 0)$ ($V_i = f_i^{-1}(0)$) are distinct each other. For the general case, refer to Sumners and Woods [4] and Yamamoto [6]. We denote the associated link for f by L . Then L is the disjoint union of K_1, \dots, K_r constructed as follows; let L^0 be the link consisting of all $K_i^0 = \mathcal{C}(V_i, 0) \cap S_\varepsilon^3$ ($i=1, \dots, r$). We note that L^0 has the same link type of the torus link of type (r, r) . Then L is the link obtained by iteration of type $\{(m_{ij}, \lambda_{ij})\}_{j=1}^{q_i}$ on each K_i^0 for $i=$

$1, \dots, r$.

Let $l = k_1 \cup \dots \cup k_r$ be a link of r components in S^3 , and l' be the link obtained from l by iteration of type (m, λ) on a component k_r , where $m > 1$. Let $\Delta(l; t_1, \dots, t_r)$ and $\Delta(l'; t_1, \dots, t_r)$ be the Alexander polynomials of l and l' respectively. Sumners and Woods [4] proved the following useful theorem.

THEOREM (Sumners and Woods [4] 5.1.) Let l and l' be as above. Then we have

$$(2.2) \quad \Delta(l'; t_1, \dots, t_r) = \Delta(l; t_1, \dots, t_r^m) Q(t_r, y; m, \lambda),$$

where $y = \sum_{i=1}^{r-1} t_i^{<k_i, k_r>}$, $<k_i, k_r>$ is the linking number of k_i and k_r

and $Q(t, s; m, \lambda)$ is the Alexander polynomial of two components formed by the torus knot k of type (m, λ) and the unknotted meridian curve on the boundary torus containing k ,

$$(2.3) \quad Q(t, s; m, \lambda) = ((t^\lambda s)^m - 1) / (t^\lambda s - 1).$$

Let L be the algebraic link associated for a reducible polynomial $f = f_1 \dots f_r$ whose tangent cones are distinct each other. Since L^0 has the same link type of the torus link of type (r, r) , we have

$$\Delta(L^0; t_1, \dots, t_r) = (t_1 \dots t_r - 1)^{r-2}.$$

Then we can compute the Alexander polynomial $\Delta(L; t_1, \dots, t_r)$ of L by (2.2). Let $\nu_{i;j,k}$ and y_{ij} be

$$\nu_{i;j,k} = \begin{cases} m_{i,j} - \dots - m_{i,k} & 1 \leq j \leq k \leq q_i \\ 1 & j > k, \end{cases}$$

and

$$y_{i,j} = \prod_{\substack{l=1 \\ l \neq i}}^r t_l^{\nu_{l;1,q} - \nu_{i;1,j-1}},$$

respectively for $i=1, \dots, q$. Then we have

LEMMA 2.1. The Alexander polynomial of L is

$$(2.4) \quad \Delta(L; t_1, \dots, t_r) = \left(\prod_{i=1}^r t_i^{\nu_{i;1,q_i} - 1} \right)^{r-2} \prod_{i=1}^r \prod_{j=1}^{q_i} Q(t_i^{\nu_{i;j+1,q_i}}, y_{ij}; m_{ij}, \lambda_{ij})$$

§3. Proof of THEOREM A.

Let $\Phi(t; m)$ be a polynomial of the form

$$(3.1) \quad \Phi(t; m) = t^{m-1} + t^{m-2} + \dots + t + 1,$$

where m is a positive integer. We call a polynomial of the form

(3.1) Φ -polynomial. Then a polynomial $Q(t, s; m, \lambda)$ is a Φ -polynomial

$$Q(t, s; m, \lambda) = \Phi(t^{\lambda} s; m).$$

Therefore the Alexander polynomial $\Delta(L; t_1, \dots, t_r)$ of the link L can be written as

$$(3.2) \quad \Delta(L; t_1, \dots, t_r) = \left(\prod_{i=1}^r t_i^{\xi_{i-1}} \right)^{r-2} \Phi \left(\prod_{i=1}^r t_i^{\xi_i}; g \right) \\ \prod_{i=1}^r \prod_{j=1}^{q_i} \Phi(t_i^{\lambda_{ij} \nu_{i;j+1, q_i}} y_{ij}; m_{ij}),$$

where $g = \text{g.c.m.} \{ \nu_{i;1, q_i} \}_{i=1}^r$ and $\xi_i = \nu_{i;1, q_i} / g$ for $i=1, \dots, r$.

Before proceeding the proof of THEOREM A, we preperate several lemmas. Let $\varphi_1 = \Phi(t_1^{\omega_1}, \dots, t_r^{\omega_r}; m)$ and $\varphi_2 = (t_1^{\omega'_1}, \dots, t_r^{\omega'_r}; m')$ for integers m and $m' \geq 2$. The following LEMMA 3.1 and LEMMA 3.2 are showed directly.

LEMMA 3.1. The product $\varphi_1 \cdot \varphi_2$ is a Φ -polynomial if and only if $\omega_i \neq \omega'_i$, and $\omega_i m = \omega'_i$ if $\omega_i < \omega'_i$ or $\omega_i = \omega'_i m'$ if $\omega_i > \omega'_i$, for all $i=1, \dots, r$.

Let $\varphi_i = \varphi_{i,1} \cdots \varphi_{i,u_i}$ ($i=1,2$) be a decomposition of φ_i by Φ -polynomials $\varphi_{i,j}$. We say that φ_1 and φ_2 are relatively prime if a product $\varphi_{1,j} \varphi_{2,j'}$ is not a Φ -polynomial for any $j=1, \dots, u_1$ and any $j'=1, \dots, u_2$.

We suppose that m is factored by prime integers as $m = a_1 \cdots a_u$. Then we have

LEMMA 3.2. A Φ -polynomial $\Phi(t; m)$ is factored by (irreducible) Φ -polynomials as

$$\Phi(t; m) = \prod_{j=1}^u \Phi(t^{\alpha_j}; a_j),$$

where $\alpha_1 = 1$ and $\alpha_j = a_1 \cdots a_{j-1}$ for $j=2, \dots, u+1$.

We note that an iterated torus knot K of type $\{(m_j, \lambda_j)\}_{j=1}^q$ is algebraic if and only if $n_{j-1}m_j < n_j$ for all $j=2, \dots, q$, where n_j are given by (2.1). Therefore, for all j and k such that $1 \leq j < k \leq q$, we have

$$(3.3) \quad \lambda_k > \lambda_j m_j \dots m_k.$$

To prove THEOREM A, it is sufficient to show that we can determine the knot types of components K_i ($i=1, \dots, r$) of L from the Alexander polynomial $\Delta(L; t_1, \dots, t_r)$ of L .

LEMMA 3.3. Let $\varphi_{ij} = (t_i^{\lambda_{i,j} \nu_{i;j+1, q_{ij} y_{ij}; m_{ij}})$ and $\varphi_{ik} = (t_i^{\lambda_{i,k} \nu_{i;k+1, q_{ik} y_{ik}; m_{ik}})$. Then φ_{ij} and φ_{ik} are relatively prime for $i=1, \dots, r$ and $1 \leq j < k \leq q_i$.

Proof. By (3.3), for all j and k such that $1 \leq j < k \leq q_i$,

$$\lambda_{ik} \nu_{i;k+1, q_i} > \lambda_{ij} \nu_{i;j+1, q_i}.$$

Then by LEMMA 3.1, φ_{ij} and φ_{ik} are relatively prime.

LEMMA 3.4. φ_{ij} and φ_{lk} are relatively prime for $i, l=1, \dots, r$, $i \neq l$, $j=1, \dots, q_i$ and $k=1, \dots, q_l$.

Proof. We suppose that m_{ij} and m_{lk} are factored by prime integers as $m_{ij} = a_1 \dots a_u$ and $m_{lk} = b_1 \dots b_v$ respectively. Let α_p and β_s be $\alpha_1 = 1$, $\alpha_p = a_1 \dots a_{p-1}$ ($2 \leq p \leq u+1$), $\beta_1 = 1$ and $\beta_s = b_1 \dots b_{s-1}$ ($2 \leq s \leq v+1$). We suppose that φ_{ij} and φ_{lk} are not relatively prime.

Then there are $p(1 \leq p \leq u)$ and $s(1 \leq s \leq v+1)$ such that

$$(3.4) \quad \lambda_{ij} \nu_{i;j+1, q_i} \alpha_p = \nu_{i;1, q_i} \nu_{l;1, k-1} \beta_s$$

or there are $p(1 \leq p \leq u+1)$ and $s(1 \leq s \leq v)$ such that

$$(3.5) \quad \nu_{i;1, q_i} \nu_{i;1, j-1} \alpha_p = \lambda_{lk} \nu_{l;k+1, q_l} \beta_s.$$

Since $\alpha_p < m_{ij}$ for $1 \leq p \leq u$ and $\beta_s < m_{lk}$ for $1 \leq s \leq v$, $\bar{\alpha}_p = m_{ij}/\alpha_p > 1$ and $\bar{\beta}_s = m_{lk}/\beta_s > 1$. Then by (3.4),

$$\lambda_{ij} = \bar{\alpha}_p \nu_{l;1, k-1} \beta_s$$

or by (3.5),

$$\lambda_{lk} = \bar{\beta}_s \nu_{i;1, j-1} \alpha_p$$

These contradict that $\text{g.c.d.}(m_{ij}, \lambda_{ij}) = 1$ or $\text{g.c.d.}(m_{lk}, \lambda_{lk}) = 1$. Then by LEMMA 3.1, φ_{ij} and φ_{lk} are relatively prime.

LEMMA 3.5. φ_{ij} and $\Phi(\prod_{i=1}^r t_i^{\xi_i}; g)$ are relatively prime for $i=1, \dots, r$, and $j=1, \dots, q_i$.

Proof. By (3.3) and since $m_{i,1} < \lambda_{i,1}$, for any i and any j ,

$$\nu_{i;1, q_i} < \lambda_{ij} \nu_{i;j+1, q_i}$$

Then by LEMMA 3.1, φ_{ij} and $\Phi(\prod_{i=1}^r t_i^{\xi_i}; g)$ are relatively prime.

By lemmas 3.3, 3.4 and 3.5, the Alexander polynomial $\Delta(L; t_1, \dots, t_r)$ of L is uniquely representd by the form (3.2).

Therefore we can uniquely determine the knot types of compo-

nents K_i of L from the exponents of t_i in (3.2). This completes the proof.

§4. Reduced Alexander polynomials.

Let l be a link of r components. Then the reduced Alexander polynomial $\Delta(l; t)$ is given by the equation $\Delta(l; t) = (t-1)\Delta(l; t, \dots, t)$ (see Milnor [3]). The following example shows that algebraic links can not be classified by reduced Alexander polynomials.

EXAMPLE 4.1. Let $f(z_0, z_1) = z_0(z_0^5 - z_1^6)$ and $f'(z_0, z_1) = (z_0^3 - z_1^2)(z_0^3 - z_1^{10})$. Then associated links L and L' (for f and f' respectively) have the same reduced Alexander polynomial $\Delta(L; t) = \Delta(L'; t) = (t-1)(t^6+1)$. But L and L' are not of the same link type.

Let Γ and Γ' be the Seifert matrices of L and L' respectively. From Amida-diagrams of L and L' (see [5]), we can compute Γ and Γ' . Then by computations with a computer, we have that the signatures $\sigma(L)$ of L and $\sigma(L')$ of L' are $\sigma(L)=19$ and $\sigma(L')=23$.

EXAMPLE 4.2. Let $g(z_0, \dots, z_n) = z_0(z_0^5 - z_1^6) + z_2^2 + \dots + z_n^2$ and $g'(z_0, \dots, z_n) = (z_0^3 - z_1^2)(z_0^3 + z_1^{10}) + z_2^2 + \dots + z_n^2$. Let K and K' be associated knots for g and g' respectively. Then Alexander polynomial $\Delta(K; t)$ and $\Delta(K'; t)$ are also equal to $(t-1)(t^6+1)$. But K and K' are not of the same knot type because signatures $\sigma(K)=19$ and $\sigma(K')=23$. Therefore algebraic knots K^{2n-1} can not be classified by Alexander polynomials.

The author wishes to thank Professors K. Nakajima and M. Takada for computation of signatures.

References

- [1] Lê Dũng Tráng; Sur les noeûds algebriques. Comp. Math. 25, 281-321 (1972).
- [2] J. Milnor: Singular points of complex hypersurfaces. Ann. of Math. Study 61, Princeton Univ. Press (1968).
- [3] F. Pham; Cours de 3ème cycle, Dep. Math. de la Faculté des Sciences de Paris, à paraître au Centre de Math. de l'Ecole Polytechnique.
- [4] D. W. Sumners and J. M. Woods: The monodromy of reducible plane curves, Inventiones Math. 40, 107+141 (1977).
- [5] M. Yamamoto: Amida-diagrams and Seifert matrices of positive iterated torus knots, (to appear).
- [6] _____: A topological classification of algebraic links, (to appear).

Department of Mathematics

School of Science and Engineering

Waseda University