

On a problem of Sakai in unbounded derivations

H. TAKAI (都立大理)

As a quantization of spaces, especially n -dimensional real lines, Sakai [7] posed the following interesting problem: are there simple C^* -algebras \mathcal{A} and n -family $\{\delta_i\}_{i=1}^n$ of non approximately bounded pregenerators of \mathcal{A} such that given a $*$ -derivation δ of \mathcal{A} with $D(\delta) = \bigcap_{i=1}^n D(\delta_i)$, there exist $k_1, k_2 \in \mathbb{R}$ and an approximately bounded $*$ -derivation δ_0 of \mathcal{A} with the property that $\delta = \sum_{i=1}^n k_i \delta_i + \delta_0$.

In this note, we show that there is at least one model for two dimensional case. It is nothing but the irrational rotation algebra, namely the C^* -crossed product $C(T) \times_{\theta} \mathbb{Z}$ of the C^* -algebra $C(T)$ of all continuous functions on the one dimensional torus T by an irrational angle θ . More precisely we have the following:

Theorem 1. Let \mathcal{A}_θ be the irrational rotation algebra. Then there exist two non approximately bounded pregenerators δ_1, δ_2 of \mathcal{A}_θ such that any

*-derivation δ of \mathcal{A}_0 with $D(\delta) = D(\delta_1) \cap D(\delta_2)$ can be expressed as $\delta = k_1\delta_1 + k_2\delta_2 + \delta_0$ for some $k_1, k_2 \in \mathbb{R}$ and an approximately bounded *-derivation δ_0 of \mathcal{A}_0 .

Remark 1. Suppose $D(\delta) = D(\delta_j)$ ($j=1$ or 2), then one can show that $\delta = k\delta_j + \delta_0$ for some $k \in \mathbb{R}$.

We now state our main theorem as follows:

Theorem 2. Let $(\mathcal{A}, \mathcal{G}, \alpha)$ be a C^* -dynamical system where \mathcal{A} is unital abelian, \mathcal{G} is discrete abelian, and α is effective. Suppose $\beta_t = \exp t\delta_0$ ($t \in T$) commuting with α , and there exists an eigenunitary u for β which generates \mathcal{A} . Then for any *-derivation δ of $\mathcal{A} \rtimes_{\alpha} \mathcal{G}$ with $D(\delta) = D(\tilde{\delta}_0) = D(\delta_0) \otimes_{\beta} \mathcal{G}$ there exist a $k \in \mathbb{R}$, pre-generator δ_1 and an approximately bounded *-derivation δ_2 of $\mathcal{A} \rtimes_{\alpha} \mathcal{G}$ such that (i) $D(\delta_j) = D(\delta)$ ($j=1,2$) $\delta_1|_{\mathcal{A}} = 0$, δ_1 commutes with $\tilde{\delta}_0$ (ii) $\delta = k\tilde{\delta}_0 + \delta_1 + \delta_2$, where $D(\delta_0) \otimes_{\beta} \mathcal{G}$ is the set of all $D(\delta_0)$ -valued function of \mathcal{G} with finite support, and $\tilde{\delta}_0(x)(\eta) = \delta_0[x(\eta)]$ ($x \in D(\delta_0) \otimes_{\beta} \mathcal{G}$).

Remark 2. If $\mathcal{G} = \mathbb{Z}$, $\delta_1 = l\delta_1'$ for some $l \in \mathbb{R}$ where δ_1' is independent of δ .

Let (\mathcal{A}, G, α) and (\mathcal{A}, H, β) be two C^* -dynamical systems where α, β commute. Then there is a C^* -dynamical system $(\mathcal{A} \rtimes_{\alpha} G, H, \tilde{\beta})$ such that $\tilde{\beta}_t(x)(g) = \beta_t[x(g)]$ ($x \in L^1(G; \mathcal{A})$). Then we have the following proposition of fixed point type:

Proposition 3. $(\mathcal{A} \rtimes_{\alpha} G)^{\tilde{\beta}} = \mathcal{A} \rtimes_{\alpha} G$

Proof. By definition, $\mathcal{A} \rtimes_{\alpha} G \subset (\mathcal{A} \rtimes_{\alpha} G)^{\tilde{\beta}}$.

Suppose the inclusion is proper, then $(\mathcal{A} \rtimes_{\alpha} G) \times_{\hat{\alpha}} \hat{G} \subsetneq (\mathcal{A} \rtimes_{\alpha} G)^{\tilde{\beta}} \times_{\hat{\alpha}} \hat{G}$ since $\tilde{\beta}$ commutes with $\hat{\alpha}$.

Since $(\mathcal{A} \rtimes_{\alpha} G)^{\tilde{\beta}} \times_{\hat{\alpha}} \hat{G} \subset ((\mathcal{A} \rtimes_{\alpha} G) \times_{\hat{\alpha}} \hat{G})^{\tilde{\beta}}$, it follows from duality [6, 8] that $\mathcal{A} \rtimes_{\alpha} \mathcal{C}(L^2(G)) \subsetneq (\mathcal{A} \rtimes_{\alpha} \mathcal{C}(L^2(G)))^{\beta \otimes \alpha}$ which is a contradiction. Q. E. D.

Comment 1. We only consider locally compact abelian groups throughout this note.

In what follows, let δ be a $*$ -derivation of $\mathcal{A} \rtimes_{\alpha} G$ such that $D(\delta) = D(\delta_0)$ where δ_0 is a generator of \mathcal{A} commuting with α . Suppose δ commutes with $\hat{\alpha}$, and G is discrete. Then $\delta(a) \in \mathcal{A}$ for $a \in D(\delta_0)$. Let $(x_n)_n \subset D(\delta)$ with $x_n \rightarrow 0$, $\delta(x_n) \rightarrow y \in \mathcal{A} \rtimes_{\alpha} G$. Since $x_n = \sum_k a_k^{(n)} \lambda(k)$ ($a_k^{(n)} \in D(\delta_0)$), using the conditional expectation ε of $\mathcal{A} \rtimes_{\alpha} G$ onto \mathcal{A}

one has $\varepsilon(x_n \lambda(g)^*) \rightarrow 0$ and $\varepsilon[(\delta(x_n) - \gamma) \lambda(g)^*] \rightarrow 0$ for each g in \mathcal{G} . Thus $a_g^{(n)} \rightarrow 0$ and $\varepsilon[\sum_k (\delta(a_k^{(n)}) \lambda(k-g) + a_k^{(n)} \delta(\lambda(k)) \lambda(g)^* - \gamma_k \lambda(k-g))] \rightarrow 0$ where $\gamma = \sum_k \gamma_k \lambda(k)$ is the Fourier expansion of γ in $\mathcal{O} \times_s \mathcal{G}$ ($\gamma_k \in \mathcal{O}$). Then $a_g^{(n)} \rightarrow 0$ and $\delta(a_g^{(n)}) \rightarrow \gamma_g$ for all g in \mathcal{G} . Since $\mathcal{D}(\delta|_{\mathcal{O}}) = \mathcal{D}(\delta_0)$, it follows from Batty's result [2] that $\delta|_{\mathcal{O}}$ is closable. So $\gamma_g = 0$ for all $g \in \mathcal{G}$. Consequently we have the following:

Lemma 4. If \mathcal{G} is discrete, any $*$ -derivation δ of $\mathcal{O} \times_s \mathcal{G}$ such that (i) $\mathcal{D}(\delta) = \mathcal{D}(\delta_0)$ and (ii) δ commutes with $\hat{\alpha}$ is closable.

Remark 3. In the above lemma, the conclusion is unclear unless the condition (ii) is added.

Now let δ be a $*$ -derivation of $\mathcal{O} \times_s \mathcal{G}$ with $\mathcal{D}(\delta) = \mathcal{D}(\delta_0)$. Define $\mathcal{J} = \{x \in \mathcal{D}(\delta_0) \mid a \mapsto \delta(ax) \text{ is continuous from } \mathcal{D}(\delta_0) \text{ into } \mathcal{O} \times_s \mathcal{G}\}$. Since $\delta(a \lambda(g) b) = \delta(\lambda(g)) \alpha_g^{-1}(a) b + \lambda(g) \delta(\alpha_g^{-1}(a) b)$ and δ_0 commutes with α , we have $x \lambda(g) b = 0$ for all $g \in \mathcal{G}$ and $b \in \mathcal{J}$ if $a_n \in \mathcal{D}(\delta_0) \rightarrow 0$ and $\delta(a_n) \rightarrow x \in \mathcal{O} \times_s \mathcal{G}$. Then $\varepsilon(x \lambda(g)) b = 0$ where ε is the projection of norm one from $\mathcal{O} \times_s \mathcal{G}$ onto \mathcal{O} . So $\varepsilon(x \lambda(g)) \in L(\mathcal{J})$, the left annihilator

of \mathcal{I} . Since \mathcal{I} is a two sided ideal of $\mathcal{D}(\delta_0)$, it follows from the same way as Longo [4] that $L(\mathcal{I}) = 0$. Thus $\varepsilon(x\lambda(f)) = 0$ for all $f \in \mathcal{G}$. Let $x = \sum_g x_g \lambda(f)$ be the Fourier expansion of x . Then $x_g = 0$. So $x = 0$. Then $\delta|_{\mathcal{A}}$ is closable from $(\mathcal{D}(\delta_0), \|\cdot\|_{\delta_0})$ into $\mathcal{A} \times \mathcal{G}$. Therefore we have the following:

Lemma 5. Let δ be a $*$ -derivation of $\mathcal{A} \times \mathcal{G}$ with $\mathcal{D}(\delta) = \mathcal{D}(\tilde{\delta}_0)$. Then δ is relatively bounded on $\mathcal{D}(\delta_0)$ with respect to δ_0 , namely $\|\delta(a)\| \leq K(\|a\| + \|\delta_0(a)\|)$ for all $a \in \mathcal{D}(\delta_0)$, with some positive constant K .

Remark 4. Since δ_0 is a pregenerator, one can not directly apply Longo's result. However the crucial part of the above proof is due to his idea [4].

By the above lemma, let $\beta_t = \exp t\delta_0$ ($t \in \mathbb{R}$). Then there exist derivations $\tilde{\delta}_f$ ($f \in L^1(\mathbb{R})$) of $\mathcal{A} \times \mathcal{G}$ such that (i) $\mathcal{D}(\tilde{\delta}_f) = \mathcal{D}(\tilde{\delta}_0)$ and (ii) $\tilde{\delta}_f = \int_{\mathbb{R}} f(t) \tilde{\beta}_t \circ \delta \circ \tilde{\beta}_{-t} dt$. In fact, since $\|\delta(a)\| \leq M(\|a\| + \|\delta_0(a)\|)$ for $a \in \mathcal{D}(\delta_0)$, $\|\delta \circ \beta_t(a) - \delta \circ \beta_s(a)\| \leq M\{\|\beta_t(a) - \beta_s(a)\| + \|\beta_t \circ \delta_0(a) - \beta_s \circ \delta_0(a)\|\}$. So $t \mapsto \delta \circ \beta_t(a)$ is continuous for each $a \in \mathcal{D}(\delta_0)$. Thus $t \mapsto \delta \circ \beta_t(x)$ is also continuous for $x \in \mathcal{D}(\tilde{\delta}_0)$ which gives

derivations \tilde{S}_f for $f \in L^1(\mathbb{R})$ of $\mathcal{O}X_\alpha \mathcal{G}$ satisfying (i) and (iii). Similarly, for each $g \in \mathcal{G}$ one has a derivation \hat{S}_g of $\mathcal{O}X_\alpha \mathcal{G}$ such that (i) $D(\hat{S}_g) = D(\tilde{\delta}_0)$ and (iii) $\hat{S}_g = \int_{\hat{\mathcal{G}}} \langle \tilde{F}, P \rangle \hat{\alpha}_P \cdot \delta \cdot \hat{\alpha}_P^{-1} dp$. Moreover suppose $P_t = \exp t \delta_0$ is periodic, then we have that $(\hat{S}_1)_0^\sim = (\tilde{S}_0)_1^\wedge$ commutes with $\hat{\alpha}$, $\tilde{\beta}$. In what follows we treat $*$ -derivations of $\mathcal{O}X_\alpha \mathcal{G}$ with the same domain as $D(\tilde{\delta}_0)$ commuting with $\hat{\alpha}$ and $\tilde{\beta}$, which are denoted by δ . Since it commutes with $\hat{\alpha}$, it follows from Lemma 4 that it is closable. Hence one may assume that it is closed. Let $x \in C^*(\mathcal{G})$, and $(x_i) \subset D(\delta)$ which converge to x . Put $y_i = \int_T \tilde{\beta}_t(x_i) dt \in \mathcal{O}X_\alpha \mathcal{G}$. Since δ commutes with $\tilde{\beta}$ and δ is closed, $y_i \in D(\delta) \cap (\mathcal{O}X_\alpha \mathcal{G})^{\tilde{\beta}}$ and $y_i \rightarrow x$ since $(\mathcal{O}X_\alpha \mathcal{G})^{\tilde{\beta}} = C^*(\mathcal{G})$ by Proposition 1. So $\delta|_{C^*(\mathcal{G})}$ is a closed $*$ -derivation of $C^*(\mathcal{G})$ since $\delta(y_i) \in C^*(\mathcal{G})$. Since $\hat{\alpha}_P \cdot \delta \cdot \hat{\alpha}_P^{-1} = \delta$ for $P \in \hat{\mathcal{G}}$ and $\tilde{F} \cdot \hat{\alpha}_P \cdot \tilde{F}^{-1} = \tau_P$ on $C(\hat{\mathcal{G}})$, $\hat{\delta} = \tilde{F} \cdot \delta \cdot \tilde{F}^{-1}$ commutes with τ on $C(\hat{\mathcal{G}})$ where \tilde{F} is the Fourier isomorphism of $C^*(\mathcal{G})$ onto $C(\hat{\mathcal{G}})$, and τ is the shift action of $\hat{\mathcal{G}}$ on $C(\hat{\mathcal{G}})$. It follows from Goodman-Nakazato [3, 5] that there exists a one parameter subgroup (P_t) of $\hat{\mathcal{G}}$ such that $\hat{\delta}(f)(P) = \lim_{t \rightarrow 0} t^{-1}(f(P_t P))$

$- f(P))$ for all $f \in \mathcal{D}(\hat{\delta})$. Since $\langle \vartheta, \cdot \rangle \in \mathcal{D}(\hat{\delta})$, one has $\delta(\lambda(\vartheta)) = \partial(\vartheta)\lambda(\vartheta)$ for all $\vartheta \in \mathcal{G}$ where $\partial(\vartheta) = \lim_{t \rightarrow 0} (\langle \vartheta, P_t \rangle - 1)$. Let $\delta_1(a\lambda(\vartheta)) = \partial(\vartheta)a\lambda(\vartheta)$ for all $a \in \mathcal{D}(\delta_0)$ and $\vartheta \in \mathcal{G}$. Then it is a pregenerator of $\mathcal{A} \rtimes_{\alpha} \mathcal{G}$ such that $\mathcal{D}(\delta_1) = \mathcal{D}(\tilde{\delta}_0)$ and $\delta_1|_{\mathcal{A}} = 0$, δ_1 commutes with $\tilde{\delta}_0$. Since δ is a closed $*$ -derivation of $\mathcal{A} \rtimes_{\alpha} \mathcal{G}$ and $\delta|_{\mathcal{A}}$ commutes with $\beta_t = \exp t\delta_0$, it follows from Batty [1] that $\delta|_{\mathcal{A}} = k\delta_0$ for some $k \in \mathbb{R}$. Therefore we have that $\delta(a\lambda(\vartheta)) = k\delta_0(a)\lambda(\vartheta) + a\partial(\vartheta)\lambda(\vartheta) = (k\tilde{\delta}_0 + \delta_1)(a\lambda(\vartheta))$, which implies the following lemma:

Lemma 6. Let $(\mathcal{A}, \mathcal{G}, \alpha)$ be a C^* -dynamical system where \mathcal{A} is unital abelian and \mathcal{G} is discrete abelian. Let $\beta_t = \exp t\delta_0$ be a periodic action of \mathbb{R} on \mathcal{A} . Suppose β is ergodic, then given a $*$ -derivation δ of $\mathcal{A} \rtimes_{\alpha} \mathcal{G}$ with the property that (i) $\mathcal{D}(\delta) = \mathcal{D}(\tilde{\delta}_0)$ and (ii) δ commutes with $\hat{\alpha}, \tilde{\beta}$, there exist a $k \in \mathbb{R}$ and a pregenerator δ_1 of $\mathcal{A} \rtimes_{\alpha} \mathcal{G}$ such that (i) $\mathcal{D}(\delta_1) = \mathcal{D}(\delta)$, $\delta_1|_{\mathcal{A}} = 0$, δ_1 commutes with $\tilde{\delta}_0$, and (ii) $\delta = k\tilde{\delta}_0 + \delta_1$ on $\mathcal{D}(\tilde{\delta}_0)$.

Remark 5. The pregenerator δ_1 defined above would

be written as $\delta_1 = \gamma \delta_1'$ for some $\gamma \in \mathbb{R}$ where δ_1' is not depending on δ . Actually if $\mathcal{G} = \mathbb{Z}$, we have $\delta_1'(a \lambda(n)) = i n a \lambda(n)$ for $a \in \mathcal{D}(\delta_0)$ and $n \in \mathbb{Z}$.

Let δ be a linear mapping from a $*$ -subalgebra $\mathcal{D}(\delta)$ of \mathcal{A} into \mathcal{A} such that $\delta(ab) = \delta(a) \alpha_g(b) + a \delta(b)$ for all $a, b \in \mathcal{D}(\delta)$ where $g \neq e \in \mathcal{G}$ is a fixed element. Suppose there is a unitary u of $\mathcal{D}(\delta)$ such that $1 \notin \text{Sp}(\alpha_g(u) u^*)$, then we have by direct computation that $\delta(u^n) = \sum_{k=0}^{n-1} \alpha_g(u^k) u^k \delta(u) u^{n-1-k}$. Since $1 \notin \text{Sp}(\alpha_g(u) u^*)$, one has that $\sum_{k=0}^{n-1} \alpha_g(u^k) u^k = (\alpha_g(u^n) u^n - 1) (\alpha_g(u) u^* - 1)^{-1}$. So $\delta(u^n) = \delta(u) u^* (\alpha_g(u) u^* - 1)^{-1} (\alpha_g - \text{id})(u^n) = \delta(u) (\alpha_g(u) - u)^{-1} (\alpha_g - \text{id})(u^n)$ for all $n \in \mathbb{Z}$ since $\delta(1) = 0$. Put $a_g = \delta(u) (\alpha_g(u) - u)^{-1} \in \mathcal{A}$. Since $a_g (\alpha_g - \text{id})$ is bounded on \mathcal{A} , the conclusion follows. Namely we have the following:

Lemma 7. Suppose \mathcal{A} is unital abelian and \mathcal{G} is discrete. Let δ be a linear mapping of a $*$ -subalgebra $\mathcal{D}(\delta)$ of \mathcal{A} into \mathcal{A} such that $\delta(ab) = \delta(a) \alpha_g(b) + a \delta(b)$ for $a, b \in \mathcal{D}(\delta)$ for a fixed $g \neq e$. Suppose there exists a unitary $u \in \mathcal{D}(\delta)$ such that $1 \notin \text{Sp}(\alpha_g(u) u^*)$, then $\delta = a_g (\alpha_g - \text{id})$ on $\mathcal{D}(\delta) \cap C^*(u)$.

for some $a_g \in \mathcal{A}$.

Remark 6. By the above lemma, there is no unbounded α_g -cocycle closed $*$ -derivation if α_g has an eigenunitary generating \mathcal{A} .

Now let $\hat{\beta}_g$ ($g \in \mathcal{G}$) be a derivation of $\mathcal{A} \rtimes_{\alpha} \mathcal{G}$ as in the previous way (following to Remark 4). Then it implies that $\delta = \sum_g \hat{\beta}_g$ on $\mathcal{D}(\delta)$. In fact, let $\delta(a) = \sum_g \delta(a)(g) \lambda(g)$ and $\delta(\lambda(h)) = \sum_g \delta(\lambda(h))(g) \lambda(g)$ be the Fourier expansion of $\delta(a)$ and $\delta(\lambda(h))$ respectively.

Then $\hat{\beta}_g(a) = \delta(a)(g) \lambda(g)$ and $\hat{\beta}_g(\lambda(h)) = \delta(\lambda(h))(g+h) \lambda(g+h)$.

Suppose δ commutes with $\tilde{\beta}$, it follows from Lemma 6

that $\hat{\beta}_e = k \tilde{\delta}_0 + \delta_1$ on $\mathcal{D}(\delta)$ where k, δ_1 are as in

Lemma 6. Let $\delta_g(a) = \hat{\beta}_g(a) \lambda(g)^*$ for $a \in \mathcal{D}(\delta_0)$ ($g \neq e$).

Then δ_g satisfy the condition of Lemma 7. Suppose

there exists a unitary $u \in \mathcal{D}(\delta_0)$ such that (i) $1 \notin \text{Sp}(\alpha_g(u)u^*)$ ($g \neq e$) and (ii) $\mathcal{A} = C^*(u)$. Since δ commutes

with $\tilde{\beta}$, and α commutes with $\beta_h = \exp t \delta_0$ which is ergodic, we have $a_g \in \mathbb{C}1$. Then $\hat{\beta}_g(a) = a_g (\alpha_g - \text{id})(a) \lambda(g) = [a_g \lambda(g), a]$. Hence $\hat{\beta}_g(a \lambda(h)) = \hat{\beta}_g(a) \lambda(h) + a \hat{\beta}_g(\lambda(h)) = [a_g \lambda(g), a \lambda(h)] + a \hat{\beta}_g(\lambda(h))$. Since $\hat{\beta}_g - \text{ad}(a_g \lambda(h))$ is a derivation on $\mathcal{D}(\delta)$, one has $\hat{\beta}_g(\lambda(h)) = 0$ for $h \in \mathcal{G}$.

(In fact, since $\hat{S}_g(\lambda(h)) = \delta(\lambda(h))(g+h)\lambda(g+h)$, we have that $\delta(\lambda(h+k))(h+k+g)u = \delta(\lambda(h))(h+g)\alpha_g(u) + \delta(\lambda(k))(k+g)u$ for all $h, k \in G$. Since $1 \in D(\delta)$, we have $\delta(1)(g) = 0$. So $\delta(\lambda(h))(h+g) = 0$ for all $h \in G$ or $\alpha_g(u) = u$. Since $1 \notin \text{Sp}(\alpha_g(u)u^*)$, we have $\delta(\lambda(h))(h+g) = 0$ for all $h \in G$. Consequently $\delta = k\tilde{\delta}_0 + \delta_1 + \sum_{g \neq e} \text{ad}(a_g \lambda(g))$ on $D(\delta)$. Set $\delta_H = \text{ad}(\sum_{g \in H} a_g \lambda(g))$ for a finite set H of $G - \{e\}$ with $H = -H$. Then δ_H are bounded $*$ -derivations of $\mathcal{O}X_\alpha G$ such that $\delta_H(\lambda(h)) = 0$ and δ_H converges to δ_2 pointwisely on $D(\delta)$ where $\delta_2(a\lambda(h)) = \sum_{g \neq e} [a_g \lambda(g), a\lambda(h)] (= (\delta - \hat{S}_e)(a)\lambda(h))$. Then $\delta = k\tilde{\delta}_0 + \delta_1 + \delta_2$ on $D(\delta)$ and $\delta_2(\lambda(g)) = 0$ for all $g \in G$, which implies the following proposition:

Proposition B. Let (\mathcal{A}, G, α) be a C^* -dynamical system where \mathcal{A} is unital abelian and G is discrete. Let $\beta_t = \exp t\tilde{\delta}_0$ be an ergodic action of T on \mathcal{A} commuting with α . Suppose there exists a unitary $u \in D(\tilde{\delta}_0)$ such that (i) $1 \notin \text{Sp}(\alpha_g(u)u^*)$ ($g \neq e$), (ii) $\mathcal{A} = C^*(u)$, then given a $*$ -derivation δ of $\mathcal{O}X_\alpha G$ such that (i) $D(\delta) = D(\tilde{\delta}_0)$ and (ii) δ commutes with $\tilde{\beta}$, there exist a $k \in \mathbb{R}$, a pregenerator δ_1 and an approximately bounded $*$ -derivation δ_2 of $\mathcal{O}X_\alpha G$ such that (i) $D(\delta_1) = D(\delta)$, $\delta_1|_{\mathcal{A}} = 0$, δ_1 commutes with $\tilde{\delta}_0$,

ii) $D(\delta_2) = D(\delta)$, $\delta_2(\lambda(g)) = 0$ for all $g \in G$, and iii) $\delta = k\tilde{\delta}'_0 + \delta_1 + \delta_2$.

Remark 7. In the case of discrete abelian groups, the Fourier expansion of any element of $\mathcal{M}X_\alpha G$ can be taken in the uniform sense. In fact, taking a net $\{f_i\}$ of positive definite functions on G with finite support converging to 1, one can show that $\sum_g f_i(g) a_g \lambda(g)$ converges to $\sum_g a_g \lambda(g) \in \mathcal{M}X_\alpha G$ uniformly.

Proof of Theorem 2: Since β commutes with α and β is ergodic, we have $\alpha_g(u)u^* \in \mathbb{C}1$. Since $\mathcal{M} = C^*(u)$ and α is effective, there are $c_g \neq 1$ ($g \neq e$) such that $\alpha_g(u) = c_g u$. So $1 \notin sp(\alpha_g(u)u^*)$ ($g \neq e$). Let $\tilde{F}_n = \int_T e^{-int} \tilde{\beta}_t \circ \delta \circ \tilde{\beta}_t^{-1} dt$ on $D(\delta)$ for $n \in \mathbb{Z}$. Since \tilde{F}_0 commutes with $\tilde{\beta}$, it follows from Proposition 8 that $\tilde{F}_0 = k\tilde{\delta}'_0 + \delta'_1 + \delta'_2$ where δ'_i, k are as in Proposition 8. Since $\tilde{\beta}_t \circ \tilde{F}_n \circ \tilde{\beta}_t^{-1} = e^{int} \tilde{F}_n$ ($n \in \mathbb{Z}$), $\tilde{\beta}_t \circ \tilde{F}_n(\lambda(g)) = e^{itn} \tilde{F}_n(\lambda(g))$. Since $\tilde{\beta}_t(u^n) = e^{itn} u^n$, we have that $u^{-n} \tilde{F}_n(\lambda(g)) \in (\mathcal{M}X_\alpha G)^{\tilde{\beta}} = C^*(G)$. So there are $b(n, g) \in C^*(G)$ such that $\tilde{F}_n(\lambda(g)) = u^n b(n, g)$. Let $\delta(\lambda(g)) = \sum_k \delta(\lambda(g))(k) \lambda(k)$ and $b(n, g) = \sum_k b(n, g)(k) \lambda(k)$ be the Fourier expansion of $\delta(\lambda(g))$ and $b(n, g)$ respectively.

Since $\mathcal{A} = C^*(U)$ and $\beta_t(U) = e^{it}U$, we have that $\delta(\lambda(\mathcal{F})(h))$
 $= a(0) + \sum_{n \neq 0} b(n, \mathcal{F})(h) U^n$ where $a(0)$ is the 0-component of
 the expansion of $\delta(\lambda(\mathcal{F})(h))$ in \mathcal{A} . Since $\tilde{\mathcal{S}}_0 = k\tilde{\delta}_0 + \delta_1' + \delta_2'$,
 one has $\tilde{\mathcal{S}}_0(\lambda(\mathcal{F})) = \partial(\mathcal{F})\lambda(\mathcal{F})$. By unicity, $\int_T \beta_t(\delta(\lambda(\mathcal{F})(h))) dt$
 $= \partial(\mathcal{F})1$ ($\mathcal{F} \neq h$), $= 0$ (otherwise), which is nothing but $a(0)$.
 Therefore we deduce that $\delta(\lambda(\mathcal{F})) = \partial(\mathcal{F})\lambda(\mathcal{F}) + \sum_h \sum_{n \neq 0} b(n, \mathcal{F})(h) \times$
 $U^n \lambda(h) = \partial(\mathcal{F})\lambda(\mathcal{F}) + \sum_{n \neq 0} U^n b(n, \mathcal{F}) = \partial(\mathcal{F})\lambda(\mathcal{F}) + \sum_{n \neq 0} \tilde{\mathcal{S}}_n(\lambda(\mathcal{F}))$.
 moreover $\delta(a) = \sum_g \hat{\mathcal{S}}_g(a)$ for all $a \in \mathcal{D}(\delta_0)$. (It follows
 from Lemma 7 that $\hat{\mathcal{S}}_g(a) = f_g(\alpha_g - id)(a)\lambda(\mathcal{F})$ for some
 $f_g \in \mathcal{A}$ ($g \neq e$). So $\hat{\mathcal{S}}_g(a) = [f_g\lambda(\mathcal{F}), a]$ for all $a \in \mathcal{D}(\delta_0)$.
 Since $\hat{\mathcal{S}}_e$ commutes with $\hat{\alpha}$, we have $\hat{\mathcal{S}}_e(a) \in \mathcal{A}$ for all
 $a \in \mathcal{D}(\delta_0)$. Since $(\hat{\mathcal{S}}_e)_0$ commutes with $\hat{\alpha}$ and $\hat{\beta}$, it means
 that $(\hat{\mathcal{S}}_e)_0 = k\tilde{\delta}_0 + \delta_1$ where k, δ_1 are as in Lemma 6.
 Then $\int_T e^{-it} \beta_t \circ \hat{\mathcal{S}}_e(U) dt = k\delta_0(U)$. Since $\beta_t(U) = e^{it}U$,
 we have $\delta_0(U) = iU$. Let $\hat{\mathcal{S}}_e(U) = \sum_n a_n U^n$ ($a_n \in \mathbb{C}$). Then
 $a_1 = ik$. Therefore $\hat{\mathcal{S}}_e(U) = k\delta_0(U) + \sum_{n \neq 1} a_n U^n$. Since $\hat{\mathcal{S}}_e$
 is a *-derivation, we deduce that $\hat{\mathcal{S}}_e(U^2) = 2\hat{\mathcal{S}}_e(U)U =$
 $2kn\delta_0(U)U^{n-1} + \sum_{m \neq 1} 2na_m U^{m+n-1} = k\delta_0(U^2) + \sum_{m \neq 1} 2na_m U^{m+n-1}$.
 Hence $\hat{\mathcal{S}}_e(U^2)\lambda(\mathcal{F}) = k\tilde{\delta}_0(U^2)\lambda(\mathcal{F}) + \sum_{m \neq 1} 2na_m U^{m+n-1}\lambda(\mathcal{F})$.
 Consequently, we have that $\delta(U^2\lambda(\mathcal{F})) = \delta(U^2)\lambda(\mathcal{F}) + U^2\delta(\lambda(\mathcal{F}))$
 $= (k\tilde{\delta}_0 + \delta_1)(U^2\lambda(\mathcal{F})) + \sum_{h \neq e} [f_h\lambda(h), U^2]\lambda(\mathcal{F}) + \sum_{m \neq 0} U^2 \tilde{\mathcal{S}}_m(\lambda(\mathcal{F}))$

$+ \sum_{m \neq 0} n a_{m+1} u^{n+m} \lambda(g)$. Since $\delta - k \tilde{\delta}_0 - \delta_1$ is a $*$ -derivation, so is $\sum_{h \neq e} [f_h \lambda(h), u^n] \lambda(g) + \sum_{m \neq 0} u^{n+m} \tilde{S}_m(\lambda(g)) + \sum_{m \neq 0} n a_{m+1} u^{n+m} \lambda(g)$. Since $\text{ad}(f_h \lambda(h))(u^n \lambda(g)) + u^n \text{ad}(f_h \lambda(h))(\lambda(g)) = \text{ad}(f_h \lambda(h))(u^n \lambda(g))$, we deduce that $u^n (\sum_{m \neq 0} \tilde{S}_m(\lambda(g)) - \sum_{h \neq e} [f_h \lambda(h), \lambda(g)]) + \sum_{m \neq 0} n a_{m+1} u^{n+m} \lambda(g)$ is a $*$ -derivation. Let $a = \sum_{m \neq 0} a_{m+1} u^m \in \mathcal{O}$. Conventionally put $\sigma(\lambda(g)) = \sum_{m \neq 0} \tilde{S}_m(\lambda(g)) - \sum_{h \neq e} [f_h \lambda(h), \lambda(g)]$. Moreover, put $\Delta(u^n \lambda(g)) = u^n \sigma(\lambda(g)) + n a u^n \lambda(g)$. Since $\delta_0(u^n) = i n u$, we see $n a u^n \lambda(g) = (-i) a \tilde{\delta}_0(u^n \lambda(g))$. Now since $\Delta(u^n \lambda(g) u^m \lambda(h)) = \Delta(u^n \lambda(g)) u^m \lambda(h) + u^n \lambda(g) \Delta(u^m \lambda(h))$, we can show that $u^{n+m} (\sigma(\lambda(g)) \lambda(h) - \lambda(g) \sigma(\lambda(h))) = m (\alpha_g(a) - a) u^{n+m} \lambda(g+h)$. Put $h=e$ and $m=1$. Then we have $u^n \sigma(\lambda(g)) = (\alpha_g(a) - a) u^n \lambda(g)$ for all $n \in \mathbb{Z}$ and $g \in G$. Therefore $\Delta(u^n \lambda(g)) = (\alpha_g(a) - a) u^n \lambda(g) + n a u^n \lambda(g) = (\alpha_g(a) + (n-1)a) u^n \lambda(g)$. Since Δ is a derivation, we get $\alpha_g(a) = a$ for all $g \in G$. So $a = \varepsilon 1$ for some $\varepsilon \in \mathbb{C}$. Then $\Delta(u^n \lambda(g)) = \varepsilon n u^n \lambda(g) = i \varepsilon \tilde{\delta}_0(u^n \lambda(g))$. Finally, we obtain that $\delta(u^n \lambda(g)) = (c \tilde{\delta}_0 + \delta_1)(u^n \lambda(g)) + \sum_{h \neq e} [f_h \lambda(h), u^n \lambda(g)]$ for some $c \in \mathbb{R}$. Let $\delta_F(a \lambda(g)) = \sum_{h \in F} [f_h \lambda(h), a \lambda(g)]$ for $a \in \mathcal{D}(\delta_0)$ and $g \in G$ where F is a finite set of $G - \{e\}$ with $F = -F$. Then δ_F is a bounded $*$ -derivation of $\mathcal{O} \times_2 G$ for all F and $\delta_F \rightarrow \delta_2$ pointwisely. Hence δ_2 is approximately bounded. This completes the proof.

References

- [1] G. J. K. Batty : *Derivations on compact spaces*, Preprint, 1978.
- [2] G. J. K. Batty : *Small perturbations of C^* -dynamical systems*, Preprint, 1978.
- [3] F. M. Goodman : *Translation invariant closed $*$ -derivations*, Preprint, 1980.
- [4] R. Longo : *Automatic relative boundedness of derivations in C^* -algebras*, *J. Func. Anal.*, 34 (1979), 21-28.
- [5] H. Nakazato : *Closed $*$ -derivations on compact groups*, Preprint, 1980.
- [6] G. K. Pedersen : *C^* -algebras and their automorphism groups*, Academic Press, 1979.
- [7] S. Sakai : *Derivations in operator algebras*, Lecture Notes, 1979.
- [8] H. Takai : *On a duality for crossed products of C^* -algebras*, *J. Func. Anal.*, 19 (1975), 25-39.