

A remark on the continuous variation of  
secondary characteristic classes for foliations

By Shigeyuki MORITA\*

§1. Introduction.

Thurston [3] has constructed a family of codimension  $n$  foliations on certain  $(2n + 1)$  manifold with continuously varying Godbillon-Vey invariant, establishing a surjection,

$$gv : H_{2n+1}(\mathbb{B}\Gamma_n ; \mathbb{Z}) \rightarrow \mathbb{R} \rightarrow 0.$$

The purpose of this note is to show that, using his results, we can show that some secondary characteristic classes other than that of Godbillon-Vey vary also continuously. We can also show that these classes are independent. Thus we can construct a surjection

$$H_{2n+1}(\mathbb{B}\Gamma_n ; \mathbb{Z}) \rightarrow \mathbb{R}^{r(n)+1} \rightarrow 0$$

where  $r(n)$  is a number depending on  $n$ . (See §3).

The foliations which we are going to construct to realize some characteristic classes are product foliations. The reason why these classes vary continuously is that they are in some sense "decomposable." The Godbillon-Vey class is "indecomposable" in the sense that it vanishes on any product foliation. In §3, we shall remark that there are infinitely many series of indecomposable classes, first of which are those of Godbillon-Vey.

To evaluate the characteristic classes on product foliations, we need the "Cartan formula" for the secondary classes. This will be done

---

\* Partially supported by N S F grant MPS 72-05055 A02.

in §2.

The author would like to express his hearty thanks to Dr. Liang for many conversations and to Dr. Lazarov for helpful suggestions. Actually the formulation of the Cartan formula in §2 is due to him.

## §2. The Cartan formula.

Suppose we are given two foliations  $(M, \mathcal{F})$  and  $(N, \mathcal{G})$  of codimensions  $p$  and  $q$  respectively. Then we can construct the product foliation  $(M \times N, \mathcal{F} \times \mathcal{G})$ . The classifying map for this foliation factors through  $B\Gamma_p \times B\Gamma_q$ ;

$$M \times N \longrightarrow B\Gamma_p \times B\Gamma_q \xrightarrow{\mu} B\Gamma_n$$

where  $n = p + q$  and  $\mu$  is induced from the natural inclusion  $\Gamma_p \times \Gamma_q \subset \Gamma_{p+q}$ .

To calculate the characteristic classes of the product foliation  $(M \times N, \mathcal{F} \times \mathcal{G})$ , it is necessary to identify the image of the classes in  $H^*(B\Gamma_n; \mathbb{R})$  under the homomorphism  $\mu^*$ . To do this, let us define characteristic classes for codimension  $n$  foliations in a slightly different way from that of Bott [2]\*. Recall that Bott used the class  $c_i$ , which is the degree  $2i$  part of  $\det(I - \frac{1}{2\pi}\Omega)$  where  $\Omega$  is the curvature matrix. We simply replace  $c_i$  by  $\Sigma_i = (\frac{-1}{2\pi})^i \text{Tr } \Omega^i$  and let  $\gamma_{2i+1}$  be the form corresponding to  $h_{2i+1}$  in the Bott's definition. Let  $W\Omega'_n$  be a differential graded algebra defined by

\* This idea is due to Lazarov.

$$WO'_n = \hat{\mathbb{R}} [\Sigma_1, \dots, \Sigma_n] \otimes E(\gamma_1, \gamma_3, \dots, \gamma_{2k+1})$$

$$\deg \Sigma_i = 2i, \quad \deg \gamma_{2i+1} = 4i + 1, \quad \text{and}$$

$$d \Sigma_i = 0, \quad d \gamma_{2i+1} = \Sigma_{2i+1}.$$

Then by exactly the same way as in [2], we obtain a homomorphism

$$H^*(WO'_n) \rightarrow H^*(B\Gamma_n; \mathbb{R}).$$

Now let us define a homomorphism of differential graded algebras

$$\alpha : WO'_n \rightarrow WO'_p \otimes WO'_q$$

by

$$\alpha(\Sigma_i) = \Sigma_i \otimes 1 + 1 \otimes \Sigma_i$$

$$\alpha(\gamma_{2i+1}) = \gamma_{2i+1} \otimes 1 + 1 \otimes \gamma_{2i+1}.$$

Then we have

Proposition 1 (Cartan formula)

The following diagram is commutative.

$$\begin{array}{ccc}
 H^*(B\Gamma_n; \mathbb{R}) & \xrightarrow{\mu^*} & H^*(B\Gamma_p \times B\Gamma_q; \mathbb{R}) \\
 \uparrow & & \uparrow \text{cross product} \\
 & & H^*(B\Gamma_p; \mathbb{R}) \otimes H^*(B\Gamma_q; \mathbb{R}) \\
 & & \uparrow \\
 H^*(WO'_n) & \xrightarrow{\alpha_*} & H^*(WO'_p) \otimes H^*(WO'_q)
 \end{array}$$

Proof. Let  $\mathcal{F}_1$  and  $\mathcal{F}_2$  be two foliations of codimension  $p$  and  $q$  respectively on a smooth manifold  $M$ . Assume that  $\mathcal{F}_1$  and  $\mathcal{F}_2$  are transversal. Then  $\mathcal{F} = \mathcal{F}_1 \cap \mathcal{F}_2$  is a codimension  $n$  ( $= p + q$ ) foliation on  $M$ . The normal bundle  $\nu(\mathcal{F})$  is canonically isomorphic to

$\nu(\mathcal{F}_1) \oplus \nu(\mathcal{F}_2)$ . Since the interior direct sum of Riemannian (resp. Bott) connections on  $\nu(\mathcal{F}_1)$  and  $\nu(\mathcal{F}_2)$  give rise to those of  $\nu(\mathcal{F})$  and since we are using the class  $\Sigma_i$ , which is, up to a scalar, just  $\text{Tr } \Omega^1$ , our Proposition is obvious for this particular case. Then the general case follows from the usual argument (cf [2]).

### §3. Main theorem.

Let  $S$  be a complex analytic surface constructed by Kodaira (cf [1]), having the following properties.

- (i)  $S$  is the total space of a fibre bundle over a curve with fibre another curve.
- (ii)  $\text{sign}(S) \neq 0$ .

Let  $\xi \subset \tau(S)$  be the tangent bundle along the fibres. From (ii) we conclude that

$$P_1(\xi) = 3 \text{ sign } S \neq 0.$$

According to Thurston [5],  $B\bar{I}_2$  is 3-connected. Therefore again by Thurston [4],  $\xi$  is homotopic to the normal bundle of a codimension 2 foliation  $\mathcal{F}$ . We have

$$P_1(\nu(\mathcal{F})) = 3 \text{ sign } S \neq 0.$$

Now let  $(M^{2n+1}, \mathcal{G}_t^n)$  be the family of codimension  $n$  foliations on a manifold  $M^{2n+1}$  constructed by Thurston [3], such that

$$\langle \text{gv}(\mathcal{G}_t^n), [M^{2n+1}] \rangle = t \in \mathbb{R}.$$

where  $t$  ranges over some open set of  $\mathbb{R}$ . We consider the product foliation

$$\underbrace{(S, \mathcal{F}) \times \dots \times (S, \mathcal{F})}_i \times (M^{2n+1-4i}, \mathcal{G}_t).$$

These are a family of codimension  $n$  foliations on  $(2n+1)$  manifold  $(S)^i \times M^{2n+1-4i}$ . We claim that some characteristic class in  $H^{2n+1}(B\Gamma_n; \mathbb{R})$  varies continuously on this family.

More precisely we have the following. Let  $r(n)$  be the greatest integer satisfying the inequality  $2n+1 - 4r(n) \geq 3$ . Then we have

Theorem. There is a surjection

$$H_{2n+1}(B\Gamma_n; \mathbb{Z}) \rightarrow \mathbb{R}^{r(n)+1} \rightarrow 0$$

for any  $n \geq 1$ .

Proof. We consider the following family of foliations.

$$(N^i, \mathcal{H}_t^i) = ((S)^i \times M^{2n+1-4i}, \mathcal{F}^i \times \mathcal{G}_t) \quad i = 0, 1, \dots, r(n).$$

Next we consider characteristic classes  $\gamma_1^{\Sigma_1^n}$ ,  $\gamma_1^{\Sigma_1^{n-2}\Sigma_2}$ ,  $\dots$ ,  $\gamma_1^{\Sigma_1^{n-2r(n)}\Sigma_2^{r(n)}}$ .

We claim that

$$(1) \quad \gamma_1^{\Sigma_1^{n-2i}\Sigma_2^i}(\mathcal{H}_t^i) = i! \cdot (6 \operatorname{sign} S)^i \cdot t \quad i = 0, \dots, r(n).$$

$$(2) \quad \gamma_1^{\Sigma_1^{n-2i}\Sigma_2^i}(\mathcal{H}_t^j) = 0 \quad \text{for } j > i.$$

Clearly our theorem follows from these two statements.

Now we first verify (1). We have a map

$$\alpha : WO_n^1 \rightarrow WO_2^1 \otimes \dots \otimes WO_2^1 \otimes WO_{n-2i}^1,$$

which is an iteration of the maps of type  $WO_n^1 \rightarrow WO_p^1 \otimes WO_q^1$  considered in §2. Let  $x \in H^*(WO_2^1 \otimes \dots \otimes WO_2^1 \otimes WO_{n-2i}^1)$  be a cohomology class, and let

$(a_1, a_2, \dots, a_{i+1})$  be an  $(i+1)$ -tuple of non-negative integers. We define  $\chi_{(a_1, a_2, \dots, a_{i+1})}$  to be the multi-degree  $(a_1, a_2, \dots, a_{i+1})$  part of  $\chi$ . Then we have, by Proposition 1,

$$\begin{aligned} & \alpha_*([\gamma_1 \Sigma_1^{n-2i} \Sigma_2^i])_{(4,4,\dots,4,2n+1-4i)} \\ &= i! [\Sigma_2] \otimes [\Sigma_2] \otimes \dots \otimes [\Sigma_2] \otimes [\gamma_1 \Sigma_1^{n-2i}]. \end{aligned}$$

Therefore

$$\langle [\gamma_1 \Sigma_1^{n-2i} \Sigma_2^i] (\chi^i t), [N^i] \rangle = i! (6 \text{ sign } S)^i \cdot t.$$

Next we prove (2).

The same calculation as above using proposition 1 yields

$$\alpha_*([\gamma_1 \Sigma_1^{n-2i} \Sigma_2^i])_{(4,4,\dots,4,2n+1-4j)} = 0 \quad \text{for } j > i.$$

This proves (2) and hence our Theorem.

Remark. Let us call an element  $x \in H^{2n+1}(W'_n)$  "indecomposable" if  $\alpha_*(x) = 0$  for any factorization  $\alpha : W'_n \rightarrow W'_{i_1} \otimes \dots \otimes W'_{i_k}$  ( $i_1 + \dots + i_k = n$ ).

Let  $x$  be such an element. Then obviously,

$$x(\text{product foliation}) = 0.$$

It is easy to show that the classes

$$\gamma_{2i+1} \Sigma_{j_1}^{k_1} \dots \Sigma_{j_\ell}^{k_\ell} \in H^{2n+4i+1}(W'_{n+2i})$$

$$(n = j_1 k_1 + \dots + j_\ell k_\ell)$$

are indecomposable if  $j_s$  is odd for every  $s = 1, \dots, \ell$ . Thus we have

infinitely many series of indecomposable elements. For example,

$$\gamma_1 \Sigma_1, \gamma_1 \Sigma_1^2, \gamma_1 \Sigma_1^3, \dots$$

$$\gamma_1 \Sigma_3, \gamma_1 \Sigma_1 \Sigma_3, \gamma_1 \Sigma_1^2 \Sigma_3, \dots$$

$$\gamma_1 \Sigma_5, \gamma_1 \Sigma_1 \Sigma_5, \gamma_1 \Sigma_1^2 \Sigma_5, \dots$$

(In Bott's definition,  $\gamma_1$  corresponds to  $h_1$ ,  $\Sigma_1$  corresponds to  $c_1$ ,  $\Sigma_2$  corresponds to  $c_1^2 - 2c_2$  and so on). We also have another type of indecomposable elements, e.g.  $\gamma_1 \Sigma_2 \in H^5(WO_2')$ .

By the argument in this note, the problem of continuous variation of characteristic classes in  $H^{2n+1}(WO_n')$  splits into the following two problems.

- (1) To construct a family of codimension  $n$  foliation on some  $M^{2n+1}$  on which indecomposable elements take values continuously and independently.
- (2) To show that the natural map  $\pi^* : H^{4k}(BGL_{2k} \mathbb{R}; \mathbb{R}) \rightarrow H^{4k}(B\Gamma_{2k}; \mathbb{R})$  is injective. (This is the case for  $k = 1$  by Thurston [5].)

## REFERENCES

- [1] M. F. Atiyah : The signature of fibre-bundles, Global Analysis, University of Tokyo Press, (1969), 73-84.
- [2] R. Bott : Lectures on characteristic classes and foliations, Springer Verlag Lecture Notes No. 279, Berlin (1972), 1-94.
- [3] W. Thurston : Variation of the Godbillon-Vey invariant in higher codimensions (unpublished).
- [4] W. Thurston : The theory of foliations of codimension greater than one, Comm. Math. Helv 49 (1974), 214-231.
- [5] W. Thurston : Foliations and groups of diffeomorphisms, Bull. Amer. Math. Soc. 80 (1974), 304-307.

Department of Mathematics

University of Tokyo

Institute for Advanced Study

Princeton