

ON DECOMPOSITION OF LATTICE Ideals of A LATTICE-ORDERED SEMIGROUP

KENTARO MURATA

Our purpose of the present note is to obtain a unique decomposition theorem of lattice ideals of l-semigroups treated in [2]. The decomposition theorem is a generalization of the unique factorization of elements in the arithmetical l-groups [7]. Applying our theorem to submodules over a maximal bounded order of a ring, we obtain a decomposition of the modules [5].

1. PRELIMINARIES. Let $L = (L, \cdot, \leq)$ be a (conditionally) complete l-semigroup with multiplicative unity e . We assume the following two conditions:

(1) L has a map $a \mapsto a^{-1}$ into itself with two properties (i) $aa^{-1}a \leq a$ and (ii) $axa \leq a$ implies $a \leq a^{-1}$.

(2) e is maximally integral: $c^2 \leq c$ and $e \leq c$ imply $c = e$.

For any a of L we define $a^* = (a^{-1})^{-1}$, and define $a^* \circ b^* = (a^*b^*)^* = (ab)^*$ [2]. Then the set $L^* = \{a^*; a \in L\}$ is a complete l-group under \circ and \leq [3]. Hence the group (L^*, \circ) is commutative by the well known theorem of l-groups. If we classify L by the quasi-equal relation $a \sim b$ defined by $a^{-1} = b^{-1}$, then the set L/\sim of all cosets forms an l-group canonically and it is isomorphic to (L^*, \circ, \leq) . We now put

the ascending chain condition in the sense of quasi-equality for integral elements of L . Then we can prove that $p^* = p$ for any prime p which is not quasi-equal to e [2]. In the following \mathbb{P} will denote the set of all primes not quasi-equal to e . Then any element a of L is factored into a finite number of primes:

$$a \sim \prod_{p \in \mathbb{P}} p^{\nu(p,a)}$$

where $\nu(p,a)$ is the p -exponent of a . We have then (1°) $\nu(p,a) = 0$ for all but finite many $p \in \mathbb{P}$, (2°) $a \sim b$ if and only if $\nu(p,a) = \nu(p,b)$ for all $p \in \mathbb{P}$, (3°) $\nu(p,a) = \nu(p,a^*)$, (4°) $\nu(p,ab) = \nu(p,a) + \nu(p,b)$, (5°) $\nu(p,a \cup b) = \min \{ \nu(p,a), \nu(p,b) \}$, (6°) $a \leq b^*$ (i.e. $a^* \leq b^*$) if and only if $\nu(p,a) \geq \nu(p,b)$ for all $p \in \mathbb{P}$.

A lattice ideal (abbr. l-ideal) J is called closed if $a \in J$ implies $a^* \in J$. Let A be any non-empty subset of L , and let A' be the join semi-lattice generated by A . Then the set-theoretical union of all principal closed l-ideals $J(a^*)$'s generated by $a \in A'$ is the closed l-ideal generated by A . Let P be any subset of \mathbb{P} . If P is non-void, the closed l-ideal generated by $\{p_1^{-1} \cdots p_n^{-1}; p_i \in P\}$ is called a P -component of the cone I and denoted by I_P . If P is void, I_P means I itself. A P -component J_P of the closed l-ideal J will be defined to be the closed l-ideal generated by $J \cdot I_P = \{xy; x \in J, y \in I_P\}$. For convenience the closed l-ideal generated by the l-ideal J will be denoted by J^* . For two l-ideals J_1 and J_2 we define quasi-equal relation by $J_1 \sim J_2 \iff J_1^* = J_2^*$. $J_1 \circ J_2$ means the closed l-ideal

generated by $\{xy; x \in J_1, y \in J_2\}$ for any two l-ideals J_1 and J_2 . Then the set of all closed l-ideals $\mathcal{J} = (\mathcal{J}, \circ, \subseteq)$ forms a complete l-semigroup which contains the cl-semigroup (L^*, \circ, \leq) isomorphically. It can be seen that (\mathcal{J}, \circ) is a commutative semigroup.

The set-theoretical union $Z_{-\infty}$ of the rational integers Z and the symbol $-\infty$ is a totally ordered additive semigroup. For any l-ideal J of L we define

$$\nu(p, J) = \inf \{ \nu(p, a); a \in J \}.$$

Fixing J and moving p over \mathbb{P} , $\nu(p, J)$ is considered as a map from \mathbb{P} into $Z_{-\infty}$. The map is written by ν_J , that is $\nu_J(p) = \nu(p, J)$.

Let now σ be a map from \mathbb{P} into $Z_{-\infty}$ such that $\sigma(p) \leq 0$ for almost all $p \in \mathbb{P}$, and let S be the set of all such maps. Then the set G of all vectors $[\sigma(p)]$ forms a complete l-semigroup under the usual addition and the order \succeq defined by $[\sigma(p)] \succeq [\sigma'(p)] \iff \sigma(p) \geq \sigma'(p)$ for all $p \in \mathbb{P}$. In symbol: $G = (G, +, \succeq)$.

2. LEMMAS AND MAIN RESULTS.

LEMMA 1. For each $\sigma \in S$, the set $K[\sigma]$ of all $x \in L$ such that $\nu(p, x) \geq \sigma(p)$ for all $p \in \mathbb{P}$ forms a closed l-ideal of L .

Proof. This is immediate by (2°), (5°) and (6°) in Section 1.

LEMMA 2. For each closed l-ideal J we have $K[\nu_J] = J$.

Proof. Similarly obtained as the proof of Lemma 3 in [7].

LEMMA 3. For each $\sigma \in S$ we have $\nu_{K[\sigma]} = \sigma$.

Proof. Similarly obtained as the proof of Lemma 4 in [7].

By using LEMMAS 2 and 3 we obtain the following

THEOREM 1. The map $f: J \mapsto f(J) = [\nu_J(p)]$ gives an l-semigroup isomorphism from $(\mathcal{J}, \circ, \subseteq)$ onto $(G, +, \leq)$.

Let $P_+(J)$, $P_0(J)$, $P_-(J)$ and $P_{-\infty}(J)$ be the sets of primes p in \mathbb{P} such that $\nu_J(p)$ is positive, zero, negative and $-\infty$, respectively.

LEMMA 4. Let J be a closed l-ideal such that both $P_+(J)$ and $P_-(J)$ are void. If $P_0(J)$ is contained in the set-theoretical union of $P_0(J(a))$ and $P_+(J(a))$, then a is contained in J and conversely.

By using Corollary to Theorem 2.3 in [2] we get the following

LEMMA 5. Let J be a closed l-ideal. If J is multiplicatively closed, the vector $f(J)$ has no integral coordinate except zero, and vice versa.

LEMMA 6. Let J be a closed l-ideal containing the cone I . If J is closed under multiplication, J is the $P_{-\infty}(J)$ -component of I .

THEOREM 2. Any l-ideal J of L is decomposed as follows:

$$(*) \quad J \sim \prod_{p \in P_+} J(p^{\nu_p}) \cdot \left(\bigvee_{p \in P_-} J(\bigcup' p^{\nu_p}) \right) \cdot I_p.$$

where $\nu_p = \nu_J(p)$, $P_+ = P_+(J^*)$, $P_- = P_-(J^*)$, \bigcup' denotes a finite join and \bigvee denotes the set-theoretical union of all $J(\bigcup' p^{\nu_p})$. Conversely, let A, B, C be any three subsets of \mathbb{P} such that they are disjoint and one of them is finite, e. g. so is A , and let α_q and $-\beta_q$ be positive and negative integers respectively such that α_q corresponds $q \in A$ and $-\beta_q$ corresponds to $q \in B$. Then

$$(**) \quad \prod_{q \in A} J(q^{\alpha_q}) \cdot \left(\bigvee_{q \in B} J(\bigcup' q^{-\beta_q}) \right) \cdot I_C$$

is an l -ideal of L . Moreover if J of (*) is quasi-equal to (**), then $P_+ = A$, $P_- = B$, $P_{-\infty} = C$, $\nu_p = \alpha_q$ ($p \in P_+$), $\nu_p = -\beta_q$ ($p \in P_-$) by suitable enumeration of p ; that is, the decomposition (*) is unique within quasi-equality.

Proof. Let J be any l -ideal of L . Firstly we suppose that J is closed. $f(J)$ is represented as $f(J) = u_+(J) + u_-(J) + u_{-\infty}(J)$, where $u_+(J)$, $u_-(J)$, $u_{-\infty}(J)$ are the vectors whose p -coordinates are $\nu_J(p)$ if p is positive-, negative-, $-\infty$ -spots (zero otherwise), respectively. It is clear that $f^{-1}(u_+(J)) = \prod_{p \in P} J(p)^{\nu_p}$. Take any element a of $f^{-1}(u_-(J))$, and let $a^* = p_1^{\lambda_1} \circ \dots \circ p_n^{\lambda_n}$, $p_i \in \mathbb{P}$. If $\lambda_i \geq 0$ for all i , then a^* is integral, hence so is the element a . Therefore a is contained in $\bigvee J(\circ p^{\nu_p})$. If $\lambda_1 < 0, \dots, \lambda_r < 0, \lambda_{r+1} > 0, \dots, \lambda_n > 0$ for r with $0 < r \leq n$, then we obtain $a \leq a^* \leq p_1^{\lambda_1} \circ \dots \circ p_r^{\lambda_r} \leq (p_1^{\nu_{p_1}} \cup \dots \cup p_r^{\nu_{p_r}})^* = p_1^{\nu_{p_1}} \circ \dots \circ p_r^{\nu_{p_r}}$. This implies $a \in J(p_1^{\nu_{p_1}} \circ \dots \circ p_r^{\nu_{p_r}})$. Hence $f^{-1}(u_-(J)) \subseteq \bigvee J(\circ p_i^{\nu_{p_i}})$. The converse inclusion is easy to see. Next, by using LEMMAS 5 and 6 we obtain $f^{-1}(u_{-\infty}(J)) = I_{P_{-\infty}(J)}$. The last part of the theorem can be proved easily.

3. APPLICATION.

1. Let R be a noncommutative ring with a bounded maximal order σ , and let \mathcal{L} be all the non-zero fractional two-sided σ -ideals (abbr. ideals) in R [4]. \mathcal{L} is then a conditionally complete l -semi-group under module-product and set-inclusion. We assume throughout this paragraph that the ascending chain condition in the sense of

quasi-equality holds for integral ideals [1]. The term submodule means a two-sided σ -submodule of R which contains a regular element of R . A submodule M of R is said to be closed if $\mathfrak{a} \subseteq M$ implies $\mathfrak{a}^* = (\mathfrak{a}^{-1})^{-1} \subseteq M$, where \mathfrak{a} is an ideal and \mathfrak{a}^{-1} is the inverse of \mathfrak{a} . The set-theoretical union M^* of \mathfrak{a}^* for the ideals \mathfrak{a} contained in M is the closed submodule generated by M . Two submodules M_1 and M_2 are said to be quasi-equal iff $M_1^* = M_2^*$. In symbol: $M_1 \sim M_2$. If we define $M_1 M_2$ of any two closed submodules M_1 and M_2 to be the set-theoretical union of all ideals $(\sum_{i=1}^n \mathfrak{a}_i \mathfrak{b}_i)^*$ where $\mathfrak{a}_i \subseteq M_1$, $\mathfrak{b}_i \subseteq M_2$, then the set $\mathfrak{M}^* = (\mathfrak{M}^*, \cdot, \subseteq)$ of all closed submodules of R forms a commutative cl-semigroup. If we classify the cl-semigroup \mathfrak{M} consisting of all submodules of R by the quasi-equal relation \sim , then \mathfrak{M}/\sim , the set of all cosets $[M_1], [M_2], \dots$, is a commutative cl-semigroup which is isomorphic to $(\mathfrak{M}^*, \cdot, \subseteq)$, where the product of two cosets is the coset containing $(M_1 M_2)^*$ and the order \leq is defined by $[M_1] \leq [M_2] \iff M_1^* \subseteq M_2^*$. Let J be any closed l-ideal of \mathcal{L} . Then the set-theoretical union $M(J)$ of all ideals in J is a closed submodule of R . Conversely the correction $J(M)$ of all ideals in the closed submodule M is an l-ideal of \mathcal{L} . Then we have $J \mapsto M(J) \mapsto J(M(J)) = J$ and $M \mapsto J(M) \mapsto M(J(M)) = M$. Let $(\mathcal{L}^*, \circ, \subseteq)$ be the cl-semigroup consisting of all closed l-ideals in \mathcal{L} , where " \circ " is defined as in the former section. Then the map $M \mapsto J(M)$ gives an l-semigroup isomorphism from $(\mathfrak{M}^*, \cdot, \subseteq)$ onto $(\mathcal{L}^*, \circ, \subseteq)$. Under that isomorphism the cl-group consisting of all ideals corresponds to the

principal lattice ideals. By using THEOREM 2 we obtain:

$$M \sim f_1^{\alpha_1} \cdots f_n^{\alpha_n} \left(\sum'_x q_x^{-\beta_x} \right) \sigma_P$$

where $P_+(J(M)) = \{f_1, \dots, f_n\}$, $\alpha_i = \nu_{f_i}$, $P_-(J(M)) = \{q_x\}$, $\beta_x = -\nu_{q_x}$, P is the complement of $P_{-\infty}(J(M))$ in the set of all prime ideals not quasi-equal to σ , \sum' denotes the restricted sum, and σ_P is the P -component of σ . Moreover the above decomposition is unique within quasi-equality. If in particular σ is Asano, each (non-zero) submodule of R is uniquely decomposed (within commutativity) as follows:

$$M = f_1^{\alpha_1} \cdots f_n^{\alpha_n} \left(\sum'_x q_x^{-\beta_x} \right) \sigma_P.$$

Furthermore the P_1 -component M_{P_1} of M is represented as follows:

$$M_{P_1} = f_1^{\alpha_1} \cdots f_r^{\alpha_r} \left(\sum'_\lambda q_\lambda^{-\beta_\lambda} \right) \sigma_{P \vee P_1}$$

where $\{f_1, \dots, f_r\} = \{f_1, \dots, f_n\} - P_1$ and $\{q_\lambda\} = \{q_x\} - P_1$ (Cf. [4] and [5].)

2. Let σ be a Dedekind domain with its quotient field K . Then any non-zero σ -submodule M of K can be decomposed as in the case of the former paragraph. By using the decomposition we can prove the following statements.

The map $\varphi: x \mapsto \varphi(x)$ from a non-zero σ -submodule M_1 to a non-zero σ -submodule M_2 is an σ -isomorphism if and only if there exists a non-zero element t of K such that $\varphi(x) = tx$ for all $x \in M_1$. Two non-zero σ -submodules M_1 and M_2 are said to have the same $-\infty$ -type iff $\sigma_{P_{-\infty}(M_1)} = \sigma_{P_{-\infty}(M_2)}$. Then in order that M_1 and M_2 have

the same $-\infty$ -type, it is necessary and sufficient that there is an ideal \mathfrak{a} such that $M_2 = M_1 \mathfrak{a}$. Let \mathfrak{m} be the ideal generated by all prime ideals in $P_{-\infty}(M)$, and let \mathfrak{a} be an ideal. Then M is σ -isomorphic to $M\mathfrak{a}$ if and only if \mathfrak{a} is represented as $\mathfrak{a} = \mathfrak{m}(a)$ for a non-zero element a of K . Any intermediate ring T of σ and K is a P -component of σ , and it is a Dedekind ring. An integral T -ideal \mathfrak{p} of T is prime if and only if $\mathfrak{p} = \mathfrak{f}T$, where \mathfrak{f} is a prime ideal in $P_0(T)$.

REFERENCES

- [1] K. Asano, Zur Arithmetik in Schieftringen I, Osaka Math. J. **1** (1949) 98-134.
- [2] K. Asano and K. Murata, Arithmetical ideal theory in semigroups, Journ. Institute of Polytec., Osaka City Univ. **4** (1953) 9-33.
- [3] L. Fuchs, Partially ordered algebraic systems, International Series of Monographs in Pure and Applied Math., **28** (1963).
- [4] N. Jacobson, The theory of rings, Mathematical Surveys II, Amer. Math. Soc. (1943).
- [5] K. Murata, On submodules over an Asano order of a ring, Proc. Japan Acad. **50** (1974) 584-588.
- [6] K. Murata, On lattice ideals in a conditionally complete lattice-ordered semigroup (Forthcoming).
- [7] K. Murata, On lattice ideals in arithmetical lattice-ordered groups, "Sugaku (Math. Soc. of Japan)" **29** No.1 (in Japanese) (1977).