数理解析研究所講究録
152巻 1972年 119-129

A MONOTONE MAP ON 2-MANIFOLDS, WHOSE IMAGE IS HOMEOMORPHIC TO ITS DOMAIN SPACE, A 2-MANIFOLD.

BY

YUTAKA DOHI

§0. The main results in this note are Theorem A (see §2) and Theorem B (see §3) which are extensions of Whyburn's theorem [1]; Any monotone mapping of a plane onto a plane is compact. In §1, extensions of a well-known Moore's theorem concerning a decomposition of a 2-sphere, are stated without proof, Theorem 1, 2, which are used to show Theorem A and B. Terminologies are explained also in §1, whose meanings will be found among Lemmata 1~10. Finally the author regrets that he has not enough informations concerning these problems: Anyone gets the same results?

§1. A map $f(X)=Y$ is <u>compact</u> <u>(connected)</u> iff the inverse image $f^{-1}(B)$ of any compact (connected) set B of Y is compact (connected).

Lemma 1. Suppose M is a connected compact n-manifold, $n>1$, and W is its connected open subspace with a totally disconnected complement $K=X-W$. Any compact and connected map $f(W)=W$, has a unique extension $g(M)=M$, which is also compact and connected, whose

restriction on K is a homeomorphism on K.

A map $f(X)=Y$ is monotone iff the inverse image $f^{-1}(y)$ of any point y of Y is always compact and connected in X. A fundamental domain of a 2-manifold is a homeomorphic image of a connected open set in a plane.

Lemma 2. A subset C in a 2-manifold M is cellular iff C is compact, connected and is contained in such a fundamental domain W of M, that the complement $W-K$ is connected.

Lemma 3. Let X be a locally compact Hausdorff space and $\{K_i \mid i=1, 2, \ldots\}$ be its countable compact covering, then for any open set U of X, there is an integer n such that $U \cap \mathrm{Int}\, K_n \neq \phi$.

Lemma 4. A separable metric n-manifold M has a countable connected open cover $M=\bigcup_{i=1}^{\infty} M_i$ such that the closure $\overline{M}_i$ is compact and $\overline{M}_i \subset M_{i+1}$ for each i.

Lemma 5. Given an injective map $f: M_1 \to M_2$ from a m-manifold without boundary M_1 into a m-manifold M_2, then the map f is an imbedding and its image $f(M_1)$ is an open set in the interior of M_2.

A map $f(X)=Y$ is <u>quasi-compact</u> iff the set B of Y, whose inverse image $f^{-1}(B)$ is closed in X, is also closed in Y. A map $f(X)=Y$ is <u>upper semi-continuous</u> (u.s.c.) iff for any open set U of X,

the set $\tilde{U}$ in X, defined by $\tilde{U}=\cup\{f^{-1}(y) \mid y\in Y,\ f^{-1}(y)\subset U\}$, is open in X.

Lemma 6. A map f(X)=Y is closed iff the map f is quassi-compact and u.s.c.

Lemma 7. Let f(X)=Y be a closed map. If X is a separable metric space, so is Y.

Lemma 8. Let the map f(X)=Y be quasi-compact such that the inverse image of any point of Y is connected. Then if X is locally connected, so is Y.

Lemma 9. If the map f(X)=Y is closed and monotone, then it is compact and connected.

Lemma 10. A monotone map f(X)=Y from a locally compact space X onto a Hausdorff space Y is U.s.c.

Lemma 11. Let f(X)=Y be a monotone map from a locally compact metric space X onto a Hausedorff space Y. For any compact set K in Y, the inverse image $f^{-1}f(K)$ is also compact in X.

A disjoint closed cover G of a space X is called <u>a decomposition of a space X</u>, and its quotient space X'=X/G is called <u>a decomposition space</u> of X by G, where the projection ϕ :X→X' is clearly quasi-compact. A decomposition G of a space X is <u>u.s.c.</u> iff the projection ϕ :X→X/G is u.s.c; and it is <u>compact</u> <u>(connected)</u> iff each element of which is

compact (connected); it is <u>non-separating</u> iff for any element $K \in G$, the complement $X-K$ is connected. A map $f(X)=Y$ induces naturally a decomposition of X, $G(f)=\{f^{-1}(y) \mid y \in Y\}$, andwe denote its decomposition space by $\phi : X \to X/G(f)$, where the combined map $h=f\phi^{-1}: X/G(f) \to Y$ is well defined, and bijective. It is clear that the map h is homeomorphism iff the map f is quasi-compact.

Moore's Theorem. Given a non-trivial decomposition $G(X)$ of a space X, where the space X is a 2-sphere or 2-plane, which is monotone, u.s.c. and non-separating, then the decomposition space X/G is homeomorphic to the space X.

Lemma 12. Given a 2-sphere X and its connected open subset W with a decomposition $G(W)$ which is u.s.c. and monotone. Let K be the complement of W in X, and $G(K)$ be its decomposition whose element is a component of K. Define the decomposition of a 2-sphere X, by $G(X)=G(W) \cup G(K)$, then the decomposition space $X/G(X)$ is a Hausdorff space.

Theorem 1. ([2]) Given a connected open subspace W of a 2-sphere S^2 and its non-trivial decomposition $G(W)$, which is monotone, u.s.c. and non-separating. Then the decomposition space $W/G(W)$ is homeomorphic to W.

Theorem 2. Let M be a separable metric 2-manifold without boundary, G be its u.s.c. cellular decomposition. Then the decomposition space M/G is homeomorphic to M.

Suppose A is a subset of a space X. A point a of A is a cut point of A, iff the complement A- a is disconnected. A subspace A is <u>a true cyclic element of X</u>, iff it is maximal in a sense that it has no cut points. <u>A cactoid</u> is a locally connected continuum every true cyclic element of which is a 2-sphere.

Theorem 3. (Moore) Every monotone image of a 2-sphere, which is a locally connected continuum, is a cactoid and every cactoid is the image under some monotone mapping.

§2. Theorem A. Let W be a connected open subspace of a 2-sphere S^2, and M be a topological 2-manifold in the most large sense. Given a monotone map $f:W\to M$, whose image f(W) has a non-vacuous interior in M, then the image f(W) is in the interior of M, and homeomorphic to W, moreover, the map $f:W\to f(W)$ is closed, compact, and connected, and the natural decomposition space by f, g(W)=W' is homeomorphic to W.

Proof. The map $f:W\to f(W)$, is monotone, u.s.c. by Lemma 10, and non-separating (see the later argument), so the natural decom-

position space by f, $g(W)=W'$ is homeomorphic to W by Theorem 1. Now the injective map $fg^{-1}:W'\to M$, is an imbedding and $fg^{-1}(W')=f(W)$ is open subspace of M, that is, f(W) is homeomorphic to W. (see Lemma 5.) Since the map $fg^{-1}:W'\to f(W)$ is a homeomorphism, the map $f:W\to f(W)$ is quasi-compact, so f is closed by Lemma 6. A closed monotone map $f(X)=Y$ is compact and connected by Lemma 9. Now we show that f is non-separating. Take a point $y\in f(W)$, then the inverse image $f^{-1}(y)=N$ is compact and connected in W, since the map f is monotone. The 2-manifold W has a connected open cover $W=\bigcup_{i=1}^{\infty}W_i$ such that $\overline{W}$ is compact and $\overline{W}_i\subset W_{i+1}$ for each i. (see Lemma 4) There is an integer n_0 such that $W_n\supset N$, $n>n_0$, because N is compact. Take an open set $U\subset \mathrm{Int}\, f(W)$, such that $\overline{U}$ is compact in Int f(W). Since $\overline{U}=\bigcup_{i=1}^{\infty} f(\overline{W}_i)\cap\overline{U}$ is a compact covering of a compact Hausdorff space, (see Lemma 3), there is an integer n_1, such that for any $n>n_1$, $f(\overline{W}_n)$ contains an open 2-cell C^2 of M. Choose an integer n, such that $n>n_0+n_1$. Since $H=f^{-1}f(\overline{W}_n)$ is compact by Lemma 11, the restriction $f:H\to f(\overline{W}_n)$ is a closed map, so H is connected by Lemma 9. Thus H is a continuum in W. There are a finite number of 2-disks in the 2-sphere S^2, say $D_1, D_2, \ldots, D_d$, whose union is denoted by $D=\bigcup_{i=1}^{d}D_i$, which satisfy that $D\cap H=\phi$, $(\bigcup_{i=1}^{d}\mathring{D}_i)\supset(S^2-W)$. We may assume that the boundary

∂D consists of a finite number of disjoint 1-spheres, say S_1, S_2, $\ldots$, S_s, that is $\partial D=\bigcup_{j=1}^{s} S_j$. For each j, the inverse set $f^{-1}f(S_j)\subset W$, is a continuum by the same argument for H, so the union $f^{-1}f\partial D=\bigcup_{j=1}^{s} f^{-1}fS_j$ is compact in W and has less than (s+1) components, Let Q be the component of the complement $W-(f^{-1}f\partial D)$, which contains a connected set H, then it is clear that $\overline{Q}$ is compact in W and ∂Q has a finite number of components, which implies that the complement (S^2-Q) has a finite number of components, say E_1, E_2, $\ldots$, E_m, $(m\leqq s)$, that is $S^2-Q=\bigcup_{i=1}^{m} E_i$. Since f has a connected decomposition, we know that $f^{-1}fQ=Q$, so the restriction $f:Q\to f(Q)$, is a quasi-compact map, because $f:\overline{Q}\to f(\overline{Q})$ is a closed map. (See Lemma 9.) Consider the decomposition space of the 2-sphere S^2, $\phi:S\to K$, by its monotone decomposition defined by $G(S^2)=\{E_1, \ldots, E_m\}\cup\{f^{-1}f(x)\mid x\in Q\}$, which is a Hausdorff space by Lemma 12. So the map $\phi(S^2)=K$ is closed by Lemma 10 and 9, which implies K is a locally connected separable metric space which is also compact and connected. (See Lemmata 7 and 8.) After all the monotone image K of a 2-sphere is a cactoid, by Theorem 3. The map $h=f\phi^{-1}:\phi(Q)\to f(Q)$ is a homeomorphism because $f:Q\to f(Q)$ is quasi-compact, so the inverse image $h^{-1}(C^2)$ of an open 2-cell $C^2\subset f(Q)$, is a non-degenerate connected subset of K, which has no cut point of

$h^{-1}(C^2)$, whence we may say that the cactoid K has at least one E_0-set, namely one 2-sphere $\Omega \subset K$. There is a point $p \in \Omega$, such that $K-\{p\}$ is connected, because generally any simple link or E_0-set in a connected set X contains at most a countable number of cut points of X. Now we define a connected subset K_0 in K, by $K_0 = \phi(Q)-\{p\}$,

$K_0 = \phi(Q)-\{p\} = K-\{\phi(E_1), \ldots, \phi(E_m), p\}$, in which the subset $Z = \Omega-\{\phi(E_1), \ldots, \phi(E_m), p\}$ is closed and open, so we know that $K_0 = Z$, that is, K is a 2-sphere Ω. The reason why the set Z is closed and open in K_0: It is clear that the closure of Z in K is contained in Ω, because the compact set Ω is closed in a Hausdorff space K and $Z \subset \Omega$, which means that $\overline{Z} \cap K_0 = Z$, that is Z is closed in K_0. Next, the map from a 2-manifold Z without a boundary into a 2-manifold M, $h: Z \to M$, is injective, the image h(Z) is open in M, by Lemma 5, so h(Z) is also open in $h(K_0)$. So $Z = h^{-1}(hz)$ is open in K_0, because $h|K_0$ is a homeomorphism. Finally the closed monotone map $\phi: S^2 \to K$ is connected by Lemma 9, so the inverse image of a connected set $K-\phi f^{-1}(y)$, where K is a 2-sphere and $\phi f^{-1}(y)$ is a point of $\phi(Q)$, is connected, that is, the complement $S^2-f^{-1}(y)$ is connected, so $W-f^{-1}(y)$ is connected.

§3. Theorem B. Let $f: M_1 \to M_2$ be a monotone map, Int $f(M_1) \neq \phi$, from a separable metric 2-manifold M_1 without a boundary into a 2-manifold

M_2. If any element K of the monotone decomposition G of X, defined by $G=\{f^{-1}(y) \mid y\in f(M_1)\}$, is contained in a fundamental domain W of M_1, Then the map f is closed, $M\approx f(M_1)$ (homeomorphic) and $f(M_1)\subset \mathring{M}_2$.

Proof. There is such a compact set A in M_1 that Int $f(A)\neq\phi$ in M_2 and $f^{-1}f(A)=A$. Take a countable open cover $N=\{N_i\}$ of M_1 such that the closure $\overline{N}_i$ is compact and $\overline{N}_i\subset N_{i+1}$ for each i=1, 2, ..., and a 2-disk D in the interior of $f(M_1)$. Since $\{D\cap f(\overline{N}_i) \mid i=1, \ldots\}$ is a compact cover of a 2-disk D, there is an integer m such that the interior of $(D\cap f\overline{N}_m)$ in D is non-vacuous, that is Int $f(\overline{N}_m)\neq\phi$ in M_2, whence we define $A=f^{-1}f\overline{N}_m$, which is a compact set in M_1. There is a fundamental domain W_0 such that the image fW_0 is open in $\mathring{M}_2$, $f^{-1}fW_0=W_0$ and for any $K\in G_{W_0}=\{K\in G \mid K\subset W_0\}$, the complement (W_0-K) is connected. For any element $K_\nu\in G$, we may choose such a fundamental domain W_ν containing K_ν as $f^{-1}fW_\nu=W_\nu$. Take a fundamental domain W of K_ν, then $\widetilde{W}=\{K_\mu\in G \mid K_\mu\subset W\}$ is an open set of M_1, because G is u.s.c. decomposition of M_1. Let W_ν be the component of $\widetilde{W}$, which contains K_ν. It is clear that W_ν be a desired one. There is an open set V_ν of M_1, such that $K_\nu\subset V_\nu\subset\overline{V}_\nu\subset W_\nu$, because a metric space M_1 is normal. Define $\widetilde{V}_\nu$ by $\widetilde{V}_\nu=\{K\in G \mid K\subset V_\nu\}$, then we have $A=\bigcup_\nu(A\cap\widetilde{V}_\nu)=\bigcup_{i=1}^{n}(A\cap\widetilde{V}_i)=\bigcup_{i=1}^{n}(A\cap cl\widetilde{V}_i)\subset\bigcup_{i=1}^{n}W_i$, because A is compact. There is an integer m, such that

Int $f(A) \cap$ Int $f(A \cap cl\widetilde{V_m}) \neq \phi$, in other words, Int $(fW_m) \neq \phi$ in M_2. Now $W_0 = W_m$ is a desired one, by the Theorem A. For any $K_\nu \in G$, the complement $(W_\nu - K_\nu)$ is connected. Take a path $\sigma : [0, 1] \to M_1$, such that $\sigma(0) \in W_0$ and $\sigma(1) \in K_\nu$. Since the set $f^{-1}f\sigma[0, 1]$ is compact and connected, there is a sequence of fundamental domain $W_0, W_1, \ldots, W_n = W_\nu$ such that $f^{-1}f\sigma[0, 1] \subset \bigcup_{i=0}^{n} W_i$ and $(\bigcup_{i=0}^{j} W_i) \cap W_{j+1} \neq \phi$ $(j=0, 1, \ldots, n-1)$. It is clear that $W_0 \cap W_1 = f^{-1}f(W_0 \cap W_1)$, so $G_{01} = \{K \in G \mid K \subset W_0 \cap W_1\}$ is a decomposition of the intersection $W_0 \cap W_1$, which is monotone, u.s.c. and non-separating. Using Theorem 1, it is known that $(W_0 \cap W_1) \approx f(W_0 \cap W_1)$ which is in $\mathring{M}_2$ and open in M_2. It implies that Int $fW_1 \neq \phi$. By Theorem A we know that $f(W_1)$ is open in $\mathring{M}_2$ and for any $K \in G_1 = \{K \in G \mid K \subset W_1\}$ $(W_1 - K)$ is connected. After the same type of n arguments, we know that $(W_\nu - K_\nu)$ is connected. Finally by Theorem 2, the decomposition space $\phi : M_1 \to M_1' = M_1/G$ is homeomorphic to M_1, and the map $h = f\phi^{-1} : M_1' \to M_2$ is a homeomorphism, which implies that $M_1 \approx f(M_1)$, $f(M_1) \subset \mathring{M}_2$ and the map $f : M_1 \to M_2$ is quasi-compact. Since the map $f : M_1 \to M_2$ is also u.s.c., the monotone map is closed, that is compact and connected.

REFERENCES

[1] "Compactness of certain mapplings," by G. T. Whyburn, American Journal of Mathematics 81 (1959), pp. 306-314

[2] "On the monotone decompositions of open sets of a 2-sphere, which induces the same spaces," by Y. Dohi, Journal of Tokyo Metropolitan Junior College of Commerce, No. 6.

[3] "Decomposition Spaces," by G. T. Whyburn in Topology of 3-Manifolds and Related Topics, pp. 1-14. (Prentice-Hall, Inc.)

Tokyo Metropolitan Junior College of Commerce
(April, 1972)