

Title	Extreme Values of an Implicit Function (不等式に関する研究)
Author(s)	MOCHIZUKI, NOZOMU
Citation	数理解析研究所講究録 (1973), 191: 84-86
Issue Date	1973-11
URL	http://hdl.handle.net/2433/107244
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

EXTREME VALUES OF AN IMPLICIT FUNCTION

NOZOMU MOCHIZUKI

Let $\varphi(x,y)$ be a C^2 -function in a domain E of \mathbb{R}^2 and $\varphi_x(a,b) = \varphi_y(a,b) = 0$ for $(a,b) \in E$. Then, as is well known, the local behavior of $\varphi(x,y)$ is examined by the sign of $\Delta_0(a,b) = \varphi_{xx}(a,b)\varphi_{yy}(a,b) - \varphi_{xy}(a,b)^2$; that is, if $\Delta_0(a,b) > 0$ $\varphi(a,b)$ is a local maximum or a local minimum according to $\varphi_{xx}(a,b) < 0$ or $\varphi_{xx}(a,b) > 0$.

Let now $f(x,y,z)$ be a C^2 -function in a domain D of \mathbb{R}^3 and let $f(a,b,c) = 0$, $f_z(a,b,c) \neq 0$ for $(a,b,c) \in D$. Then a C^2 -function $\varphi(x,y)$ is uniquely determined in a neighborhood of (a,b) , for which $z = \varphi(x,y)$ if and only if $f(x,y,z) = 0$.

As for the local behavior of such $\varphi(x,y)$, we have the following simple criterion. Although straightforward computations are sufficient for the proof, we have not found such a result in the literature.

THEOREM 1. Let $f(x,y,z)$ and $\varphi(x,y)$ be as above, and let

$$f(a,b,c) = f_x(a,b,c) = f_y(a,b,c) = 0, \quad f_z(a,b,c) \neq 0.$$

Put $\Delta(x,y,z) = f_{xx}(x,y,z)f_{yy}(x,y,z) - f_{xy}(x,y,z)^2$.

(i) If $\Delta(a,b,c) > 0$, then $c = \varphi(a,b)$ is a local maximum or a local minimum of $\varphi(x,y)$ according to $sf_{xx}(a,b,c) < 0$

or $\varepsilon f_{xx}(a,b,c) > 0$, where $\varepsilon = \text{sgn}(-f_z(a,b,c))$.

(2) If $\Delta(a,b,c) < 0$, then $c = \varphi(a,b)$ is not an extreme value of $\varphi(x,y)$.

PROOF. From $f_x + f_z \varphi_x = 0$ follows that $f_{xx} + f_z \varphi_{xx} = 0$, hence

$$\varphi_{xx}(a,b) = - \frac{f_{xx}(a,b,c)}{f_z(a,b,c)} .$$

Similarly, we have

$$\varphi_{yy}(a,b,c) = - \frac{f_{yy}(a,b,c)}{f_z(a,b,c)} , \quad \varphi_{xy}(a,b,c) = - \frac{f_{xy}(a,b,c)}{f_z(a,b,c)} .$$

Thus we have

$$\varphi_{xx}(a,b)\varphi_{yy}(a,b) - \varphi_{xy}(a,b)^2 = \frac{\Delta(a,b,c)}{f_z(a,b,c)^2} ,$$

which completes the proof.

We remark that the generalization to the case of n variables is immediate.

Let $f(x,y) = f(x_1, x_2, \dots, x_n, y)$ be a C^2 -function in a domain D of \mathbb{R}^{n+1} such that $f(a,b) = 0$ for a point $(a,b) \in D$. Let $f_y(a,b) \neq 0$. Then there exists a C^2 -function $\varphi(x) = \varphi(x_1, \dots, x_n)$ in a suitable neighborhood of $a = (a_1, \dots, a_n)$ in \mathbb{R}^n such that $y = \varphi(x)$ if and only if $f(x,y) = 0$.

Let $f_{x_j}(a,b) = 0$, $j = 1, 2, \dots, n$. We have then

$$\varphi_{x_j}(a) = 0; \quad \varphi_{x_i x_j}(a) = - \frac{f_{x_i x_j}(a,b)}{f_y(a,b)}, \quad 1 \leq i, j \leq n.$$

It is well known that $\varphi(a)$ is a local maximum if the matrix $A = [\varphi_{x_i x_j}(a)]$ is negative-definite and $\varphi(a)$ is a local minimum if A is positive-definite and, moreover, $\varphi(a)$ is not an extreme value if A is indefinite. We have thus

THEOREM 2. Let $f(x,y)$ be as above and let $f_{x_j}(a,b) = 0$, $j = 1, 2, \dots, n$. Then

(1) $b = \varphi(a)$ is a local maximum or a local minimum according as the matrix $A = [\varepsilon f_{x_i x_j}(a,b)]$ is negative-definite or positive-definite, where $\varepsilon = \text{sgn}(-f_y(a,b))$.

(2) $\varphi(x)$ does not admit an extreme value at $x = a$ if A is indefinite,

EXAMPLE. Theorem 1 is sometimes useful in computing extreme values of $\varphi(x,y)$. For example, let $\varphi(x,y) = (x+1)/\sqrt{x^2+y^2+1}$. Then, by setting $f(x,y,z) = (x+1)^2 - (x^2+y^2+1)z^2 = 0$, one can easily see that $\sqrt{2} = \varphi(1,0)$ is a local maximum of $\varphi(x,y)$.