

A CASE OF PURE LEIOMYOMA OF THE PROSTATE

Hiroshi MASUDA and Hidetoshi YAMANAKA

From the Department of Urology, Gunma University School of Medicine

Seiji NAKATA and Jin SATO

From the Department of Urology, Gunma Cancer Center

Akira OGAWA

From the Department of Pathology, Gunma Cancer Center

Nozomu KOSAKU

From the Department of Urology, Kosaku Clinic

Pure leiomyoma of the prostate is very rare. A 67-year-old man was hospitalized with gross hematuria and urinary retention. Drip infusion pyelography revealed a defect of the urinary bladder. By radiological examinations, submucosal tumor of the bladder neck was diagnosed. Transrectal biopsy suggested a benign prostatic tumor histologically. Open resection of the tumor was performed. Histological findings were pure leiomyoma of the prostate. The postoperative course was good and urinary retention was improved remarkably.

(Acta Urol. Jpn. 45 : 277-280, 1999)

Key words : Pure leiomyoma, Macrohematuria, Prostate

INTRODUCTION

Pure leiomyoma of the prostate is very rare, we could find only 12 cases in the Japanese literature since the case reported by Kojima in 1917. Recently, we encountered a patient with this disease who complained of asymptomatic gross hematuria and urinary retention. We discuss the features of this disease with a review of the literature.

CASE REPORT

A 67-year-old man presented with asymptomatic gross hematuria and urinary retention of 4 months duration. After a defect in the bladder was noted on drip infusion pyelography, he was referred to our hospital. Complete blood count was normal except for gross hematuria on urinalysis. Carcinoembryonal antigen (CEA) was 7.2 ng/ml, prostate specific antigen (PSA) was 8.8 ng/ml. Because serum CEA was slightly elevated, barium studies and gastroscopy were performed. Under a diagnosis of gastric cancer, total gastrectomy was performed. On rectal digital examination, the prostate was the size of a hen's egg, elastic soft, with a smooth surface, not nodular. A large spherical mass was recognized from the anterior wall to 12 o'clock on the bladder neck. The mucosa of the bladder was normal, not irregular. Drip infusion pyelography showed a defect in the bladder. Transrectal sonography revealed a homogeneous mass. Urethrography revealed elevation of the bladder base and elongation of the posterior urethra. Residual urine was 110 ml.




Fig. 1. Computed tomography of the pelvis revealed an elevated mass in the bladder lumen.

Computed tomography of the pelvis revealed an elevated mass in the bladder lumen. Bladder mucosa showed no irregularity and there was no lymphadenopathy in the pelvis (Fig. 1). Magnetic resonance imaging of the pelvis revealed a mass that appeared continuous from the bladder neck to inner portion of the prostate (Fig. 2). We found that the tumor had originated from prostate tissue. Therefore, under the diagnosis of prostatic tumor, a tumor biopsy was performed by transrectal sonography at first. The tumor was an encapsulated homogeneous mass. Pathological findings showed a benign prostatic tumor with few smooth muscle tissue. Under a diagnosis of benign prostatic tumor, open resection of the tumor was performed on August 3, 1994. By middle lower abdominal incision, we reached Retzius's space. We recognized that the




Fig. 2. On magnetic resonance imaging of the pelvis, the mass appeared continuous from the bladder neck to the inner portion of the prostate.



Fig. 3. Microscopic section of pure leiomyoma of prostate (H & E; $\times 33$).

tumor with a smooth surface pressed the bladder neck and the prostatic urethra. We enucleated the tumor. The specimen weighed 25 gm. It had a smooth surface and was encapsulated forming a whole homogeneous mass. On microscopic examination, the fragments were found to be composed of fasciculated bands of spindle-shaped smooth muscle. The ratio of nuclei to cytoplasm was low; mitoses and atypia were absent. The pathological diagnosis was leiomyoma of the prostate (Fig. 3). The postoperative course was good and urinary retention was improved remarkably. Postoperative residual urine was approximately 5 ml.

DISCUSSION

Pure leiomyoma of the prostate is very rare. Kaufman defined this entity as a circumscribed or encapsulated mass of smooth muscle, 1 cm or more in diameter, containing varying amounts of fibrous tissue but devoid of glandular elements, and which is either obviously prostatic or juxtaprostatic in origin and position¹⁾

Thirteen cases including our case have been reported in the Japanese literature. The pathogenesis of leiomyoma of the prostate remains unclear. Vassilakis suggested that infection and inflammation

transform glandular tissue into smooth muscle and that hypertrophy leads to myoma²⁾. Patch and Rhea indicated that leiomyomas originate from embryological analgen³⁾. The main clinical symptom is urinary disturbance as in benign prostatic hypertrophy, differing with the localization within the prostate. Nakamura and Ninman reported that rectal disturbance was a chief complaint^{4,5)}. The chief complaints in the present case were asymptomatic gross hematuria and dysuria. Concerning radiological diagnosis, there were few case reports and there seemed to be no general findings. Regan reported that in a large leiomyoma of the prostate, computed tomography of the pelvis revealed clear margins and a nodule in the inner portion⁶⁾. Nakamura reported magnetic resonance imaging relative to muscle on T1-weighted images⁴⁾. In our case, the pelvic computed tomograph showed clear margins, but there were no nodules in the inner portion. Pelvic magnetic resonance imaging in our case revealed a slight isosignal relative to muscle on T1-weighted images. Transrectal ultrasonography revealed a homogeneous mass, differing from prostatic carcinoma, which shows a hypoechoic pattern⁷⁾. This disease is benign, and we recommend a surgical method. Among 13 cases including our case, suprapubic prostatectomy was performed in 5, enucleation in 5, transurethral resection of the prostate in 2 and subtotal prostatectomy for prostatic cancer in 1⁸⁾. In our case, open resection was performed. Considering the definition, Belis indicated that true leiomyoma is distinguishable from hyperplasia by the presence of encapsulation and absence of glands⁹⁾, we recommend open resection of the tumor as the procedure of first choice.

CONCLUSION

We reported a case of pure leiomyoma of the prostate. We discussed the features of this disease with a review of the literature.

REFERENCES

- 1) Kaufman JJ and Bereneike RR: Leiomyoma of the prostate. *J Urol* **65**: 297-310, 1951
- 2) Vassilakis GB: Pure leiomyoma of prostate. *Urology* **11**: 93-94, 1978
- 3) Patch FS and Rhea LJ: Leiomyoma of the prostate gland. *Br J Urol* **17**: 213-228, 1935
- 4) Nakamura K and Shiramizu M: Pure leiomyoma of prostate presenting with rectal symptoms. *Acta Urol Jpn* **38**: 1067-1069, 1992
- 5) Hinman F and Sullivan JJ: Two cases of leiomyoma of the prostate. *J Urol* **26**: 475-483, 1931
- 6) Roberts RA, Norman RW and Mack FG: Rectal obstruction by prostatic carcinoma. *J Urol* **135**: 137-139, 1986
- 7) Hrick H and Thoeni RF: Neoplasms of the prostate

- gland. *Clinical Urography*. pp. 1396–1400, W.B. Saunders Co, Philadelphia, 1990
- 8) Nakamoto T, Fujiwara S, Ishino T, et al. : A case of pure leiomyoma of the prostate. *Acta Urol Jpn* **31** : 2261–2264, 1985
- 9) Beris JA, Post GJ, Rochman SC, et al. : Genitourinary leiomyoma. *Urology* **13** : 424–429, 1979
(Received on October 5, 1998)
(Accepted on December 22, 1998)

和文抄録

前立腺平滑筋腫の1例

群馬大学医学部泌尿器科学教室（主任：山中英壽教授）

増田 広*, 山中 英壽

群馬県立がんセンター泌尿器科（部長：鈴木孝憲）

中田 誠司**, 佐藤 仁***

群馬県立がんセンター病理部（部長：小川 晃）

小 川 晃

古作クリニック（院長：古作 望）

古 作 望

前立腺平滑筋腫は非常に稀な疾患である。67歳，男性が肉眼的血尿と尿閉を主訴に来院。DIPを含む画像診断では膀胱頸部にみられた粘膜下腫瘍と診断した。超音波下経直腸的生検では平滑筋組織を含む良性

の前立腺腫瘍であった。腫瘍摘除術を施行。病理学的診断は前立腺平滑筋腫であった。術後排尿困難の著明な改善を認めた。

(泌尿紀要 45 : 277-280, 1999)

* 現：東葛病院泌尿器科

** 現：足利赤十字病院泌尿器科

*** 現：群馬女子短期大学生活科