

Online Supplemental materials

Fig. S1. The redox response of Redoxfluor in vitro. Effect of pH upon the FRET ratio of Redoxfluor. Circles, reduced probes; squares, oxidized probes; red, C-probe; blue, A-probe.

Fig. S2. Visualization of the redox state in *P. pastoris*. The C-probe responds to various oxidants in *P. pastoris*. Bar, 2 μm .

Fig. S3. Redoxfluor response in the presence of cycloheximide. The wild-type CHO-K1 strain used in Fig. 3B was pre-treated with 20 $\mu\text{g/ml}$ cycloheximide for 1 h, incubated in medium containing 200 μM H_2O_2 and the same concentration of cycloheximide for 20 min, and transferred to cycloheximide-containing medium without H_2O_2 (H_2O_2 -washout) for the indicated periods. The cycloheximide treatment alone exerted an oxidizing effect upon the redox state, but the H_2O_2 -washout increased the FRET ratio showing the reversibility of Redoxfluor response. Bar, 10 μm .

Fig. S4. Biochemical assessment of the redox state in *pex5* cells using mPEG-maleimide. Cell lysate from wild-type (CHO-K1) or *pex5* (ZP105) cells expressing cytosolic Redoxfluor (C-probe or A-probe) was incubated with mPEG-maleimide, and subjected to immunoblot analysis. The molecular-weight distributions of the probe proteins are slightly greater in the lysate from the *pex* cells. The arrow indicates non-modified probe proteins and the asterisks show the modified forms of the protein.

Fig. S5. Detection of accumulated ROS by 2', 7'-dichlorodihydrofluorescein diacetate (DCF). Wild-type CHO-K1 cells exhibited greater levels of intracellularly accumulated ROS at 37°C than ZP105 cells mutant for peroxisome assembly. The values represent the fluorescent intensities of DCF in arbitrary units.

Table S1. Primers used for qRT-PCR

Video 1. The H₂O₂-induced FRET response in CHO-K1 cells expressing the C-probe. Time series speed was 1 frame per minute, and the images were taken for 20 minutes. Using our conventional FRET microscope, an increase in background fluorescence with both A- and C-probes was observed due to the increase of the medium volume of the object upon reagent addition.

Video 2. The ATZ-induced FRET response in CHO-K1 cells expressing the C-probe.

Time series speed was 1 frame per minute, and images were taken for 40 minutes.

ATZ was added at 20 minutes.


Video 3. The H₂O₂-induced FRET response in CHO-K1 cells expressing the C-


probe-PTS1. Time series speed was 1 frame per minute, and images were taken for


40 minutes. H₂O₂ was added at 20 minutes.


TABLE SI. Primers used in the qRT-PCR

Gene	Primers
<i>GST</i>	5' -TGGAAGGAGGAGGTGGTACTGTAG-3'
	5' -CCCATCATTACCATATCCACCAGG-3'
<i>PpCTA1</i>	5' -CGAGTATCCTTCATGGACTTGTTAC-3'
	5' -TCCTCAATGGGAAGTCTTTGTGTGG-3'
<i>PpGPX1</i>	5' -ACCAGTTTGGTCATCAGGAACCAGG-3'
	5' -ACCTTTGAATCCGAGGAGACCAGAC-3'
<i>PpSOD2</i>	5' -AACACACCTAAGGTGATCGAGCTAC-3'
	5' -ACCTGCCAACTTAGAGTTGGTAAGG-3'
<i>PpTSA1</i>	5' -CATTGTTGGCTGACACCAACCACAC-3'
	5' -TCCGACTGGCAGATCGTTGATAGTG-3'












TABLE SI. Primers used in the qRT-PCR

Gene	Primers
<i>GST</i>	5' -TGGAAGGAGGAGGTGGTACTGTAG-3'
	5' -CCCATCATTACCATATCCACCAGG-3'
<i>PpCTA1</i>	5' -CGAGTATCCTTCATGGACTTGTTAC-3'
	5' -TCCTCAATGGGAAGTCTTTGTGTGG-3'
<i>PpGPX1</i>	5' -ACCAGTTTGGTCATCAGGAACCAGG-3'
	5' -ACCTTTGAATCCGAGGAGACCAGAC-3'
<i>PpSOD2</i>	5' -AACACACCTAAGGTGATCGAGCTAC-3'
	5' -ACCTGCCAACTTAGAGTTGGTAAGG-3'
<i>PpTSA1</i>	5' -CATTGTTGGCTGACACCAACCACAC-3'
	5' -TCCGACTGGCAGATCGTTGATAGTG-3'