

Title	On the Thermoelasticity in the Semi-infinite Elastic Solid
Author(s)	SHIMA, Michiyasu
Citation	Bulletins - Disaster Prevention Research Institute, Kyoto University (1958), 25: 1-17
Issue Date	1958-08-05
URL	http://hdl.handle.net/2433/123680
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

昭和34. 6. 10



DISASTER PREVENTION RESEARCH INSTITUTE

BULLETIN No. 25

SEPTEMBER, 1958

ON THE THERMOELASTICITY IN THE SEMI-
INFINITE ELASTIC SOLID

BY

MICHIYASU SHIMA

KYOTO UNIVERSITY, KYOTO, JAPAN

R

DISASTER PREVENTION RESEARCH INSTITUTE
KYOTO UNIVERSITY
BULLETINS

Bulletin No. 25

September, 1958

On the Thermoelasticity in the Semi-infinite
Elastic Solid

By

Michiyasu SHIMA

Contents

On the Thermoelasticity in the Semi-infinite Elastic Solid

by

Michiyasu SHIMA

Geophysical Institute, Faculty of Science, Kyoto University

(Communicated by Prof. K. Sassa)

Abstract

When the spheroidal or spherical region of material of α larger than that of the surroundings, which is embedded in a semi-infinite elastic body, is heated, there appears the thermal stress. The displacements at the free boundary and the stresses round the thermal origin in such a problem of thermal elasticity are obtained, introducing the displacement function ϕ , and transforming the equations of equilibrium so that the results of potential theory and theory of centres of dilatation may be applied. Thus, the state of the thermal origin is estimated from the observed deformation of the free surface. For example, the dimension of the magmatic reservoir at the Volcano Aso is estimated at ca. 1 km from the observed crustal movement which may result from its expansion and contraction.

Nomenclature

The following nomenclature is used in the paper :

u, v, w	: cartesian components of displacement
e_{ij}	: strain ($i, j = x, y, z$)
e	: dilatation
E	: Young's modulus
μ	: rigidity
σ	: Poission's ratio
T_{ij}	: stress ($i, j = x, y, z$)
ϕ	: displacement function
α	: coefficient of linear thermal expansion
α_i	: coefficient of linear thermal expansion inside thermal region
α_o	: coefficient of linear thermal expansion outside thermal region
T	: change of temperature

θ_1 : inclination of spheroidal thermal origin.
§1.

When the temperature in an elastic body is not uniform, or the temperature in the elastic body of the non-uniform distribution of coefficient of thermal expansion changes uniformly, there appears a state of stress. Such a stress is called the thermal stress. In the case of the inclusion of material of the coefficient of thermal expansion larger than that of the surroundings in the earth's crust, heated by the convective currents of magma through the fissures, there appear the thermal stresses as the results of the increases of temperature and coefficient of thermal expansion, and we observe the deformation at the earth's surface. Particularly, if the temperature of the thermal origin is a little lower than the transition point of the heated material, the increase of coefficient of thermal expansion is remarkable. For example, the coefficient of thermal expansion of quartz increases rapidly near 573°C of α - β transition as shown in Fig. 1¹⁾.




Fig. 1 Thermal expansion of quartz
(after Jay).

In this paper, we calculate the thermal stresses round the thermal origin and the surface displacements which result from the increases of temperature and coefficient of thermal expansion in the spheroidal (or spherical) thermal origin of α_s larger than α_e in the elastically uniform semi-infinite solid.

§2

When there appears the change of temperature (x, y, z) in the infinite and free elastic solid which has the uniform elastic constants and the non-uniform linear coefficient of thermal expansion $\alpha(x, y, z)$, the free thermal expansion of every volume element is constrained partially by the surrounding material, and a state of thermal stress ensues. The difference between the actual strain and the free expansion $\int_0^T \alpha dT$ is related to the stress through

Hook's law. As the changes of elastic constants which result from that of temperature are very small, they can be neglected, but as that of coefficient of thermal expansion is fairly large, it is considered. As a uniform change of temperature of a small volume element does not create any angular distortion of the element, the shear stresses are unaffected by the term $\int_0^T adT$, that is, $T_{xy} = 2\mu e_{xy}$, However, the normal stresses are determined by the following equations.

$$T_{xx} = 2\mu \left[\frac{\sigma}{1-2\sigma} e + e_{xx} - \frac{1+\sigma}{1-2\sigma} \int_0^T adT \right] \quad (1)$$

cyclic.

Then, the three equations of equilibrium take the form.

$$\frac{\partial e}{\partial x} + (1-2\sigma)\nabla^2 u = 2(1+\sigma) \frac{\partial}{\partial x} \int_0^T adT \quad (2)$$

cyclic.

In order to solve the equation (2), the displacement function ϕ is introduced²⁾

$$\frac{\partial \phi}{\partial x} = u, \quad \frac{\partial \phi}{\partial y} = v, \quad \frac{\partial \phi}{\partial z} = w.$$

The equation (2) now can be written

$$\frac{\partial}{\partial x} \left[(1-\sigma)\nabla^2 \phi - (1+\sigma) \int_0^T adT \right] = 0.$$

Then equation (2) is all satisfied when

$$\nabla^2 \phi = \frac{1+\sigma}{1-\sigma} \int_0^T adT \quad (3)$$

As the state of stress represented by this function ϕ ordinarily requires certain surface tractions at the boundary of solid, by means of the principle of superposition, a complementary stress function must be determined so as to satisfy the boundary conditions. This is only a problem of given boundary tractions in the ordinary theory of elasticity.

By means of equation (3), equation (1) can be written in the form

$$T_{xx} = 2\mu \left[\frac{\partial^2 \phi}{\partial x^2} - \frac{1+\sigma}{1-\sigma} \int_0^T adT \right] \quad (4)$$

Equation (3) is of the same form as Poisson equation $\nabla^2 V = -4\pi\rho$ in the potential theory and a particular integral is given by the Newtonian potential of a distribution of material of the density $-(1+\sigma) \int_0^T adT / 4\pi(1-\sigma)$,

namely,³⁾

$$\phi = -\frac{1+\sigma}{4\pi(1-\sigma)} \int_0^r \frac{adT}{r'} d\xi d\eta d\zeta \quad (5)$$

$$r'^2 = (x-\xi)^2 + (y-\eta)^2 + (z-\zeta)^2.$$

This potential ϕ represents the complete solution for the infinite solid when $T=0$ outside the heated origin.

As the general equation (2) implies $\frac{\partial T}{\partial x}$, the validity of any solution at a surface of temperature distribution requires examinations. However its validity is shown by potential theory.²⁾

In order to relate the above solution to the ordinary elastic theory, the nucleus of thermoelastic strain is defined as follows. The formula (5) and the definition of ϕ shows that if a change of temperature of volume element $d\tau$ in the infinite body is, that of the remainder being zero, the displacement is the gradient of

$$-\frac{(1+\sigma)d\tau}{4\pi(1-\sigma)r'} \int_0^r adT$$

This is simply the singularity known in the ordinary theory of elasticity as the centre of dilatation⁴⁾, and (6) may be called its strength.

$$\frac{S}{4\pi} = \frac{1+\sigma}{4\pi(1-\sigma)} \int_0^r adT \quad (6)$$

Namely, the effect of heating is the same as that of a distribution of centres of dilatation of this strength $S/4\pi$ in an unheated body.

§3

Then we can obtain the stress distribution of a heated and bounded elastic body, if the formula for the same distribution $S(\xi, \eta, \zeta)/4\pi$ of centres of dilatation within the boundary is known.

In order to solve the problem of the distribution of centres of dilatation corresponding to that of the rise of temperature in the spheroidal (or spherical) region of a_i embedded in semi-infinite elastic body, firstly, we require the solution (displacement) (7) for the centre of dilatation of the strength $S/4\pi$ ⁵⁾

$$\mathbf{u} = -\frac{S}{4\pi} \left[\mathcal{V} \left(\frac{1}{R_1} \right) + \mathcal{V}_2 \left(\frac{1}{R_2} \right) \right]$$

$$\mathcal{V}_2 = (3-4\sigma)\mathcal{V} + 2\mathcal{V}z \frac{\partial}{\partial z} - 4(1-\sigma)\mathbf{k}\mathcal{V}^2z \quad (7)$$

$$R_1^2 = (x - \xi)^2 + (y - \eta)^2 + (z - \zeta)^2, \quad R_2^2 = (x - \xi)^2 + (y - \eta)^2 + (z + \zeta)^2,$$

, where the $x - y$ coordinate axes are laid on the free surface and z -axis is directed downward. If $S(\xi, \eta, \zeta)$ distributes in the region V_1 in $z \geq 0$,

$$\begin{aligned} \mathbf{u} &= \int_{V_1} \mathbf{u}_0 d\tau = -\frac{1}{4\pi} \left(\mathcal{V} \int_{V_1} \frac{S}{R_1} d\tau + \mathcal{V}_2 \int_{V_1} \frac{S}{R_2} d\tau \right) \\ &= -\frac{1}{4\pi} \left(\mathcal{V} \int_{V_1} \frac{S}{R_1} d\tau + \mathcal{V}_2 \int_{V_2} \frac{S}{R_1} d\tau \right) = \mathcal{V} \phi_1 + \mathcal{V}_2 \phi_2 \quad (8) \\ \phi_1 &= -\frac{1}{4\pi} \int_{V_1} \frac{S}{R_1} d\tau \quad \phi_2 = -\frac{1}{4\pi} \int_{V_2} \frac{S}{R_1} d\tau, \end{aligned}$$

, or writing down their components

$$\begin{aligned} u &= \frac{\partial \phi_1}{\partial x} + (3 - 4\sigma) \frac{\partial \phi_2}{\partial x} + 2z \frac{\partial^2 \phi_2}{\partial x \partial z} \\ v &= \frac{\partial \phi_1}{\partial y} + (3 - 4\sigma) \frac{\partial \phi_2}{\partial y} + 2z \frac{\partial^2 \phi_2}{\partial y \partial z} \quad (9) \\ w &= \frac{\partial \phi_1}{\partial z} + (-3 + 4\sigma) \frac{\partial \phi_2}{\partial z} + 2z \frac{\partial^2 \phi_2}{\partial z^2}. \end{aligned}$$

, where V_2 is the image of V_1 in the plane $z=0$ and ϕ_2 is simply the reflection transformation of ϕ_1 in the plane $z=0$.

Then, when the potentials ϕ_1 and ϕ_2 for a distribution of $S(\xi, \eta, \zeta)$ are known, we can obtain the displacements (u, v, w) by means of differentiation of ϕ_1 and ϕ_2 . When the spheroid and spheres are adopted as V_1 ,

spheroid

$$\begin{aligned} &\frac{x^2 \cos^2 \theta_1 + x(z-d) \sin 2\theta_1 + (z-d)^2 \sin^2 \theta_1}{a^2} + \frac{y^2}{b^2} \\ &+ \frac{x^2 \sin^2 \theta_1 - x(z-d) \sin 2\theta_1 + (z-d)^2 \cos^2 \theta_1}{c^2} = 1 \end{aligned}$$

two spheres

$$(x + b_1)^2 + y^2 + (z - d_1)^2 = a^2, \quad (x + b_2)^2 + y^2 + (z - d_2)^2 = a^2, \quad (10)$$

from the known results for their potential³⁾, ϕ outside the thermal origin are prolate spheroid $a > b = c$

$$\begin{aligned} \phi_i &= -\frac{Sac^2}{4(a^2 - c^2)} \left[\log \frac{\sqrt{a^2 + q_i} + \sqrt{a^2 - c^2}}{\sqrt{c^2 + q_i}} \left(2\sqrt{a^2 - c^2} + \frac{y^2 + Z_i - 2X_i}{\sqrt{a^2 - c^2}} \right) \right. \\ &\left. + \frac{2X_i}{\sqrt{a^2 + q_i}} - \frac{(y^2 + Z_i)\sqrt{a^2 + q_i}}{c^2 + q_i} \right] \quad (11) \end{aligned}$$

$$q_i = \frac{1}{2} \left[r_i^2 - a^2 - c^2 + \sqrt{(r_i^2 + a^2 - c^2)^2 - 4(a^2 - c^2)X_i} \right],$$

oblate spheroid $a = b > c$

$$\begin{aligned} \phi_i &= -\frac{Sac^2}{4(a^2 - c^2)} \left[\tan^{-1} \frac{\sqrt{a^2 - c^2}}{\sqrt{c^2 + q_i}} \left(2\sqrt{a^2 - c^2} - \frac{y^2 + X_i - 2Z_i}{\sqrt{a^2 - c^2}} \right) \right. \\ &\quad \left. + \frac{(y^2 + X_i)\sqrt{c^2 + q_i}}{a^2 + q_i} - \frac{2Z_i}{\sqrt{c^2 + q_i}} \right] \\ q_i &= \frac{1}{2} \left[r_i^2 - a^2 - c^2 + \sqrt{(r_i^2 - a^2 + c^2)^2 + 4(a^2 - c^2)Z_i} \right] \end{aligned} \quad (12)$$

$$\begin{aligned} X_i &= x^2 \cos^2 \theta_i + xz_i \sin 2\theta_i + z_i^2 \sin^2 \theta_i, & Z_i &= x^2 \sin^2 \theta_i - xz_i \sin 2\theta_i + z_i^2 \cos^2 \theta_i \\ r_i^2 &= x^2 + y^2 + z_i^2, & z_1 &= z - d, & z_2 &= z + d, & \theta_1 &= -\theta_2. \end{aligned}$$

two spheres

$$\phi_i = -a^3 S \left[\frac{1}{3R_{i1}} + \frac{1}{3R_{i2}} \right] \quad (13)$$

$$\begin{aligned} R_{11}^2 &= (x + b)^2 + y^2 + (z - d_1)^2, & R_{12}^2 &= (x - b)^2 + y^2 + (z - d_2)^2, \\ R_{21}^2 &= (x + b)^2 + y^2 + (z + d_1)^2, & R_{22}^2 &= (x - b)^2 + y^2 + (z + d_2)^2. \end{aligned}$$

Inserting (11), (12), (13) into (8), we obtain

$$\begin{aligned} u &= -\gamma [A_1 + (3 - 4\sigma)A_2 + 2zA] \\ v &= -\gamma [B_1 + (3 - 4\sigma)B_2 + 2zB] \\ w &= -\gamma [C_1 + (-3 + 4\sigma)C_2 + 2zC] \end{aligned} \quad (14)$$

prolate spheroid

$$\begin{aligned} r &= \frac{Sac^2}{4(a^2 - c^2)} \\ A_i &= \frac{2x(\sin^2 \theta_i - 2\cos^2 \theta_i) - 3z_i \sin 2\theta_i}{\sqrt{a^2 - c^2}} \log \frac{\sqrt{a^2 + q_i} + \sqrt{a^2 - c^2}}{\sqrt{c^2 + q_i}} \\ &\quad + \frac{4x\cos^2 \theta_i + 2z_i \sin 2\theta_i}{\sqrt{a^2 + q_i}} - \frac{(2x\sin^2 \theta_i - z_i \sin 2\theta_i)\sqrt{a^2 + q_i}}{c^2 + q_i} \\ A &= -\frac{3\sin 2\theta_2}{\sqrt{a^2 - c^2}} \log \frac{\sqrt{a^2 + q_2} + \sqrt{a^2 - c^2}}{\sqrt{c^2 + q_2}} + \frac{2\sin 2\theta_2}{\sqrt{a^2 + q_2}} + \frac{\sin 2\theta_2 \sqrt{a^2 + q_2}}{c^2 + q_2} \\ &\quad - \frac{q_{2x}(a^2 - c^2)}{\sqrt{a^2 + q_2}(c^2 + q_2)} \left\{ \frac{z_2 \sin 2\theta_2 (a^2 - c^2)}{(a^2 + q_2)(c^2 + q_2)} - \frac{2x\cos^2 \theta_2}{a^2 + q_2} - \frac{2x\sin^2 \theta_2}{c^2 + q_2} \right\}, \quad (15) \\ B_i &= 2y \left\{ \frac{1}{\sqrt{a^2 - c^2}} \log \frac{\sqrt{a^2 + q_i} + \sqrt{a^2 - c^2}}{\sqrt{c^2 + q_i}} - \frac{\sqrt{a^2 + q_i}}{c^2 + q_i} \right\} \end{aligned}$$

$$\begin{aligned}
B &= \frac{2yq_2(a^2 - c^2)}{\sqrt{a^2 + q_2}(c^2 + q_2)^2}, \\
C_i &= \frac{-3x\sin 2\theta_i + 2z_i(\cos^2\theta_i - 2\sin^2\theta_i)}{\sqrt{a^2 - c^2}} \log \frac{\sqrt{a^2 + q_i} + \sqrt{a^2 - c^2}}{\sqrt{c^2 + q_i}} \\
&\quad + \frac{2x\sin 2\theta_i + 4z_i\sin^2\theta_i}{\sqrt{a^2 + q_i}} - \frac{(-x\sin 2\theta_i + 2z_i\cos^2\theta_i)\sqrt{a^2 + q_i}}{c^2 + q_i} \\
C &= \frac{2\cos^2\theta_2 - 4\sin^2\theta_2}{\sqrt{a^2 - c^2}} \log \frac{\sqrt{a^2 + q_2} + \sqrt{a^2 - c^2}}{\sqrt{c^2 + q_2}} + \frac{4\sin^2\theta_2}{\sqrt{a^2 + q_2}} - \frac{2\cos^2\theta_2\sqrt{a^2 + q_2}}{c^2 + q_2} \\
&\quad - \frac{(a^2 - c^2)q_{2z}}{\sqrt{a^2 + q_2}(c^2 + q_2)} \left\{ \frac{x\sin 2\theta_2(a^2 - c^2)}{(a^2 + q_2)(c^2 + q_2)} - \frac{2z_2\sin^2\theta_2}{a^2 + q_2} - \frac{2z_2\cos^2\theta_2}{c^2 + q_2} \right\}
\end{aligned}$$

oblate spheroid

$$\begin{aligned}
r &= \frac{Sa^2c}{4(a^2 - c^2)} \\
A_i &= -\frac{2x(\cos^2\theta_i - 2\sin^2\theta_i) + 3z_i\sin 2\theta_i}{\sqrt{a^2 - c^2}} \tan^{-1} \frac{\sqrt{a^2 - c^2}}{\sqrt{c^2 + q_i}} \\
&\quad - \frac{4x\sin^2\theta_i - 2z_i\sin 2\theta_i}{\sqrt{c^2 + q_i}} + \frac{(2x\cos^2\theta_i + z_i\sin 2\theta_i)\sqrt{c^2 + q_i}}{a^2 + q_i} \\
A &= -\frac{3\sin 2\theta_2}{\sqrt{a^2 - c^2}} \tan^{-1} \frac{\sqrt{a^2 - c^2}}{\sqrt{c^2 + q_2}} + \frac{2\sin 2\theta_2}{\sqrt{c^2 + q_2}} + \frac{\sin 2\theta_2\sqrt{c^2 + q_2}}{a^2 + q_2} \\
&\quad + \frac{q_{2z}(a^2 - c^2)}{(a^2 + q_2)\sqrt{c^2 + q_2}} \left\{ \frac{2x\cos^2\theta_2 + z_2\sin 2\theta_2}{a^2 + q_2} + \frac{2x\sin^2\theta_2 - z_2\sin 2\theta_2}{c^2 + q_2} \right\} \\
B_i &= 2y \left\{ -\frac{1}{\sqrt{a^2 - c^2}} \tan^{-1} \frac{\sqrt{a^2 - c^2}}{\sqrt{c^2 + q_i}} + \frac{\sqrt{c^2 + q_i}}{a^2 + q_i} \right\} \\
B &= -\frac{2yq_{2z}(a^2 - c^2)}{(a^2 + q_2)\sqrt{c^2 + q_2}}, \tag{16} \\
C_i &= -\frac{3x\sin 2\theta_i + 2z_i(\sin^2\theta_i - 2\cos^2\theta_i)}{\sqrt{a^2 - c^2}} \tan^{-1} \frac{\sqrt{a^2 - c^2}}{\sqrt{c^2 + q_i}} \\
&\quad + \frac{2x\sin 2\theta_i - 4z_i\cos^2\theta_i}{\sqrt{c^2 + q_i}} + \frac{(x\sin 2\theta_i + 2z_i\sin^2\theta_i)\sqrt{c^2 + q_i}}{a^2 + q_i} \\
C &= \frac{4\cos^2\theta_2 - 2\sin^2\theta_2}{\sqrt{a^2 - c^2}} \tan^{-1} \frac{\sqrt{a^2 - c^2}}{\sqrt{c^2 + q_2}} - \frac{4\cos^2\theta_2}{\sqrt{c^2 + q_2}} + \frac{2\sin^2\theta_2\sqrt{c^2 + q_2}}{a^2 + q_2} \\
&\quad + \frac{q_{2z}(a^2 - c^2)}{(a^2 + q_2)\sqrt{c^2 + q_2}} \left\{ \frac{x\sin 2\theta_2 + 2z_2\sin^2\theta_2}{a^2 + q_2} + \frac{-x\sin 2\theta_2 + 2z_2\cos^2\theta_2}{c^2 + q_2} \right\}.
\end{aligned}$$

two spheres

$$r = \frac{Sa^3}{3},$$

$$\begin{aligned}
A_i &= -\left\{ \frac{x+b}{R_{i1}^3} + \frac{x-b}{R_{i2}^3} \right\} & A &= 3 \left\{ \frac{(x+b)(z+d_1)}{R_{21}^5} + \frac{(x-b)(z+d_2)}{R_{12}^5} \right\}, \\
B_i &= -y \left\{ \frac{1}{R_{i1}^3} + \frac{1}{R_{i2}^3} \right\} & B &= 3y \left\{ \frac{z+d_1}{R_{21}^5} + \frac{z+d_2}{R_{22}^5} \right\}, \\
C_1 &= -\left\{ \frac{z-d_1}{R_{11}^3} + \frac{z-d_2}{R_{12}^3} \right\} & C_2 &= -\left\{ \frac{z+d_1}{R_{21}^3} + \frac{z+d_2}{R_{22}^3} \right\} \\
C &= -\left\{ \frac{1}{R_{21}^5} + \frac{1}{R_{22}^5} - \frac{3(z+d_1)^2}{R_{21}^5} - \frac{3(z+d_2)^2}{R_{22}^5} \right\} \quad (17)
\end{aligned}$$

, where q_{1z} , q_{1y} , q_{1z} are the derivatives of q_i with respect to x , y , z .

Inserting the strains which are derived by differentiation of (14), (15), (16), (17) into (1), the stresses are as follows

$$\begin{aligned}
T_{xx} &= 2\mu\gamma \left(D_1 + (3-4\sigma)D_2 + 2zD - 4\sigma F_2 - \frac{1+\sigma}{1-2\sigma} \int_0^r adT \right) \\
T_{yy} &= 2\mu\gamma \left(E_1 + (3-4\sigma)E_2 + 2zE - 4\sigma F_2 - \frac{1+\sigma}{1-2\sigma} \int_0^r adT \right) \\
T_{zz} &= 2\mu\gamma \left(F_1 - F_2 + 2zF - \frac{1+\sigma}{1-2\sigma} \int_0^r adT \right) \\
T_{xy} &= \mu\gamma [G_1 + (3-4\sigma)G_2 + 2zG] \\
T_{yz} &= \mu\gamma [H_1 + H_2 + 2zH] \\
T_{zx} &= \mu\gamma [I_1 + I_2 + 2zI]. \quad (18)
\end{aligned}$$

prolate spheroid

$$\begin{aligned}
D_i &= \frac{4\cos^2\theta_i - 2\sin^2\theta_i}{\sqrt{a^2 - c^2}} \log \frac{\sqrt{a^2 + q_i} + \sqrt{a^2 - c^2}}{\sqrt{c^2 + q_i}} - \frac{4\cos^2\theta_i}{\sqrt{a^2 + q_i}} + \frac{2\sin^2\theta_i \sqrt{a^2 + q_i}}{c^2 + q_i} \\
&\quad + \frac{q_{1z}(a^2 - c^2)}{\sqrt{a^2 + q_i}(c^2 + q_i)} \left\{ \frac{(a^2 - c^2)z_i \sin 2\theta_i}{(a^2 + q_i)(c^2 + q_i)} - \frac{2x \sin^2\theta_i}{c^2 + q_i} - \frac{2x \cos^2\theta_i}{a^2 + q_i} \right\} \\
D &= \frac{q_{2x} \sin 2\theta_2 (a^2 - c^2)}{(a^2 + q_2)^{3/2} (c^2 + q_2)^2} - \frac{2q_{2z} (a^2 - c^2)}{\sqrt{a^2 + q_2} (c^2 + q_2)} \left\{ \frac{\sin^2\theta_2}{c^2 + q_2} + \frac{\cos^2\theta_2}{a^2 + q_2} \right\} \\
&\quad + \frac{q_{2xz} (a^2 - c^2)}{\sqrt{a^2 + q_2} (c^2 + q_2)} \left\{ \frac{z_2 \sin 2\theta_2 (a^2 - c^2)}{(a^2 + q_2)(c^2 + q_2)} - \frac{2x \sin^2\theta_2}{c^2 + q_2} - \frac{2x \cos^2\theta_2}{a^2 + q_2} \right\} \\
&\quad + \frac{q_{2xz} q_{2z} (a^2 - c^2)}{(a^2 + q_2)^{3/2} (c^2 + q_2)^2} \left\{ - \frac{z_2 \sin 2\theta_2 (a^2 - c^2) (2a^2 + 3c^2/2 + 7q_2/2)}{(a^2 + q_2)(c^2 + q_2)} \right. \\
&\quad \left. + \frac{2x \sin^2\theta_2 (2a^2 + c^2/2 + 5q_2/2)}{c^2 + q_2} + \frac{2x \cos^2\theta_2 (a^2 + 3c^2/2 + 5q_2/2)}{a^2 + q_2} \right\}, \\
E_i &= -\frac{2}{\sqrt{a^2 - c^2}} \log \frac{\sqrt{a^2 + q_i} + \sqrt{a^2 - c^2}}{\sqrt{c^2 + q_i}} + \frac{2\sqrt{a^2 + q_i}}{c^2 + q_i} + \frac{2y q_{1y} (a^2 - c^2)}{\sqrt{a^2 + q_i} (c^2 + q_i)^2} \\
E &= -\frac{2(q_{2z} + y q_{2zy}) (a^2 - c^2)}{\sqrt{a^2 + q_2} (a^2 + q_2)^2} + \frac{y q_{2y} q_{2z} (a^2 - c^2) (4a^2 + c^2 + 5q_2)}{(a^2 + q_2)^{3/2} (c^2 + q_2)^3}
\end{aligned}$$

$$\begin{aligned}
F_i &= \frac{4\sin^2\theta_i - 2\cos^2\theta_i}{\sqrt{a^2 - c^2}} \log \frac{\sqrt{a^2 + q_i} + \sqrt{a^2 - c^2}}{\sqrt{c^2 + q_i}} - \frac{4\sin^2\theta_i}{\sqrt{a^2 + q_i}} + \frac{2\cos^2\theta_i\sqrt{a^2 + q_i}}{c^2 + q_i} \\
&\quad + \frac{q_{iz}(a^2 - c^2)}{\sqrt{a^2 + q_i}(c^2 + q_i)} \left\{ \frac{x\sin 2\theta_i(a^2 - c^2)}{(a^2 + q_i)(c^2 + q_i)} - \frac{2z_i\sin^2\theta_i}{a^2 + q_i} - \frac{2z_i\cos^2\theta_i}{c^2 + q_i} \right\} \\
F &= -\frac{4q_{2z}(a^2 - c^2)}{\sqrt{a^2 + q_2}(c^2 + q_2)} \left\{ \frac{\sin^2\theta_2}{a^2 + q_2} + \frac{\cos^2\theta_2}{c^2 + q_2} \right\} + \frac{q_{2zz}(a^2 - c^2)}{\sqrt{a^2 + q_2}(c^2 + q_2)} \\
&\quad \times \left\{ \frac{x\sin 2\theta_2(a^2 - c^2)}{(a^2 + q_2)(c^2 + q_2)} - \frac{2z_2\sin^2\theta_2}{a^2 + q_2} - \frac{2z_2\cos^2\theta_2}{c^2 + q_2} \right\} \\
&\quad + \frac{q_{2z^2}(a^2 - c^2)}{(a^2 + q_2)^{3/2}(c^2 + q_2)^2} \left\{ -\frac{x\sin 2\theta_2(a^2 - c^2)(2a^2 + 3c^2/2 + 7q_2/2)}{(a^2 + q_2)(c^2 + q_2)} \right. \\
&\quad \left. + \frac{2z_2\sin^2\theta_2(a^2 + 3c^2/2 + 5q_2/2)}{a^2 + q_2} + \frac{2z_2\cos^2\theta_2(2a^2 + c^2/2 + 5q_2/2)}{c^2 + q_2} \right\} \\
G_i &= -\frac{4yq_{iz}(a^2 - c^2)}{\sqrt{a^2 + q_i}(c^2 + q_i)^2}, \\
G &= \frac{4(a^2 - c^2)y}{\sqrt{a^2 + q_2}(c^2 + q_2)} \left\{ -q_{2zz} + \frac{q_{2z}q_{2z}(a^2 + 4c^2 + 5q_2)}{2(a^2 + q_2)(c^2 + q_2)^2} \right\}, \\
H_i &= -\frac{4yq_{iz}(a^2 - c^2)}{\sqrt{a^2 + q_i}(c^2 + q_i)^2} \\
H &= \frac{4y(a^2 - c^2)}{\sqrt{a^2 + q_2}(c^2 + q_2)} \left\{ -q_{2zz} + \frac{q_{2z^2}(a^2 + 4c^2 + 5q_2)}{2(a^2 + q_2)(c^2 + q_2)^2} \right\}, \\
I_i &= \frac{6\sin 2\theta_i}{\sqrt{a^2 - c^2}} \log \frac{\sqrt{a^2 + q_i} + \sqrt{a^2 - c^2}}{\sqrt{c^2 + q_i}} - \frac{2\sin 2\theta_i(a^2 + 2c^2 + 3q_i)}{\sqrt{a^2 + q_i}(c^2 + q_i)} \\
&\quad + \frac{2q_{iz}(a^2 - c^2)}{\sqrt{a^2 + q_i}(c^2 + q_i)} \left\{ \frac{z_i\sin 2\theta_i(a^2 - c^2)}{(a^2 + q_i)(c^2 + q_i)} - \frac{2x\sin^2\theta_i}{c^2 + q_i} - \frac{2x\cos^2\theta_i}{a^2 + q_i} \right\} \\
I &= \frac{4q_{2z}\sin 2\theta_2(a^2 - c^2)^2}{(a^2 + q_2)^{3/2}(c^2 + q_2)^2} + \frac{2q_{2zz}(a^2 - c^2)}{\sqrt{a^2 + q_2}(c^2 + q_2)} \left\{ \frac{z_2\sin 2\theta_2(a^2 - c^2)}{(a^2 + q_2)(c^2 + q_2)} \right. \\
&\quad \left. - \frac{2x\cos^2\theta_2}{a^2 + q_2} - \frac{2x\sin^2\theta_2}{c^2 + q_2} \right\} + \frac{2q_{2z^2}(a^2 - c^2)}{(a^2 + q_2)^{3/2}(c^2 + q_2)^2} \\
&\quad \times \left\{ -\frac{z_2\sin 2\theta_2(a^2 - c^2)(2a^2 + 3c^2/2 + 7q_2/2)}{(a^2 + q_2)(c^2 + q_2)} + \frac{2x\sin^2\theta_2(2a^2 + c^2/2 + 5q_2/2)}{c^2 + q_2} \right. \\
&\quad \left. + \frac{2x\cos^2\theta_2(a^2 + 3c^2/2 + 5q_2/2)}{a^2 + q_2} \right\}.
\end{aligned}$$

oblate spheroid

$$\begin{aligned}
D_i &= \frac{2\cos^2\theta_i - 4\sin^2\theta_i}{\sqrt{a^2 - c^2}} \tan^{-1} \frac{\sqrt{a^2 - c^2}}{c^2 + q_i} + \frac{4\sin^2\theta_i}{\sqrt{c^2 + q_i}} - \frac{2\cos^2\theta_i\sqrt{c^2 + q_i}}{a^2 + q_i} \\
&\quad - \frac{q_{iz}(a^2 - c^2)}{(a^2 + q_i)\sqrt{c^2 + q_i}} \left\{ \frac{2x\cos^2\theta_i + z_i\sin 2\theta_i}{a^2 + q_i} + \frac{2x\sin^2\theta_i - z_i\sin 2\theta_i}{c^2 + q_i} \right\}
\end{aligned}$$

$$\begin{aligned}
D &= -\frac{q_{2z}(a^2-c^2)}{(a^2+q_2)\sqrt{c^2+q_2}} \left\{ \frac{2\cos^2\theta_2+\sin 2\theta_2}{a^2+q_2} + \frac{2\sin^2\theta_2-\sin 2\theta_2}{c^2+q_2} \right\} \\
&\quad - \frac{q_{2xz}(a^2-c^2)}{(a^2+q_2)\sqrt{c^2+q_2}} \left\{ \frac{2x\cos^2\theta_2+z_2\sin 2\theta_2}{a^2+q_2} + \frac{2x\sin^2\theta_2-z_2\sin 2\theta_2}{c^2+q_2} \right\} \\
&\quad + \frac{q_{2z}q_{2z}(a^2-c^2)}{(a^2+q_2)^2(c^2+q_2)^{3/2}} \left\{ \frac{(2x\cos^2\theta_2+z_2\sin 2\theta_2)(a^2/2+2c^2+5q_2/2)}{a^2+q_2} \right. \\
&\quad \left. + \frac{(3a^2/2+c^2+5q_2/2)(2x\sin^2\theta_2-z_2\sin 2\theta_2)}{c^2+q_2} \right\}, \\
E_t &= \frac{2}{\sqrt{a^2-c^2}} \tan^{-1} \frac{\sqrt{a^2-c^2}}{\sqrt{c^2+q_t}} - \frac{2\sqrt{c^2+q_t}}{a^2+q_t} - \frac{2yq_{tz}(a^2-c^2)}{(a^2+q_t)^2\sqrt{c^2+q_t}} \quad (20) \\
E &= -\frac{2(q_{2z}+yq_{2z})}{(a^2+q_2)^2\sqrt{c^2+q_2}} (a^2-c^2) + \frac{2yq_{2z}q_{2z}(a^2-c^2)(a^2/2+2c^2+5q_2/2)}{(a^2+q_2)^2(c^2+q_2)^{3/2}}, \\
F_t &= \frac{2\sin^2\theta_t-4\cos^2\theta_t}{\sqrt{a^2-c^2}} \tan^{-1} \frac{\sqrt{a^2-c^2}}{\sqrt{c^2+q_t}} + \frac{4\cos^2\theta_t}{\sqrt{c^2+q_t}} - \frac{2\sin^2\theta_t\sqrt{c^2+q_t}}{a^2+q_t} \\
&\quad - \frac{q_{tz}(a^2-c^2)}{(a^2+q_t)\sqrt{c^2+q_t}} \left\{ \frac{x\sin 2\theta_t+2z_t\sin^2\theta_t}{a^2+q_t} + \frac{2z_t\cos^2\theta_t-x\sin 2\theta_t}{c^2+q_t} \right\} \\
F &= -\frac{4q_{2z}(a^2-c^2)}{(a^2+q_2)\sqrt{c^2+q_2}} \left\{ \frac{\sin^2\theta_2}{a^2+q_2} + \frac{\cos^2\theta_2}{c^2+q_2} \right\} - \frac{q_{2xz}(a^2-c^2)}{(a^2+q_2)\sqrt{c^2+q_2}} \\
&\quad \times \left\{ \frac{x\sin 2\theta_2+2z_2\sin^2\theta_2}{a^2+q_2} + \frac{-x\sin 2\theta_2+2z_2\cos^2\theta_2}{c^2+q_2} \right\} + \frac{q_{2z}^2(a^2-c^2)}{(a^2+q_2)^2(c^2+q_2)^{3/2}} \\
&\quad \times \left\{ \frac{x\sin 2\theta_2+2z_2\sin^2\theta_2}{a^2+q_2} (a^2/2+2c^2+5q_2/2) \right. \\
&\quad \left. + \frac{(-x\sin 2\theta_2+2z_2\cos^2\theta_2)(3a^2/2+c^2+5q_2/2)}{c^2+q_2} \right\}, \\
G_t &= -\frac{4yq_{tz}(a^2-c^2)}{(a^2+q_t)^2\sqrt{c^2+q_t}}, \\
G &= \frac{4y(a^2-c^2)}{(a^2+q_2)\sqrt{c^2+q_2}} \left\{ -q_{2xz} + \frac{q_{2z}q_{2z}(a^2/2+2c^2+5q_2/2)}{(a^2+q_2)^2(c^2+q_2)} \right\}, \\
H_t &= \frac{-4yq_{tz}(a^2-c^2)}{(a^2+q_t)^2\sqrt{c^2+q_t}}, \\
H &= \frac{4y(a^2-c^2)}{(a^2+q_2)\sqrt{c^2+q_2}} \left\{ -q_{2xz} + \frac{q_{2z}^2(a^2/2+2c^2+5q_2/2)}{(a^2+q_2)^2(c^2+q_2)} \right\}, \\
I_t &= \frac{6\sin 2\theta_t}{\sqrt{a^2-c^2}} \tan^{-1} \frac{\sqrt{a^2-c^2}}{\sqrt{c^2+q_t}} - \frac{4\sin 2\theta_t(a^2+c^2/2+3q_t/2)}{(a^2+q_t)\sqrt{c^2+q_t}} \\
&\quad - \frac{2q_{tz}(a^2-c^2)}{(a^2+q_t)\sqrt{c^2+q_t}} \left\{ \frac{2x\cos^2\theta_t+z_t\sin 2\theta_t}{a^2+q_t} + \frac{2x\sin^2\theta_t-z_t\sin 2\theta_t}{c^2+q_t} \right\} \\
I &= \frac{4q_{2z}\sin 2\theta_2(a^2-c^2)^2}{(a^2+q_2)^2(c^2+q_2)^{3/2}} - \frac{2q_{2xz}(a^2-c^2)}{(a^2+q_2)\sqrt{c^2+q_2}} \left\{ \frac{2x\cos^2\theta_2+z_2\sin 2\theta_2}{a^2+q_2} \right.
\end{aligned}$$

$$\begin{aligned}
& + \frac{2x\sin^2\theta_2 - z_2\sin 2\theta_2}{c^2 + q_2} \left\} + \frac{2q_2z^2(a^2 - c^2)}{(a^2 + q_2)(c^2 + q_2)^{3/2}} \right. \\
& \times \left\{ \frac{(2x\cos^2\theta_2 + z_2\sin 2\theta_2)(a^2/2 + 2c^2 + 5q_2/2)}{a^2 + q_2} \right. \\
& \left. \left. + \frac{(2x\sin^2\theta_2 - z_2\sin 2\theta_2)(3a^2/2 + c^2 + 5q_2/2)}{c^2 + q_2} \right\}.
\end{aligned}$$

two spheres

$$\begin{aligned}
D_t &= \frac{1}{R_{11}^3} + \frac{1}{R_{12}^3} - \frac{3(x+b)^2}{R_{11}^5} - \frac{3(x-b)^2}{R_{12}^5} \\
D &= -\frac{3(z+d_1)}{R_{21}^5} - \frac{3(z+d_2)}{R_{22}^5} + \frac{15(x+b)^2(z+d_1)}{R_{21}^7} + \frac{15(x-b)^2(z+d_2)}{R_{22}^7}, \\
E_t &= \frac{1}{R_{11}^3} + \frac{1}{R_{12}^3} - \frac{3y^2}{R_{11}^5} - \frac{3y^2}{R_{12}^5} \\
E &= -\frac{3(z+d_1)}{R_{21}^5} - \frac{3(z+d_2)}{R_{22}^5} + \frac{15y^2(z+d_1)}{R_{21}^7} + \frac{15y^2(z+d_2)}{R_{22}^7} \\
F_1 &= \frac{1}{R_{11}^3} + \frac{1}{R_{12}^3} - \frac{3(z-d_1)^2}{R_{11}^5} - \frac{3(z-d_2)^2}{R_{12}^5} \\
F_2 &= \frac{1}{R_{21}^3} + \frac{1}{R_{22}^3} - \frac{3(z+d_1)^2}{R_{21}^5} - \frac{3(z+d_2)^2}{R_{22}^5} \\
F &= -\frac{9(z+d_1)}{R_{21}^5} - \frac{9(z+d_2)}{R_{22}^5} + \frac{15(z+d_1)^3}{R_{21}^7} + \frac{15(z+d_2)^3}{R_{22}^7}, \\
G_t &= -\frac{6y(x+b)}{R_{11}^5} - \frac{6y(x-b)}{R_{12}^5} \\
G &= \frac{30y(x+b)(z+d_1)}{R_{21}^7} + \frac{30y(x-b)(z+d_2)}{R_{22}^7}, \\
H_1 &= -\frac{6y(z-d_1)}{R_{11}^5} - \frac{6y(z-d_2)}{R_{12}^5} & H_2 &= -\frac{6y(z+d_1)}{R_{21}^5} - \frac{6y(z+d_2)}{R_{22}^5} \\
H &= -\frac{6yz}{R_{21}^5} - \frac{6yz}{R_{22}^5} + \frac{30y(z+d_1)^2}{R_{21}^7} + \frac{30y(z+d_2)^2}{R_{22}^7}, \\
I_1 &= -\frac{6(x+b)(z-d_1)}{R_{11}^5} - \frac{6(x-b)(z-d_2)}{R_{12}^5} \\
I_2 &= -\frac{6(x+b)(z+d_1)}{R_{21}^5} - \frac{6(x-b)(z+d_2)}{R_{22}^5} \\
I &= -\frac{6(x+b)}{R_{21}^5} - \frac{6(x-b)}{R_{22}^5} + \frac{30(x+b)(z+d_1)^2}{R_{21}^7} + \frac{30(x-b)(z+d_2)^2}{R_{22}^7},
\end{aligned}$$

§4

When

spheroid : $a=1$, $c=2$, $d=3$, $\theta_1 = -\pi/4$, $\sigma=1/4$

two spheres : $a=1$, $b=1$, $d_1=2$, $d_2=4$, $\sigma=1/4$

as shown in Fig. 2, the results of calculations of u, v, w at the free surface for the three kinds of the above thermal origin are shown in Fig. 3~6.

From the comparison of the case of prolate spheroid with that of oblate spheroid, both their states of displacements are resemble and the ratio of their quantities is nearly equal to that of their occupying




Fig. 2.




Fig. 3. Horizontal displacement for spheroid.

region 1 : 2. The ratio of the mean slope of the left side to that of the right is about 3 : 4. It may be difficult that we estimate the inclination of the thermal origin from the difference between the slopes of both sides, as it is small even at $\theta_1 = -\pi/4$. This tendency appears also in the case of two spheres, which occupy nearly the same place as the




Fig. 4 Vertical displacement for spheroid on $y=0$.




Fig. 5 Horizontal displacement for two spheres




Fig. 6 Vertical displacement for two spheres.




Fig. 7 τ_m for spheroid

that of spheroids.

Using the expressions for the above stresses (18), (19), (20), (21), we calculate the maximum value of shear stress

$$\tau_m = \sqrt{(T_{zz} - T_{xx})^2 + 4T_{zx}^2} / 2$$

in the plane $y=0$ along the intersection between the plane $y=0$ and the surface of the thermal origin of the described form. These results are shown in Fig. 7~8.

Their values are maximum about the minimums of the radius of curvature as we expect. But the difference between the value near the surface and that distant from there is small. For, toward the free surface, while the principal tension increases,

the principal compression decreases.

As the thermal origin is compressed by the surrounding material, the shear stress is very small within it, and then any fracture will not occur first in it.

It is evident from the expressions that the displacement and the stress are respectively in proportion to SD and $S\mu$, where D is the dimension of the thermal origin. We will estimate their values of rocks. Examples of coefficient of thermal expansion of rocks are shown in the following table.¹⁾




Fig. 8 τ_m for two spheroids.

Table 1 Thermal Expansion of Rocks

Rock	α (ordinary temperature)
Granite	8×10^{-6} deg ⁻¹
Basalt	5.4
Periclase	10
Andesites	7

While the values of coefficient of thermal expansion at high temperature (700°~1000°C) are ordinarily 2~4 times as much as the above, they decrease with pressure. But the effect of pressure is small and then those at about 700°C and 1000 atmosphere may be 2~3 times as much.¹⁾ Taking the rise of temperature 10°C and $\alpha_{tm} 3 \cdot 10^{-6}$ which is the mean value in its interval, assuming that the variation of α_t is linear and that D is 1 km, SD is

$$SD = \frac{(1+\sigma)D}{1-\sigma} \int_0^T \alpha dT = \frac{5}{3} D \alpha_{tm} T = \frac{5}{3} 10^5 \cdot 3 \cdot 10^{-6} \cdot 10 = 50$$

For example, the displacements w of prolate spheroid and oblate spheroid are ca. 11 cm and 20.5 cm.

Taking the rigidity of rock $3 \cdot 10^{11}$ dyne/cm² in the granite layer estimated from the seismic wave velocities, we obtain

$$S\mu = 5 \cdot 10^{-43} \cdot 10^{11} = 1.5 \cdot 10^8$$

For example, the maximum of τ_m for prolate spheroid is $2.3 \cdot 10^8$ dyne/cm².

Besides, in case that the temperature changes similarly also in the surrounding medium, we replace $\alpha_{tm} T$ with $(\alpha_{tm} - \alpha_{em}) T$.

In the above discussion, the elastic constants outside and inside the thermal origin have been assumed to be the same.

But, in fact, they differ if coefficients of thermal expansion are not equal. Nevertheless, if the configurations of the thermal origin are the same, the states of deformation may be similar, in spite of the differences between the absolute quantities. Thus, we may estimate the depth of the thermal origin and etc. from the figure of deformation at the surface.

§5.

From the results of successive levelings at Volcano Aso, the crustal move-




Fig. 9 Calculated curve of vertical displacement with observed results. (depth is 1.6km)

ment is remarkable in the region of radius of order of 2km near the craters.⁶⁾ This fact suggests that there is the thermal origin (magmatic reservoir) under the earth's surface, corresponding to this phenomena. That the depth of the upper surface of the thermal origin is ca. 0.86 km has been estimated from the records of eruption earthquakes.⁷⁾ Now assuming the thermal origin the spherical form for simplicity, we estimate the dimension. The displacement w at the surface is

$$w = \frac{\alpha^3 S \cdot d}{3(r^2 + d^2)^{3/2}} \quad (23)$$

, where d is the depth of centre of the thermal origin. The point = r where w is $1/\epsilon$ of

the maximum value at $r=0$, is

$$\frac{\alpha^3 S d}{3(r^2 + d^2)^{3/2}} / \frac{\alpha^3 S}{3d^2} = \frac{1}{\epsilon}$$

Then,

$$d = \frac{r}{\sqrt{\epsilon^{2/3} - 1}} \quad (24)$$

Evidently, r is not related to the radius a . Taking $\epsilon=4$ which corresponds to the margin of the remarkable crustal movement, we obtain $d = \text{ca. } 1.6 \text{ km}$ and then estimate the radius of thermal origin at $0.7 \sim 0.8 \text{ km}$. The vertical displacements which are calculated by inserting $d = 1.6 \text{ km}$ into (23) are shown in Fig. 9 with the observed results. Although the above inference is ambiguous in some respects, it may be said that the diameter is the order of 1 km .

The writer wishes to express his hearty thanks Prof. K. Sassa for his instructions.

Reference

- 1) F. Birch : Handbook of Physical Constants (1954)
- 2) J. N. Goodier : Phil. Mag. 23, 1017 (1937)
- 3) D. Kellogg : Foundations of Potential Theory (1929)
- 4) A. E. H. Love : Mathematical Theory of Elasticity (1927)
- 5) R. D. Mindlin : Physics 7, 195 (1936)
- 6) K. Yoshikawa : Zisin 7, 151 (1954)
- 7) K. Sassa : Mem. Coll. Soci. Kyoto Univ. Ser. A 18, 255 (1935)

Publications of the Disaster Prevention Research Institute

The Disaster Prevention Research Institute publishes reports of the research results in the form of bulletins. Publications not out of print may be obtained free of charge upon request to the Director, Disaster Prevention Research Institute, Kyoto University, Kyoto, Japan.

Bulletins :

- No. 1 On the Propagation of Flood Waves by Shoitiro Hayami, 1951.
No. 2 On the Effect of Sand Storm in Controlling the Mouth of the Kiku River by Tojiro Ishihara and Yuichi Iwagaki, 1952.
No. 3 Observation of Tidal Strain of the Earth (Part I) by Kenzo Sassa, Izuo Ozawa and Soji Yoshikawa. And Observation of Tidal Strain of the Earth by the Extensometer (Part II) by Izuo Ozawa, 1952.
No. 4 Earthquake Damages and Elastic Properties of the Ground by Ryo Tanabashi and Hatsuo Ishizaki, 1953.
No. 5 Some Studies on Beach Erosions by Shoitiro Hayami, Tojiro Ishihara and Yuichi Iwagaki, 1953.
No. 6 Study on Some Phenomena Foretelling the Occurrence of Destructive Earthquakes by Eiichi Nishimura, 1953.
No. 7 Vibration Problems of Skyscraper. Destructive Element of Seismic Waves for Structures by Ryo Tanabashi, Takuzi Kobori and Kiyoshi Kaneta, 1954.
No. 8 Studies on the Failure and the Settlement of Foundations by Sakurō Murayama, 1954.
No. 9 Experimental Studies on Meteorological Tsunamis Traveling up the Rivers and Canals in Osaka City by Shoitiro Hayami, Katsumasa Yano, Shohei Adachi and Hideaki Kunishi, 1955.
No.10 Fundamental Studies on the Runoff Analysis by Characteristics by Yuichi Iwagaki, 1955.
No.11 Fundamental Considerations on the Earthquake Resistant Properties of the Earth Dam by Motohiro Hatanaka, 1955.
No.12 The Effect of the Moisture Content on the Strength of an Alluvial Clay by Sakurō Murayama, Kōichi Akai and Tōru Shibata, 1955.
No.13 On Phenomena Forerunning Earthquakes by Kenzo Sassa and Eiichi Nishimura, 1956.
No.14 A Theoretical Study on Differential Settlements of Structures by Yoshitsura Yokoo and Kunio Yamagata, 1956.
No.15 Study on Elastic Strain of the Ground in Earth Tides by Izuo Ozawa, 1957.
No.16 Consideration on the Mechanism of Structural Cracking of Reinforced Concrete Buildings Due to Concrete Shrinkage by Yoshitsura Yokoo and S. Tsunoda, 1957.
No.17 On the Stress Analysis and the Stability Computation of Earth Embankments by Kōichi Akai, 1957.
No.18 On the Numerical Solutions of Harmonic, Biharmonic and Similar Equations by the Difference Method Not through Successive Approximations by Hatsuo Ishizaki, 1957.
No.19 On the Application of the Unit Hydrograph Method to Runoff Analysis for Rivers in Japan by Tojiro Ishihara and Akiharu Kanamaru, 1958.
No.20 Analysis of Statically Indeterminate Structures in the Ultimate State by Ryo Tanabashi, 1958.
No.21 The Propagation of Waves near Explosion and Fracture of Rock (I) by Soji Yoshikawa, 1958.
No.22 On the Second Volcanic Micro-Tremor at the Volcano Aso by Michiyasu Shima, 1958.
No.23 On the Observation of the Crustal Deformation and Meteorological Effect on It at Ide Observatory and on the Crustal Deformation Due to Full Water and Accumulating Sand in the Sabo-Dam by Michio Takada, 1958.
No.24 On the Character of Seepage Water and Their Effect on the Stability of Earth Embankments by Kōichi Akai, 1958.
No.25 On the Thermoelasticity in the Semi-infinite Elastic Solid by Michiyasu Shima

Bulletin No. 25 Published September, 1958

昭和 33 年 8 月 1 日 印刷

昭和 33 年 8 月 5 日 発行

編輯兼
発行者 京都大学防災研究所

印刷者 山代 多三郎

京都市上京区寺之内通小川西入

印刷所 山代印刷株式会社