

1 Sprout initiation and growth for three years after cutting in an abandoned secondary forest in Kyoto,  
2 Japan  
3  
4 Ayumi Imanishi<sup>1</sup>, Junko Morimoto<sup>2</sup>, Junichi Imanishi<sup>3</sup>, Shozo Shibata<sup>1</sup>, Asami Nakanishi<sup>1</sup>, Naoya  
5 Osawa<sup>4</sup>, Shinjiro Sakai<sup>1</sup>  
6  
7 1. Field Science Education and Research Center, Kyoto University, Kyoto, Japan  
8 2. Graduate School of Agriculture, Hokkaido University, Sapporo, Japan  
9 3. Graduate School of Global Environmental Studies, Kyoto University, Kyoto, Japan  
10 4. Graduate School of Agriculture, Kyoto University, Kyoto, Japan  
11  
12 Corresponding author: Ayumi Imanishi  
13 Field Science Education and Research Center, Kyoto University,  
14 Oiwake-cho, Sakyo-ku, Kyoto 606-8502, Japan  
15 E-mail: makinoa@kais.kyoto-u.ac.jp  
16 Tel.: +81-75-753-6434

1    Abstract

2            Secondary forests in Japan have been abandoned and the ecosystem has degraded since the high  
3    economic growth period. We carried out cutting in January in three small areas of a long-term  
4    abandoned secondary forest and investigated the sprout initiation and growth of woody plants for  
5    three years in order to reveal the early stage of sprout regeneration and to understand the sprouting  
6    ability and characteristics of each species for effective management. The percentage of sprouted  
7    stumps and the number of sprouting shoots was substantially maximized in autumn in the first year.  
8    These results suggested that autumn monitoring in the first year after cutting shows the maximum  
9    percentage of sprouted stumps and maximum sprout number, when cutting was conducted in the  
10    dormant season. For species characteristics, *Eurya japonica* showed low percentage of sprouted  
11    stumps in the lower plot, in which the mean DBH of the species was smaller than in the other plot.  
12    The sprout ability of *E. japonica* was deemed to be influenced by parent tree size. *Ilex pedunculosa*  
13    and *Lyonia ovalifolia* var. *elliptica* had high percentage of sprouted stumps and many sprouts. These  
14    species are useful for obtaining sprouting shoots, such as for firewood, but it is difficult to control  
15    their sprouting.

16

17    Keywords: suburban secondary forest, vegetation management, sprout ability

18

# 1 Introduction

2

3       Secondary forests in Japan were traditionally managed to obtain firewood, small timber and  
4 fertilizer (Takeuchi 2003). The sustainable management system maintained the unique biodiversity  
5 (Washitani 2003); however, the methods of obtaining energy and fertilizer have changed since the  
6 high economic growth period, and the value of secondary forests has gradually decreased (Takeuchi  
7 2003). In western Japan, most of these abandoned secondary forests have been changed from pine  
8 and deciduous broad-leaved forests to evergreen broad-leaved forests (Morimoto and Morimoto  
9 2003). The National Biodiversity Strategy of Japan, established in 2007, highlights the ecosystem  
10 degradation of the rural landscape, including the natural succession of secondary forests as one of  
11 three biodiversity crises (Ministry of the Environment 2007).

12       Recently, suburban secondary forests have been re-evaluated in terms of recreation,  
13 environmental education and the conservation of indigenous biodiversity and culture (Hattori et al.  
14 1995). At the site level, vegetation management is conducted by citizen volunteers and local  
15 governments (Shigematsu 2002), such as selective logging of evergreen shrubs and bamboos  
16 (Yamazaki et al. 2000; Yamase et al. 2005), and clearing underbrush and fallen leaves (Hosogi et al.  
17 2001; Kataoka et al. 2003); however, the number of citizen volunteers has been estimated at about  
18 20,000 and they can manage only 0.03 % of the area of secondary forests in Japan (Tsunekawa

2003). It is necessary to establish efficient adaptive vegetation management methods in secondary forests.

For vegetation management of secondary forests, it is essential to understand sprouting because secondary forests had been utilized and maintained by sprouting regeneration at 15–20-year cut intervals. The early stage of regeneration after cutting is an important period because plant species composition and density change dramatically within several years after cutting (Brokaw 1985, Breugel et al. 2006). In the early stage, sprouts generally grow earlier than seedlings and hinder seedlings from growing (Kennard et al. 2002). We reported about seedling regeneration after cutting (Imanishi et al. 2009). In the present study, we will report the early stage of sprout regeneration.

Sprouting is a major mode of regeneration of woody plants, not only after artificial disturbances such as cutting and burning (Uhl et al. 1981; Kammesheidt 1998), but also after natural disasters such as hurricanes and tornados (Glitzenstein and Harcombe 1988; Peterson and Pickett 1991; Bellingham et al. 1994) and after natural gap creation (Putz and Brokaw 1989). As a result, the factor that affects sprouting ability has attracted the interest of researchers and managers. Parent tree age (Kayll and Gimingham 1965; Johnson 1975), parent tree size (Johnson 1975; Jones and Raynal 1987), disturbance season (Blaisdell and Mueggler 1956; Malanson and Trabaud 1988; Babeux and Mauffette 1994) and site quality (Mroz et al. 1985; Forrester et al. 2003) are reported as the factors

that influence the sprouting ability and it was reported that sprouting ability differed among species (e.g. Mroz et al. 1985; Bellingham et al. 1994).

We investigated the sprouting of woody plants for three years after cutting in order to reveal the early stage of sprout regeneration and to understand the sprouting ability and characteristics of each species.

## Materials and methods

### *Study area and experimental design*

All research was conducted at Kamigamo Experimental Station (Field Science, Education and Research Center, Kyoto University, Kyoto) in western Japan (35°04'N, 135°46'E) (Fig. 1). The bedrock is sandstone and slate, and the soil type is brown forest soil. The annual mean temperature and the annual mean precipitation in the experimental forest from 1971 to 2000 were 14.6°C and 1581.8 mm, respectively (Kyoto University Forests 2002).

Three plots were set along a slope of the experimental forest to study sprouting ability and growth of woody plants under different light environments. The plots (upper, middle and lower) were 30 × 30 m, 20 × 30 m, and 40 × 15 m, respectively (Fig. 1). Cuttings were conducted after measuring stem diameters at breast height (DBHs) within these plots.

The area within and around plots became a part of the experimental forest from a national forest

in 1958. There is not an accurate data about the stand condition of the area before 1958. However, it was reported that the forest neighboring the area was cut clearly around 1925 and after that, dominated by Japanese red pine, *Pinus densiflora* Sieb. et Zucc. (Akai et al., 1986). Furthermore, it was assumed that the area might be also cut from 1920s to 1940s and the sapling of Hinoki, *Chamaecyparis obtusa* (Sieb. et Zucc.) Endl, which had naturally generated before cutting, might grow from the result of stem analyses of *C. obtusa* collected in the plots (Yanagimoto et al., 2000). After 1958, the area had been left under natural conditions. From 1970s, *P. densiflora* began to be killed by pine wilt disease and most *P. densiflora* died in early 1990s. This area is now dominated by *C. obtusa*.

To characterize the vegetation in the study area before cutting, we identified all trees taller than 1.3 m and measured DBHs in the three plots in August 1999. The percentage of total basal area (BA) was calculated for each species in the three plots. The BA of *C. obtusa* in the upper plot was 40.1 m<sup>2</sup>/ha, accounted for 88.7 % of total BA (Table 1). For shrub species, it was dominated by *Eurya japonica* Thunb. (0.97 m<sup>2</sup>/ha; 2.1 %) and *Rhododendron reticulatum* D. Don (0.85 m<sup>2</sup>/ha; 1.9 %) (Table 1). In the middle plot, the BA of *C. obtusa* was 37.7 m<sup>2</sup>/ha (93.5 %) (Table 1). For shrub species, it was dominated by *E. japonica* (0.53 m<sup>2</sup>/ha; 1.3 %) and *Camellia japonica* L. (0.46 m<sup>2</sup>/ha; 1.2 %) (Table 1). In the lower plot, it was dominated by *C. obtusa* (19.3 m<sup>2</sup>/ha; 46.1 %) and *Quercus serrata* Thunb. ex Murray (12.3 m<sup>2</sup>/ha; 29.3 %) (Table 1). For shrub species, it was dominated by

1 *Camellia japonica* (4.32 m<sup>2</sup>/ha; 10.3 %) (Table 1).

2 Almost all of the trees were cut in January 2000 in order to change the dominant species from *C.*  
3 *obtusa* to *P. densiflora* and/or deciduous broad-leaved trees; however, to improve vegetation  
4 recovery after cutting, several mother trees with a DBH greater than 5 cm were left in each of the  
5 experimental plots (Table 1).

6 *Sprout measurements*

7 We established survey areas within the three plots to acquire data on sprouts and light  
8 environment by excluding edges of the plots that were significantly shaded by the surrounding trees  
9 so that the survey areas were kept more homogeneous for sprouts. The survey areas were 20 × 20 m  
10 in the upper and middle plots and 15 × 25 m in the lower plot. 8 species, 379 stumps within the  
11 survey areas were surveyed (Table 2).

12 The number of sprouting shoots per stump within each survey area was recorded in May,  
13 September and November 2000, May and November 2001, and May and November 2002. In order  
14 to estimate the sprout growth of a stump, the height of the dominant shoot of each stump were  
15 measured in May and November 2001 and May and November 2002.

16 *Measurement of solar radiation*

17 On 29 August and 2 September 2002, hemispherical photographs were taken at 47 points (16 in  
18 the upper plot, 16 in the middle plot and 15 in the lower plot). The points were set at the center of 5

× 5 m quadrats into which each survey area was divided. Hemispherical photographs were taken 1.0 m above ground level in each quadrat using a Nikon Coolpix 995 with a fish-eye lens.

The photographs were analyzed using Gap Light Analyzer ver. 2.0 software (Frazer et al. 1999) to estimate the relative solar radiation from May to November. The radiation parameter to derive above-canopy solar radiation data was applied to modelled.

#### *Statistical analyses*

Relative solar radiation was compared among the plots using analysis of variance (ANOVA) and Tukey's HSD Test. DBHs of parent trees of each species were compared among the plots by the Mann-Whitney test with Bonferroni correction.

The percentages of sprouted stumps among the plots were tested in each species by the chi-squared test and Haberman's residual analysis. The timing of sprout initiation and death of each species was analyzed in each plot by comparing the numbers of sprouting shoots between survey times using the Wilcoxon signed rank test for matched pairs with Bonferroni correction. Then, the numbers of sprouting shoots in November 2001 were compared by the Mann-Whitney test between the groups that had significantly decreased in 2002 and that had not decreased to confirm if greater number of sprouting shoots in the previous year leads to greater rate of death. Heights of dominant shoots of each species were compared among the plots by the Mann-Whitney test with Bonferroni correction to see if sprout growth was different among the plots. The relationship between number of

sprouting shoots and height of dominant shoots in November 2001 was studied by Spearman's rank correlation analysis.

Statistical significance was set at  $p < 0.05$  and statistical analyses were performed using software (SPSS for Windows, version 13.0J; SPSS Inc. 2004).

## Results

### *Relative solar radiation*

Relative solar radiation in the upper plot was significantly higher than in the middle and lower plots (Fig. 2). In the middle plot, it was significantly higher than in the lower plot (Fig. 2).

### *DBH*

For *E. japonica*, the mean DBH in the lower plot was significantly smaller than in the upper and middle plots (Table 2). The mean DBH of *Camellia japonica* was smaller in the middle plot than the lower plot whereas that of *Cleyera japonica* Thunb. was larger in the middle plot than the lower plot (Table 2). The mean DBH of *Lyonia ovalifolia* (Wall.) Drude var. *elliptica* (Sieb. et Zucc.) Hand.-Mazz. was smaller in the middle plot than the upper plot whereas *R. reticulatum* was larger in the middle plot than the upper plot (Table 2).

### *Percentage of sprouted stumps*

The percentage of sprouted stumps in May 2000 differed among species (Fig. 3), varying from

0 % of *Camellia japonica* to 100 % of *L. ovalifolia* var. *elliptica*. In September 2000, the percentage of sprouted stumps of each species increased to more than 80 %, except for *E. japonica* in the lower plot (Fig. 3). In November 2002, there were some dead stumps that had once sprouted (Fig. 3).

For *E. japonica*, the actual percentages of sprouted stumps in the upper and middle plots were significantly higher than in the lower plot in November 2002 (Fig. 3).

#### *Number of sprouting shoots*

Almost all sprouting shoots initiated during 2000 (Fig. 4). While the two *Rhododendron* species in the upper plot continued to sprout shoots until May 2001, the number of shoots of *Ilex pedunculosa* Miq. in the upper plot and *Camellia japonica* in the lower plot had significantly decreased by May 2001 (Fig. 4). In 2002, the number of shoots of *E. japonica* in the upper and middle plots, *L. ovalifolia* var. *elliptica*, and the two *Rhododendron* species began to significantly decrease (Fig. 4). Species in which the number of sprouting shoots significantly decreased in 2002 had significantly more shoots in November 2001 than other species (Fig. 5).

#### *Height of the dominant shoot*

The height of the dominant shoot increased from May 2001 to November 2002, varying from 1.3 times in *L. ovalifolia* var. *elliptica* to 2.8 times in *Cleyera japonica* in the lower plot (Fig. 6). In November 2002, the height of *E. japonica* in the lower plot was significantly lower than in the upper and middle plots, and the heights of *R. reticulatum* in the upper plot and *Cleyera japonica* in the

lower plot were significantly lower than in the middle plot (Fig. 6).

*E. japonica* and *R. reticulatum* in the upper plot had significantly positive weak correlations between the number of sprouting shoots and the height of the dominant shoot (Table 3). *E. japonica* in the middle plot had significantly positive moderate correlation, and *I. pedunculosa* had significantly positive strong correlation (Table 3).

## Discussion

### *Sprout regeneration for three years after cutting*

More than 80 % of stumps of each species, except *E. japonica* in the lower plot, had sprouted by September in the first year after cutting (Fig. 3). The number of sprouting shoots was substantially maximized in the first growing season after cutting; the exception being two *Rhododendron* species in the upper plot in which shoots slightly increased until May 2001 (Fig. 4). Shima et al. (1989) also found that sprout initiation finished in the first year after cutting when cutting was conducted during the dormant season.

In the second or third year after cutting, sprouting shoots began to decrease (Fig. 4). Species with more shoots showed significantly reduced sprout numbers (Fig. 5). Tanaka (1989) and Rydberg (2000) reported that numerous sprout productions led to a rapid self-thinning process. However, in the present study, it is not clear that the decline of sprout number was caused by competition among

shoots within a stump or competition with the other stumps and seedlings.

Some studies have found a negative correlation between sprout number and the mean (Katagiri 1986; Manabe et al. 1991) or maximum (Brown 1994) height of species with numerous sprouts; however, *E. japonica*, *I. pedunculosa* and *R. reticulatum* in our study site had a significantly positive correlation between sprout number and the height of dominant shoots (Table 3). One possible reason is that because this study area had sufficient solar radiation, especially in the upper and middle plots (Fig. 2), competition among shoots and/or competition with the other stumps and seedlings might not be severe and vigorous stumps might have many long shoots.

#### *Species characteristics*

##### *Percentage of sprouted stumps*

For *E. japonica*, an evergreen species, the percentage of sprouted stumps in the lower plot was significantly lower than in the upper and middle plots in November 2002 (Fig. 3). The main reason for this result was considered that the mean DBH of *E. japonica* in the lower plot was significantly smaller than in the upper and middle plots (Table 2).

##### *Sprout number*

As reported, sprout number varied with species (e.g., Mroz et al. 1985; Bellingham et al. 1994). *L. ovalifolia* var. *elliptica* had an average of more than 30 sprouts per stump at maximum and *I.*

*pedunculosa* had an average of 28 sprouts per stump at maximum (Fig. 4). Yamase (2000) reported that *L. ovalifolia* var. *elliptica* had many sprouts and the sprouts of both *L. ovalifolia* var. *elliptica* and *I. pedunculosa* grew rapidly. Since in the present study, these species had a high percentage of sprouted stumps (Fig. 3) and grew rapidly (Fig. 6), sprouting regeneration seems an important strategy for their survival in highly managed forests, such as *satoyama*.

#### Height

For *E. japonica* and *Cleyera japonica*, dominant shoots were significantly lower in the lower plot than in the upper and/or middle plots in November 2002 (Fig. 6). Manabe et al. (1991) reported that the mean sprout length of *E. japonica* was shorter in darker conditions. In the present study, relative solar radiation was significantly lower in the lower plot than in the upper and middle plot (Fig. 2). However, since the mean DBH of *E. japonica* and *Cleyera japonica* was significantly smaller in the lower plot than in the upper and/or middle plots (Table 2), it is not clear that the growth increment of these species is dependent on the amount of photosynthesis or the reserve nutrient in a stump.

#### Conclusion

We investigated the stump sprouting of 8 species for three years after cutting a small area (0.06–0.09 ha). Cutting was conducted in three plots in a 50-year abandoned secondary forest in

1 January 2000. The percentage of sprouted stumps and the number of sprouting shoots was  
2 substantially maximized in autumn in the first year. The mortality of stumps sprouted by September  
3 2000 was 1.7 %, partially due to the high solar radiation in our plots. The number of sprouting  
4 shoots began to decrease in the second or third year. These results suggested that monitoring in  
5 autumn in the first year after cutting is likely to elucidate the maximum percentage of sprouted  
6 stumps and maximum sprout number when cutting was conducted in the dormant season.

7 As species characteristics, *E. japonica* had dead stumps and showed suppressed growth in the  
8 lower plot. *Cleyera japonica* was also suppressed growth in the lower plot. The sprout ability of  
9 these two species was deemed to be influenced by parent tree size because the mean DBH of these  
10 species in the lower plot was significantly smaller than in the upper and/or middle plots. However,  
11 since relative solar radiation was also significantly lower in the lower plot than in the other plot,  
12 growth increment might be influenced by the amount of photosynthesis as well as the reserve  
13 nutrient in a stump.

14 Sprouting is an important component for the management of suburban secondary forests. It is  
15 desirable to manage sprouting appropriately based on understanding species characteristics and to  
16 monitor sprout regeneration efficiently after cutting with feedback to the management.

References

Akai T, Nakai I, Okamoto N, Watanabe M (1986) Amenity management and structure of natural regenerated mixture forests of Akamatsu (*Pinus densiflora*) and Hinoki (*Chamaecyparis obtusa*) at the urban fringe of Kyoto. Bulletin of the Kyoto University Forests 57: 128-142 (in Japanese with English abstract)

Babeux P, Mauffette Y (1994) The effects of early and late spring cuts on the sprouting success of red maple (*Acer rubrum*) in northwestern Quebec. Canadian Journal of Forest Research 24: 785-791

Bellingham PJ, Tanner EVJ, Healey JR (1994) Sprouting trees in Jamaican montane forests, after a hurricane. Journal of Ecology 82: 747-758

Blaisdell JP, Mueggler WF (1956) Sprouting of bitterbrush (*Purshia tridentata*) following burning of top removal. Ecology 37(2): 365-370

Breugel MV, Martínez-Ramos M, Bongers F (2006) Community dynamics during early secondary succession in Mexican tropical rain forests. Journal of Tropical Ecology 22: 663-674

- 1 Brokaw NVL (1985) Gap-phase regeneration in a tropical forest. *Ecology* 66(3): 682-687
- 2
- 3 Brown D (1994) The development of woody vegetation in the first 6 years following clear-cutting of
- 4 a hardwood forest for a utility right-of-way. *Forest Ecology and Management* 65: 171-181
- 5
- 6 Frazer GW, Canham CD, Lertzman KP (1999) Gap Light Analyzer (GLA) Imaging
- 7 software to extract canopy structure and gap light transmission indices from true-colour
- 8 fisheye photographs, User Manual and Program Documentation, Version 2.0. Simon
- 9 Fraser University, Burnaby, British Columbia, CANADA, and Institute of Ecosystem
- 10 Studies, Millbrook.
- 11
- 12 Forrester F, Bauhus J, Connell M (2003) Competition in thinned Silvertop Ash (*Eucalyptus sieberi* L.
- 13 Johnson) stands from early coppice growth. *Forest Ecology and Management* 174: 459-475
- 14
- 15 Glitzenstein JS, Harcombe PA (1988) Effects of the December 1983 Tornado on forest vegetation of
- 16 the Big Thicket, Southeast Texas, U. S. A. *Forest Ecology and Management* 25: 269-290
- 17
- 18 Hattori T, Akamatsu H, Takeda Y, Kodate S, Kamihogi A, Yamazaki H (1995) A study on the actual

- 1 conditions of *Satoyama* (rural forests) and their management. *Humans and Nature* 6: 1-32 (in  
2 Japanese with English abstract)
- 3
- 4 Hosogi D, Kuno H, Arai K, Fukada K (2001) Effects of alternation of management to forest floor  
5 vegetation and environment of secondary coppice forests in urban area, Tokyo I. Relation between  
6 the forest vegetation management method and growth of upper layer trees or forest floor vegetation.  
7 *Journal of the Japanese Society of Revegetation Technology* 27(1): 14-19 (in Japanese with English  
8 abstract)
- 9
- 10 Imanishi A, Shibata S, Imanishi J, Terai A, Nakanishi A, Sakai S, Osawa N, Morimoto Y (2009)  
11 Change of woody-species composition in an early stage of plant succession after cutting in a  
12 suburban secondary forest dominated by *Chamaecyparis obtusa*. *Journal of the Japanese Society of  
13 Revegetation Technology* 34(4): 641-648 (in Japanese with English abstract)
- 14
- 15 Johnson PS (1975) Growth and structural development of red oak sprout clumps. *Forest Science*  
16 21(4): 413-418
- 17
- 18 Jones RH, Raynal DJ (1987) Root sprouting in American beech: production, survival, and the effect

of parent tree vigor. Canadian Journal of Forest Research 17: 539-544

Kammesheidt L (1998) The role of tree sprouts in the restoration of stand structure and species diversity in tropical moist forest after slash-and-burn agriculture in Eastern Paraguay. Plant Ecology 139: 155-165

Katagiri S (1986) Studied on sprout regeneration of Deciduous Broad-leaved trees (I) The growth of sprout shoots during five years after cutting Studies of the San'in Region. Forest resources 2: 27-36 (in Japanese)

Kataoka H, Shima K, Chiba K (2003) The change of forest environments with the conventional management in broad-leaved secondary forest and its effects on regeneration of tree species. Journal of the Japanese Society of Revegetation Technology 29(1): 297-300 (in Japanese with English abstract)

Kayll AJ, Gimingham CH (1965) Vegetative regeneration of *Calluna vulgaris* after fire. Journal of Ecology 53: 729-734

- 1 Kennard DK, Gould K, Putz FE, Fredericksen TS, Morales F (2002) Effect of disturbance intensity  
2 on regeneration mechanisms in a tropical dry forest. *Forest Ecology and Management* 162: 197-208  
3
- 4 Kyoto University Forests (2002) Meteorological observations in the Kyoto University Forests,  
5 Kyoto University Forest, Kyoto (in Japanese)  
6
- 7 Malanson GP, Trabaud L (1988) Vigour of post-fire resprouting by *Quercus coccifera* L., *Journal of*  
8 *Ecology* 76: 351-365  
9
- 10 Manabe T, Yamamoto S, Chiba K (1991) Stump sprouting of evergreen shrub, *Eurya japonica*,  
11 within a current year of the cutting, *Journal of the Japanese Society of Revegetation Technology*  
12 16(4): 1-9 (in Japanese with English abstract)  
13
- 14 Ministry of the Environment (2007) The Third National Biodiversity Strategy of Japan,  
15 <http://www.env.go.jp/en/focus/attach/071210-e.pdf> Accessed 10 December 2008  
16
- 17 Morimoto J, Morimoto Y (2003) Satoyama Landscape Transition in the Kansai Area. In: Takeuchi K,  
18 Brown RD, Washitani I, Tsunekawa A, Yokohari M (ed) *Satoyama –The Traditional Rural*

- 1 Landscape of Japan, Springer, Tokyo: pp 60-71
- 2
- 3 Mroz GD, Frederick DJ, Jurgensen MF (1985) Site and fertilizer effects on northern hardwood
- 4 stump sprouting. Canadian Journal of Forest Research 15: 535-543
- 5
- 6 Peterson CJ, Pickett STA (1991) Treefall and resprouting following catastrophic windthrow in an
- 7 old-growth hemlock-hardwoods forest. Forest Ecology and Management 42: 205-217
- 8
- 9 Putz FE, Brokaw NVL (1989) Sprouting of broken trees on Barro Colorado Island, Panama. Ecology
- 10 70(2): 508-512
- 11
- 12 Rydberg D (2000) Initial sprouting, growth and mortality of European aspen and birch after selective
- 13 coppicing in central Sweden. Forest Ecology and Management 130: 27-35
- 14
- 15 Shima K, Katagiri S, Kaneko N (1989) Dynamics of sprout shoots during the two years after cutting
- 16 a Konara (*Quercus serrata*) secondary forest, Journal of the Japanese Forest Society 71(10): 410-416.
- 17 (in Japanese)
- 18

Shigematsu T (2002) Situation and potentiality of suburban and rural *Satoyama* woodlands including fragmented urban forest reserves in present society. Design Issue of Kyusyu institute of technology 5: 1-11 (in Japanese with English abstract)

Takeuchi K (2003) *Satoyama Landscapes as Managed Nature*. In: Takeuchi K, Brown RD, Washitani I, Tsunekawa A, Yokohari M (ed) *Satoyama –The Traditional Rural Landscape of Japan*, Springer, Tokyo: pp 9-16

Tanaka S (1989) Death and Growth of sprouting shoots of *Quercus crispula* and thinning of sprouting shoots, Northern Forestry 41(6): 7-14. (in Japanese)

Tsunekawa A (2003) Strategic Management of *Satoyama* Landscapes. In: Takeuchi K, Brown RD, Washitani I, Tsunekawa A, Yokohari M (ed) *Satoyama –The Traditional Rural Landscape of Japan*, Springer, Tokyo: pp 179-191

Uhl C, Clark K, Clark H, Murphy P (1981) Early plant succession after cutting and burning in the upper Rio Negro of the Amazon Basin. Journal of Ecology 69: 631-649

- 1 Washitani I (2003) Satoyama Landscapes and Conservation Ecology. In: Takeuchi K, Brown RD,  
2 Washitani I, Tsunekawa A, Yokohari M (ed) *Satoyama* –The Traditional Rural Landscape of Japan,  
3 Springer, Tokyo: pp 16-23  
4  
5 Yamase K (2000) Stump sprouting of main woody species in secondary forest of pine mass-dieback  
6 under different thinning intensity. Journal of Japanese Institute of Landscape Architecture 63(5):  
7 465-468 (in Japanese with English abstract)  
8  
9 Yamase K, Hattori T, Mikami K, Tanaka A (2005) Species richness and species composition of the  
10 coppice forests after “Hyogo-Model” vegetation management. Journal of Japanese Institute of  
11 Landscape Architecture 68(5): 655-658 (in Japanese with English abstract)  
12  
13 Yamazaki H, Aoki K, Hattori T, Takeda Y (2000) Increase of species diversity due to management of  
14 vegetation in *Satoyama* (rural forests). Journal of Japanese Institute of Landscape Architecture 63(5):  
15 481-484 (in Japanese with English abstract)  
16  
17 Yanagimoto J, Shibata Y, Hirai T, Hashida R, Nakagawa T, Shibata S (2000)  
18 Construction of the volume table and succession for secondary forest of Hinoki in

- 1 Kamigamo experimental forest. Bulletin of the Kyoto University Forests 2000: 215-216
- 2 (in Japanese)

Table 1 Summary of cutting plots and woody plants pre- and post-cutting

		Upper plot		Middle plot		Lower plot	
Cutting area (m×m)		30×30		30×20		15×40	
Slope orientation		N 16°E		N 46°E		N 45°E	
Elevation (m)		203-214		176-190		144-168	
Slope gradient (°)		17.0		28.0		27.0	
Pre-cutting (Aug 1999)	Species richness	18		16		21	
	Population density (n/ha)	7,789		4,717		5,500	
	Woody plants (DBH ≥ 1cm)	Trees		Trees		Trees	
		BA (m <sup>2</sup> /ha)		BA (m <sup>2</sup> /ha)		BA (m <sup>2</sup> /ha)	
		<i>Chamaecyparis obtusa</i>		<i>Chamaecyparis obtusa</i>		<i>Chamaecyparis obtusa</i>	
		40.12		37.66		19.26	
Post-cutting (Jan 2000)	Remnant trees	<i>Pinus densiflora</i>		<i>Pinus densiflora</i>		<i>Quercus serrata</i>	
		1.53		0.93		12.25	
		Shrubs		Shrubs		Shrubs	
		BA (m <sup>2</sup> /ha)		BA (m <sup>2</sup> /ha)		BA (m <sup>2</sup> /ha)	
		<i>Eurya japonica</i>		<i>Eurya japonica</i>		<i>Camellia japonica</i>	
		0.97		0.53		4.32	
		<i>Rhododendron reticulatum</i>		<i>Camellia japonica</i>		<i>Wisteria floribunda</i>	
		0.85		0.46		0.19	
		5		2		5	
		111		50		400	
		Species		Species		Species	
		n		n		n	
		<i>Lyonia ovalifolia</i> var. <i>elliptica</i>		<i>Chamaecyparis obtusa</i>		<i>Quercus serrata</i>	
		3		2		13	
		<i>Pinus densiflora</i>		<i>Pinus densiflora</i>		<i>Acanthopanax sciadophylloides</i>	
		2		1		5	
		<i>Chamaecyparis obtusa</i>				<i>Chamaecyparis obtusa</i>	
		2				4	
		<i>Clethra barvinervis</i>				<i>Prunus grayana</i>	
		2				1	
		<i>Acanthopanax sciadophylloides</i>				<i>Clethra barvinervis</i>	
		1				1	

Table 2 Number and DBH of stumps of each species before the cutting

Species	Upper plot			Middle plot			Lower plot		
	n	DBH*		n	DBH*		n	DBH*	
<i>Eurya japonica</i>	71	2.44	$\pm 0.14^a$	34	2.58	$\pm 0.16^a$	28	1.53	$\pm 0.19^b$
<i>Camellia japonica</i>	7	2.75	$\pm 0.87$	10	2.53	$\pm 0.59^a$	77	4.41	$\pm 0.25^b$
<i>Cleyera japonica</i>	-	-		7	3.36	$\pm 0.70^a$	18	1.08	$\pm 0.16^b$
<i>Pieris japonica</i>	9	1.85	$\pm 0.32$	-	-		-	-	
<i>Ilex pedunculosa</i>	11	6.15	$\pm 0.74$	-	-		-	-	
<i>Lyonia ovalifolia</i> var. <i>elliptica</i>	16	2.99	$\pm 0.31^a$	7	1.81	$\pm 0.62^b$	-	-	
<i>Rhododendron reticulatum</i>	68	1.88	$\pm 0.09^a$	14	2.23	$\pm 0.21^b$	-	-	
<i>Rhododendron macrosepalum</i>	9	1.56	$\pm 0.13$	-	-		-	-	

\*: Mean  $\pm$  SE

Different letters indicate significant differences among the plots.

Table 3 Spearman's correlation coefficients between number of sprouting shoots and height of dominant shoot in November 2001

Species		n	correlation coefficients
<i>Eurya japonica</i>	upper	71	0.30 *
	middle	34	0.48 **
	lower	28	-0.05
<i>Rhododendron reticulatum</i>	upper	68	0.28 *
	middle	14	0.12
<i>Cleyera japonica</i>	middle	7	0.74
	lower	18	-0.09
<i>Lyonia ovalifolia</i> var. <i>elliptica</i>		23	0.30
<i>Ilex pedunculosa</i>		11	0.86 *
<i>Camellia japonica</i>		93	0.10

\*:  $p < 0.05$

\*\*:  $p < 0.01$

**Figure captions**

**Fig. 1** Study area and location of cutting plots

**Fig. 2** Relative solar radiation from May to November in each plot estimated from hemispherical photos

Error bars show standard errors.

Different letters indicate significant differences among plots.

**Fig. 3** Percentage of sprouted stumps of each species in May and September 2000, November 2002

Different letters indicate significant differences among plots.

(u) and (m) indicate upper plot and middle plot, respectively.

(A) and (C) indicate the actual and cumulative percentage of sprouted stumps, respectively.

**Fig. 4** Mean number of sprouting shoots per stump during the 3 years after cutting

Error bars show standard errors.

"i" or "d" shows that the number of sprouting shoots significantly increased or decreased compared to the previous survey, respectively.

(u) and (m) indicate upper plot and middle plot, respectively.

**Fig. 5** Mean number of sprouting shoots in November 2001 of species that significantly

- 1        decreased or not in 2002, comparison to the previous survey
- 2        Error bar shows standard error of the mean.
- 3        \* significant difference.
- 4        **Fig. 6**   Height of dominant shoots (cm) from May 2001 to November 2002
- 5        Error bars show standard errors.
- 6        Different letters indicate significant differences among plots.

Fig.1




Fig.2




Fig.3




Fig.4




Fig.5




Fig.6

