TITLE:

The predatory mite *Neoseiulus womersleyi* (Acari: Phytoseiidae) follows extracts of trails left by the two-spotted spider mite *Tetranychus urticae* (Acari: Tetranychidae).

AUTHOR(S):

Shinmen, Tsubasa; Yano, Shuichi; Osakabe, Masahiro

CITATION:

Shinmen, Tsubasa ...[et al]. The predatory mite *Neoseiulus womersleyi* (Acari: Phytoseiidae) follows extracts of trails left by the two-spotted spider mite *Tetranychus urticae* (Acari: Tetranychidae).. *Experimental & applied acarology* 2010, 52(2): 111-118

ISSUE DATE:

2010-10

URL:

<http://hdl.handle.net/2433/129433>

RIGHT:

The original publication is available at www.springerlink.com; This is not the published version. Please cite only the published version.; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。

**The predatory mite *Neoseiulus womersleyi* (Acari: Phytoseiidae) follows extracts
of trails left by the two-spotted spider mite *Tetranychus urticae* (Acari:
Tetranychidae)**

Tsubasa Shinmen • Shuichi Yano • Masahiro Osakabe

T. Shinmen

Laboratory of Ecological Information, Faculty of Agriculture, Kyoto University, Sakyo-ku, Kyoto
606-8502, Japan

(Present address) Pharmaceutical Research Laboratories No.2, Research & Development
Headquarters, Lion Corporation, Hirai 7-13-12, Edogawa-ku, Tokyo 132-0035; Japan

S. Yano* • Mh. Osakabe

Laboratory of Ecological Information, Graduate School of Agriculture, Kyoto University, Sakyo-ku,
Kyoto 606-8502, Japan

*Corresponding Author

e-mail: yano@kais.kyoto-u.ac.jp

Tel.: 81-75-753-6144

Fax: 81-75-753-6144

Abstract As it walks, the two-spotted spider mite *Tetranychus urticae* Koch (Acari: Tetranychidae) spins a trail of silk threads, that is followed by the predatory mite, *Neoseiulus womersleyi* Schicha (Acari: Phytoseiidae). Starved adult female *N. womersleyi* followed *T. urticae* trails laid down by five *T. urticae* females but did not follow a trail of one *T. urticae* female, suggesting that the amount of spun threads and their chemical components should correlate positively with the number of *T. urticae* individuals. To examine whether chemical components of *T. urticae* trails are responsible for the predatory mite's trail following, we collected separate *T. urticae* threads from the exuviae and eggs, and then washed the threads with methanol to separate chemical components from physical attributes of the threads. Female *N. womersleyi* did not follow *T. urticae* trails that had been washed with methanol but contained physical residues, but they did follow the direction to which the methanol extracts of the *T. urticae* trails was applied. These results suggest that the predatory mite follows chemical, not physical, attributes of *T. urticae* trails.

Key words *Tetranychus urticae* ▪ *Neoseiulus womersleyi* ▪ Silk thread ▪ Methanol extracts ▪ Trail following

36 **Introduction**

37 Predatory mites are promising biological control agents against tetranychid mites (e.g. McMurtry
38 1992; Croft and Slone 1997). For effective use of predatory mites as biological control agents, the
39 prey-searching cues of the mites must be elucidated. Although volatiles produced by
40 spider-mite-infested plants are thought to attract predatory mites (Sabelis and Van de Baan 1983;
41 Dicke et al. 1990), recent observations in open environments suggest that spider mite patches on
42 plant leaves do not attract predatory mites at a distance on the plant (Zemek et al. 2008; Yano and
43 Osakabe 2009). On the other hand, Yano and Osakabe (2009) showed that the predatory mite
44 *Phytoseiulus persimilis* Athias-Henriot (Acari: Phytoseiidae) follows trails left by adult female
45 *Tetranychus urticae*, suggesting that the spider mite trail could be a reliable prey-searching cue for
46 spider patches at a distance.
47 *Tetranychus urticae* feeds on various host plant species and is considered an important

48 agricultural pest globally (Jeppson et al. 1975). This mite often exhausts its host plant's resources,
49 and mated adult females then disperse to new hosts, primarily by walking (Margolies and Kennedy
50 1985). Ambulatory dispersing adult female *T. urticae* follow the trails left by preceding females,
51 which results in aggregation at a new colony site (Yano 2008). *Tetranychus urticae* spins a trail of
52 silk threads when walking (Saito 1977). Silk threads spun by *T. urticae* have been reported to retain
53 predatory mites and elicit typical prey-searching behavior (Hislop et al. 1978; Hoy and Smilanick
54 1981; Hislop and Prokopy 1981). Hoy and Smilanick (1981) reported that the predatory mite
55 *Metaseiulus* (= *Typhlodromus*) *occidentalis* Nesbitt (Acari: Phytoseiidae) detects silk and other
56 residuals of *T. urticae* on spider-mite-infested leaves, which can be removed by washing the leaves
57 with water. Although they suggested that water-soluble extracts of deposits are likely to serve as a
58 search cue for the predatory mite, important cues may be physically washed off. Hislop and Prokopy
59 (1981) showed that although the predatory mite *Neoseiulus fallaxis* (Garman) (Acari: Phytoseiidae)

60 preferred methanol extracts of *T. urticae* silk threads placed on filter paper, it showed a stronger
61 preference for intact silk. They concluded that *N. fallalis* may detect the chemical components of *T.*
62 *urticae* silk, although the physical stimulus of silk threads may also be involved.

63 However, these studies reported on the retention of predatory mites within a spider mite colony,
64 not on how they were guided toward the colony. Moreover, extracts of *T. urticae* silk contain
65 residuals, such as feces, exuviae, and even eggs, the latter of which are food of the above predatory
66 mites. Hence, these extracts may differ substantially from those of spider mite trails outside colonies.
67 To determine if predatory mites use the chemical components of spider mite trails to orient
68 themselves toward spider-mite-infested leaves, residual deposits that are apparently not associated
69 with trails outside the colony must be excluded.

70 In this study, we examined the prey-searching behavior of the endemic predatory mite
71 *Neoseiulus womersleyi*, which is an important predator of tetranychid mites in Japan (Hamamura

72 1986; Ehara and Shinkaji 1996). We collected *T. urticae* silk threads separately from their residuals.

73 Using extracts of the collected threads, we examined the hypothesis that *N. womersleyi* can detect

74 chemical components of *T. urticae* trails.

75

76 **Materials and methods**

77 **Mites**

78 The *T. urticae* study population was collected from a rose garden in Kyoto, Japan, and maintained on

79 individual discs of kidney bean *Phaseolus vulgaris* L. (Leguminosae) leaves pressed onto

80 water-saturated cotton in Petri dishes (90-mm diameter, 14 mm deep). The *N. womersleyi* study

81 population was also collected in Kyoto and was maintained on kidney bean leaf discs that were

82 heavily infested with *T. urticae* as prey. The leaf discs were maintained at $25 \pm 2^{\circ}\text{C}$ with $50 \pm 5\%$

83 relative humidity and a photoperiod of 16 : 8, light: dark (hereafter described as “laboratory

84 conditions”).

85

86 Preference of *N. womersleyi* for *T. urticae* trails

87 We first examined whether female *N. womersleyi* could follow trails laid down by *T. urticae*. The

88 term “trail”, as used in this study, refers to the silk threads and/or other chemical compound(s)

89 deposited by mated adult female *T. urticae* (hereafter described as “female *T. urticae*”). To conduct

90 dual-choice experiments under laboratory conditions, we connected one Parafilm square (Parafilm

91 M, American National Can Group, Chicago, IL, USA) and two bean leaf squares (10 × 10mm) with a

92 T-shaped Parafilm pathway on water-saturated cotton in Petri dishes (90-mm diameter, 14mm deep).

93 To induce a spider mite trail, we blocked a randomly selected branch with a piece of wet filter paper,

94 and then introduced either one or five 2- to 4-day-old female *T. urticae* onto the Parafilm square

95 (Fig.1a). One to five adult female *T. urticae* correspond to a typical colony size of the mite in the

96 wild (Yano, unpublished results). After 1h, when all of the *T. urticae* females had moved to the
97 available bean leaf square, we removed the wet filter paper and two bean squares together with the *T.*
98 *urticae* females from each disc, leaving only a trail on the Parafilm (Fig.1b).

99 We used 1-day-old mated adult female *N. womersleyi* (hereafter described as “female *N.*
100 *womersleyi*”) that had been previously starved since late deutonymph period. To prepare these
101 females, we isolated an old deutonymph female and an adult male *N. womersleyi* in a 1.5-ml
102 microtube (Treff AG, Degersheim, Switzerland) with a water droplet. We had previously confirmed
103 that old deutonymph females mature without feeding and that these females most intensively follow
104 spider mite trails (Yano, unpublished results). To avoid cannibalism and unsuccessful mating, we
105 used mature females only when both the female and male were alive after 48h. We introduced
106 individual female mites to the bottom of the T-shaped pathway using a fine brush (Fig.1b) and
107 recorded which branch they followed to the far end. We used each female *N. womersleyi* and

108 T-shaped Parafilm path only once. The female *N. womersleyi* that did not reach either end within
109 5min were not included in the analysis. The number of replicates was 25 for trails made by five
110 females and 30 for trails made by one female. Experimental outcomes were compared using the
111 binomial tests (Sokal and Rohlf, 1995), with the common null hypothesis that a female *N.*
112 *womersleyi* would chose either of the two branches with equal probability (i.e. 0.5).
113
114 Preference of *N. womersleyi* for *T. urticae* trails washed with methanol
115 To conduct dual-choice experiments, trails made by five *T. urticae* females were induced on a
116 Parafilm pathway in the manner described above. We then transferred the Parafilm to a Petri dish
117 filled with methanol (Wako Pure Chemical Industries Ltd, Osaka, Japan, min. 99.5%) and gently
118 shook the dish for 5min. The Parafilm was then completely dried at room temperature and placed on
119 water-saturated cotton in another Petri dish (Fig.1b). To confirm whether threads were still present

120 on Parafilm pathways after they were washed with methanol, we observed threads on the surfaces of

121 randomly sampled Parafilm pathways using a scanning electron microscope (3D Real Surface View

122 Microscope VE-8800, Keyence, Osaka, Japan).

123 We then introduced a starved 1-day-old female *N. womersleyi* to the base point of the T-shaped

124 Parafilm in the manner described above (Fig.1b) and recorded the branch that the female first

125 followed to the far end. Each female *N. womersleyi* female and T-shaped Parafilm pathway was used

126 only once. Female *N. womersleyi* that did not reach an end point within 5min were not included in

127 the analysis. The number of replicates was 59. The numbers of females were compared using a

128 binomial test in the same manner described above.

129

130 Preference of *N. womersleyi* for *T. urticae* trail extracts

131 To collect *T. urticae* silk threads separately from residuals such as exuviae and eggs, we confined 10

132 *T. urticae* females within 6h of maturation in each of ten 1.5-ml microtubes (i.e. 100 females in total)

133 with a water droplet. We had previously confirmed that *T. urticae* females within 6h of maturation do

134 not oviposit when deprived of food.

135 To allow the females to deposit silk threads inside the microtubes, the tubes were kept under

136 laboratory conditions for 48h. Each tube was then opened onto a bean leaf disc to release the females.

137 After 30min, when all the females had exited, 80μl methanol was added to each tube and the tubes

138 were shaken gently for 20min using a constant temperature shaker (Synthetech Oven SO-1G, Nippon

139 Genetics Co., Tokyo, Japan) at 30°C. Extracts from the 10 tubes (c. 800μl total) were consolidated

140 into another tube and centrifuged at 12,000 r.p.m. for 5min. To exclude residual deposits, the top

141 clear layer in the tube was collected into another tube. To evaporate the methanol, the tube was

142 opened and placed in the shaker at 30°C for 60min. The final volume of the concentrated extracts was

143 c.400μl. For the control solvent, the same amount of methanol was poured into new microtubes,

144 which were treated in the manner described above.

145 To conduct the dual-choice experiments, we constructed T-shaped pathways of filter paper
146 (35 × 35mm, 2mm wide, Fig.2). Methanol extract of about 40 spider mite's trails was applied to the
147 trunk and a randomly selected branch of each constructed pathway, with the control methanol applied
148 to the other branch. The two solutions were applied uniformly to both sides of the filter paper. Since
149 a starved *N. womersleyi* female is 0.25mm wide (1/8 of 2mm wide pathway) at most, *N. womersleyi*
150 advancing along the pathway would not contact more than 1/8 of the applied extracts. Since 1/8 of 40
151 *T. urticae* females is 5 females, this corresponds to the number of spider mites used in the first
152 experiment. Using 10μl micropipettes (Calibrated Pipets, Drummond Scientific Co., PA, USA), we
153 applied each solution little by little at the junction point to minimize mixing. The pathways were
154 completely dried at room temperature and then placed on three 15 x 15mm cardboard pieces fixed to
155 a Petri dish (Fig.2). We then introduced a starved 1-day-old female *N. womersleyi* to the base of the

156 T-shaped filter paper and recorded the branch that the females first followed to the far end. Females
157 that walked on the backside of the filter paper were observed via a 100 × 100-mm mirror placed
158 under the Petri dish (Fig.2). Each female *N. womersleyi* and T-shaped filter paper was used only once.
159 Female *N. womersleyi* that did not reach an end point within 5min were not included in the analysis.
160 The number of replicates was 54. The numbers of females were compared using a binomial test in the
161 same manner described above.

162

163 **Results**

164 Preference of *N. womersleyi* for *T. urticae* trails

165 Female *N. womersleyi* followed trails created by five female *T. urticae* (trail : control, 19 : 6;

166 $P=0.0073$, binomial test, Fig. 3a), but did not follow the trail made by one female *T. urticae* (trail :

167 control, 19 : 11; $P = 0.100$, binomial test, Fig. 3b). The latter statistically insignificant result,

168 however, may in part be due to the low number of replicates.

169

170 Preference of *N. womersleyi* for *T. urticae* trails washed with methanol

171 Although we found *T. urticae* threads on the Parafilm surfaces after they were washed with methanol,

172 the female *N. womersleyi* showed no preference for washed *T. urticae* trails (trail : control, 29 : 30; P

173 = 0.60, binomial test, Fig. 3c).

174

175 Preference of *N. womersleyi* for *T. urticae* trail extracts

176 *Neoseiulus womersleyi* females preferred trail branches with the methanol extract of *T. urticae* trails

177 to those with only methanol (trail extracts : methanol, 35 : 19; $P = 0.020$, binomial test, Fig. 3d),

178 suggesting that chemical components of *T. urticae* trails are used as a prey-searching cue. In addition,

179 we observed that some female *N. womersleyi* slowed down on the filter paper when they encountered

180 the trail extract. We also observed that some turned back from a trail-extract branch and retraced
181 their steps to the other, methanol-only branch. These behaviors were not observed during the
182 previous experiment that presented female *N. womersleyi* with *T. urticae* trails washed with
183 methanol.

184

185 Discussion

186 Not surprisingly, starved female *N. womersleyi* followed *T. urticae* trails in a density-dependent
187 fashion, as the amount of spun threads and their chemical components should correlate positively
188 with the number of *T. urticae* individuals used. Conspecific adult females (Yano 2008) and the
189 predatory mite *Phytoseiulus persimilis* (Yano and Osakabe 2009) also follow *T. urticae* trails in a
190 density-dependent fashion.

191 Because *T. urticae* spins silk threads while walking (Saito 1977), large quantities of threads can

192 be collected from long-settled mite colonies. However, residual deposits, such as exuviae, feces,
193 carcasses, and all stages of living mites, cannot be excluded from these samples. Moreover, as the
194 attractiveness of *T. urticae* threads appears to decline with time (Yano 2008), the properties of fresh
195 threads may differ from those collected at spider-mite colonies, which contain old threads. Therefore,
196 we examined *T. urticae* threads produced within 48 h, although some feces may have been included
197 in the extract.

198 The predatory mite did not follow *T. urticae* trails that had been washed with methanol, whereas
199 they did follow extracts of *T. urticae* trails. These results suggest that chemical, not physical
200 attributes of *T. urticae* trails are responsible for trail following by *N. womersleyi*, although the
201 washing treatment may have altered the physical properties of *T. urticae* threads. It is possible that
202 substances not soluble in methanol may be also responsible for the predatory mite's trail following.
203 However, *N. womersleyi* females did not follow washed *T. urticae* trails where substances not soluble

204 in methanol should remain, suggesting that such substances are unimportant. Some *N. womersleyi*
205 females slowed down and exhibited typical prey-searching behavior (Hoy and Smilanick 1981) on
206 filter paper painted with extracts of *T. urticae* trails, but no predator engaged in this behavior on
207 trails washed with methanol. These observations further suggest that the predatory mite responds
208 positively to chemical components in the *T. urticae* trails as a prey-searching cue. Therefore,
209 predatory mites most likely use the chemical cues of *T. urticae* trails to find prey colonies, and then
210 use both physical and chemical trail cues after arriving at the colonies, as suggested by Hislop and
211 Prokopy (1981).

212 To efficiently attract and retain predatory mites of tetranychid mite colonies, the chemical
213 components responsible for trail following by predatory mites must be clarified. Methods to collect
214 efficiently the unadulterated spider-mite threads should also be improved.

215

216 **Acknowledgements** We thank K. Oku of the National Agricultural Research Center, R. Uesugi

of National Institute of Vegetable and Tea Science, and H. Iida of the National Agriculture and Food Research Organization for helpful suggestions. We also thank D. E. Bowler and other members in our laboratory for valuable suggestions. This study was partly supported by the Japan Society for the Promotion of Science [Basic Research C; grant number 21580066 to S. Y.], by funding for the development of new biorational techniques in sustainable agriculture, and by a Grant-in-Aid for the 21st Century COE program for Innovative Food and Environmental Studies Pioneered by Entomomimetic Sciences, from the Ministry of Education, Culture, Sports, Science, and Technology of Japan.

References

- Croft BA, Slone DH (1997) Equilibrium densities of European red mite (Acari: Tetranychidae) after exposure to three levels of predacious mite diversity on apple. *Environ Entomol* 26:391-397
- Dicke M, Sabelis MW, Takabayashi J, Bruin J, Posthumus MA. (1990) Plant strategies of manipulating predator-prey interactions through allelochemicals: prospects for application in pest control. *J Chem Ecol* 16:3091-3118
- Ehara S, Shinkaji N (1996) Principles of Plant Acarology. Zenkoku Noson Kyoiku Kyokai, Tokyo (in Japanese).
- Hamamura T (1986) Studies on the biological control of Kanzawa spider mite, *Tetranychus kanzawai* Kishida by the chemical resistant predacious mite, *Amblyseius longispinosus* (Evans) in tea fields (Acarina: Tetranychidae, Phytoseiidae). *Bull Fruit Tree Res Stn E* 21:121-201 (in Japanese with English summary)
- Hardman JM, Smith RF, Bent E (1995) Effects of different integrated pest management programs on biological control of mites on apple by predatory mites (Acari) in Nova Scotia. *Environ Entomol* 24:125-142

- 241 Hislop RG, Alves N, Prokopy RJ (1978) Spider mite substances influencing searching behaviour of
- 242 the predator *Amblyseius fallacies* on apples. Fruit Notes 43:8-11
- 243 Hislop RG, Prokopy RJ (1981) Mite predator responses to prey and predator-emitted stimuli. J Chem
- 244 Ecol 7:895-904
- 245 Hoy MA, Smilanick JM (1981) Non-random prey location by the phytoseiid predator *Metaseiulus*
- 246 *occidentalis*: differential responses to several spider mite species. Entomol Exp Appl
- 247 29:241-253
- 248 Jeppson LR, Keifer HH, Baker EW (1975) Mites Injurious to Economic Plants. University of
- 249 California Press, Berkeley, CA, USA.
- 250 Margolies DC, Kennedy GG (1985) Movement of the two-spotted spider mite *Tetranychus urticae*
- 251 Koch (Acari: Tetranychidae), among hosts in a corn-peanut agroecosystem. Entomol Exp Appl
- 252 37:55-61
- 253 McMurtry JA (1992) Dynamics and potential impact of generalist phytoseiids in agroecosystems and
- 254 possibilities for establishment of exotic species. Exp Appl Acarol 14:371-382
- 255 Sabelis MW, Van de Baan HE (1983) Location of distant spider mite colonies by phytoseiid
- 256 predators: demonstration of specific kairomones emitted by *Tetranychus urticae* and
- 257 *Panonychus ulmi*. Entomol Exp Appl 33:303-314
- 258 Saito Y (1977) Study on spinning behavior of spider mites (Acarina: Tetranychidae). I. Method for
- 259 quantitative evaluation of the mite webbing, and the relationship between webbing and walking.
- 260 Jpn J Appl Entomol Zool 21: 27-34 (in Japanese, with English summary)
- 261 Sokal RR, Rohlf FJ (1995) Biometry. The principles and practice of statistics in biological research,
- 262 3rd edn. Freeman, San Francisco.
- 263 Yano S (2008) Collective and solitary behaviors of the two-spotted spider mite (Acari:
- 264 Tetranychidae) are induced by trail following. Ann Entomol Soc Am 101:247-252

- 265 Yano S, Osakabe Mh (2009) Do spider mite-infested plants and spider mite trails attract predatory
266 mites? Ecol Res 24:1173-1178
- 267 Zemek R, Nachman G, Ruzickova S (2008) How does *Phytoseiulus persimilis* find its prey when
268 foraging within a bean plant? In Bertrand M, Kreiter S, McCoy KD, Migeon A, Navajas M,
269 Tixier M-S, Vial L (eds) Integrative Acarology. Proceedings of the 6th European Congress.
270 European Association of Acarologists 2008, pp390-393
- 271

272 **Figure Legends**

273 **Fig.1** Experimental designs for testing the preference of *N. womersleyi* females for *T. urticae* trails

274 and trails washed with methanol. a) To guide *T. urticae* females in one direction, a piece of wet filter

275 paper blocked the other path branch; b) A 1-day-old female *N. womersleyi* was placed on the base

276 point of the T-shaped Parafilm. We recorded the branch that the female first followed to the far end.

277

278 **Fig.2** Experimental design for testing the preference of *N. womersleyi* for a methanol extract of *T.*

279 *urticae* trails. The methanol trail extract was applied to the trunk of a filter-paper path and a

280 randomly selected branch; pure methanol was applied as a control to the other branch.

281

282 **Fig.3** Summary of the dual-choice experiments. Female *N. womersleyi* followed *T. urticae* trails laid

283 down by a) five females but did not follow a trail of b) one *T. urticae* female. Moreover, female *N.*

284 *womersleyi* c) did not follow *T. urticae* trails that had been washed with methanol but d) did follow

285 the methanol extracts of the *T. urticae* trails.

Fig. 1

Fig. 2

Fig. 3