TITLE:

Berberine-induced activation of 5'-adenosine monophosphate-activated protein kinase and glucose transport in rat skeletal muscles.

AUTHOR(S):

Ma, Xiao; Egawa, Tatsuro; Kimura, Hajime; Karaike, Kouhei; Masuda, Shinya; Iwanaka, Nobumasa; Hayashi, Tatsuya

CITATION:

Ma, Xiao ...[et al]. Berberine-induced activation of 5'-adenosine monophosphate-activated protein kinase and glucose transport in rat skeletal muscles.. Metabolism: clinical and experimental 2010, 59(11): 1619-1627

ISSUE DATE:

2010-11

URL:

<http://hdl.handle.net/2433/131860>

RIGHT:

© 2010 Elsevier Inc.; This is not the published version. Please cite only the published version.; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。

Metabolism Clinical and Experimental 59 (2010) 1619–1627

Berberine-induced activation of 5'AMP-activated protein kinase and glucose transport in rat skeletal muscles

Xiao Ma, Tatsuro Egawa, Hajime Kimura, Kouhei Karaike, Shinya Masuda, Nobumasa Iwanaka, Tatsuya Hayashi*

Received 23 July 2009; accepted 17 March 2010

Laboratory of Sports and Exercise Medicine, Graduate School of Human and Environmental Studies, Kyoto University, Kyoto, 606-8501, Japan

* Corresponding author. Tel.: +81 75 753 6640; fax: +81 75 753 6640.

E-mail address: tatsuya@kuhp.kyoto-u.ac.jp (T. Hayashi).

doi:10.1016/j.metabol.2010.03.009

Abstract

Berberine (BBR) is the main alkaloid of *Coptis chinensis*, which has been used as a folk medicine to treat diabetes mellitus in Asian countries. We explored the possibility that 5'-AMP-activated protein kinase (AMPK) is involved in metabolic enhancement by BBR in skeletal muscle, the important tissue for glucose metabolism. Isolated rat epitrochlearis and soleus muscles were incubated in a buffer containing BBR, and activation of AMPK and related events were examined. In response to BBR treatment, the Thr¹⁷² phosphorylation of the catalytic α subunit of AMPK, an essential step for full kinase activation, increased in a dose- and time-dependent manner. Ser⁷⁹ phosphorylation of acetyl CoA carboxylase, an intracellular substrate of AMPK, increased correspondingly. Analysis of isoform-specific AMPK activity revealed that BBR activated both the $\alpha 1$ and $\alpha 2$ isoforms of the catalytic subunit. This increase in enzyme activity was associated with an increased rate of 3-*O*-methyl-D-glucose transport in the absence of insulin and with phosphorylation of AS160, a signaling intermediary leading to glucose transporter 4 (GLUT4) translocation. The intracellular energy status estimated from the phosphocreatine concentration was decreased by BBR. These results suggest that BBR acutely stimulates both AMPK $\alpha 1$ and AMPK $\alpha 2$ and insulin-independent glucose transport in skeletal muscle with a reduction of the intracellular energy status.

1. Introduction

Skeletal muscle plays an important role in glucose metabolism and homeostasis in humans. Skeletal muscle is the principal site of glucose transport and the rate-limiting step in glucose utilization under physiological conditions [1, 2]. Similar to insulin stimulation, exercise increases the rate of glucose transport into contracting skeletal muscle by the translocation of glucose transporter 4 (GLUT4) to the surface membranes. This phenomenon is considered to be responsible for the acute hypoglycemic effect of exercise, with glucose in the blood being taken up by contracting skeletal muscles. Indeed, the exercise-stimulated GLUT4 translocation is not impaired in insulin-resistant conditions such as type 2 diabetes mellitus and obesity [3]. Thus, the insulin-independent mechanisms of exercise have been widely used to decrease blood glucose in patients with type 2 diabetes mellitus.

It has been demonstrated that exercise and insulin use different signaling mechanisms in skeletal muscle, and 5'-AMP-activated protein kinase (AMPK) has been identified as part of the mechanisms leading to the acute enhancement of glucose transport in contracting muscles (reviewed by Kahn et al [4], Hardie et al [5], and Fuji et al [6]). AMPK is a member of a metabolite-sensing protein kinase family and acts as an energy-sensing and signaling molecule in muscle cells by monitoring cellular energy levels, such as the AMP to adenosine triphosphate (ATP) ratio. Skeletal muscle AMPK is also implicated in a variety of antidiabetic properties of exercise, including GLUT4 expression [7, 8], glycogen regulation [9, 10], fatty acid oxidation [11, 12], mitochondrial biogenesis [13, 14], and enhanced insulin sensitivity [8, 15, 16]. In addition, skeletal muscle AMPK mediates part of the action of adipokines, including leptin and adiponectin,

in glucose and lipid homeostasis, and the antidiabetic effect of metformin (MET) (reviewed by Kahn et al [4] and Hardie et al [5]). Thus, through these effects in skeletal muscle AMPK serves as a metabolic activator that may reduce the risk for type 2 diabetes mellitus.

Recent reports show that berberine (BBR), a natural plant alkaloid, stimulates glucose transport and activates AMPK in L6 cultured myocytes [17, 18]. BBR decreases intracellular energy status, represented by the AMP/ATP ratio [17]; and it has been suggested that BBR and its more biologically available derivative dihydroberberine inhibit cellular respiration by inhibiting complex 1 of the mitochondrial respiratory chain in L6 myotubes [19]. BBR is isolated from medicinal plants, such as *Coptis chinensis* (family Ranunculaceae), a Chinese herb that has been used for thousands of years in Asia to treat diabetes; and BBR treatment has beneficial effects on hyperglycemia in patients with type 2 diabetes mellitus [20, 21]. Considering that BBR and exercise can exert similar effects in stimulating AMPK accompanied with energy deprivation, skeletal muscle might be the principal site of action for the antidiabetic effect of BBR. However, to our knowledge no study has examined the direct effects of BBR on AMPK activity in skeletal muscle tissue. AMPK is a heterotrimeric kinase, consisting of a catalytic α -subunit and two regulatory subunits, β and γ . Two distinct α -isoforms (AMPK α 1 and α 2) exist in skeletal muscle. It is notable that AMPK α 1 is the predominant α -isoform and AMPK α 2 activity is undetectable in L6 cells [22].

The purpose of this study was to elucidate whether BBR acts on skeletal muscle to stimulate AMPK α 1 and α 2 and leads to metabolic enhancement, including glucose transport. We examined the short-term effects of BBR on both fast-glycolytic epitrochlearis and slow-oxidative soleus muscles using an isolated rat skeletal muscle

preparation.

2. Materials and Methods

2.1. *Experimental animals and materials*

Male Wistar rats aged 5 weeks and weighing 100 to 120 g were obtained from Shimizu Breeding Laboratories (Kyoto, Japan). The animals were housed in an animal room maintained at 23 °C with a 12:12 hour light/dark cycle and fed a standard laboratory diet and water *ad libitum*. All protocols for animal use and euthanasia were reviewed and approved by the Kyoto University Graduate School of Human and Environmental Studies and Kyoto University Radioisotope Research Center in Japan. The rats were randomly assigned to the experimental groups. BBR chloride was purchased from Sigma (Saint Louis, MO).

2.2. *Muscle incubation*

Two muscles, epitrochlearis and soleus, were used owing to their specific fiber type composition. The epitrochlearis is composed predominantly of fast-twitch glycolytic fibers (60–65% fast-twitch white, 20% fast-twitch red, 15% slow-twitch red) [23], and the soleus is composed primarily of slow-twitch oxidative fibers (0% fast-twitch white, 13% fast-twitch red, 87% slow-twitch red) [24]. Muscles were treated as described previously [25, 26] with modifications. The rats were sacrificed by cervical dislocation without anesthesia, and the muscles were rapidly and gently removed. Both ends of each muscle were tied with sutures, and the muscle was mounted on an incubation apparatus with the resting tension set to 0.5 g. To recover from AMPK activation induced during the isolating procedure, the muscles were preincubated in 7 mL of Krebs–Ringer bicarbonate buffer (KRB): 117 mmol/L NaCl, 4.7 mmol/L KCl, 2.5 mmol/L CaCl₂, 1.2 mmol/L

KH₂PO₄, 1.2 mmol/L MgSO₄, and 24.6 mmol/L NaHCO₃ containing 2 mmol/L of pyruvate (KRBP) for 30 minutes.

The muscle was then incubated in 7 mL of fresh buffer containing various concentrations (0–0.6 mmol/L) of BBR for 30 minutes. The muscles were also incubated with 2 mmol/L of 5-aminoimidazole-4-carboxamide-1- α -D-ribonucleoside (AICAR) for 30 minutes, 0.5 mmol/L 2,4-dinitrophenol (DNP) for 15 minutes or 1 μ mol/L insulin for 30 minutes to elicit maximal activation of AMPK (AICAR, DNP) or glucose transport (insulin) in isolated rat muscle [25, 27]. The muscles were also incubated with 2 mmol/L MET for 30 minutes [28]. To measure the time course of changes in AMPK phosphorylation, the muscles were preincubated for 30 minutes, and then incubated in the presence of BBR (0.3 mmol/L) for 0, 15, 30, 45 or 60 minutes. The buffer was bubbled with gas continuously using a mixture of 95% O₂–5% CO₂ and maintained at 37 °C. The muscle was then either used fresh to measure glucose uptake (see 3-*O*-methyl-D-glucose (3MG) transport) or frozen immediately in liquid nitrogen for the subsequent measurement of isoform-specific AMPK activity, adenosine triphosphate (ATP) and phosphocreatine (PCr) concentrations, and western blot analysis.

2.3. Western blot analysis

Sample preparation and western blot analysis for detection of phosphorylated AMPK α , total AMPK α and phosphorylated acetyl CoA carboxylase (ACC) were performed as described previously [26]. The muscle was homogenized in ice-cold lysis buffer (1:40 wt/vol) containing 20 mmol/L Tris–HCl (pH 7.4), 1% Triton X, 50 mmol/L NaCl, 250 mmol/L sucrose, 50 mmol/L NaF, 5 mmol/L sodium pyrophosphate, 2 mmol/L dithiothreitol, 4mg/L leupeptin, 50 mg/L trypsin inhibitor, 0.1 mmol/L benzamidine, and

0.5 mmol/L phenylmethylsulfonyl fluoride (Buffer A), and centrifuged at $16,060 \times g$ for 30 minutes at 4 °C. The supernatant (10 µg of protein) was mixed with Laemmli's sample buffer containing mercaptoethanol and boiled, and the denatured proteins were separated on either a 10% polyacrylamide gel for AMPK and Akt, or a 6% gel for ACC.

The muscle for western blot analysis of phosphorylated p38 mitogen-activated protein kinase (MAPK) and phosphorylated p44/42 MAPK was homogenized in ice-cold lysis buffer (1:40 wt/vol) containing 20 mmol/L HEPES (pH 7.4), 1.5% Triton X-100, 50 mmol/L NaF, 5 mmol/L $\text{Na}_4\text{P}_2\text{O}_7$, 50 mmol/L β -glycerophosphate (disodium-5-hydrate), 250 mmol/L sucrose, 5 mmol/L EDTA-Na and EGTA-Na, 50 mg/L soybean trypsin inhibitor, 2 mmol/L dithiothreitol, 4mg/L leupeptin, 25 mg/L aprotinin, 0.1 mmol/L benzamidine, 1 mmol/L Na_3VO_4 , and 0.5 mmol/L phenylmethylsulfonyl fluoride, and centrifuged at $16,060 \times g$ for 30 minutes at 4 °C. The supernatant (10 µg of protein) was mixed with the loading buffer containing mercaptoethanol and boiled, and the denatured proteins were separated on a 10% polyacrylamide gel.

The proteins were then transferred to polyvinylidene difluoride membranes at 100 V for 1 h. The membrane was blocked overnight at 4 °C with Block Ace (Yukijirushi Nyugyo, Tokyo, Japan) and then incubated with AMPK α (#2532; Cell Signaling Technology, Beverly, MA), phosphospecific AMPK α Thr¹⁷² (#2531; Cell Signaling Technology), phosphospecific ACC Ser⁷⁹ (#07-303; Upstate Biotechnology, Lake Placid, NY), phosphospecific Akt Ser⁴⁷³ (#9271; Cell Signaling Technology), phospho-p38 MAP Kinase Thr¹⁸⁰/Tyr¹⁸² (#9211; Cell Signaling Technology) or phospho-p44/42 MAP Kinase Thr²⁰²/Tyr²⁰⁴ (#9101; Cell Signaling Technology).

To immunoprecipitate the Akt substrate of the 160 kd (AS160), an aliquot of the supernatant of Buffer A was prepared as described above (300 µg of protein) and

incubated with anti-AS160 (Rab-GAP) (# 07-741; Upstate Biotechnology) and protein A-Sepharose CL4B beads (Amersham Biosciences, Uppsala, Sweden) overnight at 4 °C with end-over-end rotation. The immunoprecipitate was washed three times in Buffer A and then centrifuged at 4 000g for 30 seconds at 4 °C. The supernatant was removed, and the beads were mixed with the loading buffer and boiled. The denatured proteins were then separated on a 10% polyacrylamide gel. The proteins were then transferred to polyvinylidene difluoride membranes at 100 V for 1 hour. The membrane was blocked with Block Ace (Yukijirushi Nyugyo) overnight at 4 °C, and then incubated with anti-AS160 (Rab-GAP) (#30544; Upstate Biotechnology) or antiphospho-(Ser/Thr) Akt substrate (#9611; Cell Signaling Technology) as described previously [29].

The membrane was washed, reacted with anti-rabbit IgG coupled to peroxidase, and developed with enhanced chemiluminescence reagents according to the manufacturer's instructions (Amersham Biosciences, Buckinghamshire, United Kingdom). The signals on the blot were detected and quantified using a Lumino LAS-1000 System image analyzer (Fuji Photo Film, Tokyo, Japan). The mean intensity of basal samples in each membrane was used as a reference for controlling gel-to-gel variation. Equal protein loading and transfer was confirmed by Coomassie brilliant blue staining of the membranes.

2.4. Isoform-specific AMPK activity assay

The AMPK activity assay was performed as we described previously [26, 27]. Muscle was homogenized in Buffer A as described in the western blot analysis subsection and the resultant supernatant liquid (100 µg of protein) was immunoprecipitated with isoform-specific polyclonal antibodies directed against the $\alpha 1$ or $\alpha 2$ catalytic subunit of

AMPK [26] and protein A-Sepharose CL4B beads. The immunoprecipitate was washed twice in both Buffer A and wash buffer (240 mmol/L HEPES, 480 mmol/L NaCl). The kinase reaction was performed in 40 mmol/L HEPES (pH 7.0), 0.1 mmol/L SAMS peptide [26, 27], 0.2 mmol/L AMP, 80 mmol/L NaCl, 0.8 mmol/L dithiothreitol, 5 mmol/L MgCl₂, 0.2 mmol/L ATP (2 μ Ci of [γ -³²P] ATP (PerkinElmer, Wellesley, MA), in a final volume of 40 μ L for 20 minutes at 30 °C. At the end of the reaction, a 15 μ L aliquot was removed and spotted onto Whatman P81 paper (Whatman International, Maidstone, United Kingdom). The filter paper was washed at least six times in 1% phosphoric acid and once in acetone. ³²P incorporation was quantified using a scintillation counter, and kinase activity was expressed as picomoles of incorporated ATP per milligram of immunoprecipitated protein per minute.

2.5. ATP and PCr assay

ATP and PCr content were measured fluorometrically in perchloric acid extracts of epitrochlearis and soleus muscles according to the method of Lowry and Passonneau [30]. In brief, each frozen muscle was homogenized in 0.2 mol/L HClO₄ (3:25 wt/vol) in an ethanol-dry ice bath (−20 °C to −30 °C) and centrifuged at 16,000 $\times$ g for 2 minutes at −9 °C. The supernatant of the homogenate was neutralized with a solution of 2 mol/L KOH, 0.4 mol/L KCl and 0.4 mol/L imidazole, centrifuged at 16 000g for 2 minutes at −9 °C and then subjected to enzymatic analysis [30]. ATP and PCr content were expressed as nanomoles per milligram wet weight of muscle.

2.6. 3MG transport

The 3MG transport assay was performed as we described previously [25, 26]. The

muscle was transferred to 2 mL of KRB containing 1 mmol/L of 3-*O*-[methyl-³H]-D-glucose (1.5 μCi/mL) (American Radiolabeled Chemicals, St. Louis, MO) and 7 mmol/L D- [¹⁴C]-mannitol (0.3 μCi/mL) (American Radiolabeled Chemicals) at 30 °C and incubated for 10 minutes. The muscle was blotted onto filter paper, trimmed, frozen in liquid nitrogen, and stored at −80 °C. Each frozen muscle was weighed and incubated in 300 μL of 1 mol/L NaOH at 80 °C for 10 minutes. The digest was neutralized with 300 μL of 1 mol/L HCl, and the particulates were precipitated by centrifugation at 20 000g for 2 minutes. The radioactivity in aliquots of the digested protein was measured using liquid scintillation counting for dual labels, and the extracellular and intracellular spaces were calculated [31].

2.7. Statistical analysis

The data are expressed as means ± S.E. The mean values were compared using a 1-way analysis of variance (ANOVA) followed by a post hoc comparison with Fisher's protected least-significant difference method. Two mean values were compared using student's *t*-test. Differences between the groups were considered significant if $P < .05$.

3. Results

3.1. BBR increased phosphorylation of AMPK α Thr¹⁷² in skeletal muscles

The Thr¹⁷² residue in both the α 1 and α 2 catalytic subunits is the primary site responsible for AMPK activation. To determine whether BBR activates AMPK, we measured the degree of phosphorylation of AMPK α Thr¹⁷² using western blot analysis employing a phosphospecific antibody. Compared with the basal condition, both epitrochlearis (Fig. 1A) and soleus (Fig. 1B) muscles stimulated with BBR (0.3 mmol/L, 30 minutes) exhibited increased phosphorylation of Thr¹⁷² without any change in the total amount of AMPK α . ACC is a downstream target of AMPK in skeletal muscle, and the phosphorylation of the Ser⁷⁹ site of ACC reflects the total AMPK activity [32, 33]. The marked phosphorylation of ACC paralleled the increase in AMPK phosphorylation in epitrochlearis (Fig. 1A) and soleus (Fig. 1B) muscles. The AMPK activator AICAR also caused a strong phosphorylation of AMPK and ACC in epitrochlearis (Fig. 1A), but not in soleus (Fig. 1B). AICAR is taken up into skeletal muscle and metabolized by adenosine kinase to form ZMP, a monophosphorylated derivative that mimics the effects of AMP on AMPK [5]. The pharmacological inhibitor of oxidative phosphorylation, DNP, which induces a marked reduction of muscle ATP and PCr levels [27], also caused significant phosphorylation of AMPK and ACC in both muscle types (Figs. 1A and B). Metformin elicited moderate phosphorylation of AMPK and ACC in both muscles (Figs. 1A and B).

To determine the dose and time dependency of the AMPK phosphorylation, muscles were stimulated with BBR at various concentrations and for various times. The dose-response study showed a significant phosphorylation of 1.2-, 1.6- and 2.0-fold in epitrochlearis muscle (Fig. 2A), and 1.6-, 2.7- and 2.6-fold in soleus muscle (Fig. 2B) at

concentrations of 0.1, 0.3 and 0.6 mmol/L of BBR, respectively. The time-course study revealed that 0.3 mmol/L of BBR induced acute phosphorylation of AMPK in both epitrochlearis (Fig. 3A) and soleus (Fig. 3B) muscles after an incubation period as short as 15 minutes.

3.2. BBR activated both AMPK $\alpha 1$ and $\alpha 2$ in skeletal muscles

To identify which catalytic subunit is activated by BBR, isoform-specific AMPK activity was measured in anti- $\alpha 1$ and anti- $\alpha 2$ immunoprecipitates of the muscle after treatment with BBR. BBR stimulation (0.3 mmol/L, 30 minutes) significantly increased the activity of both AMPK $\alpha 1$ and AMPK $\alpha 2$ in epitrochlearis muscle, the activity of AMPK $\alpha 1$ = 14.0 ± 1.1 pmol/(mg min) in the basal condition and 50.7 ± 2.9 pmol/(mg min) after incubation with BBR; and the activity of AMPK $\alpha 2$ = 12.2 ± 1.3 pmol/(mg min) in the basal condition and 30.4 ± 2.5 pmol/(mg min) after incubation with BBR (Fig. 4A). In soleus muscle the activity of AMPK $\alpha 1$ was 54.3 ± 7.8 pmol/(mg min) in the basal condition and 104.0 ± 5.0 pmol/(mg min) after incubation with BBR, and the activity of AMPK $\alpha 2$ was 17.1 ± 1.5 pmol/(mg min) in the basal condition and 29.3 ± 1.9 pmol/(mg min) after incubation with BBR (Fig. 4B).

3.3. BBR stimulated glucose transport in skeletal muscles in the absence of insulin

We investigated whether the activation of AMPK in skeletal muscle by BBR is associated with enhanced glucose transport. In the absence of insulin, incubation with BBR (0.3 mmol/L, 30 minutes) increased the rate of 3MG transport 1.7-fold in epitrochlearis muscle (Fig. 5A) and 1.4-fold in soleus muscle (Fig. 5B) compared with the basal conditions. AICAR significantly increased 3MG transport activity in

epitrochlearis muscle (Fig. 5A), but not in soleus (Fig. 5B). Akt is a key molecule in the insulin-stimulated signaling pathway leading to enhanced glucose transport and metabolism. Recent studies suggest that AS160 is a downstream nexus for both Akt-mediated and AMPK-mediated glucose transport in skeletal muscle [34, 35], and that AS160 is an intracellular substrate of AMPK in skeletal muscle [35]. In our study, BBR significantly increased phosphorylation of AS160 (Fig. 6) without changing the phosphorylation status of Akt (Fig. 7). Additionally, BBR treatment had no effect on the phosphorylation status of p38 or p44/42 MAPK (Fig. 7).

3.4. BBR decreased energy status in skeletal muscles.

AMPK is activated in response to energy-consuming stress, such as muscle contraction, hypoxia, and inhibition of oxidative phosphorylation [27]. To clarify whether BBR increases AMPK activity by energy deprivation, we measured the muscle content of ATP and PCr. The decrease in ATP content was not statistically significant, but PCr content after the stimulation of BBR (0.3 mmol/L, 30 minutes) was significantly lower than that of the basal samples in both epitrochlearis and soleus muscles (Table 1).

4. Discussion

Our data show three novel findings relating to the metabolic effect of BBR on skeletal muscle. First, BBR had the ability to increase AMPK α Thr¹⁷² phosphorylation accompanied with both AMPK α 1 and α 2 activities (Figs. 1-4). The increased phosphorylation of endogenous substrate AS160 (Fig. 6) and enhanced 3MG transport (Fig. 5) as well as ACC phosphorylation (Fig. 1) is indicative of a substantial increase in AMPK activity *in vivo*. Second, these effects were observed in both fast-glycolytic epitrochlearis and slow-oxidative soleus muscles, suggesting that the stimulatory effect of BBR on AMPK is not specific to a particular muscle type. Third, BBR-stimulated AMPK activation was associated with a reduction in the fuel status of skeletal muscle (Table 1), as with contraction-stimulated AMPK activation.

We used an isolated muscle preparation incubated *in vitro* to eliminate systemic confounders, such as circulatory, humoral and neuronal factors. In particular, we considered the possibility that oral administration of BBR, which acts as an α -galactosidase inhibitor [36, 37], could evoke indirect metabolic changes in skeletal muscle by suppressing glucose absorption in intestines. In addition, because AMPK α 1 is activated immediately as a post mortem artifact during the dissection procedure [38], we exposed muscles to BBR after a preincubation period sufficient to stabilize AMPK α 1 activity. Thus, our method made it possible to evaluate the direct effect of BBR on muscle tissue and both AMPK α 1 and AMPK α 2 activities.

Incubation of epitrochlearis and soleus muscles with BBR (≥ 0.1 mmol/L, ≥ 15 minutes) increased AMPK phosphorylation in dose- and time-dependent manners, with peak levels of phosphorylation at ≥ 0.3 mmol/L and ≥ 30 minutes (Figs. 2 and 3). We have

recently reported that incubation of rat epitrochlearis and soleus muscles with caffeine (≥ 3 mmol/L, ≥ 15 minutes) increases AMPK phosphorylation in dose- and time-dependent manners [39]. We have also shown that activation of AMPK α 1 and α 2 by caffeine is associated with a reduction in phosphocreatine content and an increased rate of 3MG transport in the absence of insulin [39]. These results suggest that BBR has similar actions to caffeine, by acutely stimulating both AMPK α isoforms and insulin-independent glucose transport with a reduction of the intracellular energy status. It is noteworthy that consumption of beverages containing caffeine including coffee and green tea is associated with a reduced risk of type 2 diabetes mellitus [40, 41].

We examined the effects of BBR in fast-glycolytic and slow-oxidative muscles because activation of AMPK and its effects may vary by muscle type. In the studies using rat skeletal muscle, repeated AICAR administration elicited the greatest increases in GLUT4 and glycogen content in fast-twitch muscles [42-44]. Daily, subcutaneous injections of AICAR for 4 weeks increased GLUT4 and glycogen content in the red and white quadriceps, but not in the soleus muscle in rats [43]. Furthermore, AICAR exposure stimulated glucose transport in rat white muscles, but had no effect in soleus muscle [45, 46]. We also found significant increases in AMPK phosphorylation and 3MG transport activity by AICAR in epitrochlearis muscle, but not in soleus (Figs. 1 and 5). In contrast, incubation with BBR markedly increased AMPK phosphorylation and 3MG transport in both epitrochlearis and soleus muscles (Figs. 1 and 5). Considering that caffeine stimulates AMPK activity with a reduction of the intracellular energy status in both incubated epitrochlearis and soleus muscles [39], energy deprivation per se may be important in activating AMPK in rat soleus. Because of the non-muscle type-specific stimulation by BBR, we believe that most skeletal muscles respond to BBR.

The finding that BBR increased AMPK α Thr¹⁷² phosphorylation provides evidence that BBR induces covalent modification via upstream kinases. The LKB1 complex is constitutively active and the binding of AMP to AMPK facilitates the phosphorylation of AMPK by the LKB1 complex [47, 48]. Thus, LKB1 is believed to be a crucial AMPK kinase in the response to energy deprivation in skeletal muscle. In the present study, AMPK activation was accompanied by a decrease in PCr content (Table 1), raising the possibility that LKB1 is involved in AMPK activation by BBR.

Although we did not examine the long-term effect of BBR on muscle metabolism, Kim et al [49] have recently shown that intraperitoneal administration of BBR once a day for three weeks increases AMPK and ACC phosphorylation in gastrocnemius muscle of the *db/db* mouse (they did not examine the acute effect of BBR). In our earlier study, repeated activation of AMPK by intraperitoneal injection of AICAR, three times a day for seven days, increased GLUT4 protein and insulin-stimulated 2-deoxyglucose transport activity in mouse soleus muscle [8]. Similarly, in ZDF rats, which are characterized by a progressive β -cell dysfunction and a leptin receptor defect, AICAR treatment over a period of eight weeks increased whole-body insulin sensitivity, which was associated with the level of muscle GLUT4 expression [50]. These data lead us to speculate that repeated activation of AMPK by BBR may achieve long-term metabolic benefits in skeletal muscle and protect against the development of diabetes. In this context, further studies are needed to clarify whether oral administration of BBR at a physiologic dose (1.0~1.5 g daily) [20, 21] results in AMPK activation in human skeletal muscle.

Cheng et al [17] have shown that BBR stimulates glucose transport and promotes AMPK phosphorylation and p38 MAPK phosphorylation, and that the BBR-stimulated glucose transport is partially inhibited by an AMPK inhibitor Compound C or p38 MAPK

inhibitor SB202190 in L6 myotubes. In contrast, Zhou et al [51] have reported that in 3T3-L1 adipocytes BBR-stimulated glucose transport is not affected by SB203580 but is partially inhibited by PD98059, a p44/42 MAPK kinase (MEK) inhibitor that leads to inactivation of p44/42 MAPK. In the present study, BBR stimulation (0.3 mmol/L, 30 minutes) did not induce an increase in p38 MAPK phosphorylation or p44/42 MAPK phosphorylation although the BBR stimulation significantly increased the rate of 3MG transport in skeletal muscles. Our data suggest that p38 and p44/42 MAPK signaling does not play a major role in activating glucose transport in intact skeletal muscle.

In summary, we demonstrated for the first time that BBR increases both AMPK α 1 and α 2 activities in fast- and slow-twitch skeletal muscles, and that this activation is accompanied by insulin-independent glucose transport and a reduction of muscle energy status. We propose that BBR can activate muscle glucose metabolism, at least in part, by stimulating AMPK.

Acknowledgments

We thank Licht Miyamoto and Taro Toyoda for suggestions. We also thank Department of Medicine and Clinical Science Kyoto University Graduate School of Medicine and Radioisotope Research Center of Kyoto University for instrumental support. Tatsuya Hayashi was supported by research grants from the Japan Society for the Promotion of Science (20500576), Japan Vascular Disease Research Foundation, All Japan Coffee Association, and Nestle Nutrition Council Japan.

References

- [1] Kubo K and Foley JE, Rate-limiting steps for insulin-mediated glucose uptake into perfused rat hindlimb. *Am J Physiol*, 1986;250:E100-2.
- [2] Cline GW, Petersen KF, Krssak M, et al., Impaired glucose transport as a cause of decreased insulin-stimulated muscle glycogen synthesis in type 2 diabetes. *N Engl J Med*, 1999;341:240-6.
- [3] Kennedy JW, Hirshman MF, Gervino EV, et al., Acute exercise induces GLUT4 translocation in skeletal muscle of normal human subjects and subjects with type 2 diabetes. *Diabetes*, 1999;48:1192-7.
- [4] Kahn BB, Alquier T, Carling D, et al., AMP-activated protein kinase: ancient energy gauge provides clues to modern understanding of metabolism. *Cell Metab*, 2005;1:15-25.
- [5] Hardie DG, Hawley SA, and Scott JW, AMP-activated protein kinase--development of the energy sensor concept. *J Physiol*, 2006;574:7-15.
- [6] Fujii N, Jessen N, and Goodyear LJ, AMP-activated protein kinase and the regulation of glucose transport. *Am J Physiol Endocrinol Metab*, 2006;291:E867-77.
- [7] Zheng D, MacLean PS, Pohnert SC, et al., Regulation of muscle GLUT-4 transcription by AMP-activated protein kinase. *J Appl Physiol*, 2001;91:1073-83.
- [8] Nakano M, Hamada T, Hayashi T, et al., α 2 Isoform-specific activation of 5'-adenosine monophosphate-activated protein kinase by 5-aminoimidazole-4-carboxamide-1- β -D-ribose at a physiological level activates glucose transport and increases glucose transporter 4 in mouse skeletal muscle. *Metabolism*, 2006;55:300-8.

- [9] Jorgensen SB, Nielsen JN, Birk JB, et al., The alpha2-5'AMP-activated protein kinase is a site 2 glycogen synthase kinase in skeletal muscle and is responsive to glucose loading. *Diabetes*, 2004;53:3074-81.
- [10] Miyamoto L, Toyoda T, Hayashi T, et al., Effect of acute activation of 5'-AMP-activated protein kinase on glycogen regulation in isolated rat skeletal muscle. *J Appl Physiol*, 2007;102:1007-13.
- [11] Winder WW and Hardie DG, Inactivation of acetyl-CoA carboxylase and activation of AMP-activated protein kinase in muscle during exercise. *Am J Physiol*, 1996;270:E299-304.
- [12] Vavvas D, Apazidis A, Saha AK, et al., Contraction-induced changes in acetyl-CoA carboxylase and 5'-AMP-activated kinase in skeletal muscle. *J Biol Chem*, 1997;272:13255-61.
- [13] Jager S, Handschin C, St-Pierre J, et al., AMP-activated protein kinase (AMPK) action in skeletal muscle via direct phosphorylation of PGC-1alpha. *Proc Natl Acad Sci U S A*, 2007;104:12017-22.
- [14] Garcia-Roves PM, Osler ME, Holmstrom MH, et al., Gain-of-function R225Q mutation in AMP-activated protein kinase gamma3 subunit increases mitochondrial biogenesis in glycolytic skeletal muscle. *J Biol Chem*, 2008;283:35724-34.
- [15] Fiedler M, Zierath JR, Selen G, et al., 5-aminoimidazole-4-carboxy-amide-1-beta-D-ribofuranoside treatment ameliorates hyperglycaemia and hyperinsulinaemia but not dyslipidaemia in KKAY-CETP mice. *Diabetologia*, 2001;44:2180-6.
- [16] Tanaka S, Hayashi T, Toyoda T, et al., High-fat diet impairs the effects of a single bout of endurance exercise on glucose transport and insulin sensitivity in rat skeletal

muscle. *Metabolism*, 2007;56:1719-28.

[17]Cheng Z, Pang T, Gu M, et al., Berberine-stimulated glucose uptake in L6 myotubes involves both AMPK and p38 MAPK. *Biochim Biophys Acta*, 2006;1760:1682-9.

[18]Lee YS, Kim WS, Kim KH, et al., Berberine, a natural plant product, activates AMP-activated protein kinase with beneficial metabolic effects in diabetic and insulin-resistant states. *Diabetes*, 2006;55:2256-64.

[19]Turner N, Li JY, Gosby A, et al., Berberine and its more biologically available derivative, dihydroberberine, inhibit mitochondrial respiratory complex I: a mechanism for the action of berberine to activate AMP-activated protein kinase and improve insulin action. *Diabetes*, 2008;57:1414-8.

[20]Yin J, Xing H, and Ye J, Efficacy of berberine in patients with type 2 diabetes mellitus. *Metabolism*, 2008;57:712-7.

[21]Zhang Y, Li X, Zou D, et al., Treatment of type 2 diabetes and dyslipidemia with the natural plant alkaloid berberine. *J Clin Endocrinol Metab*, 2008;93:2559-65.

[22]Lessard SJ, Chen ZP, Watt MJ, et al., Chronic rosiglitazone treatment restores AMPK α 2 activity in insulin-resistant rat skeletal muscle. *Am J Physiol Endocrinol Metab*, 2006;290:E251-7.

[23]Nesher R, Karl IE, Kaiser KE, et al., Epitrochlearis muscle. I. Mechanical performance, energetics, and fiber composition. *Am J Physiol*, 1980;239:E454-60.

[24]Armstrong RB and Phelps RO, Muscle fiber type composition of the rat hindlimb. *Am J Anat*, 1984;171:259-72.

[25]Hayashi T, Hirshman MF, Kurth EJ, et al., Evidence for 5' AMP-activated protein kinase mediation of the effect of muscle contraction on glucose transport. *Diabetes*, 1998;47:1369-73.

- [26] Toyoda T, Hayashi T, Miyamoto L, et al., Possible involvement of the α 1 isoform of 5'AMP-activated protein kinase in oxidative stress-stimulated glucose transport in skeletal muscle. *Am J Physiol Endocrinol Metab*, 2004;287:E166-73.
- [27] Hayashi T, Hirshman MF, Fujii N, et al., Metabolic stress and altered glucose transport: activation of AMP-activated protein kinase as a unifying coupling mechanism. *Diabetes*, 2000;49:527-31.
- [28] Zhou G, Myers R, Li Y, et al., Role of AMP-activated protein kinase in mechanism of metformin action. *J Clin Invest*, 2001;108:1167-74.
- [29] Arias EB, Kim J, Funai K, et al., Prior exercise increases phosphorylation of Akt substrate of 160 kDa (AS160) in rat skeletal muscle. *Am J Physiol Endocrinol Metab*, 2007;292:E1191-200.
- [30] Lowry OH and Passonneau JV, eds. *A Flexible System of Ebzymatic Analysis*. 1972, Academic Press: New York. 151-6.
- [31] Young DA, Uhl JJ, Cartee GD, et al., Activation of glucose transport in muscle by prolonged exposure to insulin. Effects of glucose and insulin concentrations. *J Biol Chem*, 1986;261:16049-53.
- [32] Davies SP, Sim AT, and Hardie DG, Location and function of three sites phosphorylated on rat acetyl-CoA carboxylase by the AMP-activated protein kinase. *Eur J Biochem*, 1990;187:183-90.
- [33] Park SH, Gammon SR, Knippers JD, et al., Phosphorylation-activity relationships of AMPK and acetyl-CoA carboxylase in muscle. *J Appl Physiol*, 2002;92:2475-82.
- [34] Kramer HF, Witczak CA, Taylor EB, et al., AS160 regulates insulin- and contraction-stimulated glucose uptake in mouse skeletal muscle. *J Biol Chem*, 2006;281:31478-85.

- [35] Treebak JT, Glund S, Deshmukh A, et al., AMPK-mediated AS160 phosphorylation in skeletal muscle is dependent on AMPK catalytic and regulatory subunits. *Diabetes*, 2006;55:2051-8.
- [36] Pan GY, Huang ZJ, Wang GJ, et al., The antihyperglycaemic activity of berberine arises from a decrease of glucose absorption. *Planta Med*, 2003;69:632-6.
- [37] Liu L, Deng Y, Yu S, et al., Berberine attenuates intestinal disaccharidases in streptozotocin-induced diabetic rats. *Pharmazie*, 2008;63:384-8.
- [38] Toyoda T, Tanaka S, Ebihara K, et al., Low-intensity contraction activates the alpha1-isoform of 5'-AMP-activated protein kinase in rat skeletal muscle. *Am J Physiol Endocrinol Metab*, 2006;290:E583-90.
- [39] Egawa T, Hamada T, Kameda N, et al., Caffeine acutely activates 5'adenosine monophosphate-activated protein kinase and increases insulin-independent glucose transport in rat skeletal muscles. *Metabolism*, 2009;58:1609-17.
- [40] Salazar-Martinez E, Willett WC, Ascherio A, et al., Coffee consumption and risk for type 2 diabetes mellitus. *Ann Intern Med*, 2004;140:1-8.
- [41] Iso H, Date C, Wakai K, et al., The relationship between green tea and total caffeine intake and risk for self-reported type 2 diabetes among Japanese adults. *Ann Intern Med*, 2006;144:554-62.
- [42] Holmes BF, Kurth-Kraczek EJ, and Winder WW, Chronic activation of 5'-AMP-activated protein kinase increases GLUT-4, hexokinase, and glycogen in muscle. *J Appl Physiol*, 1999;87:1990-5.
- [43] Winder WW, Holmes BF, Rubink DS, et al., Activation of AMP-activated protein kinase increases mitochondrial enzymes in skeletal muscle. *J Appl Physiol*, 2000;88:2219-26.

- [44]Buhl ES, Jessen N, Schmitz O, et al., Chronic treatment with 5-aminoimidazole-4-carboxamide-1-beta-D-ribofuranoside increases insulin-stimulated glucose uptake and GLUT4 translocation in rat skeletal muscles in a fiber type-specific manner. *Diabetes*, 2001;50:12-7.
- [45]Balon TW and Jasman AP, Acute exposure to AICAR increases glucose transport in mouse EDL and soleus muscle. *Biochem Biophys Res Commun*, 2001;282:1008-11.
- [46]Ai H, Ihlemann J, Hellsten Y, et al., Effect of fiber type and nutritional state on AICAR- and contraction-stimulated glucose transport in rat muscle. *Am J Physiol Endocrinol Metab*, 2002;282:E1291-300.
- [47]Hawley SA, Boudeau J, Reid JL, et al., Complexes between the LKB1 tumor suppressor, STRAD alpha/beta and MO25 alpha/beta are upstream kinases in the AMP-activated protein kinase cascade. *J Biol*, 2003;2:28.
- [48]Sakamoto K, Goransson O, Hardie DG, et al., Activity of LKB1 and AMPK-related kinases in skeletal muscle: effects of contraction, phenformin, and AICAR. *Am J Physiol Endocrinol Metab*, 2004;287:E310-7.
- [49]Kim WS, Lee YS, Cha SH, et al., Berberine improves lipid dysregulation in obesity by controlling central and peripheral AMPK activity. *Am J Physiol Endocrinol Metab*, 2009;296:E812-9.
- [50]Pold R, Jensen LS, Jessen N, et al., Long-term AICAR administration and exercise prevents diabetes in ZDF rats. *Diabetes*, 2005;54:928-34.
- [51]Zhou L, Yang Y, Wang X, et al., Berberine stimulates glucose transport through a mechanism distinct from insulin. *Metabolism*, 2007;56:405-12.

Legends

Fig. 1. Berberine (BBR) increases the phosphorylation of AMPK α and acetyl CoA carboxylase (ACC) in rat skeletal muscle. Isolated epitrochlearis (A) and soleus (B) muscles were incubated in the absence (basal), or presence of BBR (0.3 mmol/L), AICAR (2 mmol/L) or metformin (MET) (2 mmol/L) for 30 minutes, or in the presence of 2,4-dinitrophenol (DNP) (0.5 mmol/L) for 15 minutes. The tissue lysate was subjected to Western blot analysis with antibodies to phosphorylated AMPK α , AMPK α , phosphorylated ACC or ACC. Representative immunoblots are shown.

Fig. 2. BBR increases phosphorylation of AMPK α in a dose-dependent manner in rat skeletal muscles. Isolated epitrochlearis (A) and soleus (B) muscles were incubated in the presence of 0 (basal), 0.1, 0.3 or 0.6 mmol/L of BBR for 30 minutes. The tissue lysate was subjected to Western blot analysis with antibodies to phosphorylated AMPK α or AMPK α . Fold increases are expressed relative to the level in muscles of the basal group. The values are means $\pm$ S.E. The number of muscles in each group is as follows: basal (8), BBR 0.1 mmol/L (6), BBR 0.3 mmol/L (8), BBR 0.6 mmol/L (6). * P < .05 and ** P < .01 vs. basal; # P < .05 and ## P < .01 vs. BBR 0.1 mmol/L.

Fig. 3. BBR increases phosphorylation of AMPK α in a time-dependent manner in rat skeletal muscles. Isolated epitrochlearis (A) and soleus (B) muscles were incubated in the presence of BBR (0.3 mmol/L) for 0 (basal), 15, 30, 45 or 60 minutes. The tissue lysate was subjected to Western blot analysis with antibodies to phosphorylated AMPK α or AMPK α . Fold increases are expressed relative to the level in muscles of the basal group.

The values are means $\pm$ S.E. The number of muscles in each group is as follows: basal (8), BBR 15 minutes (6), BBR 30 minutes (8), BBR 45 minutes (6), BBR 60 minutes (6). * $P < .05$ and ** $P < .01$ vs. basal; # $P < .05$ and ## $P < .01$ vs. BBR 15 minutes.

Fig. 4. BBR activates both AMPK α 1 and α 2 activities in rat skeletal muscles. Isolated epitrochlearis (A) and soleus (B) muscles were incubated in the absence (basal) or presence of BBR (0.3 mmol/L) for 30 minutes, or in the presence of DNP (0.5 mmol/L) for 15 minutes. The isoform-specific AMPK activity was determined in the anti-AMPK α 1 and anti-AMPK α 2 immunoprecipitates. Fold increases are expressed relative to the level in muscles of the basal group. The values are means $\pm$ S.E. The number of muscles in each group is as follows: basal (7), BBR (7), DNP (4). ** $P < .01$ vs. basal.

Fig. 5. BBR increases 3MG transport in rat skeletal muscles. Isolated epitrochlearis (A) and soleus (B) muscles were incubated in the absence (basal), or presence of BBR (0.3 mmol/L), AICAR (2 mmol/L) or insulin (1 μ mol/L) for 30 minutes, and then 3MG transport activity was determined. The values are means $\pm$ S.E. The number of muscles in each group is as follows: basal (7), BBR (7), AICAR (4), Insulin (4). ** $P < .01$ vs. basal.

Fig. 6. BBR increases phosphorylation of AS160 in rat skeletal muscles. Isolated epitrochlearis (A) muscle was incubated in the absence (basal), or presence of BBR (0.3 mmol/L) or AICAR (2 mmol/L) for 30 minutes. Isolated soleus (B) muscle was incubated in the absence (basal), or presence of BBR (0.3 mmol/L) for 30 minutes or DNP (0.5 mmol/L) for 15 minutes. The immunoprecipitate with anti-AS160 (Rab-GAP) was

subjected to Western blot analysis with antiphospho-(Ser/Thr) Akt substrate or anti-AS160 (Rab-GAP). Representative immunoblots are shown.

Fig. 7. BBR does not increase phosphorylation of Akt, p38 MAPK or p42/44 MAPK in rat skeletal muscles. Isolated epitrochlearis (A) and soleus (B) muscles were incubated in the absence (basal) or presence of BBR (0.3 mmol/L) for 30 minutes. The tissue lysate was subjected to Western blot analysis with antibodies to phosphorylated Akt, p38 MAPK or p44/42 MAPK. Representative immunoblots are shown.

Table 1

BBR decreases phosphocreatine (PCr) levels in rat skeletal muscles.

		Basal	BBR
epitrochlearis	ATP	5.7±0.5	5.2±0.4
	PCr	19.7±0.7	12.9±1.6*
soleus	ATP	2.5±0.3	2.1±0.3
	PCr	9.8±1.2	5.5±0.5*

Isolated epitrochlearis and soleus muscles were incubated in the absence (basal) or presence of BBR (0.3 mmol/L) for 30 minutes, and the intracellular ATP and PCr contents (in nmol/mg wet weight of muscle) were measured. The values are means ± S.E. The number of muscles in each group is as follows: epitrochlearis; ATP (8), PCr (8), soleus; ATP (6), PCr (8). *P < .01 vs. basal.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7