

Title	On the principal series representation of $SU(2,2)$ (Automorphic representations, automorphic L -functions and arithmetic)
Author(s)	Bayarmagnai, G.
Citation	数理解析研究所講究録 (2009), 1659: 157-165
Issue Date	2009-07
URL	http://hdl.handle.net/2433/140926
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

On the principal series representation of $SU(2, 2)$

G. Bayarmagnai

1 Introduction

Let G denote the special unitary group $SU(2, 2)$. In the paper, we will deal with the principal series representations of G which are parabolically induced by the minimal parabolic subgroup P_{min} with Langlands decomposition $P_{min} = MAN$;

$$\pi_{\sigma, \nu} = \text{Ind}_{P_{min}}^G (\sigma \otimes e^{\nu+\rho} \otimes 1_N),$$

where ρ is the half sum associated to the root system of the pair (G, A) , ν is a complex valued real linear form on $\mathfrak{a} = \text{Lie}(A)$, σ is a unitary character of M .

Let η be a continuous unitary character of N . We then have the Jacquet functional $J_{\sigma, \nu}$ on the space of differentiable functions of $L^2_{\sigma}(K)$, the representation space of $\pi_{\sigma, \nu}$, such that $J_{\sigma, \nu}(\pi_{\sigma, \nu}(n)f) = \eta(n)J_{\sigma, \nu}(f)$ for any $n \in N$. The functional defines an intertwiner J from $\pi_{\sigma, \nu}|_K$ to $\mathcal{A}_{\eta}(N \backslash G)$ by sending any $v \in \pi_{\sigma, \nu}|_K$ to the function $J_v(g) := J_{\sigma, \nu}(\pi_{\sigma, \nu}(g)v)$, ($g \in G$). Here the subspace of all K -finite vectors of $\pi_{\sigma, \nu}$ is denoted by $\pi_{\sigma, \nu}|_K$ and $\mathcal{A}_{\eta}(N \backslash G)$ is the subspace of $C^{\infty}(G)$ consisting of all moderate growth functions $f(g)$ such that $f(ng) = \eta(n)f(g)$ for $n \in N$ and $g \in G$. In fact, J is an intertwiner of K and \mathfrak{g} -equivariant, and hence the study of the image of J (the Whittaker model) leads us to the problem of the investigations of the (\mathfrak{g}, K) -module structure and the functions $J_v(g)$ for certain K -types of $\pi_{\sigma, \nu}$.

The main goal of this paper is to describe the above mentioned objects in terms of parameters of the principal series representation $\pi_{\sigma, \nu}$ explicitly. Note that our results are quite similar to that of Ishii [4] and Oda [5], for both $Sp(2, \mathbb{R})$ and $SU(2, 2)$ have the same restricted root system.

We also consider a matrix representations of the Knapp-Stein intertwining operator which have been motivated by a result of Goodman-Wallach [2].

2 Preliminaries

Let K be the compact group $S(U(2) \times U(2))$. Then K is the maximal compact subgroup of G fixed by the Cartan involution θ for G given by

$$\theta(g) = {}^t \bar{g}^{-1}, \quad g \in G.$$

We fix the following basis for the 7 dimensional Lie algebra $\mathfrak{k}_{\mathbb{C}}$, the complexification of $\mathfrak{k} = \text{Lie}(K)$:

$$\begin{aligned} h^1 &= \begin{pmatrix} h & 0 \\ 0 & 0 \end{pmatrix}, & h^2 &= \begin{pmatrix} 0 & 0 \\ 0 & h \end{pmatrix}, & I_{2,2} &= \begin{pmatrix} 1_2 & 0 \\ 0 & -1_2 \end{pmatrix}, \\ e_{\pm}^1 &= \begin{pmatrix} e_{\pm} & 0 \\ 0 & 0 \end{pmatrix}, & e_{\pm}^2 &= \begin{pmatrix} 0 & 0 \\ 0 & e_{\pm} \end{pmatrix}, \end{aligned}$$

where $h = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$, $e_+ = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ and $e_- = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$.

For every K -module V , it is clear that $I_{2,2} \in \mathfrak{k}_{\mathbb{C}}$ commutes with the action of K on V . If V is irreducible, then by Schur's lemma, the operator is a scalar of the identity map.

Lemma 2.1 *Let m_1, m_2 be positive integers and l be an integer. If $m_1 + m_2 + l$ is an even integer, then there is an irreducible K -module $(\tau_{[m_1, m_2; l]}, V_{m_1, m_2})$ with a basis $\{f_{pq} \mid 0 \leq p \leq m_1, 0 \leq q \leq m_2\}$ of V_{m_1, m_2} such that $I_{2,2}f_{pq} = lf_{pq}$ and*

$$\begin{aligned} h^1(f_{pq}) &= (2p - m_1)f_{pq}, & e_+^1(f_{pq}) &= (m_1 - p)f_{p+1,q}, & e_-^1(f_{pq}) &= pf_{p-1,q}, \\ h^2(f_{pq}) &= (2q - m_2)f_{pq}, & e_+^2(f_{pq}) &= (m_2 - q)f_{p,q+1}, & e_-^2(f_{pq}) &= qf_{p,q-1}. \end{aligned}$$

It follows from the fact that $SU(2) \times SU(2) \times \mathbb{C}^{(1)}$ is a twofold covering of K with the projection given by

$$pr(g_1, g_2; u) = \text{diag}(ug_1, u^{-1}g_2), \quad g_1, g_2 \in SU(2), \quad u \in \mathbb{C}^{(1)}.$$

3 K -finite vectors in the principal series

In this section, for each simple K -module $\tau \in \hat{K}$, we associate a matrix function $\mathbf{S}^{(\tau)}(k)$, $k \in K$, whose entries give a basis for the τ -isotypic component of $\pi_{\sigma, \nu}$. The main feature of this basis is that the both \mathfrak{g} and K -actions on $\pi_{\sigma, \nu} \upharpoonright_K$ have simple expressions in terms of parameters of given representation. For more details about this theme, we refer to [5] which is our main reference.

Proposition 3.1 *Let $H(\tau)$ be the τ -isotypic component of $L^2(K)$, and put $\dim(\tau) = n$. There exists a unique square matrix function $\mathbf{S}^{(\tau)}(k)$, $k \in K$, of size n with entries in $H(\tau)$,*

$$\mathbf{S}^{(\tau)}(k) = \begin{bmatrix} f_{11}(k) & \cdots & f_{n1}(k) \\ \vdots & \ddots & \vdots \\ f_{1n}(k) & \cdots & f_{nn}(k) \end{bmatrix} = \{f_{ij}(k)\}_{1 \leq i, j \leq n},$$

satisfying the following two conditions:

1. $\mathbf{S}^{(\tau)}(1_K) = \text{diag}(1, \dots, 1) \in M_n(\mathbb{C})$,
2. For each α ($1 \leq \alpha \leq n$), the set $\{f_{\alpha 1}(k), \dots, f_{\alpha n}(k)\}$ is a basis for τ as in Lemma 2.1. Moreover, we have

$$H(\tau) = \bigoplus_{\alpha} W_{\alpha},$$

where W_{α} denotes the space spanned by $f_{\alpha 1}(k), \dots, f_{\alpha n}(k)$.

Proof. The existence of the matrix function is similar to that of [5]. We consider the uniqueness. Assume that there exist two matrices $\mathbf{F}^{(\tau)}(k) = \{f_{ij}(k)\}$ and $\mathbf{G}^{(\tau)}(k) = \{g_{ij}(k)\}$ as required. Denote by F_{α} the isomorphism between τ and the space spanned by $\{f_{\alpha j}(k), \dots, f_{\alpha n}(k)\}$. Similarly, we define G_{α} for the α -th column of $\mathbf{G}^{(\tau)}(k)$. As a result, we obtain two ordered bases $\{F_{\alpha}\}_{\alpha}$ and $\{G_{\alpha}\}_{\alpha}$ for the n -dimensional vector space $\text{Hom}_K(\tau, H(\tau))$. Then we have the n by n matrix $A = \{a_{\alpha\beta}\}$, the change of coordinate matrix, such that

$$F_{\alpha} = \sum_{\beta} a_{\alpha\beta} G_{\beta}.$$

For a basis $\{f_{\gamma}\}$ of τ , one obtains

$$f_{\alpha\gamma}(k) = F_{\alpha}(f_{\gamma}) = \sum_{\beta} a_{\alpha\beta} G_{\beta}(f_{\gamma}) = \sum_{\beta} a_{\alpha\beta} f_{\beta\gamma}(k).$$

Evaluation at the point 1_K shows that

$$a_{\alpha\gamma} = \delta_{\alpha\gamma}.$$

If $v \neq 0 \in W_\alpha \cap W_\beta$, then $Kv = W_\alpha = W_\beta$. Schur's lemma and second condition imply that $\alpha = \beta$. Assume there is a matrix $\mathbf{S}^{(\tau)}(k)$ as required, we then have the direct sum decomposition of $H(\tau)$. \square

For each $\tau_m = \tau_{[m_1, m_2; l]} \in \hat{K}$, define a finite set $I(\tau_m)$ to be the collection of indices α such that W_α occurs in $\pi_{\sigma, \nu} |_K$ as a K -module. Thus, the cardinality of $I(\tau_m)$ is the K -multiplicity of τ_m in $\pi_{\sigma, \nu}$. Let s be the integer parameter corresponding to $\sigma \in \hat{M}$. By setting $n = (m_1 + m_2 + s)/2$, one can see that $p + q = n$ if $\alpha \in I(\tau_m)$ with $\alpha = (m_2 + 1)p + q + 1, (q \leq m_2)$. We identify the index α with the pair (p, q) defined by α .

We define a matrix function $\mathbf{S}_{\sigma, \nu}^{(\tau_m)}(k)$ attached to the τ -isotypic component of $\pi_{\sigma, \nu}$ by eliminating all the α -th columns of $\mathbf{S}^{(\tau_m)}(k)$ when $p + q \neq n$ and change the α -th columns by 0 if $\alpha \notin I(\tau_m)$ and $p + q = n$.

4 The (\mathfrak{g}, K) -module structure on $\pi_{\sigma, \nu}$

Let $\mathfrak{g} = \mathfrak{k} + \mathfrak{p}$ be the Cartan decomposition of $\mathfrak{g} = Lie(G)$ corresponding to θ . In this section, we explicitly describe $\mathfrak{p}_\mathbb{C}$ -action on the space

$$\pi_{\sigma, \nu} |_K \cong \bigoplus_{\tau_m \in \hat{K}} \bigoplus_{\alpha \in I(\tau_m)} W_\alpha.$$

Since the adjoint representation of K on $\mathfrak{p}_\mathbb{C}$ splits into two irreducible components, the antiholomorphic part \mathfrak{p}_- and the holomorphic part \mathfrak{p}_+ , it is enough to investigate the \mathfrak{p}_+ -action for our purpose. Let E_{ij} be the matrix unit of $M_4(\mathbb{R})$ with 1 in the (i, j) -entry and zero elsewhere. Then the set $\{E_{ij} \mid i = 1, 2, j = 3, 4\}$ forms a basis for \mathfrak{p}_+ . For a fixed pair $(e_1, e_2), e_j \in \{\pm 1\}$ with $j = 1, 2$, we define \mathbf{c}_t^j by

$$\mathbf{c}_t^j = \frac{t}{m_j + 1} \quad (0 \leq t \leq m_j + e_j).$$

Let (τ_m, V_m) be an irreducible representation of K with parametrization $m = [m_1, m_2; l]$. By the well known Clebsch-Gordan theorem, the irreducible components in the K -module $\mathfrak{p}_+ \otimes_{\mathbb{C}} \tau_m$ are precisely the K -representations

$$T = \{ \tau_{[m_1+e_1, m_2+e_2; l+2]} \mid e_1, e_2 \in \{\pm 1\} \},$$

and we will denote these by $\tau_{[\pm, \pm; +]}$ or $\tau_{[e_1, e_2; +]}$.

For each K -isomorphism between τ_m and W_α in Proposition 3.1, we have the following surjective homomorphism $\mathfrak{p}_+ \otimes_{\mathbb{C}} \tau_m \rightarrow \mathfrak{p}_+ W_\alpha$ of K -modules. Therefore, we obtain an injection

$$\mathfrak{p}_+ H_{\sigma, \nu}(\tau_m) \hookrightarrow \bigoplus_{\tau_{m'} \in T} H_{\sigma, \nu}(\tau_{m'})$$

which implies the following theorem. Here $H_{\sigma, \nu}(\tau_m)$ stands for the τ_m -isotypic component of $\pi_{\sigma, \nu}$.

Theorem 4.1 *Let $\tau_{[e_1, e_2; +]}$ be a simple K -submodule of the K -module $\mathfrak{p}_+ \otimes_{\mathbb{C}} \tau_m$ for a given simple K -module τ_m and the K -module (Ad, \mathfrak{p}_+) . Then we have that*

$$\mathbf{C}_{[e_1, e_2; +]} \mathbf{S}_{\sigma, \nu}^{(\tau_m)}(k) = \mathbf{S}_{\sigma, \nu}^{(\tau_{[e_1, e_2; +]})}(k) \Gamma_{[e_1, e_2; +]},$$

where the product of matrices of the left hand side is the differential operation. Here, $r = (s + l)/2$ and

1. $\Gamma_{[-,-,+]} = \{a_{ij}\}_{0 \leq i \leq n-1, 0 \leq j \leq n}$ is a matrix whose all non zero entries are given by

$$\begin{aligned} a_{t-1,t} &= \frac{1}{2}(\nu_2 + 1 + m_1 + r - 2t) && \text{if } (t, n-t) \in I(\tau_m), (t-1, n-t) \in I(\tau_{m'}), \\ a_{t,t} &= -\frac{1}{2}(\nu_1 - 1 - m_2 + r - 2t) && \text{if } (t, n-t) \in I(\tau_m), (t, n-t-1) \in I(\tau_{m'}). \end{aligned}$$

and $C_{[-,-,+]} = \{C_{ij}\}$ is a matrix of size $(m_1 m_2) \times (m_1 + 1)(m_2 + 1)$ with entries given by

$$\begin{aligned} C_{m_2 p+q+1, (m_2+1)p+q+1} &= -E_{14}, \\ C_{m_2 p+q+1, (m_2+1)p+q+2} &= -E_{13}, \\ C_{m_2 p+q+1, (m_2+1)(p+1)+q+1} &= E_{24}, \\ C_{m_2 p+q+1, (m_2+1)(p+1)+q+2} &= E_{23}, \end{aligned}$$

for each $0 \leq p \leq m_1 - 1$ and $0 \leq q \leq m_2 - 1$, but all other entries are 0.

2. $\Gamma_{[+,+,+]} = \{a_{ij}\}_{0 \leq i \leq n+1, 0 \leq j \leq n}$ is a matrix whose all non zero entries are given by

$$\begin{aligned} a_{t,t} &= \frac{1}{2}(\nu_2 + 1 + m_1 + r - 2t)(1 - c_t^1) c_{\nu-t+1}^2 && \text{if } (t, n-t) \in I(\tau_m), (t, n-t+1) \in I(\tau_{m'}), \\ a_{t+1,t} &= \frac{1}{2}(\nu_1 + 3 + 2m_1 + m_2 + r - 2t) c_{t+1}^1 (c_{\nu-t}^2 - 1) && \text{if } (t, n-t) \in I(\tau_m), (t+1, n-t) \in I(\tau_{m'}). \end{aligned}$$

and $C_{[+,+,+]} = \{C_{ij}\}$ is a matrix of size $(m_1 + 2)(m_2 + 2) \times (m_1 + 1)(m_2 + 1)$ with entries given by

$$\begin{aligned} C_{(m_2+2)p+q+1, (m_2+1)p+q+1} &= -(1 - c_p^1)(1 - c_q^2) E_{23}, \\ C_{(m_2+2)p+q+1, (m_2+1)p+q} &= (1 - c_p^1) c_q^2 E_{24}, \\ C_{(m_2+2)p+q+1, (m_2+1)(p-1)+q+1} &= -c_p^1 (1 - c_q^2) E_{13}, \\ C_{(m_2+2)p+q+1, (m_2+1)(p-1)+q} &= c_p^1 c_q^2 E_{14}, \end{aligned}$$

for each $0 \leq p \leq m_1 + 1$ and $0 \leq q \leq m_2 + 1$, but all other entries are 0.

3. $\Gamma_{[-,+,+]} = \{a_{ij}\}_{0 \leq i \leq n, 0 \leq j \leq n}$ is a square matrix whose all non zero entries are given by

$$\begin{aligned} a_{t-1,t} &= \frac{1}{2}(\nu_2 + 1 + m_1 + r - 2t) c_{\nu-t+1}^2 && \text{if } (t, n-t) \in I(\tau_m), (t-1, n-t+1) \in I(\tau_{m'}), \\ a_{t,t} &= \frac{1}{2}(\nu_1 + 1 + m_2 + r - 2t)(1 - c_{\nu-t}^2) && \text{if } (t, n-t) \in I(\tau_m), (t, n-t) \in I(\tau_{m'}). \end{aligned}$$

and $C_{[-,+,+]} = \{C_{ij}\}$ is a matrix of size $m_1(m_2 + 2) \times (m_1 + 1)(m_2 + 1)$ with entries given by

$$\begin{aligned} C_{(m_2+2)p+q+1, (m_2+1)p+q+1} &= (1 - c_q^2) E_{13}, \\ C_{(m_2+2)p+q+1, (m_2+1)p+q} &= -c_q^2 E_{14}, \\ C_{(m_2+2)p+q+1, (m_2+1)(p+1)+q+1} &= -(1 - c_q^2) E_{23}, \\ C_{(m_2+2)p+q+1, (m_2+1)(p+1)+q} &= c_q^2 E_{24}, \end{aligned}$$

for $0 \leq p \leq m_1 + 1$ and $0 \leq q \leq m_2 - 1$, but all other entries are 0.

4. $\Gamma_{[+,-,+]} = \{a_{ij}\}_{0 \leq i \leq n, 0 \leq j \leq n}$ is a square matrix whose all non zero entries are given by

$$\begin{aligned} a_{t,t} &= \frac{1}{2}(\nu_2 + 1 + m_1 + r - 2t)(1 - c_t^1) && \text{if } (t, n-t) \in I(\tau_m), (t, n-t) \in I(\tau_{m'}), \\ a_{t+1,t} &= \frac{1}{2}(\nu_1 + 1 + 2m_1 - m_2 + r - 2t) c_{t+1}^1 && \text{if } (t, n-t) \in I(\tau_m), (t+1, n-t-1) \in I(\tau_{m'}). \end{aligned}$$

and $C_{[+, -, +]} = \{C_{ij}\}$ is a matrix of size $(m_1 + 2)m_2 \times (m_1 + 1)(m_2 + 1)$ with entries given by

$$\begin{aligned} C_{m_2 p + q + 1, (m_2 + 1)p + q + 1} &= (1 - c_p^1)E_{24}, \\ C_{m_2 p + q + 1, (m_2 + 1)p + q + 2} &= (1 - c_p^1)E_{23}, \\ C_{m_2 p + q + 1, (m_2 + 1)(p - 1) + q + 1} &= c_p^1 E_{14}, \\ C_{m_2 p + q + 1, (m_2 + 1)(p - 1) + q + 2} &= c_p^1 E_{13}, \end{aligned}$$

for $0 \leq p \leq m_1 + 1$ and $0 \leq q \leq m_2 - 1$, but all other entries are 0.

4.0.1 The Knapp-Stein operator

In this subsection, we consider a matrix representation of the Knapp-Stein operator with respect to the basis for $\pi_{\sigma, \nu} |_K$. This is motivated by Theorem 6.7 in the paper of Goodman-Wallach [2].

Let us recall the Knapp-Stein intertwining operator $A_{\sigma, \nu}^s$ from the space of all C^∞ -vectors of $\pi_{\sigma, \nu}$ to that of $\pi_{s(\sigma), s(\nu)}$ defined by

$$(A_{\sigma, \nu}^s f)(k) = \int_{\tilde{N}_s} a(n_s s^* k)^{\nu + \rho} f(k(n_s s^* k)) dn_s, \quad (f \in \pi_{\sigma, \nu}^\infty).$$

Here $s^* \in K$ such that $s := Ad(s^*) \in W(A)$, $\tilde{N}_s = N \cap s^* N s^{*-1}$ and $s(\sigma)$ is a character of M given by $s(\sigma)(m) = \sigma(s^* m s^{*-1})$, $m \in M$. Since it is a linear map from $\pi_{\sigma, \nu}$ to $\pi_{s(\sigma), s(\nu)}$ satisfying

$$A_{\sigma, \nu}^s \pi_{\sigma, \nu}(x) f = \pi_{s(\sigma), s(\nu)}(x) A_{\sigma, \nu}^s f, \quad x \in G \text{ (or } U(\mathfrak{g})),$$

we have a linear map

$$A^s(\tau) : \text{Hom}_K(\tau, \pi_{\sigma, \nu} |_K) \rightarrow \text{Hom}_K(\tau, \pi_{s(\sigma), s(\nu)} |_K).$$

for any $\tau \in \hat{K}$.

Let $[\alpha_i]$ be the K -isomorphism from τ to W_{α_i} for $\alpha_i \in I(\tau)$. We equip the space $\text{Hom}_K(\tau, \pi_{\sigma, \nu} |_K)$ with the basis consisting of the K -homomorphisms $[\alpha_i]$. Similarly, we choose a basis for the space $\text{Hom}_K(\tau, \pi_{s(\sigma), s(\nu)} |_K)$. Then we want to compute all entries a_{ij} of the matrix $A^s(\tau) = (a_{ij})$ such that

$$A^s(\tau)[\alpha_i] = \sum_{\alpha_j^s \in I} a_{ij} \cdot [\alpha_j^s]$$

where $I = \{\alpha^s \mid W_{\alpha^s} \hookrightarrow \pi_{s(\sigma), s(\nu)} |_K\}$. For each basis vector f_{pq} of τ as in Lemma 2.1, we have that

$$(A^s(\tau)[\alpha_i])(f_{pq}) = \sum_{\alpha_j^s \in I} a_{ij} \cdot [\alpha_j^s](f_{pq}) = \sum_{\alpha_j^s \in I} a_{ij} \cdot f_{\alpha_j^s, pq}^{(\tau)}(k).$$

On the other hand, by definition of the map $A^s(\tau)$, one has

$$(A^s(\tau)[\alpha_i])(f_{pq}) = (A_{\sigma, \nu}^s f_{\alpha_i, pq}^{(\tau)})(k), \quad \alpha_i \in I(\pi_{\sigma, \nu}, \tau).$$

Thus we have the following formula for the coefficients a_{ij} of the matrix $A^s(\tau)$ for each $\tau \in \hat{K}$.

Lemma 4.2 *Let α_i be in $I(\pi_{\sigma, \nu}, \tau)$ and α_j^s be in $I(\pi_{s(\sigma), s(\nu)}, \tau)$. Then the (i, j) -th coefficient of $A^s(\tau)$*

$$a_{ij} = (A_{\sigma, \nu}^s f_{\alpha_i, \alpha_j^s}^{(m)})(1_4).$$

Example 4.3

Let s be a generator of $W(A)$ whose image is the matrix $\text{diag}(1, -1)$ under the representation of $W(A)$ on \mathfrak{a}^* . Then we choose the corresponding $s^* \in K$ as the matrix $\text{diag}(1, -i, 1, i)$ and hence

$$\bar{N}_s = \exp(\mathfrak{g}_{-2\lambda_2}) = \left\{ n_s(t) = \kappa^{-1} \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & t & & 1 \end{pmatrix} \kappa : t \in \mathbb{R} \right\}.$$

Since $n_s \in \bar{N}_s$, one has ${}^t n_s I_{2,2} n_s = I_{2,2}$ and hence $n_s s^* = I_{2,2} {}^t n_s^{-1} I_{2,2} s^*$. Thus, we have the following.

Assume $n_s = n_s(t) \in \bar{N}_s$. Let $n' \in N$, $a(n_s s^*) \in A$ and $k(n_s s^*) \in K$ be so that $n_s s^* = n' a(n_s s^*) k(n_s s^*)$. Then

$$\begin{aligned} a(n_s s^*)^{\nu+\rho} &= (1+t^2)^{-\frac{\nu_2+1}{2}}, \\ k(n_s s^*) &= \text{diag}(1, -iu, -1, -iu^{-1}) \end{aligned}$$

where $u = ((1-it)/(1+it))^{\frac{1}{2}}$.

For a fixed $\tau_m \in \hat{K}$, therefore

$$f_{\gamma_i, \beta_j}(k(n_s s^*)) = 0 \text{ when } \gamma_i \neq \beta_j$$

If $\tau = \tau_{[m_1, m_2; l]}$ then we have

$$A^s(\tau) = 2\pi 2^{-\nu_2} \Gamma(\nu_2) \text{diag} \left[\frac{(-1)^{(m_1+m_2)/2-p+1} i^{m_2+r}}{\Gamma(\frac{1}{2}\nu_2 + \frac{1}{2} + d) \Gamma(\frac{1}{2}\nu_2 + \frac{1}{2} - d)} \right]_p$$

where $d = \frac{1}{2}(m_1 + r - 2p)$ for $(p, (m_1 + m_2)/2 - p) \in I(\pi_{\sigma, \nu}, \tau_m)$.

5 Whittaker functions

The main focus of this section is on the integral expressions of Whittaker functions on G related to certain principal series. The results of the section 4.1 lead us to the study of Whittaker functions related to some K -types. For this purpose, we focus our investigation on the principal series representations which contain one dimensional K -types and apply the method used in [4] to evaluate such Whittaker functions. More precisely, in this setting, the character σ of M factors through a character χ of μ_2 . Let $(\pi_{\chi, \nu}, L_\chi^2(K))$ denote the principal representation series corresponding to such character σ .

For an integer u , define a function $f_u(k)$ on K by $f_u(k) := \det(k_2)^u$, $k = \begin{pmatrix} k_1 & 0 \\ 0 & k_2 \end{pmatrix} \in K$.

Lemma 5.1 *Let $f_u(k)$ be as above. Then $\tau_{[0,0;2u]} \cong \mathbb{C} f_u(k)$ as K -modules. Moreover, if $\chi(-1) = (-1)^u$ then $f_u(k) \in L_\chi^2(K)$ and $[\pi_{\chi, \nu} : \tau_{[0,0;2u]}] = 1$.*

5.1 The Jacquet integral.

Let $J_{\chi, \nu}$ be the Jacquet functional on the subspace of differentiable functions of $L_\chi^2(K)$ given by

$$J_{\chi, \nu}(f) = \int_N \eta(n)^{-1} a(s^* n)^{\nu+\rho} f(k(s^* n)) dn$$

for a differentiable function f in $L_\chi^2(K)$ and the longest element $s \in W(A)$. Here $W(A)$ is the Weyl group defined as the quotient of $M^* = N_K(\mathfrak{a})$, the normalizer of \mathfrak{a} in K , by M and s^* is an element of M^* mapping to the longest element $s \in W(A)$.

Then one has $J_{\chi,\nu}(\pi(n)f) = \eta(n)J_{\chi,\nu}(f)$ and hence

$$J \in \text{Hom}_{(\mathfrak{g},K)}(\pi_{\chi,\nu}|_K, \mathcal{A}_\eta(N \setminus G)), \quad (1)$$

where J associates $v \in \pi_{\chi,\nu}|_K$ to the function $J_v(g) := J_{\chi,\nu}(\pi_{\chi,\nu}(g)v)$, ($g \in G$). We want to have an explicit formula for the A -radial part:

$$J_{f_u}(a) = J_{\chi,\nu}(\pi_{\chi,\nu}(a)f_u) = a^{-\nu+\rho} \int_N \eta(ana)^{-1} a(s^*n)^{\nu+\rho} f_u(k(s^*n)) dn.$$

In our case, we can choose $I_{2,2} \in K$ for $s^* \in K$.

5.1.1 The first modification

Let E_{ij} be the usual matrix with 1 in (i, j) -entry and zero elsewhere. Put

$$\begin{aligned} E_0 &= \kappa^{-1}(E_{12} - E_{43})\kappa, & E_1 &= i\kappa^{-1}(E_{12} + E_{43})\kappa, & E_2 &= \kappa^{-1}E_{24}\kappa, \\ F_0 &= \kappa^{-1}(E_{14} + E_{23})\kappa, & F_1 &= i\kappa^{-1}(E_{14} - E_{23})\kappa, & F_2 &= \kappa^{-1}E_{13}\kappa, \end{aligned}$$

by setting $i = \sqrt{-1}$ and

$$\kappa = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ -i & 0 & i & 0 \\ 0 & -i & 0 & i \end{pmatrix}.$$

Then the corresponding root spaces of positive roots in $\Phi(\mathfrak{g}, \mathfrak{a})$ are given by

$$\begin{aligned} \mathfrak{g}_{\lambda_1 - \lambda_2} &= E_0 \cdot \mathbb{R} \oplus E_1 \cdot \mathbb{R}, & \mathfrak{g}_{2\lambda_2} &= E_2 \cdot \mathbb{R}, \\ \mathfrak{g}_{\lambda_1 + \lambda_2} &= F_0 \cdot \mathbb{R} \oplus F_1 \cdot \mathbb{R}, & \mathfrak{g}_{2\lambda_1} &= F_2 \cdot \mathbb{R}. \end{aligned}$$

Let \mathfrak{n} be a subalgebra defined by $\mathfrak{n} = \sum_{\alpha \in \Phi_+} \mathfrak{g}_\alpha$. We now describe elements of a maximal nilpotent subgroup N of G given by $N = \exp(\mathfrak{n})$.

The Killing form $B(X, Y) = \text{tr}(\text{ad}X \cdot \text{ad}Y)$, ($X, Y \in \mathfrak{g}$) and Cartan involution θ of \mathfrak{g} induce an inner product \langle, \rangle of \mathfrak{g} via

$$\langle X, Y \rangle = -B(X, Y^\theta), \quad (X, Y \in \mathfrak{g}).$$

Then one has that $\langle \mathfrak{g}_\alpha, \mathfrak{g}_\beta \rangle = 0$ if $\alpha \neq \beta$, because of the involution θ . Moreover, one can see that the set $\{E_i, F_i \mid i = 0, 1, 2\}$ is an \langle, \rangle -orthogonal basis for \mathfrak{n} such that a each element $n = n(n_0, n_1, n_2, n_3)$ in the maximal unipotent group $N = \exp(\mathfrak{n})$ is expressed in the form:

$$\kappa^{-1} \begin{pmatrix} 1 & n_0 & & \\ & 1 & & \\ & & 1 & \\ & & -\bar{n}_0 & 1 \end{pmatrix} \begin{pmatrix} 1 & n_1 & n_2 \\ & 1 & \bar{n}_2 & n_3 \\ & & 1 & \\ & & & 1 \end{pmatrix} \kappa$$

for $n_1, n_3 \in \mathbb{R}$, $n_0, n_2 \in \mathbb{C}$.

Lemma 5.2 *We have*

1. Set $N_1 = \begin{pmatrix} n_1 & n_2 \\ \bar{n}_2 & n_3 \end{pmatrix}$ for $n = n(n_0, n_1, n_2, n_3) \in N$. Then

$$f_u(k(I_{2,2}n)) = \left(\frac{\det(1 - \sqrt{-1}N_1)}{\det(1 + \sqrt{-1}N_1)} \right)^{\frac{1}{2}}.$$

2. Let η be a character of N determined by a real number c_2 and $c = c_0 + \sqrt{-1}c_1 \in \mathbb{C}$. Then

$$\eta(ana^{-1}) = \exp(2\sqrt{-1} \left(\frac{a_1}{a_2} \operatorname{Re}(\bar{c}n_0) + c_2 a_2^2 n_3 \right)),$$

where $a_i = \exp(t_i)$, ($i = 1, 2$) for $a = a(t_1, t_2) \in A$.

3. For $\nu = (\nu_1, \nu_2) \in \operatorname{Hom}_{\mathbb{R}}(\mathfrak{a}, \mathbb{C})$, we have that $a(I_{2,2}n)^{\nu+\rho} = \Delta_1^{-\frac{\nu_1-\nu_2}{2}-1} \Delta_2^{-\frac{\nu_2+1}{2}}$ where
 $\Delta_1 = 1 + n_1^2 + \bar{n}_2 n_2 + (\bar{n}_0 n_2 + n_0 \bar{n}_2)(n_1 + n_3) + \bar{n}_0 n_0(1 + \bar{n}_2 n_2 + n_3^2)$,
 $\Delta_2 = 1 + n_1^2 + 2n_2 \bar{n}_2 + n_3^2 + (n_1 n_3 - n_2 \bar{n}_2)^2$ for $n = n(n_0, n_1, n_2, n_3) \in N$.

For future convenience, we choose a new coordinate for A by

$$y = (y_1, y_2) = \left(\frac{a_1}{a_2}, a_2^2 \right).$$

Since $f \rightarrow J_f(g)$ is the Whittaker realization of $\pi_{\chi, \nu}$, $J_{f_u}(a)$ is the radial part of a Whittaker function on G belonging to π_{ν} . Thus, in the new coordinate system, we can summarize that the radial part of Whittaker function associated with the K -type τ_u can be written in the form

$$y_1^{-\nu_1+3} y_2^{-\frac{\nu_1+\nu_2}{2}+2} \int_N \Delta_1^{-\frac{\nu_1-\nu_2}{2}-1} \Delta_2^{-\frac{\nu_2+1}{2}} \times \exp(-2\sqrt{-1}(y_1 \operatorname{Re}(\bar{c}n_0) + c_2 y_2 n_3)) f_u(k(I_{2,2}n)) dn,$$

where dn is a multiplicative Haar measure on N . Now we shall give a normalization of Haar measure of N . Since the exponential map of \mathfrak{n} onto N is an analytic isomorphism, there exists a unique Haar measure dn on N that corresponds to Lebesgue measure on \mathfrak{n} .

Lemma 5.3 *The radial part of the moderate growth Whittaker function $W_{(\nu_1, \nu_2)}(y_1, y_2; u)$ (up to constant) associated with the K -type τ_u can be written in the form*

$$y_1^{-\nu_1+3} y_2^{-\frac{\nu_1+\nu_2}{2}+2} \int_{\mathbb{R}^6} \Delta_1^{-\frac{\nu_1-\nu_2}{2}-1} \Delta_2^{-\frac{\nu_2+1}{2}} \exp(-2\sqrt{-1}(c_0 z_0 y_1 + c_1 t_0 y_1 - n_3 y_2)) f_u(k(I_{2,2}n))$$

with respect to $dz_0 dt_0 dn_1 dz_2 dt_2 dn_3$. Here $n_i = z_i + \sqrt{-1}t_i$ ($i = 0, 2$).

In fact, it suffices to consider the cases $u = 0$ and 1 for our purposes. Let $K_{\mu}(z)$ be the Bessel function.

Theorem 5.4 *Let $\pi_{\chi, \nu}$ be irreducible and η be a nondegenerated unitary character N . Then we have the following assertions on the A -radial part of the primary Whittaker function $W_{(\nu_1, \nu_2)}(y_1, y_2; u)$.*

If χ is trivial then the Whittaker function $W_{(\nu_1, \nu_2)}(y_1, y_2; 0)$ is identified with $y_1^3 y_2^2$ times

$$\int_0^{\infty} \int_0^{\infty} T_{\nu_1, \nu_2}(y_1, y_2, t_1, t_2) \frac{dt_1}{t_1} \frac{dt_2}{t_2}.$$

If χ is non-trivial then the Whittaker function $\bar{W}_{(\nu_1, \nu_2)}(y_1, y_2; 1)$ is identified with $y_1^4 y_2^3/4$ times

$$\int_0^{\infty} \int_0^{\infty} T_{\nu_1, \nu_2}(y_1, y_2, t_1, t_2) (\sqrt{t_1/t_2} - 1/\sqrt{t_1 t_2}) \frac{dt_1}{t_1} \frac{dt_2}{t_2}$$

where $T_{\nu_1, \nu_2}(y_1, y_2, t_1, t_2)$ is the function

$$K_{\frac{\nu_1+\nu_2}{2}}(2\sqrt{t_2/t_1}) K_{\frac{\nu_2-\nu_1}{2}}(2\sqrt{t_1 t_2}) \exp\left(-|c_2|y_2 t_1 - \frac{|c_2|y_2}{t_1} - \frac{t_2}{|c_2|y_2} - (c_0^2 + c_1^2)|c_2| \frac{y_1^2 y_2}{t_2}\right)$$

Note here that, we need the following formula to reduce the number of integral symbols corresponding to the root spaces $\mathfrak{g}_{\lambda_1-\lambda_2}$ and $\mathfrak{g}_{\lambda_1+\lambda_2}$:

Formula 5.5 *Let $a, c \in \mathbb{R}_+^*$ and $b, \alpha, \beta \in \mathbb{R}$ such that $\alpha^2 + \beta^2 = 1$. Then we have*

$$\int_{\mathbb{R}} \int_{\mathbb{R}} \exp(-c(x^2 + y^2) - a(\alpha x + \beta y)^2 + 2\sqrt{-1}b(\alpha x + \beta y)) dx dy = \frac{\pi}{(c^2 + ac)^{\frac{1}{2}}} \exp\left(\frac{-b^2}{a+c}\right).$$

References

- [1] G. Bayarmagnai, Explicit evaluation of certain Jacquet integrals on $SU(2, 2)$, Preprint, 2008.
- [2] Goodman, R. Wallach, N. R., Whittaker vectors and conical vectors. *J. Funct. Anal.*, 39 (1980), 199-279.
- [3] T. Hayata, Differential equations for principal series Whittaker functions on $SU(2, 2)$. *Indag. Math. (N.S.)* 8 (1997), no.4, 493–528.
- [4] T. Ishii, On principal series Whittaker functions on $Sp(2, \mathbb{R})$, *J. Func. Anal.* 225 (2005) 1-32.
- [5] T. Oda, The standard (\mathfrak{g}, K) -modules of $Sp(2, \mathbb{R})$, Preprint Series, UTMS 2007-3, Graduate School of Mathematical Sciences, University of Tokyo.
- [6] T. Miyazaki and T.Oda, Principal series Whittaker functions on $Sp(2, \mathbb{R})$, – Explicit formulae of differential equations, Proceeding of the 1993 Workshop, Automorphic forms and related topics, The Pyungsan Institute for Mathematical Sciences, pp. 55-92.
- [7] David A. Vogan, Representations of real reductive Lie groups. *Progress in Mathematics*, vol. 15, Birkhauser, Boston, Basel, Stuttgart, 1981.
- [8] N.R. Wallach, Real reductive groups. I, *Pure and Applied Mathematics* vol. 132, Academic Press Inc., Boston, MA, 1988.
- [9] Yamashita, H., Embedding of discrete series into induced representations of semisimple Lie groups, I, General theory and the case of $SU(2, 2)$, *Japan J. Math.*, 16 (1990), 31-95.
- [10] Yamashita, H., Embedding of discrete series into induced representations of semi-simple Lie groups, II, Generalized Whittaker models for $SU(2, 2)$, *J. Math. Kyoto Univ.*, 31-1 (1991), 543-571.