



IUGG 2011, Melbourne (Australia), 2 July 2011
- JA05 Data rescue, digitisation and metadata
requirements in geophysics -

3761

Inter-university Upper atmosphere Global Observation NETwork (IUGONET)

**H. Hayashi, Y. Koyama, T. Hori, Y. Tanaka, M. Kagitani,
A. Shinbori, S. Abe, T. Kouno, D. Yoshida, S. UeNo,
N. Kaneda, M. Yoneda, and IUGONET project team**


The IUGONET project aims at building “e-infrastructure” for researchers to effectively find, get, and analyze various kinds of upper atmospheric data spread over universities and institutes.

- *To distribute ground-based observational data accumulated over 50 years since IGY (both digital and analogue data)*
- *To promote analyses of multi-disciplinary data, which will lead to comprehensive studies of mechanisms of long-term variations in the upper atmosphere*

Participating universities and research institutes

- Planetary Plasma and Atmospheric Research Center, [Tohoku University](#)
- [National Institute of Polar Research](#)
- Solar Terrestrial Environment Laboratory, [Nagoya University](#)
- Research Institute for Sustainable Humanosphere, [Kyoto University](#)
- World Data Center for Geomagnetism, [Kyoto University](#)
- Kwasan and Hida Observatories, [Kyoto University](#)
- Space Environment Research Center, [Kyushu University](#)

Observations by IUGONET institutions



Various observation parameters (wind, geomagnetism, aurora, sunspot etc.) taken by various techniques in various time periods at various locations and altitudes

Such observational data not necessarily well used in scientific researches so far
→ What's the problem?

SuperDARN Radar

Syowa Station
SuperDARN radar x2
MF radar
aurora imagers
magnetometer
ELF/VLF receiver
riometer

Peru Ica University




- ▲ MST radar
- ◆ MF/meteor radar
- ◆ MAGDAS magnetometer
- ▲ FM - CW radar
- OMTI imager
- ◆ WDC magnetometer
- Magnetic Equator (IGRF2005, Height 100km)

Project Schematic


PROBLEM: Various kind, huge amount of data spread over institutes and universities



SOLUTION: Create a metadata database for cross-search of these distributed data



Promote new types of upper atmospheric research by analysis of multi-disciplinary data




● Many metadata formats available in Earth and planetary sciences !

- Dublin Core
- ISO 19115 / 19139
- GCMD DIF
- FGDC CSDGM
- IPY Metadata Profile
- ISTP Standards
- SPASE
- ...

● IUGONET adopted **SPASE** as our base format

originally developed to describe research resources regarding heliospheric and magnetospheric satellite observations

- closely related to STP and upper atmospheric researches (➔ **easy to use as a base format**)
- new metadata elements & words appendable (➔ **customizable according to our data**)
- widely-used in existing Virtual Observatories (➔ **possible to exchange metadata**)



(<http://www.spase-group.org>)

- IUGONET uses **DSpace** as the metadata DB platform

- ✓ Free software, widely used by digital repositories in many universities over the world.
- ✓ Including fundamental functions to register, search, provide, and harvest metadata written in the IUGONET metadata format.



registered



```
<?xml version="1.0" encoding="UTF-8"?>
<spase lang="en" xmlns:xsi="http://www.w3.org/2001/XMLSchema-
instance" xmlns="http://www.iugonet.org/data/schema"
xsi:schemaLocation="http://www.iugonet.org/data/schema
http://www.iugonet.org/data/schema/iugonet.xsd ">
  <Version>1.0.0</Version>
  <NumericalData>

    <ResourceID>spase://IUGONET/NumericalData/STEL/SuperDARN/
HOK/HFradar/sd_hok_common_ergsccdf</ResourceID>
    <ResourceHeader>
      <ResourceName>SuperDARN Hokkaido HF radar, common
mode data distributed by
ERG-SC</ResourceName>
      <ReleaseDate>2009-04-01T00:00:00</ReleaseDate>
      <ExpirationDate>2199-12-31T23:59:59</ExpirationDate>
      <Description>Common mode data generated by SuperDARN
Hokkaido HF radar. Data
. Data files are distributed in the CDF format through ERG-
SC</Description>
      ...
      ...
    </NumericalData>
```

**keyword
search**

**spatial coverage
search**

**time range
search**



IUGONET Metadata DB >

This DB is now in beta testing. We appreciate your [feedback](#).

IUGONET
Metadata DB for Upper Atmosphere

☒ **Free Word:**

 (e.g. ionosphere, troposphere, magnetosphere, heliosphere.....)

☒ **Time:**
 from to [UTC]

☒ **Spatial Coverage/Map:**
 Latitude: Southernmost , Northernmost [degree]
 Longitude: Westernmost , Easternmost [degree]
 OR [\[close map\]](#)

☒ **Data Types:**
 Data Set (☒ Numerical ☒ Plot / Movie) ☐ Data File / Plot ☐ Instrument ☐ Observatory ☒

RSS Feeds
[RSS](#) [RSS](#) [RSS](#)

Search/Retrieve URL Service
[GOSS](#) [Share](#)

Latest Update Sat
 May 21 09:56:34
 JST 2011
[more...](#)

[WSC](#) XHTML 1.0

Datasets that match input keyword(s) are listed !

Relative Stop Date: 14 days ago (-P14D)
<http://gemsissc.stelab.nagoya-u.ac.jp/erg/>
 Repository: <spase://IUGONET/Repository/STEL/ERG-SC>
 Instrument: <spase://IUGONET/Instrument/STEL/SuperDARN/HOK>

The common time fitacf CDF data of SuperDARN King Salmon HF radar distributed by ERG-SC

NumericalData

Common mode data obtained by SuperDARN King Salmon HF radar. Data files are distributed in the CDF format through the ERG-SC repository

Start Date: 2006-12-02T00:00:00

Relative Stop Date: 180 days ago (-P180D)

<http://gemsissc.stelab.nagoya-u.ac.jp/erg/>

Repository: <spase://IUGONET/Repository/STEL/ERG-SC>

Instrument: <spase://IUGONET/Instrument/STEL/SuperDARN/KSR/HFradar>

Standard observation data of the troposphere and lower stratosphere taken by the MU radar (NetCDF format)

NumericalData

The 10-minute average observation data in the NetCDF (Network Common Data Form) format taken by the MU radar at Shigaraki in the Shiga prefecture, Japan (34.85N, 136.10E, 385m MSL), which has been operated in the standard observation mode of the troposphere and stratosphere. The observation data are stored in the NetCDF files of each day. The file name is (year)(month)(day).nc. The NetCDF data include range, height, time, three components of wind velocity, radial Doppler velocity, echo power, spectral width and noise level for each beam number and so on. The azimuth and zenith angles of beam 1, 2, 3, 4 and 5 are (0, 0), (0, 10), (90, 10), (180, 10) and (270, 10), respectively, in unit of degree. The value of 1.0e+10 means missing data.

Start Date: 1986-03-16T15:05:00

Relative Stop Date: 14 days ago (-P14D)

<http://www.rish.kyoto-u.ac.jp/radar-group/mu/data/>

Repository: <spase://IUGONET/Repository/RISH/RISHDB>

Instrument: <spase://IUGONET/Instrument/RISH/misc/SGK/MUradar>

Field-aligned irregularity (FAI) observation data of the ionosphere taken by the EAR (NetCDF format)

NumericalData

The field-aligned irregularity (FAI) observation data in the NetCDF (Network Common Data Form) format taken by the equatorial atmosphere radar (EAR) at Kototabang, Indonesia (0.20S, 100.32E, 865m MSL). This FAI observation mode covers a wide altitude range from 80 to 600 km in the ionosphere (D-region (below 90 km), E-region (90-150 km), and F-region (above 150 km)). The observation data are stored in the NetCDF files of each day and observation parameter. The file name is (year)(month)(day).(observation parameter).nc. The NetCDF data include range, height, time, radial Doppler velocity, echo power, spectral width and noise level for each beam number and so on. Details of the observation parameter are described in the EAR-FAI homepage (<http://www.rish.kyoto-u.ac.jp/ear/data-fai/index.html>). The value of 1.0e+10 means missing data.

Description:

The 10-minute average observation data in the NetCDF (Network Common Data Form) format taken by the equatorial atmosphere radar (EAR) at Kototabang, Indonesia (0.203S, 100.320E, 865m MSL), which has been operated in the standard observation mode of the troposphere and stratosphere. The observation data are stored in the NetCDF files of each day. The file name is (year)(month)(day).nc. The NetCDF data include range, height, time, three components of wind velocity, radial Doppler velocity, echo power, spectral width and noise level for each beam number and so on. The azimuth and zenith angles of beam 1, 2, 3, 4 and 5 are (0, 0), (0, 10), (90, 10), (180, 10) and (270, 10), respectively, in unit of degree. The value of 1.0e+10 means missing data.

Acknowledgement:

If you acquire EAR data, we ask that you acknowledge us in your use of the data. This may be done by including text such as EAR data provided by Research Institute for Sustainable Humanosphere of Kyoto University.

ReleaseDate:

2011-05-06T00:00:00

Contact PersonID:

0: space://IUGONET/Person/Hiroiyuki.Hashiguchi
 1: space://IUGONET/Person/EAR.Management.Group
 2: space://IUGONET/Person/Noriko.Hashiguchi
 3: space://IUGONET/Person/RISH.Metadata.Management.Group

Contact Role:

0: PrincipalInvestigator
 1: GeneralContact
 2: DataProducer
 3: MetadataContact

AccessInformation RepositoryID:

space://IUGONET/Repository/RISH/RISHDB

AccessInformation AccessURL URL:

<http://www.rish.kyoto-u.ac.jp/ear/data/index.html>

AccessInformation Availability:

Online

AccessInformation AccessRights:

Open

AccessInformation Format:

NetCDF

Description of data

Jump to metadata of “person”
 (e-mail address, etc.)


Information of data access

IUGONET data analysis software (UDAS) = IDL + TDAS


- To help users easily download, visualize, and analyze various data provided from us

THEMIS Data Analysis Software suite
– a set of IDL libraries

Able to make stacked plots of time series to compare various kind of data



Easy to handle data even for those who are not familiar with the data by using GUI



Project timeline

ITEMS		FY2009	FY2010	FY2011	FY2012	FY2013	FY2014	REMARKS
Virtual Information Center	Installation & stable operation	Install system —————→			Update system			Construct the integrated research environment (video and/or web conference system, etc.)
	Extension to other disciplines						—————→	Wrap up the project and discuss further extension of the system to other discipline
Metadata DB system	Development	Make prototype —————→	Develop regular system	Release product to public				Design and build the IUGONET metadata DB system on the basis of DSpace
	Stable operation				Update computers		—————→	Conduct regular operation of the metadata DB and customize it as needed
Metadata	Design of metadata format	Release ver.1 format —————→	Prepare documents —————→	Update format as needed ----->				Formulate the IUGONET common metadata format and keep updating it if necessary
	Creation of metadata			Open metadata thru metadata DB	Target relatively old, undatabased items		—————→	Create metadata in the designated format and register them in the metadata DB system
Analysis Software	Survey & Specification of analysis software	Specification —————→	Prepare documents —————→					Design an integrated analysis software to download, visualize, and analyze data provided from the IUGONET institutions
	Programming			Release product to public	Target relatively old, undatabased items		—————→	Develop the IUGONET analysis software by using TDAS (a set of IDL subroutines)
Others	Rearrangement of observational DBs		Rearrange DBs corresponding to metadata & software development				—————→	Rearrange existing observational DBs and newly compile DBs of undatabased items
	Scientific researches			Conduct scientific researches with the IUGONET products				Do interdisciplinary researches using various data from the IUGONET institutions
	Management of project website	Build project homepage —————→					—————→	Provide project information to the public through the website

Today

The IUGONET metadata database and analysis software have just been **beta-released** !

<http://search.iugonet.org/iugonet>



The screenshot shows the IUGONET Metadata DB search interface. It includes a navigation menu on the left with links to Home, IUGONET MDB Search Help, Browse Data, Browse Service, and UDAS. The main search area features a search bar with a "Free Word" input field, a "Time" range selector, and a "Spatial Coverage/Map" section with latitude and longitude coordinates. A map of Japan is displayed below the spatial coverage section. The interface also includes a "Data Types" section with checkboxes for Numerical, Plot / Movie, Data File / Plot, Instrument, and Observatory. A "Search" button is located at the bottom of the search area.

<http://www.iugonet.org/en/software.htm>



The screenshot shows the IUGONET Data Analysis Software (UDAS) page. It includes a "Topics" section with a link to "UDAS v1.00.b1" and a "What is UDAS?" section. The "What is UDAS?" section describes the software as a plug-in for THEMIS Data Analysis Software Suite (TDAS) and lists its features, including data access, export, and GUI. A "Getting started" section with links to "View screenshots" and "List of load procedures and corresponding IUGONET observations" is also present. A "Data Policy" section and a "Collaborations" section are at the bottom. A "Download UDAS" button is located on the right side of the page.

➤ We welcome your feedback

- The IUGONET project (<http://www.iugonet.org>) builds **metadata database** and **analysis software** to promote effective use of upper atmospheric data taken by various ground-based observations.
- The IUGONET products have been beta-released!
 - Metadata database : <http://search.iugonet.org/iugonet/>
 - Analysis software : <http://www.iugonet.org/en/software.html>
- The IUGONET project plans to expand this system to other types of data (satellite & simulation) and to other countries. We also would like to collaborate with other disciplines to build more comprehensive system.