

1 **Odd-numbered perfluorocarboxylates predominate over perfluorooctanoic acid in**
2 **serum samples from Japan, Korea and Vietnam**

3

4 Kouji H. Harada^a, Toshiaki Hitomi^a, Tamon Niisoe^a, Katsunobu Takanaka^b, Sigetosi
5 Kamiyama^c, Takao Watanabe^d, Chan-seok Moon^e, Hye-ran Yang^a, Nguyen Ngoc Hung^f,
6 Akio Koizumi^{a,*}

7 ^a Department of Health and Environmental Sciences, Kyoto University Graduate School
8 of Medicine, Yoshida, Kyoto 606-8501, Japan

9 ^b Department of Neurosurgery, Takayama Red Cross Hospital, Takayama 506-8550,
10 Japan

11 ^c Morinomiya Industrial Health Association, Miyagino, Sendai 983-0031, Japan

12 ^d Tohoku Bunkyo College, Katayachi, Yamagata 990-2316, Japan

13 ^e Department of Industrial Health, Catholic University of Pusan, Busan 609757, Korea

14 ^f Department of Science and Technology, Hanoi Medical University, Ton That Tung,
15 Hanoi, S.R. Vietnam

16 *Correspondence to: Akio Koizumi M.D., Ph.D.

17 Department of Health and Environmental Sciences, Kyoto University Graduate School
18 of Medicine, Yoshida Konoe, Sakyo, Kyoto 606-8501, Japan

19 Tel: +81-75-753-4456; Fax: +81-75-753-4458

20 E-mail: koizumi.akio.5v@kyoto-u.ac.jp

21

22 **Abstract**

23 Perfluorooctanoic acid (PFOA) has recently attracted attention as a potential health risk
24 following environmental contamination. However, information detailing exposure to
25 perfluorinated carboxylic acids (PFCAs) other than PFOA is limited. We measured the
26 concentrations of PFCAs (from perfluorohexanoic acid to perfluorotetradecanoic acid)
27 in serum samples obtained from patients in Japan (Sendai, Takayama, Kyoto and Osaka)
28 between 2002 and 2009, Korea (Busan and Seoul) between 1994 and 2008 and Vietnam
29 (Hanoi) in 2007/2008. Total PFCAs levels (geometric mean) were increased from 8.9 ng
30 mL⁻¹ to 10.3 ng mL⁻¹ in Japan; from 7.0 ng mL⁻¹ to 9.2 ng mL⁻¹ in Korea; and were
31 estimated at 4.7 ng mL⁻¹ in Vietnam. PFCAs of greater length than PFOA were
32 significantly increased in Sendai, Takayama and Kyoto, Japan, and levels of long-chain
33 PFCAs exceeded PFOA levels in serum. Among these PFCAs, perfluoroundecanoic
34 acid (PFUnDA) was the predominant component (28.5%), followed by
35 perfluorononanoic acid (PFNA 17.5%), perfluorodecanoic acid (PFDA 7.9%),
36 perfluorotridecanoic acid (PFTrDA 6.1%) and perfluorododecanoic acid (PFDoDA
37 1.8%). Odd-numbered PFCAs (PFNA, PFUnDA and PFTrDA) were also observed in
38 Korea and Vietnam and their presence increased significantly in Korea between 1994
39 and 2007/2008. The proportion of long-chain PFCAs in serum was relatively high
40 compared to reports in Western countries. Further investigations into the sources and
41 exposure routes are needed to predict the future trajectory of these serum PFCA levels.

42 Key words: perfluorocarboxylate; perfluorooctanoic acid; serum; temporal trend;
43 East Asia

44 **Abbreviations**

45 PFCA: perfluorinated carboxylic acids

46 PFOS: perfluorooctane sulfonate

47 PFOA: perfluorooctanoic acid

48 PFHxA: perfluorohexanoic acid

49 PFHpA: perfluoroheptanoic acid

50 PFNA: perfluorononanoic acid

51 PFDA: perfluorodecanoic acid

52 PFUnDA: perfluoroundecanoic acid

53 PFDoDA: perfluorododecanoic acid

54 PFTrDA: perfluorotridecanoic acid

55 PFTeDA: perfluorotetradecanoic acid

56 IDLs: instrumental detection limits

57 MDLs: method detection limits

58 RSD: relative standard deviation

59 SD: standard deviation

60 GM: geometric mean

61 GSD: geometric standard deviation

62

63 1. Introduction

64 Perfluorinated compounds such as perfluorooctane sulfonate (PFOS) and
65 perfluorooctanoic acid (PFOA) have recently attracted attention owing to widespread
66 contamination of the environment, wildlife and humans by these chemicals (Houde et
67 al., 2006). In 2002, after 50 years of production, 3M Company phased out their
68 manufacture of PFOS (Renner, 2001). PFOA is considered to be a major component of
69 perfluorocarboxylates (PFCAs) emission. However, in Japan, PFCA emissions consisted
70 of not only PFOA but also perfluorononanoic acid (PFNA) and perfluoroundecanoic
71 acid (PFUnDA) (of which 25 and 7 metric tons, respectively, were emitted in 2000)
72 (Prevedouros et al., 2006). These odd-numbered PFCAs (PFNA, PFUnDA and
73 perfluorotridecanoic acid (PFTrDA)) were detected at higher concentrations in samples
74 from local wildlife than similar even-numbered PFCAs (PFOA, perfluorodecanoic acid
75 (PFDA) and perfluorododecanoic acid (PFDoDA), respectively) (Furdui et al., 2008).
76 Although studies using human samples from Western countries showed that PFOA was
77 the most prevalent (followed by PFNA, PFDA and PFUnDA) (Haug et al., 2009;
78 Joensen et al., 2009; Kato et al., 2009), our previous study of Japanese women in the
79 Miyagi prefecture showed that PFNA and PFUnDA (average: 2.8 and 5.4 ng mL⁻¹,
80 respectively) were found at broadly similar serum concentrations to PFOA (average: 4.9
81 ng mL⁻¹) (Kärman et al., 2009).

82 PFCAs with longer chains than PFOA have higher bio-concentration factors
83 suggesting persistency in the environment (Martin et al., 2003). Temporal trends in
84 serum levels after 2002 showed no apparent decline of PFNA, PFDA or PFUnDA in
85 Norway (Haug et al., 2009), although serum levels of PFOA and PFOS both decreased
86 in the United States, Norway and Japan (Harada and Koizumi, 2009; Harada et al.,

87 2010; Haug et al., 2009; Olsen et al., 2008). These findings suggest a possibility that the
88 origin and source of exposure to long-chain PFCAs could differ from those of PFOA
89 and PFOS.

90 In the present study, we investigated current serum concentrations of PFCAs in three
91 Asian countries (Japan, Korea and Vietnam). We selected the cities of Busan and Seoul
92 in Korea because they are comparably urban and industrialized to Osaka, Japan. To
93 confirm the temporal trends in Japan and Korea, we used archived historical serum
94 samples stored in the human specimen bank (Koizumi et al., 2009; Koizumi et al.,
95 2005). Hanoi in Vietnam was selected to evaluate the development of PFCA
96 contamination following recent industrialization.

97

98 **2. Material and methods**

99 *2.1. Experimental design and study population*

100 To evaluate geographical differences and temporal trends in Asian countries, we
101 compared 521 samples collected from Japan (Sendai, Takayama, Kyoto and Osaka)
102 between 2002 and 2009; Korea (Busan and Seoul) between 1994 and 2008; and in
103 Hanoi, Vietnam between 2007 and 2008. Samples from Sendai and Takayama in 2008,
104 Osaka, Busan, Seoul and Hanoi are identical to a previous analysis of PFOS and PFOA
105 (Harada et al., 2010; Kärroman et al., 2009). A total of 521 serum samples with
106 information on donor age, sex and residential history (>5 yr in each area) were selected
107 from the archived samples in Kyoto Human Specimen Bank (Koizumi et al., 2009;
108 Koizumi et al., 2005) (Table 1). Serum was separated from cellular components and
109 stored at -30 °C until analysis.

110 The study population in Osaka and Kyoto consisted of residents that had been

111 intensely exposed to PFOA from a local industrial source (the fluoropolymer
112 manufacturer, Daikin Company) (Harada et al., 2004, 2007, 2010; Kärman et al., 2009;
113 Niisoe et al., 2010). In contrast, there is no known potential industrial source of PFCAs
114 that would affect sample populations in the other cities studied.

115 For historical comparisons, samples were selected so that age and gender were
116 matched among time points, except for Busan in 2000 and Osaka (Table 1).

117 The research protocol for the present study was reviewed and approved by the
118 Ethics Committee of the Kyoto University Graduate School of Medicine on 14
119 November 2003 (E25).

120

121 *2.2. Reagents*

122 Ammonium acetate (purity: >99% by HPLC) was purchased from Aldrich
123 (Steinheim, Germany). Acetonitrile (LC-MS grade) and water (distilled LC-MS grade)
124 were obtained from Kanto Chemicals (Tokyo, Japan). Acetic acid and benzyl bromide
125 were purchased from Wako pure chemicals (Osaka, Japan). Mixture of native PFCAs,
126 ¹³C₄-labeled PFOA and ¹³C₅-labeled PFNA were obtained from Wellington Laboratories
127 (Guelph, Ontario, Canada).

128

129 *2.3. Determination of PFCAs in serum*

130 Perfluorohexanoic acid (PFHxA), perfluoroheptanoic acid (PFHpA), PFOA, PFNA,
131 PFDA, PFUnDA, PFDoDA, PFTrDA and perfluorotetradecanoic acid (PFTeDA) were
132 analyzed. Serum samples were subjected to a clean-up procedure using a dispersive
133 carbon method described by Powley et al. (2005). Briefly, the serum samples (0.5 mL,
134 except for Korean samples between 1994 and 2000, which were 0.25 mL) together with

135 an internal standard ($^{13}\text{C}_4\text{-PFOA}$, 1 ng) were extracted with 5 mL of acetonitrile,
136 followed by centrifugation at $1600 \times g$ for 15 min. The supernatants were transferred
137 into new tubes with 25 mg of ENVI-Carb and 50 μL of acetic acid, and the solutions
138 were mixed by vortexing for 30 s. After centrifugation at $1600 \times g$ for 15 min, the
139 extracts were dried under a nitrogen stream. The residue was then re-dissolved in 100 μL
140 of 100 mM benzyl bromide acetone containing the recovery performance standard
141 $^{13}\text{C}_5\text{-PFNA}$ (1 ng) for 1 hour at 80 $^\circ\text{C}$ and transferred to an autosampler vial. Extracts
142 were analyzed using gas chromatography-mass spectrometry (Agilent
143 6890GC/5973MSD, Agilent Technologies Japan, Ltd., Tokyo, Japan) in electron impact
144 ionization mode using single ion monitoring. PFCA benzyl esters were separated on a
145 DB-5MS column (30 m length, 0.25 mm i.d., 1 μm film thickness) with a helium carrier
146 gas. Split-less injections (1 μL) were performed with the injector set at 220 $^\circ\text{C}$, and the
147 split was opened after 1.5 min. The initial oven temperature was 70 $^\circ\text{C}$ for 2 min,
148 ramped at 20 $^\circ\text{C min}^{-1}$ to 100 $^\circ\text{C}$, and then at 30 $^\circ\text{C min}^{-1}$ to 280 $^\circ\text{C}$. Ion fragments
149 ($[\text{M}]^+$) were monitored and used as quantification ions (Table 2).

150 Instrumental detection limits (IDL) were defined as the mass of analyte producing a
151 peak with a signal-to-noise ratio of 3, and ranged from 1 pg (PFTeDA) to 0.25 pg (other
152 PFCAs) (Table 2). Since blank samples (0.5 mL distilled water) contain no detectable
153 concentrations, the method detection limits (MDL) value was considered to be equal to
154 the IDL corresponding to 0.2 ng mL^{-1} for PFTeDA and 0.025 ng mL^{-1} for other PFCAs
155 (Table 2).

156

157 2.4. Quality assurance

158 Quantification was performed using an internal standard method with the external

159 standards dissolved in 10% methanol in water. $^{13}\text{C}_4$ -PFOA was used as the internal
160 standard for PFCAs. $^{13}\text{C}_5$ -PFNA was used to calculate a recovery rate of $^{13}\text{C}_4$ -PFOA.
161 All samples were quantified using a seven-point calibration curve with a relative
162 standard deviation (RSD) of the relative response factors <15% for all compounds. The
163 recoveries were evaluated by five replicate fortifications (fortified by 10 times the
164 original concentration of serum) of a human serum sample with low contamination
165 (Table 2). The procedural blank levels were evaluated in duplicate for 11 samples each
166 using 0.5 mL distilled water.

167 Using the above method, we reanalyzed 361 samples originally tested in a previous
168 study by HPLC-MS/MS (Harada et al., 2010; Kärrman et al., 2009). The reanalyzed
169 samples showed $5.14\pm 11.60\text{ ng mL}^{-1}$ for PFOA, which equates to 101.7% of the levels
170 obtained in the previous study ($5.05\pm 11.16\text{ ng mL}^{-1}$, $p=0.478$ by paired t -test). Pearson's
171 correlation coefficient, r and slope were 0.9882 and 1.128, respectively ($p<0.0001$).
172 Levels (mean \pm SD) of PFHpA, PFNA, PFDA and PFUnDA in Osaka in 2004 were also
173 confirmed in this study (HPLC-MS/MS vs GC-MS: $0.26\pm 0.14\text{ ng mL}^{-1}$ vs $0.24\pm 0.09\text{ ng}$
174 mL^{-1} , $6.68\pm 1.78\text{ ng mL}^{-1}$ vs $6.16\pm 1.91\text{ ng mL}^{-1}$, $2.55\pm 0.99\text{ ng mL}^{-1}$ vs $2.74\pm 1.32\text{ ng}$
175 mL^{-1} , $5.80\pm 2.13\text{ ng mL}^{-1}$ vs $5.12\pm 2.69\text{ ng mL}^{-1}$, respectively; $p>0.05$ by paired t -test).
176 RSDs of difference between methods were 33.1%, 9.8%, 13.6% and 11.5% for PFHpA,
177 PFNA, PFDA and PFUnDA, respectively and average RSD was 17.0%.

178 To assess potential interlaboratory difference in analysis, NIST standard reference
179 material (SRM) 1957 was analyzed (Table 2). The values from PFHpA to PFUnDA
180 were comparable to those from interlaboratory comparison exercises (Lindstrom et al.,
181 2009; Keller et al., 2010).

182 Mean recovery rate (RSD) of $^{13}\text{C}_4$ -PFOA in 521 samples was 96.5% (8.8%). To

183 evaluate possible matrix effect in serum sample, we further analyzed 100 samples
184 extracts fortified with 1 ng of PFHpA, PFNA, PFDA, PFUnDA, PFDoDA and PFTrDA
185 standards. Recoveries of fortified standards were 98.7%, 104.6%, 102.0%, 97.2%,
186 102.2% and 96.3% for PFHpA, PFNA, PFDA, PFUnDA, PFDoDA and PFTrDA,
187 respectively. It is therefore considered that there was no substantial suppression or
188 enhancement of target ions, if any.

189

190

191 2.5. *Statistical analysis*

192 All statistical analyses were carried out using the JMP software (Version 4; SAS
193 Institute Inc., Cary, NC). Values of $p < 0.05$ were considered to indicate statistical
194 significance. Concentrations of less than the detection limit were all approximated to
195 'half of detection limit' for statistical analyses. Serum levels of PFCAs were assumed to
196 distribute lognormally because the serum levels of PFCAs in the samples displayed
197 right-skewed patterns and geometric means were comparable to medians. Statistical
198 analyses were conducted after logarithmic transformation of the serum concentrations.
199 Differences between mean values were tested by Tukey-Kramer's honestly significant
200 difference (HSD) test after ANOVA. Correlation was tested by Spearman's rank
201 correlation coefficient (ρ). Factor analysis was used to transform a number of
202 contaminants into a smaller number of potential factors of sources. Factor analysis was
203 conducted *via* correlation matrix. In essence, the factor analysis is a model which
204 presumes the existence of a smaller set of factors that can reproduce exactly the
205 correlation in the larger set of variable. To achieve this goal, the linear combinations of
206 factors (i.e., principal component) will be generated in such a manner that each

207 composite variate will account a smaller portion of the total variation i.e., variance.
208 Eigenvalues of a principal component is a measure how much this principal component
209 can account for the variation and eigenvector indicates an associated set of coefficients
210 with a principal component for each factor. Eigenvectors were employed through the
211 analysis when eigenvalues were close to or greater than 1 which means its eigenvector
212 can account the equivalent of one or more original variables. Normalized varimax
213 rotation (an orthogonal rotation of the factor axes) was applied to these eigenvectors to
214 simplify them into a few variables with high correlations.

215

216 **3. Results**

217 *3.1. Temporal changes in PFCA concentrations in Japan*

218 The descriptive statistics for PFCAs are presented in Table 3. Most samples
219 contained PFOA, PFNA, PFDA, PFUnDA and PFTrDA at both time points. No samples
220 contained PFHxA and PFTeDA at concentrations above MDL. PFHpA levels were
221 significantly decreased in all sampling sites in Japan between 2002/2004 and 2008/2009
222 ($p < 0.05$ by Student's *t*-test). PFOA was relatively high in Osaka and Kyoto although
223 levels of this compound nevertheless significantly decreased in this period ($p < 0.05$ by
224 Student's *t*-test). In Sendai and Takayama, PFOA levels also decreased but this
225 difference was not statistically significant. In contrast, PFCAs longer than PFOA
226 showed significant increases in Sendai, Takayama and Kyoto with few exceptions.
227 Among these PFCAs, PFUnDA was the predominant component, followed by PFNA,
228 PFDA, PFTrDA and PFDoDA. These odd-numbered PFCAs (i.e. PFUnDA, PFNA and
229 PFTrDA) were detected at higher concentrations than neighboring, even-numbered
230 PFCAs (PFDA and PFDoDA).

231 In Osaka, levels of PFNA, PFDA and PFUnDA, as with PFOA, significantly
232 decreased from 2004 to 2008. PFDoDA and PFTrDA levels did not change. Among four
233 sampling sites in 2008/2009, Osaka and Kyoto had higher PFOA, PFNA and PFDA
234 levels than Sendai and Takayama ($p < 0.05$ by Tukey's HSD test) but PFUnDA, PFDoDA
235 and PFTrDA showed no regional differences ($p > 0.05$ by ANOVA).
236 As a consequence of the increase in long-chain PFCAs, the proportion of PFOA in the
237 total PFCA content became less than 50% in all locations except Osaka.

238

239 *3.2. Temporal trends in the serum concentrations of PFCAs in Korea*

240 The PFCA concentrations in the serum samples collected in Busan and Seoul
241 between 1994 and 2008 are shown in Table 4. As is the case with Japan, PFOA, PFNA,
242 PFDA, PFUnDA and PFTrDA were frequently detected in 2007/2008. PFHxA and
243 PFTeDA were not detected in any samples at concentrations above MDL. In agreement
244 with the previous report by Harada et al. (2010), PFOA levels were stable from 1994 to
245 2008 in Busan and Seoul ($p > 0.05$ by ANOVA). In contrast, odd-numbered PFCAs
246 (PFNA, PFUnDA and PFTrDA) were significantly increased during this period ($p < 0.05$
247 by Tukey's HSD test or Student's *t*-test). The PFCA levels had the following order of
248 prevalence in 1994: PFOA > PFNA ~ PFUnDA > PFDA > PFTrDA > PFHpA ~ PFDoDA.
249 However, by 2007/2008 the order had changed to:
250 PFOA > PFUnDA > PFNA > PFDA ~ PFTrDA > PFDoDA > PFHpA. Between 1994 and
251 2007/2008, total PFCA levels were significantly increased by 1.31- and 1.53-fold in
252 Busan and Seoul, respectively ($p < 0.05$ by Tukey's HSD test or Student's *t*-test).
253 Samples from Busan contained higher concentrations of PFHpA, PFOA, PFNA, PFDA,
254 PFUnDA and PFTrDA than did those from Seoul in both 1994 and 2007/2008 ($p < 0.05$

255 by Student's *t*-test).

256

257 3.3. PFCAs concentrations in Hanoi, Vietnam in 2008-2009

258 PFOA, PFNA, PFDA and PFUnDA were detected in all samples, and PFDoDA and
259 PFTrDA were also detected, albeit less frequently (Table 5). PFHxA, PFHpA and
260 PFTeDA were not detected in any samples from Hanoi. The concentration of PFUnDA
261 was highest among the PFCAs studied, followed by PFNA, PFDA, PFOA, PFTrDA and
262 PFDoDA. The proportion of PFOA relative to total PFCAs was only 12.9%.

263

264 3.4. Correlations among PFCA levels and factor analysis

265 Correlation coefficients among PFCAs in 521 samples are listed in Table 6. PFHpA
266 was relatively less correlated with other PFCAs, except for PFOA ($\rho=0.398$). PFOA also
267 significantly correlated with PFNA and PFDA (ρ coefficient >0.5) but was less well
268 correlated with PFUnDA, PFDoDA and PFTrDA. In general, PFCA concentrations
269 indicated a strong correlation between PFCAs of similar (i.e. adjacent) chain length.

270 To delineate potential patterns in the data, PFCA concentrations were examined
271 using factor analysis. The contributions of factors 1 and 2 to the total variance were
272 49.72% and 19.40% (with an eigenvalue >1), respectively (Table 7). After varimax
273 rotation, the first factor indicated a higher eigenvector for longer-chain PFCAs than
274 PFNA. The second factor had a more positive eigenvector for shorter-chain PFCAs than
275 PFDA. Since there is a point source of PFCAs in both Osaka and Kyoto, we evaluated
276 whether this predominant source may perturb the results of the factor analysis.
277 Eliminating Osaka and Kyoto samples, however, did not alter a correlation matrix
278 among PFCAs with changes in eigenvalues being less than 5% (data not shown),

279 indicating that the dominant point source had no substantial influence on the
280 interpretation of factor 1 and factor 2.

281 Factor 1 is characterized by PFUnDA dominance (factor loading: 0.858) and another
282 by PFOA dominance (0.819), respectively. This characteristic pattern indicates
283 fingerprints of PFCAs sources in Asia. Temporal transition of factor scores is
284 demonstrated by score plots shown in Supplemental Fig. 1. In sampling sites in Japan
285 and Korea (except for Osaka), centers of score plot moved rightwards and downwards,
286 indicating that the factor 1 score increased and factor 2 score decreased during these
287 periods. Mean factor scores of each sampling site are also shown in Table 7. In Japan,
288 factor 1 scores significantly increased from 2002/2003 to 2008/2009 ($p < 0.05$ by
289 Student's *t*-test), except for Osaka which already had a high factor 1 score (0.92) in
290 2004. This increase in factor 1 scores was also observed in Busan and Seoul from 1994
291 to 2007/2008 ($p < 0.05$ by Tukey's HSD test or Student's *t*-test). Although the factor 1
292 score in Hanoi was lower than those in other sites in 2007–2009, it surpassed scores in
293 Sendai and Kyoto in 2002/2003 and in Busan and Seoul in 1994. Contrary to factor 1,
294 factor 2 scores in all sampling sites in Japan significantly declined between 2002/2004
295 and 2008/2009 ($p < 0.05$ by Student's *t*-test) and also in Busan and Seoul from 1994 to
296 2007/2008 ($p < 0.05$ by Tukey's HSD test or Student's *t*-test). Factor 2 in Hanoi was the
297 lowest among all sampling sites.

298

299 **4. Discussion**

300 In the present study, we uncovered two major fingerprints (factor 1 and factor 2) by
301 analyzing serum samples from three countries in East Asia. Characteristic PFCA
302 composition was observed for odd-numbered PFCAs such as PFUnDA and PFTrDA

303 with residual PFDoDA and PFDA, which can correspond to factor 1. Even in
304 populations exposed to low levels of PFOA, notably Hanoi, PFUnDA showed
305 substantial serum levels. Moreover, levels of those PFCAs with longer chain lengths
306 than PFOA were significantly elevated in Japan and Korea in recent years. In the
307 late-2000s, consequently, long-chain PFCA levels exceeded PFOA levels in most
308 sampling sites. This finding suggests an emergence of specific sources of exposure in
309 East Asia.

310 In several countries, serum PFOA has reportedly decreased (Harada et al., 2010;
311 Olsen et al., 2008). In contrast, PFCAs of longer chain lengths than PFOA were
312 frequently detected in serum samples in this study. Total levels of long-chain PFCAs
313 were comparable to or greater than PFOA levels (except in Osaka) and showed trends
314 towards increases in Japan and Korea. Correlation between PFOA and long-chain
315 PFCAs was not strong which suggests that the sources of long-chain PFCA
316 contamination have different exposure route than PFOA. Indeed, factor analysis
317 demonstrated two major factors as sources of PFCAs. The first factor had loading on
318 longer-chain PFCAs than PFOA and the second factor on PFHpA, PFOA and PFNA.
319 Temporal trends of these factors were opposite and contamination derived from factor 1
320 might be expected to emerge in around a decade. This transition of factor scores was
321 similar in Japan, Korea and Hanoi. Contamination derived from factor 1 may have been
322 prevailing in East Asian countries.

323 Among long-chain PFCAs, odd-numbered PFCAs accounted for the major
324 proportion. Serum or blood levels of PFCAs reported from populations in China, Sri
325 Lanka, Australia, Norway, Sweden, Denmark, Poland, Belgium, Spain and USA are
326 summarized in Table 8 (Ericson et al., 2007; Falandysz et al., 2006; Guruge et al., 2005;

327 Haug et al., 2009; Joensen et al., 2009; Kärrman et al., 2006; Kuklennyik et al., 2004;
328 Pan et al., 2010; Roosens et al., 2010; Toms et al., 2009). The PFCA composition in our
329 current study, which was characterized by a large proportion of PFUnA, was apparently
330 different from Western countries (Table 8). Although PFOA levels in these countries
331 were comparable, long-chain PFCAs were not major components in Western countries,
332 except for Antwerp, Belgium and Atlanta, USA. Therefore, this composition can be
333 considered as a clear fingerprint for East Asian countries and is implicated in the
334 origination of factor 1.

335 However, their source remains unclear due to insufficient monitoring data of PFCAs.
336 Interestingly, a review by Prevedouros et al. (2006) indicated that PFNA has been
337 manufactured in Japan *via* oxidation of fluorotelomer olefins together with PFUnDA
338 and PFTrDA. Industrial application of these odd-numbered PFCAs, namely Surflon
339 S-111, might contribute to the East Asian-specific pattern of serum body burdens. The
340 temporal increase in long-chain PFCAs warrants further investigations of the sources
341 and exposure routes to assist in predicting future changes in the serum levels of these
342 contaminants.

343 In this study, there was a limitation in chemical analysis. $^{13}\text{C}_4$ -PFOA was used for
344 internal standard for PFCAs (C_7 - C_{14}). Chemical properties of PFCAs may, however, be
345 different even though they have similar structures. Matrix effects also might affect
346 quantification of PFCAs other than PFOA. Thus it is logically possible that recovery
347 rates of $^{13}\text{C}_4$ -PFOA might be extensively deviated from those of other PFCAs.
348 Nevertheless, such a possibility is unlikely because recovery rates of PFCAs were higher
349 than 90% and RSD were within 10%, indicating that there was no substantial difference
350 in recoveries among PFCAs in this method. Furthermore, a good agreement of results

351 in SRM analysis by interlaboratory comparisons assured that our analytical method in
352 this study is sound. Collectively, these findings consistently support that analytical
353 method in this study was sufficiently qualified.

354 Recent epidemiological investigations have raised concern regarding developmental
355 effects of PFOA on children (Steenland et al., 2010). In contrast, few studies have been
356 conducted on the effects of PFCAs of different chain length. Even though PFCAs have
357 similar structure, their chemical properties and biological activity are likely different. In
358 several *in vitro* studies, long-chain PFCAs caused biological responses at lower doses
359 than PFOA (Liao et al., 2009; Matsubara et al., 2006; Upham et al., 1998). The
360 toxicokinetics of long-chain PFCAs are also unclear, especially in humans. These
361 uncertainties necessitate more comprehensive toxicological studies on PFCAs.

362

363 **Acknowledgments**

364 We are grateful to Dr. Anna Kärrman (Örebro University, Sweden) for valuable
365 instructions for quality assurance. This study was mainly supported by Special
366 Coordination Funds for Promoting Science and Technology (No. 1300001) sponsored
367 by the Japan Science and Technology Agency, and a Grant-in-Aid for Health Sciences
368 Research from the Ministry of Health, Labour and Welfare of Japan (H21-Food-003).
369 The authors have no financial conflicts of interest.

370

371 **References**

- 372 Ericson, I., Gomez, M., Nadal, M., van Bavel, B., Lindstrom, G., Domingo, J.L., 2007.
373 Perfluorinated chemicals in blood of residents in Catalonia (Spain) in relation to
374 age and gender: A pilot study. *Environ Int.*
- 375 Falandysz, J., Taniyasu, S., Gulkowska, A., Yamashita, N., Schulte-Oehlmann, U., 2006.
376 Is fish a major source of fluorinated surfactants and repellents in humans living
377 on the Baltic Coast? *Environ Sci Technol* 40, 748-751.
- 378 Furdui, V.I., Helm, P.A., Crozier, P.W., Lucaciu, C., Reiner, E.J., Marvin, C.H., Whittle,
379 M.D., Mabury, S.A., Tomy, G., 2008. Temporal trends of perfluoroalkyl
380 compounds with isomer analysis in lake trout from Lake Ontario (1974-2004).
381 *Environ. Sci. Technol.* 42, 4739-4744.
- 382 Guruge, K.S., Taniyasu, S., Yamashita, N., Wijeratna, S., Mohotti, K.M., Seneviratne,
383 H.R., Kannan, K., Yamanaka, N., Miyazaki, S., 2005. Perfluorinated organic
384 compounds in human blood serum and seminal plasma: a study of urban and
385 rural tea worker populations in Sri Lanka. *J Environ Monit* 7, 371-377.
- 386 Harada, K., Koizumi, A., Saito, N., Inoue, K., Yoshinaga, T., Date, C., Fujii, S., Hachiya,
387 N., Hirose, I., Koda, S., Kusaka, Y., Murata, K., Omae, K., Shimbo, S.,
388 Takenaka, K., Takeshita, T., Todoriki, H., Wada, Y., Watanabe, T., Ikeda, M.,
389 2007. Historical and geographical aspects of the increasing perfluorooctanoate
390 and perfluorooctane sulfonate contamination in human serum in Japan.
391 *Chemosphere* 66, 293-301.
- 392 Harada, K., Saito, N., Inoue, K., Yoshinaga, T., Watanabe, T., Sasaki, S., Kamiyama, S.,
393 Koizumi, A., 2004. The influence of time, sex and geographic factors on levels
394 of perfluorooctane sulfonate and perfluorooctanoate in human serum over the

395 last 25 years. *J Occup Health* 46, 141-147.

396 Harada, K.H., Koizumi, A., 2009. Environmental and biological monitoring of
397 persistent fluorinated compounds in Japan and their toxicities. *Environ Health*
398 *Prev Med* 14, 7-19.

399 Harada, K.H., Yang, H.R., Moon, C.S., Hung, N.N., Hitomi, T., Inoue, K., Niisoe, T.,
400 Watanabe, T., Kamiyama, S., Takenaka, K., Kim, M.Y., Watanabe, K., Takasuga,
401 T., Koizumi, A., 2010. Levels of perfluorooctane sulfonate and
402 perfluorooctanoic acid in female serum samples from Japan in 2008, Korea in
403 1994-2008 and Vietnam in 2007-2008. *Chemosphere* 79, 314-319.

404 Haug, L.S., Thomsen, C., Becher, G., 2009. Time trends and the influence of age and
405 gender on serum concentrations of perfluorinated compounds in archived human
406 samples. *Environ. Sci. Technol.* 43, 2131-2136.

407 Houde, M., Martin, J.W., Letcher, R.J., Solomon, K.R., Muir, D.C., 2006. Biological
408 monitoring of polyfluoroalkyl substances: A review. *Environ Sci Technol* 40,
409 3463-3473.

410 Joensen, U.N., Bossi, R., Leffers, H., Jensen, A.A., Skakkebaek, N.E., Jorgensen, N.,
411 2009. Do perfluoroalkyl compounds impair human semen quality? *Environ*
412 *Health Perspect* 117, 923-927.

413 Kärman, A., Harada, K.H., Inoue, K., Takasuga, T., Ohi, E., Koizumi, A., 2009.
414 Relationship between dietary exposure and serum perfluorochemical (PFC)
415 levels--a case study. *Environ Int* 35, 712-717.

416 Kärman, A., van Bavel, B., Jarnberg, U., Hardell, L., Lindstrom, G., 2006.
417 Perfluorinated chemicals in relation to other persistent organic pollutants in
418 human blood. *Chemosphere* 64, 1582-1591.

419 Kato, K., Calafat, A.M., Wong, L.Y., Wanigatunga, A.M., Caudill, S.P., Needham, L.L.,
420 2009. Polyfluoroalkyl Compounds in Pooled Sera from Children Participating in
421 the National Health and Nutrition Examination Survey 2001-2002. *Environ. Sci.*
422 *Technol.* 43, 2641-2647.

423 Koizumi, A., Harada, K., Inoue, K., Hitomi, T., Yang, H.-R., Moon, C.-S., Wang, P.,
424 Hung, N., Watanabe, T., Shimbo, S., Ikeda, M., 2009. Past, present, and future of
425 environmental specimen banks. *Environmental Health and Preventive Medicine*
426 14, 307-318.

427 Koizumi, A., Yoshinaga, T., Harada, K., Inoue, K., Morikawa, A., Muroi, J., Inoue, S.,
428 Eslami, B., Fujii, S., Fujimine, Y., Hachiya, N., Koda, S., Kusaka, Y., Murata, K.,
429 Nakatsuka, H., Omae, K., Saito, N., Shimbo, S., Takenaka, K., Takeshita, T.,
430 Todoriki, H., Wada, Y., Watanabe, T., Ikeda, M., 2005. Assessment of human
431 exposure to polychlorinated biphenyls and polybrominated diphenyl ethers in
432 Japan using archived samples from the early 1980s and mid-1990s. *Environ Res*
433 99, 31-39.

434 Kuklennyik, Z., Reich, J.A., Tully, J.S., Needham, L.L., Calafat, A.M., 2004. Automated
435 solid-phase extraction and measurement of perfluorinated organic acids and
436 amides in human serum and milk. *Environ Sci Technol* 38, 3698-3704.

437 Liao, C., Wang, T., Cui, L., Zhou, Q., Duan, S., Jiang, G., 2009. Changes in Synaptic
438 Transmission, Calcium Current, and Neurite Growth by Perfluorinated
439 Compounds Are Dependent on the Chain Length and Functional Group.
440 *Environmental Science & Technology* 43, 2099-2104.

441 Martin, J.W., Mabury, S.A., Solomon, K.R., Muir, D.C.G., 2003. Bioconcentration and
442 tissue distribution of perfluorinated acids in rainbow trout (*Oncorhynchus*

443 mykiss). *Environmental Toxicology and Chemistry* 22, 196-204.

444 Matsubara, E., Harada, K., Inoue, K., Koizumi, A., 2006. Effects of perfluorinated
445 amphiphiles on backward swimming in *Paramecium caudatum*. *Biochem*
446 *Biophys Res Commun* 339, 554-561.

447 Niisoe, T., Harada, K.H., Ishikawa, H., Koizumi, A., 2010. Long-term simulation of
448 human exposure to atmospheric perfluorooctanoic acid (PFOA) and
449 perfluorooctanoate (PFO) in the Osaka urban area, Japan. *Environ Sci Technol*
450 44, 7852-7857.

451 Olsen, G.W., Mair, D.C., Church, T.R., Ellefson, M.E., Reagen, W.K., Boyd, T.M.,
452 Herron, R.M., Medhdizadehkashi, Z., Nobiletti, J.B., Rios, J.A., Butenhoff, J.L.,
453 Zobel, L.R., 2008. Decline in perfluorooctanesulfonate and other
454 polyfluoroalkyl chemicals in American Red Cross adult blood donors,
455 2000-2006. *Environ Sci Technol* 42, 4989-4995.

456 Pan, Y., Shi, Y., Wang, J., Cai, Y., Wu, Y., 2010. Concentrations of perfluorinated
457 compounds in human blood from twelve cities in China. *Environ Toxicol Chem*
458 29, 2695-2701.

459 Powley, C.R., George, S.W., Ryan, T.W., Buck, R.C., 2005. Matrix effect-free analytical
460 methods for determination of perfluorinated carboxylic acids in environmental
461 matrixes. *Anal Chem* 77, 6353-6358.

462 Prevedouros, K., Cousins, I.T., Buck, R.C., Korzeniowski, S.H., 2006. Sources, fate and
463 transport of perfluorocarboxylates. *Environ Sci Technol* 40, 32-44.

464 Renner, R., 2001. Scotchgard scotched - Following the fabric protector's slippery trail to
465 a new class of pollutant. *Sci Am* 284, 18.

466 Roosens, L., D'Hollander, W., Bervoets, L., Reynders, H., Van Campenhout, K.,

467 Cornelis, C., Van Den Heuvel, R., Koppen, G., Covaci, A., 2010. Brominated
468 flame retardants and perfluorinated chemicals, two groups of persistent
469 contaminants in Belgian human blood and milk. *Environ Pollut* 158, 2546-2552.

470 Steenland, K., Fletcher, T., Savitz, D.A., 2010. Epidemiologic evidence on the health
471 effects of perfluorooctanoic acid (PFOA). *Environ Health Perspect* 118,
472 1100-1108.

473 Toms, L.M., Calafat, A.M., Kato, K., Thompson, J., Harden, F., Hobson, P., Sjodin, A.,
474 Mueller, J.F., 2009. Polyfluoroalkyl chemicals in pooled blood serum from
475 infants, children, and adults in Australia. *Environ Sci Technol* 43, 4194-4199.

476 Upham, B.L., Deocampo, N.D., Wurl, B., Trosko, J.E., 1998. Inhibition of gap
477 junctional intercellular communication by perfluorinated fatty acids is dependent
478 on the chain length of the fluorinated tail. *Int J Cancer* 78, 491-495.

479
480
481

482 **Figure captions**

483 **Supplemental Fig. 1.**

484 Plot of first- and second factor scores of 521 samples in Osaka (A), Kyoto (B),
485 Takayama (C), Sendai (D), Busan (E) and Seoul and Hanoi (F). Overall, 50% of the
486 values locate within the boundary circles for each sampling site and time period.

487

488

489

Table 1
Study area and study population

Sampling site	Population (x10 ³)	Latitude and longitude	Year	n (%female)	Age ^a	(range)
Japan						
Sendai	1,031	38°17'04" N 140°55'46" E	2008	50 (100)	37.5±9.44	(21-53)
	1,023	-	2003	50 (100)	36.6±10.1	(20-59)
Takayama	94 (65) ^b	36°08'13" N 137°15'16" E	2008	50 (100)	40.5±4.78	(29-49)
	67	-	2003	50 (100)	39.9±4.5	(31-45)
Kyoto	1,466	35°01'18" N 135°46'38" E	2009	30 (50)	33.2±14.7	(21-68)
	1,469	-	2002	30 (50)	35.4±11.3	(21-58)
Osaka	2,652	34°45'31" N 135°31'52" E	2008	50 (100)	45.9±8.92 ^{A*}	(30-63)
	2,619	-	2004	10 (100)	60.9±6.3 ^B	(49-69)
Korea						
Busan	3,711	35°14'39" N 129°05'54" E	2008	35 (100)	40.1±6.44 ^{A*}	(18-49)
	3,732	-	2000	30 (100)	35.4±4.27 ^B	(28-45)
	3,961	-	1994	39 (100)	42.3±4.65 ^A	(34-52)
Seoul	10,421	37°27'52" N 127°01'56" E	2007	36 (100)	34.5±8.24	(20-54)
	10,798	-	1994	24 (100)	38.0±7.41	(24-51)
Vietnam						
Hanoi	6,232	21°00'08" N 105°49'50" E	2007-2008	37 (100)	30.2±5.76	(20-40)

* Means of age with different letters differed significantly ($p < 0.05$ by Tukey's HSD test). For example, the letters A and B indicate that the corresponding values differ significantly at $p < 0.05$.

a Data are presented as mean \pm standard deviation.

b Takayama city merged with neighboring cities in 2005. Numbers in parentheses denote populations areas corresponding to those used in 2003.

Table 2

Recovery, detection limits and QA for PFCA analysis in human serum samples

Compound	Quantification (confirmation)	Recovery and (reproducibility) % (RSD%) ^a (n=5)	Instrument detection limit ^b (pg)	detection limit ^c (ng mL ⁻¹)	SRM1957 ^d (ng mL ⁻¹)
PFHxA	404 (385)	92.2 (8.41)	0.25	0.05	<0.05
PFHpA	454 (435)	94.5 (4.12)	0.25	0.05	0.27
PFOA	504 (485)	101.7 (6.99)	0.25	0.05	4.77
¹³ C ₄ PFOA	508 (489)	102.8 (5.47)	-	-	-
PFNA	554 (535)	97.4 (7.61)	0.25	0.05	0.96
¹³ C ₅ PFNA	559 (540)	-	-	-	-
PFDA	604 (585)	91.9 (8.63)	0.25	0.05	0.26
PFUnDA	654 (635)	94.1 (7.22)	0.25	0.05	0.16
PFDoDA	704 (685)	95.7 (4.87)	0.5	0.1	<0.1
PFTTrDA	754 (735)	98.6 (9.41)	0.5	0.1	<0.1
PFTeDA	785 (786)	92.4 (8.18)	1	0.2	<0.2

^a RSD: relative standard deviation^b 1 µl injection^c 0.5 mL serum sample^d 0.5 mL serum sample of NIST SRM 1957 was analyzed.

Table 3
Serum concentrations of PFCAs in Japan

Sampling site	Year	n		Concentration (ng mL ⁻¹)							ΣPFCAs
				PFHpA	PFOA	PFNA	PFDA	PFUnDA	PFDODA	PFTTrDA	
Sendai	2008	50	GM(GSD)	0.06(2.17)*	2.44(1.56)	1.80(1.40)*	0.72(1.46)*	3.00(1.59)*	0.17(1.99)*	0.60(2.00)*	9.13(1.41)*
			range	<0.05-0.37	0.85-6.05	0.90-3.58	0.31-1.58	1.15-8.08	<0.1-0.52	<0.1-1.43	3.81-17.52
			detection%	58	100	100	100	100	82	96	
Sendai	2003	50	GM(GSD)	0.15(3.75)	2.65(1.61)	1.01(1.85)	0.52(1.71)	1.68(1.75)	0.10(1.85)	0.31(2.12)	6.92(1.51)
			range	<0.05-1.25	0.87-7.59	0.21-4.94	0.09-1.57	0.32-5.70	<0.1-0.37	<0.1-1.13	2.74-17.94
			detection%	72	100	100	100	100	58	100	
Takayama	2008	50	GM(GSD)	0.04(2.29)*	2.51(1.84)	1.78(1.42)*	0.85(1.51)*	3.12(1.51)	0.20(2.15)*	0.60(2.66)	9.87(1.39)
			range	<0.05-0.49	0.82-11.25	1.01-4.50	0.26-2.68	1.28-7.13	<0.1-0.61	<0.1-2.46	5.44-22.09
			detection%	38	100	100	100	100	82	94	
Takayama	2003	50	GM(GSD)	0.11(2.35)	3.19(1.62)	1.30(1.73)	0.65(1.63)	2.74(1.60)	0.14(1.88)	0.55(1.72)	9.18(1.49)
			range	<0.05-1.72	1.36-20.28	0.64-9.88	0.18-2.26	0.77-7.81	<0.1-0.51	0.16-1.98	4.49-37.04
			detection%	88	100	100	100	100	80	100	
Kyoto	2009	30	GM(GSD)	0.11(1.98)*	5.28(1.57)*	2.78(1.42)*	1.10(1.45)	3.20(1.64)*	0.24(1.87)*	0.45(1.57)*	13.67(1.42)
			range	<0.05-0.31	2.60-16.52	1.34-4.40	0.60-2.25	1.20-11.26	<0.1-0.99	0.22-1.15	6.60-26.81
			detection%	96.7	100	100	100	100	93.3	100	
Kyoto	2002	30	GM(GSD)	0.23(1.89)	7.12(1.54)	2.09(1.67)	0.91(1.66)	1.89(1.65)	0.12(2.04)	0.31(1.83)	12.98(1.52)
			range	0.08-1.25	2.69-19.64	0.81-5.37	0.35-2.54	0.72-5.44	<0.1-0.37	<0.1-1.00	5.38-33.75
			detection%	100	100	100	100	100	66.7	96.7	
Osaka	2008	50	GM(GSD)	0.07(3.11)*	13.46(1.79)*	3.54(1.62)*	1.11(1.60)*	3.05(1.73)*	0.16(2.55)	0.52(2.62)	23.08(1.64)*
			range	<0.05-1.11	5.59-201.68	0.85-14.57	0.36-2.80	1.01-8.79	<0.1-0.75	<0.1-1.95	10.77-220.07
			detection%	48	100	100	100	100	68	94	
Osaka	2004	10	GM(GSD)	0.21(2.00)	29.54(1.29)	6.41(1.38)	2.38(1.48)	5.45(1.46)	0.25(2.28)	0.44(2.40)	45.42(1.27)
			range	0.05-0.45	20.60-45.20	3.07-9.22	1.41-4.17	3.19-9.01	<0.1-0.51	<0.1-1.02	31.67-65.57
			detection%	100	100	100	100	100	90	90	

GM: Geometric mean; GSD: Geometric standard deviation

* GMs between time points are significantly different in each sampling site ($p < 0.05$ by Student's t test after log transformation).

Table 4
Serum concentrations of PFCAs in Korea

Sampling site	Year	n		Concentration (ng mL ⁻¹)							ΣPFCAs
				PFHpA	PFOA	PFNA	PFDA	PFUnDA	PFDoDA	PFTTrDA	
Busan	2008	35	GM(GSD)	0.04(1.92) ^{*A}	4.67(1.40)	1.91(1.45) ^{*A}	0.91(1.38)	2.91(1.54) ^{*A}	0.20(2.03) ^{*AB}	0.94(1.92) ^{*A}	11.87(1.38) ^{*A}
			range	<0.05-0.16	2.77-9.80	1.02-3.89	0.44-1.76	1.03-7.62	<0.1-0.81	<0.1-2.74	6.70-24.13
			detection%	40	100	100	100	100	85.7	97.1	
	2000	30	GM(GSD)	0.10(1.59) ^B	3.69(1.47)	1.77(1.41) ^A	0.84(1.45)	2.06(1.66) ^{AB}	0.14(1.61) ^A	0.72(1.67) ^A	9.58(1.39) ^B
			range	<0.1-0.28	1.19-7.33	0.89-3.61	0.32-1.47	0.58-3.95	<0.20-0.39	<0.20-1.73	4.31-16.00
			detection%	30	100	100	100	100	33.3	100	
	1994	39	GM(GSD)	0.10(1.58) ^B	4.11(1.43)	1.35(1.96) ^B	0.89(1.65)	1.37(2.81) ^B	0.11(1.61) ^B	0.36(2.90) ^B	9.05(1.46) ^B
			range	<0.1-0.32	1.72-9.63	<0.10-5.20	0.25-2.98	<0.20-13.16	<0.20-1.03	0.10-2.89	4.08-32.50
			detection%	35.9	100	97.4	100	92.3	7.7	69.2	
Seoul	2007	36	GM(GSD)	0.03(1.48)	2.29(1.34)	1.13(1.32) [*]	0.58(1.38)	2.18(1.48) [*]	0.12(2.03)	0.59(2.10) [*]	7.10(1.35) [*]
			range	<0.05-0.12	1.22-4.64	0.74-2.01	0.32-1.00	1.10-5.62	<0.10-0.38	<0.10-1.54	3.94-12.55
			detection%	13.9	100	100	100	100	66.7	97.2	
	1994	24	GM(GSD)	0.08(1.00)	2.09(1.54)	0.65(2.01)	0.45(2.06)	0.54(3.89)	0.10(1.26)	0.16(2.40)	4.63(1.49)
			range	<0.1	0.89-4.09	<0.1-1.73	<0.1-1.18	<0.20-3.59	<0.20-0.31	<0.20-1.08	2.56-10.69
			detection%	0	100	95.8	95.8	70.8	4.2	25	

GM: Geometric mean; GSD: Geometric standard deviation

* GMs among different time points are significantly different in each sampling sites ($p < 0.05$ by Student's t test or Tukey's HSD test after log transformation). Alphabetic suffix was used for comparisons among three groups. For example, the letters A and B indicate that the corresponding values differ significantly at $p < 0.05$, while A and AB or AB and B indicated that the corresponding values do

Table 5
 Serum concentrations of PFCAs in Hanoi, Vietnam

Sampling site	Year	n		Concentration (ng mL ⁻¹)							ΣPFCAs
				PFHpA	PFOA	PFNA	PFDA	PFUnDA	PFDODA	PFTTrDA	
Hanoi	2007-2008	37	GM(GSD)	0.03(1.00)	0.61(1.55)	0.89(1.47)	0.82(1.67)	1.55(1.53)	0.09(1.85)	0.36(3.38)	4.73(1.38)
			range	<0.05	0.20-1.43	0.35-1.65	0.19-2.03	0.57-3.95	<0.10-0.26	<0.10-1.99	2.58-9.43
			detection%	0	100	100	100	100	51.4	86.5	

GM: Geometric mean; GSD: Geometric standard deviation

Table 6

Correlation between different chain length PFCAs

Combination		ρ	p value
PFOA	PFHpA	0.398	<0.001
PFNA	PFHpA	0.223	<0.001
PFNA	PFOA	0.734	<0.001
PFDA	PFHpA	0.165	<0.001
PFDA	PFOA	0.534	<0.001
PFDA	PFNA	0.727	<0.001
PFUnDA	PFHpA	0.019	0.660
PFUnDA	PFOA	0.323	<0.001
PFUnDA	PFNA	0.646	<0.001
PFUnDA	PFDA	0.689	<0.001
PFDoDA	PFHpA	0.055	0.208
PFDoDA	PFOA	0.235	<0.001
PFDoDA	PFNA	0.462	<0.001
PFDoDA	PFDA	0.563	<0.001
PFDoDA	PFUnDA	0.740	<0.001
PFTTrDA	PFHpA	-0.117	0.008
PFTTrDA	PFOA	0.063	0.151
PFTTrDA	PFNA	0.264	<0.001
PFTTrDA	PFDA	0.360	<0.001
PFTTrDA	PFUnDA	0.552	<0.001
PFTTrDA	PFDoDA	0.471	<0.001

 ρ : Spearman's rank correlation coefficient

Table 7
Factor analysis among PFCAs

	Initial solution		Varimax rotated	
	F1	F2	F1	F2
Eigenvalue	3.48	1.36		
Contribution (%)	49.72	19.40		
Eigenvector				
PFHpA	0.092	0.618	-0.198	0.713
PFOA	0.365	0.480	0.327	0.819
PFNA	0.474	0.179	0.673	0.610
PFDA	0.469	0.036	0.745	0.459
PFUnDA	0.446	-0.230	0.858	0.168
PFDoDA	0.374	-0.266	0.760	0.066
PFTTrDA	0.274	-0.481	0.719	-0.244
Factor score (mean±standard deviation)				
Sendai	2008		0.31±0.78*	-0.41±0.67*
	2003		-0.84±0.90	0.17±0.92
Takayama	2008		0.50±0.67*	-0.49±0.85*
	2003		-0.10±0.76	0.02±0.81
Kyoto	2009		0.44±0.68*	0.68±0.52*
	2002		-0.46±0.85	1.29±0.50
Osaka	2008		0.91±1.03	1.17±0.78*
	2004		0.92±0.71	2.42±0.48
Busan	2008		0.68±0.67 ^{†A}	-0.28±0.56 ^{†A}
	2000		0.06±0.61 ^B	0.15±0.46 ^A
	1994		-0.48±0.95 ^C	0.42±0.63 ^B
Seoul	2007		0.02±0.69 ^{†A}	-0.92±0.36 ^{†A}
	1994		-1.49±0.90 ^B	-0.22±0.48 ^B
Hanoi	2007- 2008		-0.27±0.74	-1.48±0.65

F1: 1st factor; F2: 2nd factor

* Means between time points are significantly different ($p < 0.05$ by Student's t test)


† Means among different time points are significantly different ($p < 0.05$ by Student's t test or Tukey's HSD test). For example, the letters A and B indicate that the corresponding values differ significantly at $p < 0.05$, while A and A or B and B indicated that the corresponding values do not differ significantly.

Table 8
Comparison of serum or whole blood concentrations of PFCAs with reported data

Sampling site	Year	n	Sex	Sample	Concentration (ng mL ⁻¹)	Concentration (ng mL ⁻¹)							reference
						PFHpA	PFOA	PFNA	PFDA	PFUnDA	PFDoDA	PFTTrDA	
Japan													
Sendai	2008	50	F	median serum	0.07	2.36	1.82	0.73	2.97	0.19	0.74	This study	
	2003	50	F	median serum	0.22	2.59	1.07	0.55	1.79	0.11	0.37	This study	
Takayama	2008	50	F	median serum	<0.05	2.08	1.72	0.80	3.11	0.24	0.77	This study	
	2003	50	F	median serum	0.13	3.21	1.29	0.69	2.69	0.15	0.56	This study	
Kyoto	2009	15/15	M/F	median serum	0.11	5.52	2.69	1.01	3.15	0.25	0.45	This study	
	2002	15/15	M/F	median serum	0.23	7.20	2.15	0.90	1.72	0.14	0.30	This study	
Osaka	2008	50	F	median serum	<0.05	12.80	3.32	1.10	2.98	0.19	0.72	This study	
	2004	10	F	median serum	0.23	28.90	6.87	2.53	5.83	0.35	0.51	This study	
Korea Busan	2008	35	F	median serum	<0.05	4.64	1.98	0.92	3.00	0.22	0.92	This study	
	2000	30	F	median serum	<0.1	3.98	1.92	0.87	2.27	<0.2	0.76	This study	
	1994	39	F	median serum	<0.1	3.98	1.28	0.94	1.82	<0.20	0.49	This study	
Seoul	2007	36	F	median serum	<0.05	2.21	1.11	0.57	2.37	0.14	0.74	This study	
	1994	24	F	median serum	<0.1	2.31	0.76	0.50	0.89	<0.20	<0.20	This study	
Vietnam Hanoi	2007-2008	37	F	median serum	<0.05	0.63	0.91	0.85	1.58	0.11	0.65	This study	
Norway	2006	>20	M	median serum	0.078	2.7	0.55	0.22	0.14	<0.05	0.071	Haug et al., 2009	
Sri Lanka Colombo	2003	10	M	median serum	0.146	9.32	0.299	0.18	0.186	0.015	-	Guruge et al., 2005	
China Ningbo	2006-2008	8/12	M/F	median pooled serum	<0.1	3.28	0.984	0.718	0.917	<0.18	-	Pan et al., 2010	
Spain Catalonia	2002-2007	24/24	M/F	median whole blood	<0.78	1.65	0.41	0.24	0.2	-	-	Ericson et al., 2007	
Poland Gdańsk	2003	10/5	M/F	median whole blood	0.086	2.8	0.49	0.17	0.078	0.012	-	Falandysz et al., 2006	
Belgium Antwerp	2002-2005	182	F	pooled serum	-	3.18	2.41	1.86	-	-	-	Roosens et al., 2010	
Australia Queensland	2006-2007	42/42	M/F	mean pooled serum	-	6.4	0.8	0.29	-	-	-	Toms et al., 2009	
Denmark Copenhagen	2003	105	M	median serum	0.2	4.9	0.8	0.9	0.1	0.08	<0.1	Joensen et al., 2009	
Sweden Stockholm	1997-2000	40/26	M/F	median whole blood	-	2.5	0.3	0.2	0.2	-	-	Karrman et al., 2006	
Atlanta USA	2003	10/10	M/F	median serum	-	4.35	2.35	0.35	0.7	-	-	Kuklenyik et al., 2004	


Supplemental Figure 1

A




— x Osaka 2004 — + Osaka 2008

B




— o Kyoto 2002 — • Kyoto 2009

C




— Δ Takayama 2003 — ◆ Takayama 2008

D




— □ Sendai 2003 — ■ Sendai 2008

E



— z Busan 1994 — y Busan 2000
— x Busan 2008

F



— o Seoul 1994 — • Seoul 2007
— * Hanoi