TITLE:

The functional effect of Gly209 and Ile213 substitutions on lysozyme activity of family 19 chitinase encoded by cyanophage Ma-LMM01

AUTHOR(S):

Hosoda, Naohiko; Kurokawa, Yoichi; Sako, Yoshihiko; Nagasaki, Keizo; Yoshida, Takashi; Hiroishi, Shingo

CITATION:

Hosoda, Naohiko ...[et al]. The functional effect of Gly209 and Ile213 substitutions on lysozyme activity of family 19 chitinase encoded by cyanophage Ma-LMM01. Fisheries Science 2011, 77(4): 665-670

ISSUE DATE:

2011-07

URL:

<http://hdl.handle.net/2433/143681>

RIGHT:

The final publication is available at www.springerlink.com; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。 ; This is not the published version. Please cite only the published version.

Title

The functional effect of Gly209 and Ile213 substitutions on lysozyme activity of family 19 chitinase encoded by cyanophage Ma-LMM01

Authors

Naohiko Hosoda • Yoichi Kurokawa • Yoshihiko Sako • Keizo Nagasaki • Takashi Yoshida • and Shingo Hiroishi

N. Hosoda

Department of Marine Bioscience, Fukui Prefectural University, 1-1 Gakuen-cho, Obama, Fukui 917-0003, Japan

E-mail: s0894004@s.fpu.ac.jp

Y. Kurokawa

Department of Bioscience, Fukui Prefectural University, 4-1-1 Matsuoka-Kenjojima, Eihei-cho, Yoshida-gun, Fukui 910-1195, Japan

E-mail: kurokawa@fpu.ac.jp

Y. Sako

Graduate School of Agriculture, Kyoto University, Kyoto 606-8502, Japan

E-mail: sako@kais.kyoto-u.ac.jp

K. Nagasaki

National Research Institute of Inland Sea, Fisheries Research Agency, 2-17-5 Maruishi, Hatsukaichi, Hiroshima 739-0452, Japan

E-mail: nagasaki@affrc.go.jp

S. Hiroishi

Department of Marine Bioscience, Fukui Prefectural University, 1-1 Gakuen-cho, Obama, Fukui 917-0003, Japan

E-mail: hiroishi@fpu.ac.jp

Corresponding author

T. Yoshida

Mailing address: Laboratory of Marine Microbiology, Graduate School of Agriculture, Kyoto University, Kyoto 606-8502, Japan.

Tel: +81-75-753-6218; Fax: +81-75-753-6226

E-mail: yoshiten@kais.kyoto-u.ac.jp

36 **Abstract**

37 ORF69 in the cyanophage infecting *Microcystis aeruginosa*, Ma-LMM01,
38 shows homology to the family 19 chitinases where the catalytic domain has structural
39 similarity to lysozyme. Chitinases hydrolyze chitin, a β -1, 4-linked monopolymer of
40 *N*-acetylglucosamine (GlcNAc); whereas lysozymes hydrolyzes peptidoglycan,
41 alternating β -1, 4-linked copolymers of *N*- acetylmuramic acid (MurNAc) and GlcNAc.
42 Using amino acid sequence comparison to ORF69, the putative sugar binding
43 residues, Gln162 and Lys165, from the barley chitinase (the model enzyme for the
44 family 19 chitinases) corresponding to subsites -4 and -3 were found to be replaced with
45 Gly209 and Ile213, respectively, in ORF69. To analyze their contribution to substrate
46 binding affinity, ORF69 was cloned into *Escherichia coli*; and two mutant proteins
47 G209Q and I213K were prepared using site-directed mutagenesis. The wild-type gene
48 product (gp69) showed both lysozyme and chitinase activities. In contrast, the I213K
49 mutant showed a decrease (70%) in lysozyme activity and a significant increase (50%)
50 in chitinase activity; whereas, the G209Q mutant almost completely abolished both
51 enzyme activities. The data suggest the Ile213 residue is involved in recognizing the
52 substrate MurNAc; and Gly209 has significant contribution in chitinase and lysozyme
53 activities for the wild-type gp69.

54 **Keywords**55 *Microcystis*, cyanophage, family 19 chitinase, site-directed mutagenesis

56 Introduction

57 Throughout the world, *Microcystis aeruginosa* is one of the common
58 bloom-forming species in eutrophic freshwaters. Some strains produce cyclic peptide
59 toxins called microcystins that cause serious health problems in water management [1].
60 Previously we isolated a cyanophage, Ma-LMM01, infecting the toxic *M. aeruginosa*
61 strain NIES298 [2]. The genome of Ma-LMM01 contains 184 ORFs [3]. The majority
62 of the predicted genes have no detectable homologues in present databases including
63 other *Myoviridae*; and thus Ma-LMM01 was assigned as a member of a new lineage of
64 the *Myoviridae* family [3, 4]. Of the ORFs, ORF69 is predicted to encode for a member
65 of the family 19 chitinases whose catalytic domain has structural similarity to lysozyme
66 [5].

67 Chitinase (EC 3.2.1.14) is a glycoside hydrolase that hydrolyzes chitin, a linear
68 β -1, 4-linked monopolymer of *N*-acetylglucosamine (GlcNAc). Based on structures and
69 catalytic mechanisms, the chitinases are classified into two families, 18 and 19 [6, 7].
70 Family 18 chitinases are widely distributed in a variety of organisms such as bacteria,
71 fungi, bacteriophages, animals and higher plants (classes III and V); whereas family 19
72 chitinases are found only in higher plants (classes I, II and IV). Recently, however,
73 some members of the family 19 chitinases have been found in genomes of

74 actinobacteria, proteobacteria, nematodes and bacteriophages [8]. Based on amino acid
75 sequence comparisons, phylogenetic analysis shows the family 19 chitinases are
76 separated into five clusters (clusters I to V) [8]. Of these, cluster III of the family 19
77 chitinases are most distantly related to the other clusters [8]. The cluster III family 19 of
78 chitinases consists only of those from proteobacteria and bacteriophages and the genes
79 in proteobacteria are often found within the phage-related regions. The cluster III family
80 19 chitinase genes in PA0629 from *Pseudomonas aeruginosa* and PFL_1227 from *P.*
81 *fluorescens* Pf-5 are located within a region of one of the variants of a defective phage
82 (pyocin) and prophage, respectively. Their recombinant proteins have lysozyme activity
83 that hydrolyses peptidoglycans, alternating β -1, 4-linked residues of *N*- acetylmuramic
84 acid (MurNAc) and GlcNAc [9, 10].

85 The family 19 chitinases are shown to have highly conserved catalytic residues
86 and substrate-binding residues using crystal structure analyses, e.g. those from barley
87 (cluster I) [11], Jack bean (cluster I) [12], ChiC of *Streptomyces griseus* HUT6037
88 (cluster II) [13] and ChiG of *S. coelicolor* A3(2) (cluster II) [14]. We found two
89 residues were replaced in the putative substrate-binding residues of ORF69 when
90 compared to the other family 19 chitinases. Here, we determined the lysozyme and
91 chitinase activities of ORF69 gene products from cyanophage Ma-LMM01; and

examined the function of the two residues in recognition of the substrates, chitin and peptidoglycan, using site-directed mutagenesis.

Materials and methods

Cloning of ORF69 and derivatives

The genomic DNA of Ma-LMM01 was purified as described previously [2]. To isolate the full sequence of the ORF69 gene, a PCR reaction was performed with a forward primer MaPOrf69InF and a reverse primer MaPOrf69InR1 containing *EcoRI* site (Table 1). The PCR was performed in a 50 μ l containing 200 ng Ma-LMM01 DNA, 10 μ M primers, 250 μ M each dNTPs, 1X PCR buffer for KOD-plus- and 1U KOD-Plus- (TOYOBO, Osaka, Japan). The reaction conditions were: 2 min initial denaturing at 94 °C followed by 35 cycles: 94 °C for 15 s, 45 °C for 30 s and 68 °C for 90 s. The reaction mixture was purified using a Wizard[®] SV Gel and PCR Clean-Up System (Promega, Madison, WI). The pTrc-OmpA vector was fused with the OmpA signal sequences upstream of a multiple cloning site; thus the resulting expressed protein is transported to the periplasm in the transformants [15]. The purified DNA fragments were digested with *EcoRI* (TOYOBO) and cloned into expression vector pTrc-OmpA. The coding region of the ORF69 was inserted downstream of the OmpA

signal sequence yielding pTrc-OmpA-ORF69. *E. coli* JM109 (TOYOBO) was transformed with the pTrc-OmpA-ORF69; and the transformant was selected on LB plates containing 100 µg/ml carbenicillin disodium salt (Nacalai Tesque, Kyoto, Japan) and 0.5% glucose. The DNA sequence of the resultant plasmids was verified using the primers Trc-F and Trc-R2 (Table 1).

Site-directed mutagenesis

Three mutant proteins (G209Q, I213K and E122A) were constructed using PCR-based site-directed mutagenesis. The mutagenesis primers are shown in Table 1. The PCR reaction mixtures contained 100 ng plasmid DNA template, 10 pmol each of the forward and reverse primers containing the desired mutation, 2 mM dNTPs, reaction buffer and PfuUltra High-Fidelity DNA polymerase (Stratagene, La Jolla, CA). The reaction mixture was subjected to 20 cycles of PCR (95 °C for 30 s; 55 °C for 1min; and 68 °C for 12 min); then, the resultant plasmids were digested with *Dpn* I. Finally, the mutated plasmids were transformed into *E. coli* JM109 and expressed.

Preparation of culture supernatant from transformants.

E. coli JM109 cells containing the plasmid pTrc-OmpA-ORF69 or mutated plasmids were independently grown overnight at 30 °C in LB liquid medium containing 100 µg ml⁻¹ carbenicillin disodium salt and 0.5% glucose. Two-ml of the culture was

128 diluted to 100 ml of fresh LB medium and grown with shaking at 30 °C until the
129 $OD_{660nm} = 0.5$. Protein expression was induced adding
130 isopropyl- β -D-thiogalactopyranoside (IPTG) to a final concentration of 0.1 mM. After
131 growth for another 6 h, the supernatant of two consecutive centrifugations (15,000g, 5
132 min, at 4 °C) was stored as the culture supernatant at -20 °C until used. The pelleted
133 cells were resuspended in 50 mM sodium phosphate buffer (pH 6.2) and disrupted using
134 sonication. After centrifugation, the crude extracts were also stored at -20 °C. Amount
135 of total protein was measured by method of Bradford with BSA as a standard [16].
136 **Enzymatic activities of culture supernatant from *E. coli* cells expressing the**
137 **wild-type ORF69 and its mutants.**

138 **(i) Lysozyme activity**

139 Lyophilized cells of *Micrococcus lysodeikticus* ATCC4698 (Nacalai) were
140 re-suspended at a concentration of 0.25 mg/ml in 10 ml of 50 mM sodium phosphate
141 buffer (pH 6.2); then, 100 μ g of a culture supernatant (see above) was added to 2 ml of
142 the *M. lysodeikticus* cell suspension. Cell lysis was measured continuously by
143 monitoring the decrease in turbidity (OD_{660nm}) using a Ultraspec 2100 *pro* (GE
144 Healthcare, Buckinghamshire, UK) for 15 minutes at 37 °C. Lysozyme activity was
145 calculated from the linear portion of the digestion graph representing absorbance versus

time. One unit of enzyme activity was defined as the amount of enzyme causing an

absorbance decrease of 0.01 OD_{660nm} per min at 37 °C [17].

(ii) Chitinase activity

The chitinase assay was performed using *p*-nitrophenyl chitooligosaccharide *p*NP-(GlcNAc)_n (n= 2 to 5) (Yaizu Suisan Chemical Co. Ltd., Shizuoka, Japan) as the substrate [18]. The reaction mixture (200μl) contained 2.5 mM of substrate in 50 mM sodium phosphate buffer (pH 6.2). The reaction was initiated adding 100 μg of a culture supernatant to the above reaction mixture pre-incubated for 5 min; and incubated for 15 min after addition at 37 °C. The reaction was stopped by adding 250 μl 0.2 M Na₂CO₃ and the released *p*-nitrophenol was measured at OD_{420nm}. One unit of chitinase activity was defined as the amount of enzyme causing 1μmol releasing of *p*-nitrophenol per min at 37 °C.

Results

The amino acid sequence of ORF69

A phylogenetic analysis showed the amino acid sequence of ORF69 was clustered within the cluster III family 19 chitinases (data not shown). Comparison of the conserved domain database analysis [19] showed the C-terminal region of ORF69 (residues 110-251) contains a domain similar to a glycoside hydrolase family 19

164 chitinase (cd00325). There are two catalytic residues and seven putative sugar binding
165 residues in the family 19 chitinases.

166 The position of the 161-166 residues (the 161-166 loop) in the cluster I barley
167 family 19 chitinase contains two polar amino acids (Gln162 and Lys165) (Fig. 1) [20].
168 Gln162 and Lys165 are thought to form the substrate-binding site, namely subsite -4
169 and -3, respectively (subsites are numbered according to the standard nomenclature;
170 cleavage occurs between the sugar units bound in subsites -1 and +1 [21]). These
171 residues in ORF69 and other cluster III family 19 chitinases are replaced with the
172 non-polar amino acids, Gly209 and Ile213. Therefore, we predicted mutations in
173 residues Gly209 and Ile213 of ORF69 would affect the recognition of substrates. To
174 confirm this hypothesis, site-directed mutagenesis was performed.

175 **Lysozyme activity of gp69 and its mutants**

176 Based on the vector pTrc-OmpA-ORF69 encoding the wild-type ORF69
177 enzyme (gp69), two vectors encoding mutants in the putative sugar binding site (G209Q,
178 I213K) and one in the catalytic site (E122A) were constructed.

179 The growth of transformants was monitored measuring the change in OD_{660nm}.
180 After induction with IPTG, the OD value of both transformants containing
181 pTrc-OmpA-ORF69 and pTrc-OmpA-ORF69-I213K declined about 30% from 4 to 8 h

(data not shown). In contrast, normal growths were observed in transformants

containing pTrc-OmpA-ORF69-G209Q and pTrc-OmpA-ORF69-E122A.

In transformants containing either pTrc-OmpA-ORF69 or pTrc-OmpA-ORF69-I213K, induction of protein expression caused cell lysis and most of the lysozyme activities were observed in the culture supernatant fraction (data not shown). In the transformants containing pTrc-OmpA-ORF69-G209Q, no lysozyme activity was observed in both culture supernatant fractions and crude protein from pelleted cells (data not shown). In addition, a 10-fold concentration of the crude extracts had no impact on lysozyme activity. The activities of wild-type gp69, G209Q and I213K for *M. lysodeikticus* were 61.3 ± 17.2 , 1.1 ± 1.2 and 18.1 ± 12.9 U/mg, respectively (Table 2) suggesting mutations in these residues have effects on the lysozyme activity. Lysozyme activity from the mutant protein, E122A (having a mutation in the catalytic site) was not detected.

Chitinase activity of gp69 and its mutants digesting chitooligosaccharides.

To determine the chitinase activity of wild-type gp69 and its mutants, we used the culture supernatant fraction to measure the release of *p*-nitrophenol, using several chitooligosaccharides [*p*NP-(GlcNAc)_n (n= 2 to 5)] as substrates. The highest hydrolyzing activity of wild-type gp69 was observed when *p*NP-(GlcNAc)₃ was used as

200 the substrate (Fig. 2). Whereas, the hydrolytic activity to $p\text{NP}-(\text{GlcNAc})_5$ was
201 approximately one-half compared to $p\text{NP}-(\text{GlcNAc})_3$; and the hydrolysis activity for
202 $p\text{NP}-(\text{GlcNAc})_4$ was not detected (Fig. 2). Compared to the gp69, the hydrolytic activity
203 of mutant protein I213K had approximately a 1.3-fold increase using $p\text{NP}-(\text{GlcNAc})_3$
204 and the hydrolytic activity to $p\text{NP}-(\text{GlcNAc})_4$ was increased. Thus, the I213K mutation
205 increased the hydrolyzing activity towards $p\text{NP}-(\text{GlcNAc})_3$ and $p\text{NP}-(\text{GlcNAc})_4$ even
206 though the lysozyme activity of I213K was the 30% of that of wild-type (Table 2). In
207 contrast, the mutant protein, G209Q in both fractions, showed no detectable activity
208 towards any substrate. In addition, a 10-fold concentration of the crude extracts of
209 G209Q did not show chitinase activities (data not shown).

210 Discussion

211 We tried to construct various vectors for the expression of ORF69 where
212 spontaneous mutations were observed in the cloned sequences. This was possibly due to
213 the toxicity of the gene products for *E. coli* [22]. We obtained only one clone with the
214 correct sequence of ORF69 using the pTrc-OmpA vector that allows expression of
215 OmpA signal-fused protein that was guided to the periplasmic space of the *E. coli* cell.
216 For this reason, crude extracts from the transformants were used in the enzymatic
217 experiments.

218 In general, the family 19 chitinases are endo-type chitinases that generate
219 various sizes of chitooligomers [23]. However, the release of *p*-nitrophenol from the
220 gp69 using the chitooligosaccharides as substrate suggests the wild-type gp69 is not
221 typically an endo-like enzyme. Chi19 from *Vibrio proteolyticus* hydrolyzes colloidal
222 chitin to release small oligosaccharides at the early stage of the reaction; and it is
223 thought to be an exo-like family 19 chitinase [18]. Hen egg white lysozyme (HEWL)
224 has six subsites (from -4 to +2) [24]. Also, the subsite structures of higher plant family
225 19 chitinases (cluster I) are assumed to be represented by (-4)(-3)(-2)(-1)(+1)(+2) [20]
226 or (-3)(-2)(-1)(+1)(+2)(+3) [25] except for those from *Carica papaya* [26] and *Picea*
227 *abies* (Norway spruce) [27]. In HEWL, the corresponding subsite -3 is known to be
228 responsible for interaction with MurNAc [24]. Combined, our data suggests the Ile213
229 residue corresponding to subsite -3 in ORF69 contributes to the interaction with
230 MurNAc and the replacement of Ile213 with Lys may emphasize the affinity for
231 *p*NP-(GlcNAc)₃ and *p*NP-(GlcNAc)₄ than for *M. lysodeikticus*. Whereas, the I213K
232 mutation decreased the hydrolyzing activity to *p*NP-(GlcNAc)₂ and *p*NP-(GlcNAc)₅
233 where one possible explanation is interference of substrate access to the catalytic site
234 due to the small oligosaccharides but we do not have any useful data concerning this
235 hypothesis. The substitution of Gly209 to Gln caused a significant decrease in both

236 lysozyme and chitinase activities. There is a possibility the single amino acid
237 replacement altered the conformation of recombinant protein G209Q [28]. Further study
238 is required to determine the effect of the Gly209 residue in conformational changes.

239 Nineteen cluster III family 19 chitinases (17 genes in bacterial genomes and 2
240 genes in bacteriophages) were in the current database [8, 29]. Two residues, Asn124 and
241 Lys165 (according to the barley family 19 chitinase numbering), are presumed to be
242 responsible for the subsite -3 activity. These residues are highly conserved among
243 family 19 chitinases [20]. In ORF69 of Ma-LMM01, the amino acids corresponding to
244 the Lys165 residue in 15 genes of the 19 cluster III family 19 chitinases are replaced
245 with non-polar amino acids. In the remaining four genes, the amino acids corresponding
246 to Lys165 is replaced with a Tyr residue. Further research focusing on the role of the Tyr
247 residue is necessary to determine its contribution in sugar binding.

248 Family 19 chitinases are hypothesized to be horizontally transferred from
249 higher plants to bacteria [29, 30]. In the evolutionary history of the family 19 chitinases,
250 mutation in key residues (ex. corresponding to subsite -3) may lead to alternation of
251 affinity for substrates; and may have spread to bacteria and phages as a lytic enzyme.

252 **Acknowledgement**

253 This study was supported by a Grant-in-Aid for Science Research (No. 20310045). NH
254 is a research fellow supported by the Japan Society for the Promotion of Science for
255 Young Scientists (No. 218877).

References

1. Codd GA, Lindsay J, Young FM, Morrison LF, Metcalf JS (2005) Harmful Cyanobacteria. In: Huisman J et al (eds) Harmful Cyanobacteria. Springer, Netherlands, pp 1-23
2. Yoshida T, Takashima Y, Tomaru Y, Shirai Y, Takao Y, Hiroishi S, Nagasaki K (2006) Isolation and characterization of a cyanophage infecting the toxic cyanobacterium *Microcystis aeruginosa*. Appl Environ Microbiol 72:1239-1247
3. Yoshida T, Nagasaki K, Takashima Y, Shirai Y, Tomaru Y, Takao Y, Sakamoto S, Hiroishi S, Ogata H (2008) Ma-LMM01 infecting toxic *Microcystis aeruginosa* illuminates diverse cyanophage genome strategies. J Bacteriol 190:1762-1772
4. Lavigne R, Darius P, Summer EJ, Seto D, Mahadevan P, Nilsson AS, Ackermann HW, Kropinski AM (2009) Classification of Myoviridae bacteriophages using protein sequence similarity. BMC Microbiol 9:224
5. Monzingo AF, Marcotte EM, Hart PJ, Robertus JD (1996) Chitinases, chitosanases, and lysozymes can be divided into procaryotic and eucaryotic families sharing a conserved core. Nat Struct Biol 3:133-140
6. Henrissat B (1991) A classification of glycosyl hydrolases based on amino acid sequence similarities. Biochem J 280:309-316
7. Henrissat B, Bairoch A (1993) New families in the classification of glycosyl hydrolases based on amino acid sequence similarities. Biochem J 293:781-788
8. Kawase T, Saito A, Sato T, Kanai R, Fujii T, Nikaidou N, Miyashita K, Watanabe T (2004) Distribution and phylogenetic analysis of family 19 chitinases in *Actinobacteria*. Appl Environ Microbiol 70:1135-1144
9. Nakayama K, Takashima K, Ishihara H, Shinomiya T, Kageyama M, Kanaya S, Ohnishi M, Murata T, Mori H, Hayashi T (2000) The R-type pyocin of *Pseudomonas aeruginosa* is related to P2 phage, and the F-type is related to lambda phage. Mol Microbiol 38:213-231
10. Mavrodi DV, Loper JE, Paulsen IT, Thomashow LS (2009) Mobile genetic elements in the genome of the beneficial rhizobacterium *Pseudomonas fluorescens* Pf-5. BMC Microbiol 9:8
11. Hart PJ, Pfluger HD, Monzingo AF, Hollis T, Robertus JD (1995) The refined crystal structure of an endochitinase from *Hordeum vulgare* L. seeds at 1.8 Å resolution. J Mol Biol 248:402-413
12. Hahn M, Hennig M, Schlesier B, Hohne W (2000) Structure of jack bean chitinase. Acta Cryst D56:1096-1099

- 279 13. Kezuka Y, Ohishi M, Itoh Y, Watanabe J, Mitsutomi M, Watanabe T, Nonaka T (2006) Structural studies of a two-domain chitinase
280 from *Streptomyces griseus* HUT6037. J Mol Biol 358:472-484
- 281 14. Hoell IA, Dalhus B, Heggset EB, Aspmo SI, Eijsink VG (2006) Crystal structure and enzymatic properties of a bacterial family 19
282 chitinase reveal differences from plant enzymes. FEBS J 273:4889-4900
- 283 15. Kurokawa Y, Yanagi H, Yura T (2001) Overproduction of bacterial protein disulfide isomerase (DsbC) and its modulator (DsbD)
284 markedly enhances periplasmic production of human nerve growth factor in *Escherichia coli*. J Biol Chem 276:14393-14399
- 285 16. Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of
286 protein-dye binding. Anal Biochem 72:248-254
- 287 17. Wang SL, Chang WT (1997) Purification and characterization of two bifunctional chitinases/lysozymes extracellularly produced by
288 *Pseudomonas aeruginosa* K-187 in a shrimp and crab shell powder medium. Appl Environ Microbiol 63:380-386
- 289 18. Honda Y, Taniguchi H, Kitaoka M (2008) A reducing-end-acting chitinase from *Vibrio proteolyticus* belonging to glycoside
290 hydrolase family 19. Appl Microbiol Biotechnol 78:627-634
- 291 19. Marchler-Bauer A, Anderson JB, Chitsaz F, Derbyshire MK, DeWeese-Scott C, Fong JH, Geer LY, Geer RC, Gonzales NR, Gwadz
292 M, He S, Hurwitz DI, Jackson JD, Ke Z, Lanczycki CJ, Liebert CA, Liu C, Lu F, Lu S, Marchler GH, Mullokandov M, Song JS,
293 Tasneem A, Thanki N, Yamashita RA, Zhang D, Zhang N, Bryant SH (2009) CDD: specific functional annotation with the Conserved
294 Domain Database. Nucleic Acids Res 37:205-210
- 295 20. Brameld KA, Goddard WA, 3rd (1998) The role of enzyme distortion in the single displacement mechanism of family 19 chitinases.
296 Proc Natl Acad Sci USA 95:4276-4281
- 297 21. Davies GJ, Wilson KS, Henrissat B (1997) Nomenclature for sugar-binding subsites in glycosyl hydrolases. Biochem J 321:557-559
- 298 22. Perry LJ, Heyneker HL, Wetzel R (1985) Non-toxic expression in *Escherichia coli* of a plasmid-encoded gene for phage T4
299 lysozyme. Gene 38: 259-264
- 300 23. Robertus JD, Monzingo AF (1999) The structure and action of chitinases. EXS 87:125-135
- 301 24. Fukamizo T (2000) Chitinolytic enzymes: catalysis, substrate binding, and their application. Curr Protein Pept Sci 1:105-124

- 302 25. Honda Y, Fukamizo T (1998) Substrate binding subsites of chitinase from barley seeds and lysozyme from goose egg white.
303 Biochim Biophys Acta 1388:53-65
- 304 26. Huet J, Rucktooa P, Clantin B, Azarkan M, Looze Y, Villeret V, Wintjens R (2008) X-ray structure of papaya chitinase reveals the
305 substrate binding mode of glycosyl hydrolase family 19 chitinases. Biochemistry 47:8283-8291
- 306 27. Ubhayasekera W, Rawat R, Ho SW, Wiweger M, Von Arnold S, Chye ML, Mowbray SL (2009) The first crystal structures of a
307 family 19 class IV chitinase: the enzyme from Norway spruce. Plant Mol Biol 71:277-289
- 308 28. Gilbert HJ, Drabble WT (1980) Complementation *in vitro* between *guaB* mutants of *Escherichia coli* K12. J Gen Microbiol
309 117:33-45
- 310 29. Udaya Prakash NA, Jayanthi M, Sabarinathan R, Kanguane P, Mathew L, Sekar K (2010) Evolution, homology conservation, and
311 identification of unique sequence signatures in GH19 family chitinases. J Mol Evol 70:466-478
- 312 30. Watanabe T, Kanai R, Kawase T, Tanabe T, Mitsutomi M, Sakuda S, Miyashita K (1999) Family 19 chitinases of *Streptomyces*
313 species: characterization and distribution. Microbiology 145:3353-3363
- 314 31. Tamura K, Dudley J, Nei M, Kumar S (2007) MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0.
315 Mol Biol Evol 24:1596-1599

Tables

Table 1. Plasmids and primers used in this study

Plasmids or primers	Characteristics or sequences (5' to 3')	Sources or references
Plasmids		
pTrc-OmpA	Amp ^R , lacI ^q , ori (pBR322), trcP, rrmB T1T2 terminator	Kurokawa, <i>et al.</i> , [15]
pTrc-OmpA-ORF69	pTrc-OmpA with an insert of nucleotides 52314 to 53069 from cyanophage Ma-LMM01	This study
pTrc-OmpA-ORF69-G209Q	Substitute of ORF69 Gly209 to Gln209	This study
pTrc-OmpA-ORF69-I213K	Substitute of ORF69 Ile213 to Lys213	This study
pTrc-OmpA-ORF69-E122A	Substitute of ORF69 Glu122 to Ala122	This study
Primers		
MaPOrf69InF	CTA AGA AAC ATT GAT AGA GC	This study
MaPOrf69InR1 ^{a)}	CGG <u>AAT TCT</u> CAT GTC AGC ACC GCC TGT A (<i>EcoRI</i>)	This study
ORF69-E122A-F ^{b)}	CAG CTG ATG CAC GCG TCA <i>GGG</i> AAC CTA CG	This study
ORF69-E122A-R ^{b)}	CGT AGG TTC CCT GAC GCG TGC ATC AGC TG	This study
ORF69-G209Q-F ^{b)}	AGC GCA CTA CTG GAG AAC ACC AGG GCT AAA TGA AAT AGC AGA C	This study
ORF69-G209Q-R ^{b)}	GTC TGC TAT TTC ATT TAG CCC <i>TGG</i> TGT TCT CCA GTA GTG CGC T	This study
ORF69-I213K-F ^{b)}	GAA CAC GGG GGC TAA ATG AAA <i>AGG</i> CAG ACA AGA ATG ATA TAA A	This study
ORF69-I213K-R ^{b)}	TTT ATA TCA TTC TTG TCT <i>GCC</i> TTT TCA TTT AGC CCC CGT GTT C	This study
Trc-F	ACA TCA TAA CGG TTC TGGC	Kurokawa, <i>et al.</i> , [15]
Trc-R2	CAA ATTC TGT TTT ATC AGA CC	Kurokawa, <i>et al.</i> , [15]

a): The restriction site is underlined.

b): The mutation sites are in italics.

Table 2. Lysozyme activity of the culture supernatant of *E. coli* cells expressing the wild-type gp69 and its mutants.

Protein	Activity (U/ mg)	Relative activity (%)
Gp69	61.3 ± 17.2	100
G209Q	1.1 ± 1.2	1.8
I213K	18.1 ± 12.9	29.5
E122A	ND	-
Vector	ND	-
HEWL*	1.6 ± 0.01	2.6

The activity was measured by the decrease in absorbance at OD_{660nm}. The reaction mixtures were incubated in 50 mM sodium phosphate buffer, pH6.2 at 37 °C. The data are the means of three independent assays ± SD. -, Activity not detected. *, Activities of hen egg white lysozyme (HEWL; 0.6μg) was used as positive control. One unit of enzyme activity was defined as the amount of enzyme that causes an absorbance decrease of 0.01.

Figure legends

Figure 1. Alignment of amino acid sequences for the catalytic domains of family 19 chitinases. Sequence alignment was performed using MEGA version 4 software [31]. Residues conserved in the sequences are indicated by black backgrounds, whereas residues conserved in >80 and >60% of the proteins examined are indicated by white type on a dark gray background and by black type on a light gray background, respectively. The two catalytic amino acid residues from the family 19 chitinases are indicated by closed triangles. Residues predicted to interact with saccharide molecules in theoretical models of the barley chitinase/ (GlcNAc)₆ complex [19] are indicated by closed circles. The 161-166 residues (using the numbering of the barley enzyme; see text) are indicated by dots above the sequence. Based on higher plant and *Streptomyces* chitinase, the putative secondary structure assignments are indicated with “h” for α -helix and “s” in the β -strand. ORF69, Ma-LMM01 ORF69 (accession no. YP_851083); PA0629, lytic enzyme of *P. aeruginosa* PAO1 (NP_249320); putative lytic enzyme of ORF_PML14, *P. aeruginosa* (YP_788803); PFL_1227, lytic enzyme of *P. fluorescens* Pf-5 (YP_258358); ChiG, ChiG of *Streptomyces coelicolor* A3(2) (BAA75648); ChiC, ChiC of *S. griseus* HUT 6037 (BAA23739); Jackbean, class II chitinase of *Canavalia ensiformis* (CAA07413); and Barley, class II chitinase of *Hordeum vulgare* (P23951).

375 Figure 2. Hydrolytic activity for various chitooligosaccharides using the culture
376 supernatant from *E. coli* cells expressing the wild-type ORF69 and its mutants. The
377 reaction mixture was 2.5 mM substrate, 50 mM sodium phosphate buffer (pH6.2) and
378 mixed with 100 µg of the culture supernatant. pNP-(GlcNAc)₂, white bars;
379 pNP-(GlcNAc)₃, pale-grey bars; pNP-(GlcNAc)₄, dark-grey bars; and pNP-(GlcNAc)₅,
380 black bars. The reaction mixtures were incubated for 15 min in 50 mM sodium
381 phosphate buffer (pH6.2) at 37 °C. The bars are an average of three independent
382 measurements. N.D: Not detected.

383 Fig.1

385 Fig. 2

386