TITLE:

Light-Induced Conformational Changes in Full-Length *Arabidopsis thaliana* Cryptochrome.

AUTHOR(S):

Kondoh, Masato; Shiraishi, Chiaki; Müller, Pavel; Ahmad, Margaret; Hitomi, Kenichi; Getzoff, Elizabeth D; Terazima, Masahide

CITATION:

Kondoh, Masato ...[et al]. Light-Induced Conformational Changes in Full-Length *Arabidopsis thaliana* Cryptochrome.. *Journal of molecular biology* 2011, 413(1): 128-137

ISSUE DATE:

2011-10-14

URL:

<http://hdl.handle.net/2433/147966>

RIGHT:

© 2011 Elsevier Ltd.; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。; This is not the published version. Please cite only the published version.

Title:**Light-induced conformational changes in full-length *Arabidopsis thaliana*
Cryptochrome 1**

Author names and affiliations:

Masato Kondoh¹, Chiaki Shiraishi¹, Pavel Müller², Margaret Ahmad², Kenichi Hitomi³,
Elizabeth D. Getzoff³, Masahide Terazima^{1*}

¹*Department of Chemistry, Graduate School of Science, Kyoto University, Kyoto, 606-8502,
Japan*

²*Université Paris VI, UMR-CNRS 7632, Paris, France*

³*Department of Molecular Biology and The Skaggs Institute for Chemical Biology, The
Scripps Research Institute, La Jolla, CA 92037, USA.*

* Corresponding author.

E-mail: mterazima@kuchem.kyoto-u.ac.jp, TEL: 81-75-753-4026, FAX: 81-75-753-4000

Abstract

Cryptochromes are widespread flavoproteins with homology to photolyases, a class of blue light activated DNA repair enzymes. Unlike photolyases, both plant and animal cryptochromes have a C-terminal domain. This Cryptochrome C-terminal (CCT) domain mediates interactions with other proteins, while the photolyase-like domain converts light energy to a signal via reduction and radical formation of the flavin adenine dinucleotide (FAD) cofactor. However, the mechanism by which the photolyase-like domain regulates the CCT domain is not known. Here, we applied the pulsed laser induced transient grating (TG) method to detect conformational changes induced by blue-light excitation of full-length *Arabidopsis thaliana* Cryptochrome 1 (*AtCRY1*). A significant reduction in the diffusion coefficient of *AtCRY1* was observed upon photoexcitation, indicating that a large conformational change occurs in this monomeric protein. *AtCRY1* containing a single mutation (W324F) that abolishes an intra-protein electron transfer cascade did not exhibit this conformational change. Moreover, the conformational change was much reduced in protein lacking the CCT domain. Thus, we conclude that the observed large conformational changes triggered by light excitation of the photolyase-like domain result from C-terminal domain rearrangement. This inter-domain modulation would be critical for CRYs' ability to transduce a blue-light signal into altered protein-protein interactions for biological activity. Lastly, we demonstrate that the TG technique provides a powerful method for the direct observation and understanding of photoreceptor dynamics.

Key words: blue light receptor; flavin; photoreduction; diffusion; conformational dynamics

Abbreviations: *AtCRY1*, *Arabidopsis thaliana* Cryptochrome 1; Cryptochrome, CRY; Photolyase, PHR; CCT, Cryptochrome C-terminal; FAD, flavin adenine dinucleotide; TG, transient grating; TA, Transient Absorption; FTIR, Fourier Transform Infrared; *D*, molecular diffusion coefficients; CD, circular dichroism

Introduction

Cryptochromes (CRY) are flavoproteins found in all kingdoms of life: plants ^{1,2}, animals ³⁻⁵, and bacteria ⁶. CRYs share amino acid sequence similarities, a common structural fold and the non-covalently bound flavin adenine dinucleotide (FAD) cofactor with DNA photolyase (PHR) enzymes, which catalyze the light-dependent repair of UV-damaged DNA. CRYs have little or no DNA repair activity ⁷, but instead control a wide variety of life activities ^{1,8,9}; light-dependent growth and development in plants, the circadian clock in animals, and sensitivity to the magnetic field (light-dependent magnetoreceptor) in flies ⁵.

Among plant CRYs, *Arabidopsis thaliana* Cryptochrome 1 (*AtCRY1*) has been most extensively studied ^{1,2,10-15}, while recent publications expand the fundamental knowledge about the others such as *Arabidopsis thaliana* Cryptochrome 2 (*AtCRY2*) ¹⁶ or algal Cryptochrome ¹⁷. *AtCRY1* regulates many aspects of photomorphogenesis ^{1,2,18}. Plant CRYs, represented by *AtCRY1*, contain not only the N-terminal PHR-like domain (ca. 500 amino acids), but also an additional characteristic Cryptochrome C-terminal (CCT) domain ^{1,2}. While the crystal structure of the PHR-like domain of *AtCRY1* has been reported ¹⁰, the isolated CCT domain was found to be unstructured by NMR analysis ¹¹. For plant CRYs, the CCT domain plays an important functional role; genetic and physiological analysis of truncated forms of CRYs has demonstrated that the CCT domain mediates signal transduction by interaction with protein partners and can function in isolation when fused to a reporter gene ^{12,13}. The mechanism and kinetics for CRY signal transduction from the light excitation of FAD in the N-terminal PHR-like domain to protein-protein interactions of the CCT domain remain central unanswered questions.

For CRY signaling, the FAD-containing PHR-like domain most likely initiates light signal conversion through the cofactor, regulating the activity of CCT domain in a light dependent manner ^{12,13}. Transient Absorption (TA) ¹⁴ and Fourier Transform Infrared (FTIR)

¹⁵ studies on *AtCRY1* *in vitro* have both revealed that light-induced intra-protein electron transfer from an aromatic residue (Trp or Tyr) to the chromophore takes place during formation of the FAD radical (FADH[•]), which has been shown to be the signaling state in plant CRYs ^{16,19}, following photoexcitation of oxidized flavin. This electron transfer is presumed to take place within the PHR-like domain of CRYs by analogy with that during photoactivation (photoreduction, not photoreactivation (DNA repair)) reaction of *E. coli* DNA photolyase ²⁰. Subsequent to light absorption and flavin reduction, additional changes in the protein have been probed by FTIR and partial proteolysis experiments, which suggested the possibility of light dependent conformational rearrangement of *AtCRY1* ^{11,15}. However, the conformational changes observed by FTIR are relatively weak and have not been well assigned ¹⁵. Similarly, indirect proteolysis analysis ¹¹ showed only minor effects and gave poor information in time-resolution. Therefore, the nature and reaction kinetics for the overall signal transduction process associated with plant CRYs, including potential rearrangement of the CCT domain still remains unclear.

To reveal the kinetics and mechanism of signal transduction in *AtCRY1*, it is essential to monitor time-resolved changes in protein conformation and/or intermolecular interactions. To date, the TA method has been one of the main techniques to monitor reaction kinetics of photoreceptor proteins. However, because structural changes affecting absorption are generally localized to the vicinity of the chromophore, this method is not suitable for monitoring the conformational changes elsewhere in the protein. To observe time-resolved conformational dynamics throughout the protein, we complemented TA with time-resolved detection of changes in diffusion by the pulsed laser induced transient grating (TG) technique ²¹⁻²⁹. The unique advantage of this method is its possibility to detect changes in molecular diffusion coefficients (*D*) during chemical reactions with a very short time-resolution. *D* is a valuable and important physical property to characterize a molecule in solution. This value is

sensitive to not only the protein conformational changes, but also intermolecular interaction between the protein and the solvent. If the friction between the diffusing protein and the solvent molecules increases, D should decrease. In fact, it was reported that D of a protein molecule in the aqueous solution is largely decreased upon the unfolding ²¹. Therefore, time-resolved measurement of D by the TG technique provides a powerful way to study the kinetics of both spectrally active and spectrally silent changes in molecular conformation and/or intermolecular interactions. Conformation changes observed by the TG method so far are consistent with results from other methods and kinetics of the changes has been revealed by this method; e.g., N-terminal helix unfolding of PYP ²², (consistent with circular dichroism (CD) study ³⁰), N-terminal helix formation of phytochrome ²³ (consistent with the time-resolved CD data ³¹), J α -helix unfolding of the linker region of phototropins LOV2 domain ^{24,25} (consistent with CD ³² and NMR ³³ experiments), and so on. These previous studies support the validity of the TG method.

In this study, we investigated the photochemical reactions of full-length *AtCRY1* using the TG method. We successfully detected large conformational change of full-length *AtCRY1* as the change in D and determined its kinetics. This D -change was not observed in the *AtCRY1* W324F mutant protein carrying a single mutation in the Trp triad of the N-terminal PHR-like domain. We also found that the conformational change in truncated *AtCRY1* lacking the CCT domain is very small. These observations clearly indicate that both the PHR-like and the CCT domains are responsible for the overall conformational change of plant CRYs. Thus light-dependent modulation between these two domains would likely be critical for activation of the receptor and subsequent biological activity by CRYs.

Results and discussion

Detection of conformational change in full-length *AtCRY1*

The TG signal of the full-length *AtCRY1* was measured in buffer solution at 12 °C at a protein concentration of 80 μM, and the signal at a grating wavenumber of $q^2 = 4.1 \times 10^{11} \text{ m}^{-2}$ is depicted in Fig. 1a. In principle, the TG signal intensity is proportional to the square of the photo-induced refractive index change (δn), which mainly derives from thermal energy releasing (thermal grating) and the creation and depletion of the chemical species (species grating)³⁴⁻³⁶. The observed TG signal (Fig. 1a) mainly consisted of two components; after the pulsed laser light irradiation, the TG signal quickly rose followed by a decay to the baseline within a few microseconds, and then by a rise-decay profile in the hundreds millisecond time range. If the kinetics of the signal depends on grating wavenumbers (q^2), this phase should be attributed to a diffusion process. If not, it should be due to reaction kinetics. For the assignment of the signal components, the TG signals of the *AtCRY1* sample were measured at various q^2 (Fig. 1b). The rate of the initial decay indeed depended on q^2 (Fig. 1b (inset)). The dependence indicates that this phase represent diffusion process, not reaction kinetics. Furthermore, the profile was expressed by an exponential function and the rate constant agreed well with the rate of the thermal diffusion $D_{th}q^2$ (D_{th} : thermal diffusivity), which was determined independently from the TG signal of the calorimetric reference sample (BCP) under the same condition. Hence this initial decay component was clearly attributed to the thermal grating signal ($\delta n_{th}(t)$).

$$\delta n_{th}(t) = \delta n_{th} \exp(-D_{th}q^2t) \quad (1)$$

where the δn_{th} is the initial refractive index change of the thermal grating signal. The time scales of the last rise-decay curve are also q^2 -dependent (Fig. 1 b). This phase indicates a protein diffusion process (diffusion signal: $\delta n_P(t) - \delta n_R(t)$), which reflects the decay process of the species grating signal due to the reactant (δn_R) and product (δn_P) by their diffusion processes. If we ignore the time-dependence of the molecular diffusion coefficient D , the

diffusion signal should be expressed as a following bi-exponential function^{37,38}.

$$\delta n_P(t) - \delta n_R(t) = \delta n_P \exp(-D_P q^2 t) - \delta n_R \exp(-D_R q^2 t) \quad (2)$$

where the D_R and D_P are the molecular diffusion coefficients of the reactant and product, respectively. The rise-decay feature of the observed diffusion signal indicates two diffusion components having opposite signs of δn (eq. (2)). Because the thermal grating signal had decayed to the baseline before the appearance of the diffusion signal and the sign of δn_{th} is negative at this temperature, the signs of δn for the rise and decay components were unambiguously determined to be negative and positive, respectively. The rate of the rise component is faster than that of the decay component, indicating that D was decreased by the photoreaction. D is a physical property that reflects not only the molecular size but also the intermolecular interactions including the large conformational rearrangement. The significant reduction in D following photoexcitation will be discussed later.

What are the kinetics of this D -change? The kinetics can be obtained from the q^2 dependence of the signal. In Fig. 1b, the TG signals are normalized by the thermal grating intensity, which is a measure of the concentration of photoexcited molecules. If the D -change completes before the time scale of the diffusion signal, the diffusion signal should not depend on q^2 . Contrary to this, the diffusion signal in a fast time scale is weak and its intensity dramatically increased at longer time (Fig. 1b). This time dependence of the diffusion signal implies that the D -change reaction of *A*tCRY1 took place within this observation time. Qualitatively, this change can be explained as follows: when the protein diffusion rate (Dq^2) is more rapid than its rate of D -change reaction (i.e, at a large q^2 condition), the diffusion coefficients of the reactant and product are similar and the diffusion signal should be weak, due to the cancellation of refractive index changes of the product and the reactant (δn_P and δn_R , respectively) of eq. (2). With decreasing q^2 , the diffusion signal appeared in a longer time region, and D_P also decreases significantly. Hence, the diffusion signal intensity

increased with decreasing q^2 . A similar time dependent feature of the diffusion signal was observed previously for other photo-sensor proteins²⁴⁻²⁹ and the time-dependence was analyzed by the following two state model^{37,38}.

$$R \xrightarrow{h\nu} I \xrightarrow{k} P$$

$$\delta n_P(t) - \delta n_R(t) = \left(\delta n_I - \frac{k}{(D_I - D_P)q^2 + k} \delta n_P \right) \exp\{-(D_I q^2 + k)t\} + \frac{k}{(D_I - D_P)q^2 + k} \delta n_P \exp(-D_P q^2 t) - \delta n_R \exp(-D_R q^2 t) \quad (3)$$

where R, I, P and k represent the reactant, an initial product (intermediate), the final product and the rate constant of the change, respectively. Furthermore, δn_I and D_I are the refractive index changes due to the creation of the I species and the diffusion coefficient of the intermediate, respectively.

The diffusion signals for *AtCRY1* were analyzed by the time-dependent model above. We first determined D_R and D_P , respectively, from the diffusion signal in the smallest q^2 , i.e., in the long time scale where the D -change reaction should be almost completed. We fitted the slower part (> 2 s) of the diffusion signal at the smallest q^2 by a bi-exponential function to obtain $D_R = 1.46 (\pm 0.05) \times 10^{-11} \text{ m}^2 \text{ s}^{-1}$ and $D_P = 0.62 (\pm 0.03) \times 10^{-11} \text{ m}^2 \text{ s}^{-1}$. Using these values and $D_I = D_R$, the observed TG signals at all q^2 can be reproduced well based on eq. (3). The time constant (k^{-1}) for the D -change was determined to be $k^{-1} = 0.4 (\pm 0.1)$ s.

Between the time scales for the thermal grating and diffusion signals, another weak species grating signal was observed (Fig. 1c, arrows). The precise determination of its rate constant was difficult, because of the interference from the thermal grating and the diffusion signal contributions. We analyzed this weak species grating component as follows. The total TG signal of *AtCRY1* should be fitted by the following equation, which includes thermal grating contribution $\delta n_{th}(t)$, protein diffusion contribution $\delta n_P(t) - \delta n_R(t)$, and weak species grating contributions.

$$I_{TG}(t) = \alpha \left[\delta n_{th}(t) + \sum_i a_i \exp(-k_i t) + \delta n_P(t) - \delta n_R(t) \right]^2 \quad (4)$$

The thermal and molecular diffusion contributions are given by eq. (1) and eq. (3), respectively. Additional exponential terms were added to the equation to reproduce weak species grating contributions. We conducted single exponential and bi-exponential analyses for the species grating signal (eq. (4) with $i=1$ and $i=2$). Fig.2 depicts the fitting results by single exponential (solid curve) and bi-exponential fitting (broken curve). During the fitting process, we used $D_{th}q^2$ determined by the decay rate of the signal of the calorimetric reference sample. We also used the parameters for the protein diffusion terms, which were determined in the diffusion signal analysis above. Apparently, the signal was reproduced well in the bi-exponential analysis. We could determine the time constants of 1 ms and 7.5 ms for this weak species signal. According to the previous TA studies¹⁴, absorption changes showed the creation of the radical form of *AtCRY1* after the photoexcitation. Following radical formation, a 1 ms phase (lifetime of 1.4 ms) was observed and attributed to the intra-protein electron transfer from Trp[•] to Tyr. Therefore, the 1 ms dynamics observed by the TG method is attributed to the population grating signal reflecting the absorption change involved in this electron transfer reaction. This flavin radical observed by TA returned to the original ground state with half lifetimes of 5 ms and >100 ms (corresponding to time constants of 7.2 ms and >140 ms). Therefore, we attributed the 7.5 ms dynamics observed in the TG signal to this dark reversion process of the chromophore from the radical form to the ground state. (In the present study, the phase of >100 ms was not observed, probably because this component was masked by the strong diffusion signal.) The fact that these time constants are similar to those reported in previous study¹⁴ clearly indicates that the full-length *AtCRY1* sample was stable and exhibited a proper photoreaction under our experimental conditions. Remarkably, even after the 7.5 ms decay, the TG signal did not disappear but showed the strong diffusion signal. This fact indicates that the light-induced conformational changes in the protein were retained

even after the recovery of the chromophore.

Assignment of origin of the molecular diffusion coefficient change in AtCRY1.

To date, reported photo-induced decreases in molecular diffusion coefficient D of photosensor proteins have been classified as originating from either oligomer formation²⁷⁻²⁹ or conformational change²⁴⁻²⁶. Those are referred as diffusion sensitive conformation change (DSCC). It seems rather straightforward to conclude that an oligomization reaction of AtCRY1 might cause the D reduction upon photoreaction, because, according to the Stokes-Einstein relationship, D is inversely proportional to the radius of a molecule. If the difference in D between the reactant and product ($D_P/D_R = 2.3$) was interpreted only in terms of the difference in molecular radius, the molecular volume of the product would be $(2.3)^3 = 12$ times larger than that of the reactant. (Although D -value depends on the shape of molecule, the dependence is rather weak according to the theoretical equation for the molecule with a non-spherical shape³⁹.) However, D is determined not only by the molecular size, but also by the intermolecular interaction between the protein and solvent molecules, therefore, an increase in protein/solvent intermolecular interaction must be also taken into account. If protein conformational changes produce an increase in protein/solvent intermolecular interactions, then D would be reduced too. To distinguish between these two possibilities, the concentration dependence of the signal profile should be examined. If this D -change is caused by oligomer formation, the reaction rate for the D -change would be sensitive to protein concentration, and this should result in a concentration dependent time profile of the signal. Photosensor proteins, BLUF (sensors of blue light using FAD) domains of AppA, YcgF, and LOV2 (light oxygen voltage sensing) domains of phototropins all exhibit concentration-dependent TG signals²⁷⁻²⁹. In contrast, if intramolecular protein conformational change is directly responsible for the D -change, this reaction rate should be

independent of the protein concentration, and the time profile of the TG signal should not depend on the concentration, except for the absolute intensity.

Therefore, we examined the TG signal at various sample concentrations at relatively large q^2 condition ($4.3 \times 10^{11} \text{ m}^{-2}$) (Fig. 3a and 3b). At this grating wavenumber, the diffusion signal appears in several 100 milliseconds time range, which covers the D -change ($< 1 \text{ s}$). Hence, if the rate of the D -change depends on the concentration, the profile of the signal should be sensitive to the concentration. Fig. 3a depicts the concentration dependence of the TG signal. Naturally, the signal intensity decreased with decreasing concentration. However, the time profile normalized at the peak intensity did not depend on the concentration except for a small temporal shift. (This shift could be due to a viscosity change by the dilution of the solution.) Indeed, by fitting the present signals with the present model, we determined that D_P and D_R increased slightly with decreasing the concentration of the protein ($D_R = 1.46 \times 10^{-11} \text{ m}^2 \text{ s}^{-1}$ and $D_P = 0.62 \times 10^{-11} \text{ m}^2 \text{ s}^{-1}$ at $80 \text{ } \mu\text{M}$, and $D_R = 1.59 \times 10^{-11} \text{ m}^2 \text{ s}^{-1}$ and $D_P = 0.68 \times 10^{-11} \text{ m}^2 \text{ s}^{-1}$ at $39 \text{ } \mu\text{M}$), but the rate constant for the D -change reaction did not change. Since the rate constant is concentration-independent, the origin of D -change is due to a significant and large conformational change of AtCRY1 protein, rather than light-dependent aggregation or oligomerization.

Furthermore, this large D -change cannot be explained by only one or two hydrogen bond rearrangements between protein residues and water, because of the following reason. Previously, we estimated an increase of the excess friction by unfolding of α -helix to be $0.3 \times 10^{-12} \text{ kg/s}$ ⁴⁰. (Here, the friction coefficient of species i is defined by $f_i = k_B T / D_i$ ($i = P$ or R), where k_B is the Boltzmann constant and T is the temperature. The change in the friction Δf , upon the transformation from the reactant to the product is calculated by $\Delta f = f_P - f_R$.) In the present AtCRY1 case, the increase of friction was calculated to be $3.75 \times 10^{-10} \text{ kg/s}$. This value is very large. If Δf of PYP⁴⁰ is used and only unfolding occurs for this conformation change,

this increase means about 100 residues are exposed to the buffer. Although this number is just a rough estimation of the extent of the conformation change, the significant reduction in D observed here, at least, implies that a large conformational rearrangement occurs after photoexcitation of *AtCRY1*.

Photoreduction-induced conformational change in *AtCRY1*

All structurally characterized members of the CRY/PHR family including *AtCRY1* conserve a common motif consisting of a chain of three Trp residues extending from the FAD cofactor to the protein surface^{6,20,41}. In *E. coli* PHR, this Trp chain is responsible for electron transfer to FAD during photoactivation (photoreduction of the oxidized form of PHR). After a single photo-induced electron transfer step from a nearby Trp reduces the FAD, the resulting radical state on the Trp is subsequently transferred to the terminal Trp (Trp306 in *E. coli* PHR). Because of the high structural conservation, a similar electron transfer mechanism has been assumed for CRYs^{6,14,42}. TA and FTIR studies on *AtCRY1* also suggested the involvement of an aromatic residue (Trp or Tyr) in the photoexcitation reaction^{14,15}. To decipher correlations between electron transfer and the observed conformational change, we made TG measurements for an *AtCRY1* mutant (W324F) lacking the terminal surface-exposed Trp residue of the proposed electron transfer pathway (Fig. 4). The W324F mutant shows almost the same absorption spectrum as wild-type protein, but shows no light-induced absorption change, indicating the mutant has no photoreduction reactions⁴³. For the W324F mutant, we detected no diffusion signal, although the thermal grating signals for the mutant and the wild-type were similar. These results indicate that the D -change reaction is likely linked to the electron transfer event in *AtCRY1*. Thus, the conformational change observed in the full-length wild-type protein is triggered in the N-terminal PHR-like domain.

Contribution of the CCT domain to the conformational change

Cryptochrome proteins are generally larger than photolyases, due to C-terminal extension of their sequences. In particular, plant CRYs have an additional characteristic C-terminal domain, named the Cryptochrome C-terminal (CCT) domain. The plant CCT domains have been implicated in signal transduction and shown to function as a scaffold for other proteins^{12,13}. NMR studies coupled with proteolysis indirectly suggested that some conformational change occurs in the CCT domain upon light irradiation¹¹. To directly examine and distinguish light-induced conformational changes in the N- and C-terminal domains of *AtCRY1*, we measured TG signals of truncated *AtCRY1* protein lacking the CCT domain (*trAtCRY1*) under the same experimental conditions as full-length *AtCRY1* (Fig. 4). In contrast to the full-length protein, which shows the characteristic large diffusion signal, *trAtCRY1* does not show this component, but only a very weak diffusion signal. These observations implicate that major component of the DSCC in *AtCRY1* is derived from the CCT domain.

Using a binding assay coupled with proteolysis for *AtCRY1*, the PHR-like domain was shown to interact with the CCT domain in the dark, inducing stable tertiary structure in the C-terminal domain¹¹. This interaction was destabilized in the light due to some conformational change in the CCT domain¹¹. We have reproduced these proteolysis experiments using a slightly modified protocol using the current *AtCRY1* sample and a high intensity blue LED to promote greater photoreduction (Fig. 5). Arrows indicate bands that are differentially sensitive to proteolysis in dark and light treated samples. At earlier time points, the upper indicated band is significantly more stable in dark as compared to light samples even after double the time of digestion, consistent with previous reports¹¹. At longer digestion times lower bands appear in the dark sample but not in the irradiated sample,

indicative of further differential cleavage in the digestion products. This differential resistance to proteolysis in light and dark samples indicates considerable changes in protein surface properties and susceptibility to cleavage as a result of irradiation, indicating that the light-induced structural changes are likely to be quite large. Furthermore, a proteolysis experiment for trAtCRY1 did not show any significant difference between dark and light illuminated samples (data not shown). Taken together, these results confirm the large conformational change directly observed by the TG method and are all consistent with the following interpretation; in the dark, inter-domain interactions between the PHR-like and CCT domains produce a relatively packed overall structure. Light excitation and reduction of FAD in the N-terminal PHR-like domain would affect the adjacent Trp triad and could propagate to the interface between the PHR-like and CCT domains, resulting in their dissociation. This dissociation would increase the surface area between AtCRY1 and the solvent, producing a *D*-value reduction, and simultaneously would expose interaction sites for other molecules, thereby activating signals in the photomorphogenesis pathways. The conformational changes in the CCT domain have been suggested to be important for a long time ¹². However, there has been no information on the dynamics of the proposed conformational changes. Here, our time-resolved measurement revealed not only the existence of the conformational change in the CCT domain but also the rate of the conformational change for the first time.

Interestingly, a weak diffusion signal with a small *D*-change was observed in trAtCRY1 lacking the CCT domain (Fig. 4b (blue)). The observation is consistent with a small conformational change in the vicinity of the FAD binding site, as previously observed by vibrational spectral analysis ¹⁵. TG signals are sensitive to both spectral changes and conformational changes that are spectrally silent. Therefore, the relatively large DSCC observed here is not conflicted with the conclusions from FTIR studies ¹⁵, because the

diffusion coefficient monitored by TG techniques would be sensitive to dissociation of the PHR-like and CCT domains. We propose that PHRs undergo relatively small conformational changes upon light illumination, whereas CRYs have evolved their C-terminal extensions to increase dynamics for signaling. The schematic illustrations of the possible reactions are shown in Fig. 6.

Conclusions

This work for the first time demonstrates a significant light-dependent conformational change in a CRY that may provide the basis for signaling. Although the suggestion that conformational changes in CRYs may provide a mechanism for signaling has been made since many years⁴¹, direct experimental evidence for such a dynamic transformation has been missing. Here we report a significant conformational change in full-length *AtCRY1* that is dependent on light activation of the N-terminal PHR-like domain through flavin reduction by intra-protein electron transfer involving conserved Trp residues. We further show that the C-terminal CCT domain is critical to this conformational change subsequent to the initial light activation event. We propose that electron transfer from the flavin chromophore triggers the dissociation between the CCT and PHR-like domain in the protein, which are connected to each other to form a stable tertiary structure by inter-domain interaction in the dark state. This dissociation leads to exposure of the previously buried interfaces, and the consequent increase in interaction between *AtCRY1* and solvent, resulting in the observed large *D*-change. These events associated with light-induced conformational changes most likely represent essential processes that underlie *AtCRY1* signaling in plants.

Materials and methods

Measurement

The experimental setup for the TG experiment was similar to conditions reported for other photoreceptors in previous papers^{44,45}. For excitation of *AtCRY1*, a XeCl excimer laser-pumped dye laser beam (Lamda Physik Compex 102xc, Lumonics Hyper Dye 300; $\lambda = 465$ nm) was used. The time-resolution of the instrument was ~ 20 ns. The laser beam was split into two by a beam splitter and crossed inside a quartz sample cell (optical path-length = 2 mm). The spot size of the excitation beams at the sample position was ca. 1 mm diameter. The created refractive index modulation in the sample was probed by a diode laser (835 nm) as a Bragg diffracted signal (TG signal). The TG signal was detected by a photomultiplier tube (Hamamatsu R1477), and fed into a digital oscilloscope (Tetronix 2430A). The TG signals were transferred to a microcomputer and averaged 6-36 pulse to improve a signal to noise (S/N) ratio. After every shot of the excitation pulse, the sample solution (200 μ L) was stirred so that the photo-product was not excited again. The grating wavenumber (q^2) was determined with the decay rate of the thermal grating signal of a calorimetric reference solution; bromocresol purple (BCP) which gives rise to only the thermal grating signal due to the nonradiative transition within the pulse width of the excitation laser. The various grating numbers were adjusted by changing the crossing angle of the excitation pulses and the probe beam. All measurements were carried out at room temperature (12°C).

In the proteolysis measurement, samples were 100 μ L of 10 μ M *AtCRY1* in 50 mM Tris pH 7.5, 10 mM DTT, 20% glycerol and 2 μ M trypsin placed in a 1 x 10mm glass cuvette. In the blue light condition, samples were irradiated on ice by a blue LED at 150 mmolm²sec⁻¹ through an IR filter to eliminate heating effects and temperature was continuously checked in the immediate vicinity of the cuvette throughout the whole course of the experiment. A 10 minute pre-irradiation was carried out before the addition of trypsin (T0). Aliquots were removed at T0 (before addition of trypsin), 0, 5, 10, 15, 20, 30, 45 and 60 minutes after addition of trypsin, boiled in SDS sample buffer and resolved on 13% acrylamide SDS gels.

In the dark condition, samples were harvested under the identical conditions and time points as above but without undergoing irradiation.

Sample preparation

AtCRY1 was isolated from a recombinant baculovirus expression system and purified by nickel column affinity chromatography as described previously⁴⁶. The W324F mutant of *AtCRY1* was also prepared as reported previously⁴³. For the truncated proteins, gene coding photolyase-like domain¹⁰ was inserted into pGEX-6P-1. The protein lacking the CCT domain was expressed in *E. coli* and purified with Glutathione Sepharose followed by Mono Q and Sephacryl S-300 after tag cleavage. The proteins were dissolved into the buffer solution containing 50 mM Tris-HCl, 500 mM NaCl, and 30 % (v/v) glycerol (pH 7.5). Dusts in the sample solution were removed with a centrifuge before measurement. The TG signal was measured mostly at a concentration of 80 μ M. Dependence on the concentration was examined in the concentration range from 39 μ M to 80 μ M.

Acknowledgements

This work was supported by the Grant-in-Aid for Scientific Research (A) (No.18205002), Grant-in-Aid for Scientific Research on Innovative Areas (20107003), and Scientific Research on Innovative Areas (Research in a proposed research area) (20107003) from the Ministry of Education, Science, Sports and Culture in Japan (to M.T.), NSF grant 0848311 (to M.A.) and National Institutes of Health grant GM37684 (to E.D.G.). M.K. was supported by research fellowship of Global COE program, International Center for Integrated Research and Advanced Education in Material Science, Kyoto University, Japan. K.H. was supported by The Skaggs Institute for Chemical Biology.

References

1. Cashmore, A. R., Jarillo, J. A., Wu, Y. & Liu, D. (1999). Cryptochromes: Blue light receptors for plants and animals. *Science* **284**, 760-765.
2. Ahmad, M. & Cashmore, A. R. (1993). HY4 gene of *A. thaliana* encodes a protein with characteristics of a blue-light photoreceptor. *Nature* **366**, 162-166.
3. Hsu, D. S., Zhao, X., Zhao, S., Kazantsev, A., Wang, R., Todo, T., Wei, Y. & Sancar, A. (1996). Putative Human Blue-Light Photoreceptors hCRY1 and hCRY2 Are Flavoproteins. *Biochemistry* **35**, 13871-13877.
4. Song, S., Öztürk, N., Denaro, T. R., Arat, N. Ö., Kao, Y., Zhu, H., Zhong, D., Reppert, S. M. & Sancar, A. (2007). Formation and Function of Flavin Anion Radical in Cryptochrome 1 Blue-Light Photoreceptor of Monarch Butterfly. *J. Biol. Chem.* **282**, 17608-17612.
5. Gegear, R. J., Casselman, A., Waddell, S. & Reppert, S. M. (2008). Cryptochrome mediates light-dependent magnetosensitivity in *Drosophila*. *Nature* **454**, 1014-1018.
6. Brudler, R., Hitomi, K., Daiyasu, H., Toh, H., Kucho, K., Ishiura, M., Kanehisa, M., Roberts, V. A., Todo, T., Tainer, J. A. & Getzoff, E. D. (2003). Identification of a New Cryptochrome Class: Structure, Function, and Evolution. *Molecular Cell* **11**, 59-67.
7. Sancar, A. (2008). Structure and Function of Photolyase and in Vivo Enzymology: 50th Anniversary. *J. Biol. Chem.* **283**, 32153-32157.
8. Losi, A. (2007). Flavin-based Blue-light Photosensors: A Photobiophysics Update. *Photochemistry and Photobiology* **83**, 1283-1300.
9. Sancar, A. Structure and Function of DNA Photolyase and Cryptochrome Blue-Light Photoreceptors. (2003). *Chem. Rev.* **103**, 2203-2238.
10. Brautigam, C. A., Smith, B. S., Ma, Z., Palnitkar, M., Tomchick, D. R., Machius, M. & Deisenhofer, J. (2004). Structure of the photolyase-like domain of cryptochrome 1

- from *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. USA* **101**, 12142-12147.
11. Partch, C. L., Clarkson, M. W., Özgür, S., Lee, A. L. & Sancar, A. (2005). Role of Structural Plasticity in Signal Transduction by the Cryptochrome Blue-Light Photoreceptor. *Biochemistry* **44**, 3795-3805.
 12. Yang, H., Wu, Y., Tang, R., Liu, D., Liu, Y. & Cashmore, A. R. (2000). The C Termini of Arabidopsis Cryptochromes Mediate a Constitutive Light Response. *Cell* **103**, 815-827.
 13. Yang, H., Tang, R. & Cashmore, A. R. (2001). The Signaling Mechanism of Arabidopsis CRY1 Involves Direct Interaction with COP1. *The Plant Cell* **13**, 2573-2587.
 14. Giovani, B., Byrdin, M., Ahmad, M. & Brettel, K. (2003). Light-induced electron transfer in a cryptochrome blue-light photoreceptor. *Nat. Struct. Biol.* **10**, 489-490.
 15. Kottke, T., Batschauer, A., Ahmad, M. & Heberle, J. (2006). Blue-Light-Induced Changes in Arabidopsis Cryptochrome 1 Probed by FTIR Difference Spectroscopy. *Biochemistry* **45**, 2472-2479.
 16. Banerjee, R., Schleicher, E., Meier, S., Munoz, V. R., Pokorny, R., Ahmad, M., Bittl, R. & Batschauer, A. (2007). The Signaling State of Arabidopsis Cryptochrome 2 Contains Flavin Semiquinone. *J. Biol. Chem.* **282**, 14916-14922.
 17. Langenbacher, T., Immeln, D., Dick, B. & Kottke, T. (2009). Microsecond light-induced proton transfer to flavin in the blue light sensor plant cryptochrome. *J. Am. Chem. Soc.* **131**, 14274-14280.
 18. Li, Q. & Yang, H. (2007). Cryptochrome signaling in plants. *Photochem. Photobiol.* **83**, 94-101.
 19. Bouly, J. P., Schleicher, E., Dionisio-Sese, M., Vandenbussche, F., Van Der Straeten, D., Bakrim, N., Meier, S., Batschauer, A., Galland, P., Bittl, R. & Ahmad, M. (2007).

- Cryptochrome Blue Light Photoreceptors Are Activated through Interconversion of Flavin Redox States. *J. Biol. Chem.* **282**, 9383-9391.
20. Aubert, C., Vos, M. H., Mathis, P., Eker, A. P. M. & Brettel, K. (2000). Intraprotein radical transfer during photoactivation of DNA photolyase. *Nature* **405**, 586-590.
 21. Inoue, K., Baden, N. & Terazima, M. (2005). Diffusion coefficient and the Secondary Structure of Poly-L-glutamic Acid in Aqueous solution. *J. Phys. Chem. B* **109**, 22623-22628.
 22. Hoshihara, Y., Imamoto, Y., Kataoka, M., Tokunaga, F. & Terazima, M. (2008). Conformational Changes in the N-Terminal Region of Photoactive Yellow Protein: A Time-Resolved Diffusion Study. *Biophys. J.* **94**, 2187-2193.
 23. Eitoku, T., Zarate, X., Kozhukh, G. V., Kim, J. I., Song, P. S. & Terazima, M. (2006). Time-Resolved Detection of Conformational Changes in Oat Phytochrome A: Time-Dependent Diffusion. *Biophys. J.* **91**, 3797-3804.
 24. Nakasone, Y., Eitoku, T., Matsuoka, D., Tokutomi, S. & Terazima, M. (2007). Dynamics of Conformational Changes of Arabidopsis Phototropin 1 LOV2 with the Linker Domain. *J. Mol. Biol.* **367**, 432-442.
 25. Eitoku, T., Nakasone, Y., Matsuoka, D., Tokutomi, S. & Terazima, M. (2005). Conformational Dynamics of Phototropin 2 LOV2 Domain with the Linker upon Photoexcitation. *J. Am. Chem. Soc.* **127**, 13238-13244.
 26. Tanaka, K., Nakasone, Y., Okajima, K., Ikeuchi, M., Tokutomi, S. & Terazima, M. (2009). Oligomeric-State-Dependent Conformational Change of the BLUF Protein TePixD (Til0078). *J. Mol. Biol.* **386**, 1290-1300.
 27. Nakasone, Y., Ono, T., Ishii, A., Masuda, S. & Terazima, M. (2007). Transient Dimerization and Conformational Change of a BLUF Protein: YcgF. *J. Am. Chem. Soc.* **129**, 7028-7035.

28. Hazra, P., Inoue, K., Laan, W., Helligwerf, K. J. & Terazima, M. (2006). Tetramer Formation Kinetics in the Signaling State of AppA Monitored by Time-Resolved Diffusion. *Biophys. J.* **91**, 654-661.
29. Nakasone, Y., Eitoku, T., Matsuoka, D., Tokutomi, S. & Terazima, M. (2006). Kinetic Measurement of Transient Dimerization and Dissociation Reactions of Arabidopsis Phototropin 1 LOV2 Domain. *Biophys. J.* **91**, 645-653.
30. Harigai, M., Imamoto, Y., Kamikubo, H., Yamazaki, Y. & Kataoka, M. (2003). Role of an N-terminal loop in the secondary structural change of photoactive yellow protein. *Biochemistry* **42**, 13893-13900.
31. Chen, E., Lapko, V. N., Song, P. S. & Kliger, D. S. (1997). Dynamics of the N-Terminus α -Helix Unfolding in the Photoreversion Reaction of Phytochrome A. *Biochemistry* **36**, 4903-4908.
32. Corchnoy, S. B., Swartz, T. E., Lewis, J. W., Szundi, I., Briggs, W. R. & Bogomolni, R. A. (2003). Intramolecular Proton Transfers and Structural Changes during the Photocycle of the LOV2 Domain of Phototropin 1. *J. Biol. Chem.* **278**, 724-731.
33. Harper, S. M., Christie, J. & Gardner, K. H. (2004). Disruption of the LOV-J α helix interaction activates phototropin kinase activity. *Biochemistry* **43**, 16184-16192.
34. Terazima, M. (1998). Photothermal studies of photophysical and photochemical processes by the transient grating method. *Adv. Photochem.* **24**, 255-338.
35. Terazima, M. & Hirota, N. (1993). Translational diffusion of a transient radical studied by the transient grating method; pyrazinyl radical in 2-propanol. *J. Chem. Phys.* **98**, 6257-6262.
36. Hara, T., Hirota, N. & Terazima, M. (1996). A new application of the transient grating method to a photochemical reaction; the entropy, reaction volume and partial molar volume measurements. *J. Phys. Chem.* **100**, 10194-10200.

37. Nishida, S., Nada, T. & Terazima, M. (2004). Kinetics of intermolecular interaction during protein folding of reduced cytochrome c. *Biophys. J.* **87**, 2663-2675.
38. Terazima, M. (2006). Diffusion coefficients as a monitor of reaction kinetics of biological molecules. *Phys. Chem. Chem. Phys.* **8**, 545-557.
39. Cussler, E. L. (2009). Diffusion (3rd edition) chap. 5. pp129, Cambridge University Press.
40. Khan, J. S., Imamoto, Y., Harigai, M., Kataoka, M. & Terazima, M. (2006) Conformational changes of PYP monitored by diffusion coefficient: Effect of N-terminal α -helices. *Biophys. J.* **90**, 3686-3693.
41. Busza, A., Emery-Le, M., Rosbash, M. & Emery, P. (2004). Roles of the Two Drosophila CRYPTOCHROME Structural Domains in Circadian Photoreception. *Science* **304**, 1503-1506.
42. Biskup, T., Schleicher, E., Okafuji, A., Link, G., Hitomi, K., Getzoff, E. D. & Weber, S. (2009) Direct Observation of a Photoinduced Radical Pair in a Cryptochrome Blue-Light Photoreceptor. *Angew. chem. int. ed.* **48**, 404-407.
43. Zeugner, A., Byrdin, M., Bouly, J., Bakrim, N., Giovani, B., Brettel, K. & Ahmad, M. (2005). Light-induced Electron Transfer in Arabidopsis Cryptochrome-1 Correlates with in Vivo Function. *J. Biol. Chem.* **280**, 19437-19440.
44. Terazima, M. & Hirota, N. (1991). Measurement of the quantum yield of triplet formation and short triplet lifetimes by the transient grating technique. *J. Chem. Phys.* **95**, 6490-6495.
45. Terazima, M. (2000). Is the Translational Diffusion of Organic Radicals Different from That of Closed-Shell Molecules? *Acc. Chem. Res.* **33**, 687-694.
46. Lin, C., Robertson, D. E., Ahmad, M., Raibekas, A. A., Jorns, M. S., Dutton, P. L. & Cashmore, A. R. (1995). Association of Flavin Adenine Dinucleotide with the

Arabidopsis Blue Light Receptor CRY1. *Science* **269**, 968-970.

Figure captions

Fig. 1. (a) TG signal of the full-length *AtCRY1* after blue-light excitation at 465nm in the buffer solution containing 50 mM Tris, 500 mM NaCl, and 30 % Glycerol (pH 7.5) at $q^2 = 4.1 \times 10^{11} \text{ m}^{-2}$ at the excitation wavelength of 465 nm (Red line). The signal was measured at a concentration of 80 μM , at 12 °C, and at an excitation laser power of $\sim 10 \text{ }\mu\text{J}$. (inset) Magnification of the signal in the time region of 2 μs - 40 ms. (b). TG signals of full-length *AtCRY1* at various q^2 values; $4.1 \times 10^{11} \text{ m}^{-2}$ (red), $1.9 \times 10^{11} \text{ m}^{-2}$ (orange), $7.3 \times 10^{10} \text{ m}^{-2}$ (green), and $3.5 \times 10^{10} \text{ m}^{-2}$ (blue). Signals were normalized by thermal grating intensity. (c) Magnification of the Figure (b) in the time region of 100 μs -150 ms. The q^2 -independent decay components were observed in the milliseconds time range (arrows). These components indicate reaction kinetics (absorption changes). The black broken curves in these figures show the best fitted curves by eq. (4).

Fig. 2. The comparison of single exponential (black solid curve) and bi-exponential (black dashed curve) analysis for the weak species grating components observed in the TG signal of full-length *AtCRY1* at $q^2 = 7.3 \times 10^{10} \text{ m}^{-2}$. Residual errors are also shown in the figure top by dark green solid curve (single exponential analysis) and green broken curve (bi-exponential analysis).

Fig. 3. (a) TG signals for the full-length *AtCRY1* at the protein concentration of 80 μM (red), 64 μM (orange), 50 μM (green), and 39 μM (blue) at $q^2 = 4.3 \times 10^{11} \text{ m}^{-2}$. The TG signals were normalized by the diffusion peak. (b) *AtCRY1* absorption spectra at matching concentrations.

Fig. 4. TG signal of the wild-type full-length (black), the CCT domain truncated (blue), and W324F mutant (red) of full-length *AtCry1* at $q^2 = 2.3 \times 10^{12} \text{ m}^{-2}$ under the same experimental

conditions. The signals were measured at protein concentrations of 50 μ M, at 12 °C.

Fig. 5. Partial proteolysis of the full-length *AtCRY1* in dark (left) and light (right) conditions. The bands showing differential proteolysis are indicated (arrows). The topmost band is less stable subsequent to illumination.

Fig. 6. The schematic illustrations of the proposed photoreaction of *AtCRY1*. In the dark, *AtCRY1* takes a relatively compact structure by inter-domain interactions. Blue light induces the reduction of FAD chromophore to the radical form ($\text{FADH}\cdot$) through the intra-protein electron transfer from the conserved tryptophan triad ¹⁴. The change in the PHR-like domain could propagate to the interface between the PHR-like and CCT domains, resulting in their dissociation. This dissociation would increase the surface area (colored in pink) between *AtCRY1* and the solvent, producing a reduction of the molecular diffusion coefficient and the protein stability to the protease. This simultaneously would expose interaction sites for other molecules, thereby activating signals in the photomorphogenesis pathways.

Fig.1

Fig.2

Fig.3

Fig.4

Fig.5

Fig.6