

Title	<Note> Chimpanzees in Bandafassi Arrondissement, Southeastern Senegal: Field Surveys as a Basis for the Sustainable Community-Based Conservation
Author(s)	Gašperšič, Maja; Pruetz, Jill D.
Citation	Pan Africa News (2011), 18(2): 23-25
Issue Date	2011-12
URL	<a href="http://hdl.handle.net/2433/152160">http://hdl.handle.net/2433/152160</a>
Right	Copyright © Pan Africa News.
Type	Article
Textversion	publisher

12. Fa JE, Juste J, Perez del Val J, Castroviejo J 1995. Impact of market hunting on mammal species in Equatorial Guinea. *Conserv Biol* **9**:1107–1115.
13. Salasfky N 2011. Integrating development with conservation. *Biol Conserv* **144**:973–978.
14. Tutin CEG, Parnell RJ, White LJT, Fernandez M 1995. Nest building by lowland gorillas in the Lopé Reserve, Gabon: environmental influences and implications for censusing. *Int J Primatol* **16**:53–76.
15. World Heritage (2011) *Convention Concerning the Protection of the World Cultural and Natural Heritage*. 35Com (<http://whc.unesco.org/en/decisions/4409>)
16. Vermeulen C, Julve C, Doucet J-L, Monticelli D 2009. Community hunting in logging concessions: Towards a management model for Cameroon's dense forests. *Biodivers Conserv* **18**:2705–2718.
17. Roe D, Nelson F, Sandbrook C (eds) 2009. *Community Management of Natural Resources in Africa: Impacts, Experiences and Future Directions*. IIED Natural Resource Issues No. 18.
18. Van Vliet N, Milner-Gulland EJ, Bousquet F, Saqalli M, Nasi R 2010. Effect of small-scale heterogeneity of prey and hunter distributions on the sustainability of bushmeat hunting. *Conserv Biol* **24**:1327–1337.
19. Sandbrook C, Roe D 2010. Linking conservation and poverty alleviation: the case of great apes. An overview of current policy and practice in Africa. *Conservation and Poverty Learning Group Report*.
20. Brooks TM, Wright J, Sheil D 2009. Evaluating the success of conservation actions in safeguarding tropical forest biodiversity. *Conserv Biol* **23**:1448–1457.
21. Nepal S, Spiteri A 2011. Linking livelihoods and conservation: An examination of local residents' perceived linkages between conservation and livelihoods benefits around Nepal's Chitwan National Park. *Environ Manage* **47**:727–738.
22. Williamson EA, Usongo L 1995. I. Survey of primate populations and large mammal inventory ; II. Survey of elephants, gorillas and chimpanzees; Reserve de Faune du Dja, Cameroun. *Projet ECOFAC—Composante Cameroun*.
23. Ferraro PJ, Patanayak SK 2006. Money for nothing? A call for empirical evaluation of biodiversity conservation investments. *PLOS Biol* **4**:482–488; e105.
24. Pullin AS, Knight TM 2009. Doing more good than harm—Building an evidence-base for conservation and environmental management. *Biol Conserv* **142**:931–934.
25. Sutherland WJ, Pullin AS, Dolman PM, Knight TM 2004. The need for evidence-based conservation. *Trends Ecol Evol* **19**(8):305–308.
26. Matthews E, Matthews A 2004. Survey of gorillas (*Gorilla gorilla gorilla*) and chimpanzees (*Pan troglodytes troglodytes*) in Southwestern Cameroon. *Primates* **45**:15–24.
27. Kühl H, Maisels F, Ancrenaz M, Williamson EA 2008. *Best Practice Guidelines for Surveys and Monitoring of Great Ape Populations*. Gland, Switzerland: IUCN SSC Primate Specialist Group.
28. Willie J 2006. *Contribution a l'évaluation de l'incidence de la chasse sur les populations de cephhalophes de la peripherie nord de la reserve de biosphere du Dja (Est-Cameroun)*. Master's thesis, University of Dschang, Cameroon.
29. Campbell G, Kuhl H, N'Goran Kouame P, Boesch C 2008. Alarming decline of West African chimpanzees in Cote d'Ivoire. *Curr Biol* **18**:R903/04.
30. Kormos R, Boesch C, Bakarr MI, Butynski TM 2003. *West African Chimpanzees: Status Survey and Conservation Action Plan*. IUCN/SSC Primate Specialist Group, Gland, Switzerland.
31. Stokes EJ, Strindberg S, Bakabana PC, Elkan PW, Iyenguet FC, et al. 2010. Monitoring great ape and elephant abundance at large spatial scales: Measuring effectiveness of a conservation landscape. *PLoS ONE* **5**:e10294.
32. Posa MRC, Diesmos AC, Sodhi NS, Brooks TM 2008. Hope for threatened tropical biodiversity: lessons from the Philippines. *BioScience* **58**:231–240.

## <NOTE>

# Chimpanzees in Bandafassi Arrondissement, Southeastern Senegal: Field Surveys as a Basis for the Sustainable Community-Based Conservation

Maja Gašperšič<sup>1</sup> & Jill D. Pruett<sup>2</sup>

<sup>1</sup> Department of Biology, University of Ljubljana, Slovenia

<sup>2</sup> Department of Anthropology, Iowa State University, USA  
(E-mail: [maja.g.cisse@gmail.com](mailto:maja.g.cisse@gmail.com))

## INTRODUCTION

The western chimpanzee (*Pan troglodytes verus*) is considered as one of the most threatened ape species, facing a dramatic decline over the last decade<sup>1,2</sup>. The latest conservation action plans classified Senegal as “an exceptionally important priority area” for chimpanzee protection, which demands immediate attention<sup>3</sup>. Chimpanzees have been expatriated from at least two African countries and IUCN estimated the Senegalese population to be almost extinct, numbering between 200 and 400<sup>4</sup>. Most apes range in small isolated communities in intense sympatry with local ethnic groups. Major threats include human encroachment, deforestation for crops, gold and iron digging, along with limited pet trade<sup>5</sup>. Additionally, this population lives at the northern edge of species' distribution, in extremely hot, dry and open savanna landscape that characterized an important transitional period in human evolution<sup>6</sup>. Chimpanzees though have a mythical relation with Senegalese people; therefore local folklore and taboos allows them to share space. This project is part of the investigation “Conservation of chimpanzees in south-eastern Senegal: the human element” supervised by JD Pruett. Initially, we identified ape communities in Bandafassi Arrondissement, their ranging patterns, key water and food sources, and particularly chimpanzees' relation to humans via an ethnoprimateological approach<sup>7</sup>.

Due to the importance of water sources with gallery forests, specific food-rich areas, and conflicts with humans encountered we focused on three priority field-sites: mountain slopes above Bandafassi village, where chimpanzees were known to attack goats; the stream of Angafou (12°34'N, 12°24'W), its rich gallery forest and mango orchards, being crop-raided by apes, supposedly after commercial harvesting of baobab and Saba fruits increased; and the surroundings of Nathia (12°29'N, 12°22'W), which was not identified in previous surveys<sup>2</sup> and shows no conflict between species. Two buffer-zones are surveyed sporadically to provide basic information about the presence of chimpanzees in the remaining forest patches between the Niokolo Koba national park and neighboring Guinea-Conakry. Several sites along Gambia River indicate various conflicts between chimpanzees and humans (e.g. palm-wine harvesting, raphia or bamboo-cutting, artisanal gold-digging sites). Chimpanzees are seen there only when people are not common, and it appears that their seasonal activity affects the ranging patterns of apes.

## METHODS

Habitat composition at study sites was sampled by walking several km in random directions and assessing the general habitat within a 50 m radius each 30 m. The classification of vegetation types was adapted after Bogart & Pruetz<sup>8</sup>. In March 2011 a phenology transect (1 km long, 10 m wide) was established at each study site to monitor tree production monthly throughout the year. Chimpanzee and human usage is assessed here on a total of 507 woody plants. Key food sources (e.g. *Adansonia digitata*, *Cola cordifolia*, *Parkia biglobosa*, etc.), and termite mounds and beehives were marked as well. We also noted encounters with large mammals and humans to estimate the presence of predators, competitors and potential prey-species for chimpanzees. Data were systematically collected upon sightings, nesting<sup>9</sup> and feeding sites, while signs of material culture<sup>10</sup> were recorded opportunistically. Diet composition was assessed through observations, feeding traces, and fecal-sample analyses<sup>11</sup>. Identifying ranging behavior included recording signs of chimpanzee activity and, if indices were positive (fresh debris or vocalizations), we tried to approach them.

## RESULTS

From April 2010 through June 2011 we conducted 292 field surveys and identified 13 villages along several dry season water sources associated with chimpanzees. In total we recorded 157 nesting sites, 85 feeding sites, several cases of tool-use (e.g. baobab cracking, termite-fishing, ant-dipping, tuber-digging), hand-dug wells and caves/shelters. Chimpanzees of Bandafassi were encountered more than 160 times, while data on demography, activity budget and reactions to observers stems from 146 observations at three study sites.

### Encounters with chimpanzees

Despite the rarity of closed-canopy habitat<sup>12</sup>, chimpanzees were most often observed in gallery and ecotone forest (38.3% and 0.8% respectively; see Figure 1), followed by different woodland types (closed 26.3%, open 19.1% and bamboo 9.2%), on plateaus (3.5%), in grassland and along fields (1.4% both types). These results indicate chimpanzees' preference for forested habitats, especially for nesting. This habitat type is also most heavily exploited by humans, particularly nomadic pastoralists<sup>13</sup>.

The average party size for various chimpanzee groups sighted was 5.8 (range 1–30). The largest daily parties were encountered at Nathia (mean 7.8), followed by Angafou (5.6), while the smallest groups ranged at Bandafassi (4.5). Foraging parties were on average composed of 2.4 males, 2.1 females, 1.6 juveniles, and 1.3 infants. The com-

munity at Angafou regularly nests along the stream and has indicated some acceptance of observers through time, while Nathia chimpanzees seem to have a vast home-range (> 70 km<sup>2</sup>) and use it seasonally.

As expected, apes were most often feeding (63.4% overall) upon sighting, foraging and traveling (17.9%), drinking (8.3%) resting (4.8%), nesting (2.1%) and other (in 3.4% mostly social behaviors). The most common reaction of chimpanzees in general was to leave the location (38% overall; see Figure 2), but positively they ignored us and indicated curiosity more often over time (24% and 20% respectively). However, important differences in reactions occurred among three sites with different human-chimpanzee relations. The apes indicated fear in only 14.3% of all episodes, but more significantly at Angafou. Nathia chimpanzees exhibited less fear than other communities and often ignored observers during prolonged contacts, while only apes at Bandafassi also displayed at human observers (4.3%).

### Competition with humans

So far, chimpanzees consumed fruits (73.2%), pods (7.6%), nuts (4.9%), pith (4.9%), bark (2.9%), leaves (1.5%; partly used for medicinal purposes) and flowers (0.4%) from 39 plant species of at least 17 families. Social insects (termites and weaver ants together in 2.6%), tubers (wild yam and unidentified underground storage organs), honey




Figure 1. Habitat composition/available vegetation types compared with chimpanzees' usage (nests, contacts)




Figure 2. Comparison of chimpanzee reactions to humans among three main sites

(each in 0.4% of cases) and traces of geophagy were detected in their food repertoire. Many ape food sources are also used by local humans for various purposes (Pruetz listed 17 species<sup>14</sup>), and at least five of their key feeding items are exploited commercially (e.g. *Adansonia digitata*, *Saba senegalensis*, *Tamarindus indica*, *Parkia biglobosa*, *Cola cordifolia*). Among tree species used for food or nest construction several were also cut by shepherds; e.g. *Bombax costatum*, *Khaya senegalensis*, *Acacia* spp., *Ficus* spp., and *Zizyphus* spp.<sup>13</sup>. Fulbe herders from North of the country were in fact most often encountered in the field, pruning trees for their large flocks of sheep (100–300), while harvesting of *Saba* fruits occurred as a key activity by local people in June.

### DISCUSSION: A Vision For The Sustainable Community-Based Conservation

Chimpanzees' responses to observers appear to be influenced by their experience with local people, who are the crucial element for their conservation<sup>15</sup>. Nature-culture tourism programs, including forest guards, could be an alternative solution to the complex relationship between the species<sup>16</sup>, while habituating the apes is not recommended<sup>17</sup>. The incentive could include local traditional customs related to primates (Bedik and Bassari initiation rites) and nature hiking with the opportunistic chance of chimpanzee viewing. This particular research project covers the widest geographical area in Senegal (> 500 km<sup>2</sup>), including several chimpanzee communities in rare forest patches. As of December 2011, five field assistants, serving as eco-rangers, monitor the most important chimpanzee sites. We established a limited "surveillance system" similar to J. Carter's in Guinea or further east in Senegal. Identified eco-guards monitor the ranging behavior of chimpanzees and at the same time limit crop-raiding and attacks on domestic animals. The two bordering-zones are important in terms of recognizing the existing forest-corridors available for possible gene-transfer within the population. It appears that with the continuous presence of researchers the apes ceased to capture goats, similarly as mango-raiding reportedly decreased. Encounters with shepherds, wine-collectors or hunters in the field are reported to local Eaux et forêts authorities. Preliminary estimates from surveys in Senegal by S Ndiaye in May 2011 are encouraging and indicate the Senegalese population is stronger than thought and numbers up to 500 chimpanzees<sup>18</sup>. In fact, two of our main study areas (Angafou and Nathia) were recognized as priority sites for chimpanzee conservation<sup>18</sup>. To conclude, this project has laid solid foundations for the sustainable community-based conservation in Bandafassi arrondissement. However, the prospects for the future co-existence of savanna chimpanzees and humans depend on needed support from governmental agencies as well as investment in terms of research and funding.

### ACKNOWLEDGEMENTS

First we want to thank Senegalese department of Eaux et forêts at the Ministry of environment and the Arrondissement du Bandafassi to give us the permission for research. The project was generously supported by the Slovenian Research Agency ARRS (Z6-3676). We are sincerely grateful to local

authorities and villagers for accepting our presence in their forests, USAID organization Wula Nafaa for initial funding of the project, *Fongoli savanna chimpanzee project* for logistics and Dondo Kanté for partly managing the project. Collaboration with Janis Carter and Souleye Ndiaye was invaluable. The research though would not be possible without dedicated field assistants / forest guardians: Gilbert Camara at Bandafassi mountains, Nambura Diallo at Nathia, Khadim Diallo at park borders, Samba Faye Kanté at Angafou, and Omar Diallo at Marewa (along with various tasks). Finally, special thanks goes to Seth Cissé for his overall help, data-collection and supervising the project in my absence.

### REFERENCES

1. Plumptre *et al.* 2010. Eastern chimpanzees (*Pan troglodytes schweinfurthii*): status survey and conservation action plan 2010-2020. IUCN Species Survival Commission.
2. Boesch C 2008. Why do chimpanzees die in the forest? The challenges of understanding and controlling for wild ape health. *Am J Primatol* **70**:722–726.
3. Kormos R, Boesch C, Bakarr MI, Butynski TM (eds) 2003. *West African Chimpanzees: Status Survey and Conservation Action Plan*. IUCN—World Conservation Union, Gland, Switzerland.
4. Carter J, Ndiaye S, Pruetz J, McGrew WC 2003. Senegal. In: *West African Chimpanzees: Status Survey and Conservation Action Plan*. Kormos R, Boesch C, Bakarr MI, Butynski TM (eds), IUCN, pp. 31–39.
5. Pruetz JD, Kante D 2010. Successful return of a wild infant chimpanzee (*Pan troglodytes verus*) to its natal group after capture by poachers. *Afr Primates* **7**(1):35–41.
6. Sept JM 2002. Chimpanzees on the edge: The implications of chimpanzee ecology in "savanna" landscapes for hominin evolution. In: *Primates: evolución, cultura y diversidad*. Contreras JM, Veá JJ (eds), CEFPSVLT, México.
7. Lee PC 2010. Sharing space: Can ethnoprimateology contribute to the survival of nonhuman primates in human-dominated globalized landscapes? *Am J Primatol* **72**:925–931.
8. Bogart SL, Pruetz JD 2011. Insectivory of savanna chimpanzees (*Pan troglodytes verus*) at Fongoli, Senegal. *Am J Phys Anthropol* **145**:11–20.
9. Pruetz JD, Marchant LF, Arno J, McGrew WC 2002. Survey of savanna chimpanzees (*Pan troglodytes verus*) in southeastern Senegal. *Am J Primatol* **58**:35–43.
10. McGrew WC, Baldwin PJ, Marchant LF, Pruetz JD, Scott SE, Tutin CEG 2003. Ethoarchaeology and elementary technology of unhabituated wild chimpanzees at Assirik, Senegal. *PaleoAnthropology* **2003**:1–20.
11. McGrew WC, Marchant LF, Phillips CA 2009. Standardized protocol for faecal analysis. *Primates* **50**:363–366.
12. Pruetz, JD, Bertolani P 2009. Chimpanzee (*Pan troglodytes verus*) behavioral responses to stresses associated with living in a savanna-mosaic environment: implications for hominin adaptations to open habitats. *PaleoAnthropology* **2009**:252–262.
13. Massa BE 2011. *Predicting Conflict over Scarce Resources: Chimpanzees (Pan troglodytes verus) and Fulbe Pastoralists*. Unpublished Master's thesis, Duke University.
14. Pruetz JD 2002. Competition between savanna chimpanzees and humans in southeastern Senegal. *Am J Phys Anthropol* **117**(S):128 (AAPA abstracts).
15. Ancrenaz M, Dabek L, O'Neil S 2007. The costs of exclusion: Recognizing a role for local communities in biodiversity conservation. *PLoS Biol* **5**:2443–2448.
16. Hockings K, Humle T 2009. *Best Practice Guidelines for the Prevention and Mitigation of Conflict between Humans and Great Apes*. Occasional paper 37, IUCN Species Survival Commission.
17. Macfie EJ, Williamson EA 2010. *Best Practice Guidelines for Great Ape Tourism*. Occasional paper 38, IUCN Species Survival Commission.
18. Ndiaye S 2011. *Conservation du chimpanzé au Sénégal. Etat des connaissances et réactualisation du plan d'actions*. Rapport d'étude pour USAID/Wula Nafaa.