

Title	Classical integrability of Schrödinger sigma models and q-deformed Poincaré symmetry
Author(s)	Kawaguchi, Io; Yoshida, Kentaroh
Citation	Journal of High Energy Physics (2011), 2011(11)
Issue Date	2011-11
URL	http://hdl.handle.net/2433/152371
Right	The final publication is available at www.springerlink.com ; This is not the published version. Please cite only the published version. この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。
Type	Journal Article
Textversion	author

Classical integrability of Schrödinger sigma models and q -deformed Poincaré symmetry

Io Kawaguchi and Kentaroh Yoshida

Department of Physics, Kyoto University, Kyoto 606-8502, Japan

E-mail:

io@gauge.scphys.kyoto-u.ac.jp, kyoshida@gauge.scphys.kyoto-u.ac.jp

ABSTRACT: We discuss classical integrable structure of two-dimensional sigma models which have three-dimensional Schrödinger spacetimes as target spaces. The Schrödinger spacetimes are regarded as null-like deformations of AdS_3 . The original AdS_3 isometry $SL(2, \mathbb{R})_L \times SL(2, \mathbb{R})_R$ is broken to $SL(2, \mathbb{R})_L \times U(1)_R$ due to the deformation. According to this symmetry, there are two descriptions to describe the classical dynamics of the system, 1) the $SL(2, \mathbb{R})_L$ description and 2) the enhanced $U(1)_R$ description. In the former 1), we show that the Yangian symmetry is realized by improving the $SL(2, \mathbb{R})_L$ Noether current. Then a Lax pair is constructed with the improved current and the classical integrability is shown by deriving the r/s -matrix algebra. In the latter 2), we find a non-local current by using a scaling limit of warped AdS_3 and that it enhances $U(1)_R$ to a q -deformed Poincaré algebra. Then another Lax pair is presented and the corresponding r/s -matrices are also computed. The two descriptions are equivalent via a non-local map.

KEYWORDS: Integrable Field Theory, Sigma Models, AdS-CFT Correspondence

ARXIV EPRINT: [1109.0872\[hep-th\]](https://arxiv.org/abs/1109.0872)

Contents

1	Introduction	1
2	Schrödinger sigma models	3
2.1	The action of Schrödinger sigma models	5
3	Left description based on $SL(2, \mathbb{R})_L$	6
3.1	Yangian symmetry	6
3.2	Lax pair, monodromy matrix, r/s -matrices	7
4	Right description based on enhanced $U(1)_R$	9
4.1	q -deformed Poincaré symmetry	9
4.2	Lax pair, monodromy matrix. r/s -matrices	12
5	Conclusion and Discussion	14
A	Derivation of Lax pair in the right description	15

1 Introduction

The AdS/CFT correspondence [1, 2] is the most concrete realization of dualities between gauge theories and gravitational (string) theories. In the recent progress the integrable structure played an important role in checking it at non-BPS region (For a comprehensive review, see [3]). The integrability of sigma models on AdS spaces and spheres [4, 5] is closely related to the integrable structure behind AdS/CFT and a key ingredient is that AdS spaces and spheres are represented by symmetric cosets. This feature was utilized to classify the coset geometries which can potentially be studied holographically [6]. It is well known that symmetric coset sigma models are classically integrable and infinite-dimensional symmetries, which are often called Yangian [7, 8], are realized (For early works and a review, see [9, 10] and [11]).

Recently, gravity duals for non-relativistic CFTs [12, 13] are intensively studied in relation to the holographic condensed matter scenario. Schrödinger spacetimes are proposed as gravity duals for non-relativistic CFTs [14] possessing non-relativistic conformal symmetry called Schrödinger symmetry [15, 16]. It is an interesting issue to

consider the integrability of sigma models which have the Schrödinger spacetimes as target spaces¹. The Schrödinger spacetimes are represented by non-symmetric coset [20], and hence the standard argument for symmetric cosets is not applicable directly to Schrödinger spacetimes.

In this paper we will reveal the classical integrable structure of two-dimensional sigma models defined on three-dimensional Schrödinger spacetimes, which are called “Schrödinger sigma models” here. The metric of the Schrödinger spacetime is

$$ds^2 = L^2 [d\rho^2 - 2e^{-2\rho}dudv - Ce^{-4\rho}dv^2] , \quad (1.1)$$

where C is a constant deformation parameter. When $C = 0$, (1.1) becomes AdS_3 with the radius L . When $C \neq 0$, we can put $C = \pm 1$ with the Lorentz boost,

$$u \rightarrow \lambda u , \quad v \rightarrow \lambda^{-1}v \quad (\lambda : \text{const.}).$$

The AdS_3 isometry $SO(2, 2) = SL(2, \mathbb{R})_L \times SL(2, \mathbb{R})_R$ is broken to $SL(2, \mathbb{R})_L \times U(1)_R$ due to the non-vanishing C . According to $SL(2, \mathbb{R})_L \times U(1)_R$, there are two descriptions to describe the classical dynamics of the system, 1) the left description based on $SL(2, \mathbb{R})_L$ and 2) the right description based on enhanced $U(1)_R$.

In the former description 1), the Yangian symmetry is shown to be realized as a hidden symmetry by improving the $SL(2, \mathbb{R})_L$ Noether current so that it satisfies the flatness condition, following [21]. Then a Lax pair is constructed with the improved current and the classical integrability is shown by deriving the r/s -matrix algebra. The universality class of this system is rational as is expected from the presence of Yangian. The argument here is quite similar to the previous works for squashed spheres and warped AdS spaces [21, 22].

In the latter description 2), a non-local conserved current is presented by applying a non-local map to the improved current as in the case of squashed sphere [23]. It enhances $U(1)_R$ to a q -deformed Poincaré algebra [24, 25]. Then another Lax pair, which also leads to the classical equations of motion exactly, is presented by taking a scaling limit from the Lax pair in the case of warped AdS_3 and the corresponding r/s -matrices show that the system is rational. This means that the left and right descriptions are equivalent at classical level. In fact, the two descriptions are equivalent via a non-local map.

This paper is organized as follows. In section 2 three-dimensional Schrödinger spacetimes are rewritten in terms of $SL(2, \mathbb{R})$ group element. Then the action of

¹It is worth mentioning about the very recent paper [17] in which an intimate connection between the Schrödinger spacetimes [18] and the Kerr/CFT correspondence [19] is discussed.

Schrödinger sigma models is introduced. In section 3 we study the classical integrability in the left description based on the $SL(2, \mathbb{R})_L$ symmetry. In section 4 the classical integrability is discussed in the right description based on enhanced $U(1)_R$. Section 5 is devoted to conclusion and discussion. In Appendix A we explain the derivation of Lax pair in the right description in detail.

2 Schrödinger sigma models

Schrödinger spacetime in any dimensions is homogeneous and can be described as a coset [20]. Although the coset is non-reductive in general, an exception is the three-dimensional case and the coset becomes reductive. We are confined to this case hereafter.

For the later convenient, let us introduce an $SL(2, \mathbb{R})$ group element represented by

$$g = e^{2vT^+} e^{2\rho T^2} e^{2uT^-} . \quad (2.1)$$

The $SL(2, \mathbb{R})$ generators T^a ($a = 0, 1, 2$) are expressed in terms of the standard Pauli matrices like

$$T^0 = \frac{i}{2}\sigma_2 , \quad T^1 = \frac{1}{2}\sigma_1 , \quad T^2 = \frac{1}{2}\sigma_3 ,$$

and the light-cone notation is defined as

$$T^\pm = \frac{1}{\sqrt{2}} (T^0 \pm T^1) . \quad (2.2)$$

They satisfy the relations

$$[T^a, T^b] = \varepsilon^{ab}{}_c T^c , \quad \text{Tr} (T^a T^b) = \frac{1}{2} \gamma^{ab} , \quad (2.3)$$

where the anti-symmetric tensor $\varepsilon^{ab}{}_c$ is normalized $\varepsilon^{012} = +1$ and the metric on $\mathbb{R}^{1,2}$ is $\gamma^{ab} = (-1, +1, +1)$. The group indices are raised and lowered with γ^{ab} and its inverse.

As a result, the metric (1.1) can be rewritten as

$$\begin{aligned} ds^2 &= \frac{L^2}{2} \left[\text{Tr} (J^2) - 2C (\text{Tr} (T^- J))^2 \right] \\ &= \frac{L^2}{4} \left[-2J^- J^+ + (J^2)^2 - C (J^-)^2 \right] \end{aligned} \quad (2.4)$$

in terms of the left-invariant one-form J defined as

$$J \equiv g^{-1}dg, \quad J^a = 2\text{Tr}(T^a J). \quad (2.5)$$

It is easy to see that the metric (2.4) is invariant under the $SL(2, \mathbb{R})_L \times U(1)_R$ transformation:

$$g \rightarrow g^L \cdot g \cdot e^{-\alpha T^-}. \quad (2.6)$$

The infinitesimal $SL(2, \mathbb{R})_L$ and $U(1)_R$ transformations are given by, respectively,

$$\delta^{L,a} g = \epsilon T^a g, \quad (2.7)$$

$$\delta^{R,-} g = -\epsilon g T^-. \quad (2.8)$$

Here it should be noted that AdS_3 has three kinds of anisotropic deformations, i) space-like, ii) time-like and iii) null-like deformations. The Schrödinger spacetimes correspond to null-like deformations of AdS_3 . The metric of space-like warped AdS_3 is realized with a deformation term on T^1 as

$$ds^2 = \frac{L^2}{2} \left[\text{Tr}(J^2) - 2\tilde{C} [\text{Tr}(T^1 J)]^2 \right]. \quad (2.9)$$

The metric of time-like warped AdS_3 is obtained with a deformation term on T^0 as

$$ds^2 = \frac{L^2}{2} \left[\text{Tr}(J^2) - 2\tilde{C} [\text{Tr}(T^0 J)]^2 \right]. \quad (2.10)$$

The null-like warped AdS_3 is obtained from both space-like and time-like warped AdS_3 geometries by taking a scaling limit [18]. As an example, let us consider a space-like warped AdS_3 with a deformation parameter \tilde{C} . The metric can be rewritten in terms of T^\pm as

$$ds^2 = \frac{L^2}{2} \left[\text{Tr}(J^2) - \tilde{C} [\text{Tr}(T^+ J)]^2 + 2\tilde{C} \text{Tr}(T^+ J) \text{Tr}(T^- J) - \tilde{C} [\text{Tr}(T^- J)]^2 \right]. \quad (2.11)$$

By rescaling T^\pm as

$$T^- \rightarrow \sqrt{\frac{2C}{\tilde{C}}} T^-, \quad T^+ \rightarrow \sqrt{\frac{\tilde{C}}{2C}} T^+ \quad (2.12)$$

and taking $\tilde{C} \rightarrow 0$ limit with C fixed, the metric (2.4) is reproduced. The above argument is applicable to time-like warped AdS_3 as well.

2.1 The action of Schrödinger sigma models

The action of Schrödinger sigma models is

$$S = - \iint dt dx \eta^{\mu\nu} [\text{Tr}(J_\mu J_\nu) - 2C \text{Tr}(T^- J_\mu) \text{Tr}(T^- J_\nu)] . \quad (2.13)$$

The base space is a two-dimensional Minkowski spacetime with the coordinates $x^\mu = (t, x)$ and the metric $\eta_{\mu\nu} = (-1, +1)$. Although we have applications to string theory in our mind, we do not impose periodic boundary conditions and the Virasoro conditions for simplicity here, but the boundary condition that the group element variable $g(x)$ approaches a constant element very rapidly as it goes to spatial infinities:

$$g(t, x) \rightarrow g_{(\pm)} : \text{const.} \quad (x \rightarrow \pm\infty) . \quad (2.14)$$

Thus the left-invariant current J_μ vanishes as it approaches spatial infinities,

$$J_\mu(t, x) \rightarrow 0 \quad (x \rightarrow \pm\infty) . \quad (2.15)$$

The equations of motion obtained from (2.13) are

$$\partial^\mu J_\mu - 2C \text{Tr}(T^- \partial^\mu J_\mu) T^- - 2C \text{Tr}(T^- J_\mu) [J^\mu, T^-] = 0 . \quad (2.16)$$

By multiplying T^a to (2.16) and taking the trace operation, the T^a component of the equations of motion can be obtained. The T^- component leads to the conservation law of the $U(1)_R$ current,

$$\partial^\mu J_\mu^- = 0 . \quad (2.17)$$

The T^2 and T^+ components are, respectively,

$$\partial^\mu J_\mu^2 - C J_\mu^- J^{-,\mu} = 0 , \quad (2.18)$$

$$\partial^\mu J_\mu^+ - C J_\mu^- J^{2,\mu} = 0 . \quad (2.19)$$

The equations of motion (2.16) are equivalent to (2.17)-(2.19).

In addition, the equations of motion (2.16) are equivalent to the conservation law of the $SL(2, \mathbb{R})_L$ current,

$$\partial^\mu [g J_\mu g^{-1} - 2C \text{Tr}(T^- J_\mu) g T^- g^{-1}] = 0 . \quad (2.20)$$

According to this observation on the equations of motion, one may expect that there should be two ways to describe the classical dynamics of this system. Indeed, this is the case. One description is based on the $SL(2, \mathbb{R})_L$ symmetry and the other is on the enhanced $U(1)_R$ symmetry.

3 Left description based on $SL(2, \mathbb{R})_L$

In this section we consider the classical integrability of Schrödinger sigma models in the left description based on the $SL(2, \mathbb{R})_L$ symmetry. First, we show that $SL(2, \mathbb{R})_L$ symmetry is enhanced to an infinite-dimensional symmetry, the $SL(2, \mathbb{R})_L$ Yangian, by improving the $SL(2, \mathbb{R})_L$ Noether current. Then we construct Lax pair and monodromy matrix, and show the classical integrability by deriving the classical r/s -matrix algebra.

3.1 Yangian symmetry

The action (2.13) is invariant under the $SL(2, \mathbb{R})_L$ transformation (2.7). The corresponding conserved $SL(2, \mathbb{R})_L$ current is

$$j_\mu^L = gJ_\mu g^{-1} - 2C\text{Tr}(T^- J_\mu) gT^- g^{-1} + \epsilon_{\mu\nu} \partial^\nu f. \quad (3.1)$$

The first two terms can be obtained by the Noether procedure but the last term is the ambiguity of the conserved current. The anti-symmetric tensor $\epsilon_{\mu\nu}$ on the base space is normalized as $\epsilon_{tx} = +1$ and f is an arbitrary function. When f is taken as

$$f = -\sqrt{C} gT^- g^{-1}, \quad (3.2)$$

then the current j_μ^L satisfies the flatness condition,

$$\epsilon^{\mu\nu} (\partial_\mu j_\nu^L - j_\mu^L j_\nu^L) = 0. \quad (3.3)$$

Thus the flat and conserved $SL(2, \mathbb{R})_L$ current has been obtained in Schrödinger sigma models. This improved current enables us to construct an infinite number of conserved “non-local” charges, for example, by following the BIZZ construction [26]. The first two of them are

$$\begin{aligned} Q_{(0)}^{L,a} &= \int_{-\infty}^{\infty} dx j_t^{L,a}(x), \\ Q_{(1)}^{L,a} &= \frac{1}{4} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} dx dy \epsilon(x-y) \varepsilon^a{}_{bc} j_t^{L,b}(x) j_t^{L,c}(y) - \int_{-\infty}^{\infty} dx j_x^{L,a}(x), \end{aligned} \quad (3.4)$$

where $\epsilon(x-y) \equiv \theta(x-y) - \theta(y-x)$ and $\theta(x)$ is a step function.

The next is to compute the Poisson brackets of the charges. For this purpose, the current algebra of the flat and conserved $SL(2, \mathbb{R})_L$ current is needed. It can be computed by evaluating the standard Poisson brackets of the dynamical variables contained in the classical action and is written down in terms of the component of the current, e.g. $j_\mu^{L,a} = 2\text{Tr}(T^a j_\mu^L)$,

$$\left\{ j_t^{L,a}(x), j_t^{L,b}(y) \right\}_P = \varepsilon^{ab}{}_c j_t^{L,c}(x) \delta(x-y),$$

$$\begin{aligned} \left\{ j_t^{L,a}(x), j_x^{L,b}(y) \right\}_{\mathbb{P}} &= \varepsilon^{ab} j_x^{L,c}(x) \delta(x-y) + \gamma^{ab} \partial_x \delta(x-y), \\ \left\{ j_x^{L,a}(x), j_x^{L,b}(y) \right\}_{\mathbb{P}} &= 0. \end{aligned} \quad (3.5)$$

Note that the current algebra does not contain C explicitly and is exactly the same as the $SL(2, \mathbb{R})_{\mathbb{L}}$ current algebra in sigma models defined on undeformed AdS_3 . It may be natural if one notices that C can be absorbed into the normalization of the generators by a simple rescaling,

$$T^- \rightarrow \frac{1}{\sqrt{|C|}} T^-, \quad T^+ \rightarrow \sqrt{|C|} T^+. \quad (3.6)$$

Thus the Poisson brackets of the conserved charges are also the same as the AdS_3 case,

$$\begin{aligned} \left\{ Q_{(0)}^{L,a}, Q_{(0)}^{L,b} \right\}_{\mathbb{P}} &= \varepsilon^{ab} Q_{(0)}^{L,c}, \\ \left\{ Q_{(1)}^{L,a}, Q_{(0)}^{L,b} \right\}_{\mathbb{P}} &= \varepsilon^{ab} Q_{(1)}^{L,c}, \\ \left\{ Q_{(1)}^{L,a}, Q_{(1)}^{L,b} \right\}_{\mathbb{P}} &= \varepsilon^{ab} \left[Q_{(2)}^{L,c} + \frac{1}{12} (Q_{(0)}^L)^2 Q_{(0)}^{L,c} \right], \end{aligned} \quad (3.7)$$

and therefore an infinite number of conserved charges satisfy the $SL(2, \mathbb{R})_{\mathbb{L}}$ Yangian algebra, as a matter of course.

There is another way to reproduce the current algebra (3.5). The flat conserved $SL(2, \mathbb{R})_{\mathbb{L}}$ current is found in sigma models on space-like warped AdS_3 (2.9) via a double Wick rotation as discussed in [21]. The current algebra in the case of space-like warped AdS_3 is

$$\begin{aligned} \left\{ j_t^{L,a}(x), j_t^{L,b}(y) \right\}_{\mathbb{P}} &= \varepsilon^{ab} j_t^{L,c}(x) \delta(x-y), \\ \left\{ j_t^{L,a}(x), j_x^{L,b}(y) \right\}_{\mathbb{P}} &= \varepsilon^{ab} j_x^{L,c}(x) \delta(x-y) + (1 + \tilde{C}) \gamma^{ab} \partial_x \delta(x-y), \\ \left\{ j_x^{L,a}(x), j_x^{L,b}(y) \right\}_{\mathbb{P}} &= -\tilde{C} \varepsilon^{ab} j_t^{L,c}(x) \delta(x-y). \end{aligned}$$

The rescaling (2.12) does not change the algebra at all. Thus, by taking the limit $\tilde{C} \rightarrow 0$, the current algebra (3.5) is reproduced.

3.2 Lax pair, monodromy matrix, r/s -matrices

The improved $SL(2, \mathbb{R})_{\mathbb{L}}$ current enables us to construct a Lax pair,

$$L_t^L(x; \lambda) = \frac{1}{1 - \lambda^2} [j_t^L(x) - \lambda j_x^L(x)], \quad L_x^L(x; \lambda) = \frac{1}{1 - \lambda^2} [j_x^L(x) - \lambda j_t^L(x)]. \quad (3.8)$$

Here λ is a spectral parameter. The commutation relation

$$[\partial_t - L_t^L(\lambda), \partial_x - L_x^L(\lambda)] = 0 \quad (3.9)$$

reproduces the conservation law of the improved current (equivalently equations of motion) and the flat condition.

Now let us introduce the monodromy matrix $U^L(\lambda)$ defined as

$$U^L(\lambda) \equiv \text{P exp} \left[\int_{-\infty}^{\infty} dx L_x^L(x; \lambda) \right]. \quad (3.10)$$

The symbol P denotes the path ordering. It is easy to see that the monodromy matrix is conserved,

$$\frac{d}{dt} U^L(\lambda) = 0. \quad (3.11)$$

Thus it can be regarded as a generating function of conserved charges. The expression of the conserved quantities depend on the expansion point. For example, when the monodromy matrix is expanded around $\lambda = \infty$, the Yangian charges we have discussed so far are reproduced.

The Poisson bracket of $L_x^{L,a}(x; \lambda)$ is evaluated as

$$\begin{aligned} \{L_x^{L,a}(x; \lambda), L_x^{L,b}(y; \mu)\}_{\text{P}} &= \frac{1}{\lambda - \mu} \varepsilon^{ab}{}_c \left[\frac{\mu^2}{1 - \mu^2} L_x^c(x; \lambda) - \frac{\lambda^2}{1 - \lambda^2} L_x^{L,c}(x; \mu) \right] \delta(x - y) \\ &\quad - \frac{\lambda + \mu}{(1 - \lambda^2)(1 - \mu^2)} \gamma^{ab} \partial_x \delta(x - y). \end{aligned} \quad (3.12)$$

With the tensor product notation, it can be rewritten as follows:

$$\begin{aligned} \{L_x^L(x; \lambda), \otimes L_x^L(y; \mu)\}_{\text{P}} &= [r^L(\lambda, \mu), L_x^L(x; \mu) \otimes 1 + 1 \otimes L_x^L(x; \mu)] \delta(x - y) \\ &\quad - [s^L(\lambda, \mu), L_x^L(x; \mu) \otimes 1 - 1 \otimes L_x^L(x; \mu)] \delta(x - y) \\ &\quad - 2s^L(\lambda, \mu) \partial_x \delta(x - y). \end{aligned} \quad (3.13)$$

Here we have introduced classical r -matrix $r^L(\lambda, \mu)$ and s -matrix $s^L(\lambda, \mu)$ [27], respectively, defined as

$$\begin{aligned} r^L(\lambda, \mu) &\equiv \frac{1}{2(\lambda - \mu)} \left(\frac{\mu^2}{1 - \mu^2} + \frac{\lambda^2}{1 - \lambda^2} \right) (-T^+ \otimes T^- - T^- \otimes T^+ + T^2 \otimes T^2), \\ s^L(\lambda, \mu) &\equiv \frac{\lambda + \mu}{2(1 - \lambda^2)(1 - \mu^2)} (-T^+ \otimes T^- - T^- \otimes T^+ + T^2 \otimes T^2). \end{aligned} \quad (3.14)$$

It is easy to show the extended classical Yang-Baxter equation is satisfied,

$$\begin{aligned} & [(r+s)_{13}^L(\lambda, \nu), (r-s)_{12}^L(\lambda, \mu)] + [(r+s)_{23}^L(\mu, \nu), (r+s)_{12}^L(\lambda, \mu)] \\ & + [(r+s)_{23}^L(\mu, \nu), (r+s)_{13}^L(\lambda, \nu)] = 0, \end{aligned} \quad (3.15)$$

where the subscripts denote the vector spaces on which the r - and s -matrices act. Thus the classical integrability has been shown in the left description.

4 Right description based on enhanced $U(1)_R$

In this section, we describe the classical dynamics of Schrödinger sigma models in the right description based on the enhanced $U(1)_R$ symmetry. We will show that the broken components of $SL(2, \mathbb{R})_R$ are realized as non-local symmetries². The algebra of the corresponding conserved charges is found to be q -deformed two-dimensional Poincaré algebra [24, 25]. In addition, the Lax pair related to the q -deformed Poincaré symmetry is constructed. The resulting classical r/s -matrices also satisfies the classical Yang-Baxter equation.

4.1 q -deformed Poincaré symmetry

We first show that q -deformed Poincaré symmetry is realized as a non-local symmetry in Schrödinger sigma models.

Now the $SL(2, R)_R$ symmetry of the original AdS_3 is broken to $U(1)_R$ due to the deformation. This is generated by T^- as in (2.8) and the conserved $U(1)_R$ current is

$$j_\mu^{R,-} = -2\text{Tr}(T^- J_\mu) = -J_\mu^- .$$

In contrast to the T^- component, the other components generated by T^2 and T^+ are not the isometry of the Schrödinger spacetime. However, an important observation is that there should be a non-local symmetry even in the case of Schrödinger spacetime, in analogy with our previous work [23] on squashed spheres and warped AdS spaces. By following the procedure in [23] and applying a simple non-local map³ to the flat and conserved $SL(2, \mathbb{R})_L$ current, the non-local current is given by

$$\begin{aligned} j_\mu^{R,2} &= -2e^{\sqrt{C}\chi} \text{Tr}(T^2 g^{-1} j_\mu^L g) , \\ j_\mu^{R,+} &= -2e^{\sqrt{C}\chi} \text{Tr}(T^+ g^{-1} j_\mu^L g) , \\ j_\mu^{R,-} &= -2\text{Tr}(T^- g^{-1} j_\mu^L g) . \end{aligned} \quad (4.1)$$

²For an earlier argument on non-locality of the right symmetry, based on a T-duality, see [28].

³This map is analogous to the Seiberg-Witten map [29].

The non-locality comes through the non-local field χ defined as

$$\chi(x) \equiv -\frac{1}{2} \int_{-\infty}^{\infty} dy \epsilon(x-y) j_t^{R,-}(y). \quad (4.2)$$

The boundary conditions (2.14) ensure the convergence of the integral for an arbitrary value of x .

The (2, +)-components of the non-local current are explicitly written down as

$$\begin{aligned} j_\mu^{R,2} &= -2e^{\sqrt{C}\chi} \left[\text{Tr}(T^2 J_\mu) + \sqrt{C} \epsilon_{\mu\nu} \text{Tr}(T^- J^\nu) \right] \\ &= -e^{\sqrt{C}\chi} \left(J_\mu^2 + \sqrt{C} \epsilon_{\mu\nu} J^{-,\nu} \right), \\ j_\mu^{R,+} &= -2e^{\sqrt{C}\chi} \left[\text{Tr}(T^+ J_\mu) + C \text{Tr}(T^- J_\mu) + \sqrt{C} \epsilon_{\mu\nu} \text{Tr}(T^2 J^\nu) \right] \\ &= -e^{\sqrt{C}\chi} \left(J_\mu^+ + C J_\mu^- + \sqrt{C} \epsilon_{\mu\nu} J^{2,\nu} \right). \end{aligned} \quad (4.3)$$

Note that χ satisfies the following relation,

$$\epsilon_{\mu\nu} \partial^\nu \chi = -j_\mu^{R,-}. \quad (4.4)$$

To show the conservation of the non-local currents, we need to use the relations (2.17)-(2.19) and (4.4).

The standard Noether charge

$$Q^{R,-} = \int_{-\infty}^{\infty} dx j_t^{R,-}(x) \quad (4.5)$$

generates the right action of $U(1)_R$,

$$\delta^{R,-} g = \{g, Q^{R,-}\}_P = -g T^-. \quad (4.6)$$

Similarly, non-local charges

$$Q^{R,2} = \int_{-\infty}^{\infty} dx j_t^{R,2}(x), \quad Q^{R,+} = \int_{-\infty}^{\infty} dx j_t^{R,+}(x)$$

generate non-local transformations,

$$\begin{aligned} \delta^{R,2} g &= \{g, Q^{R,2}\}_P = -g \left[T^2 e^{\sqrt{C}\chi} - \sqrt{C} T^- \xi^2 \right], \\ \delta^{R,+} g &= \{g, Q^{R,+}\}_P = -g \left[T^+ e^{\sqrt{C}\chi} - \sqrt{C} T^- \xi^+ \right]. \end{aligned} \quad (4.7)$$

Here we have introduced new non-local fields,

$$\xi^2(x) = -\frac{1}{2} \int_{-\infty}^{\infty} dy \epsilon(x-y) j_t^{R,2}(y), \quad \xi^+(x) = -\frac{1}{2} \int_{-\infty}^{\infty} dy \epsilon(x-y) j_t^{R,+}(y).$$

Note that ξ^2 and ξ^+ are well defined under the boundary conditions (2.14). We can directly check that the action (2.13) is invariant under the transformations (4.7). To show the invariance, we need to use the equations of motion (2.16) and thus the non-local transformations (4.7) are the on-shell symmetry.

The Poisson brackets of $j_t^R(x)$ are

$$\begin{aligned} \left\{ j_t^{R,+}(x), j_t^{R,-}(y) \right\}_{\text{P}} &= -j_t^{R,2}(x) \delta(x-y), \\ \left\{ j_t^{R,+}(x), j_t^{R,2}(y) \right\}_{\text{P}} &= -e^{\sqrt{C}\chi} j_t^{R,+}(x) \delta(x-y) - \frac{\sqrt{C}}{2} \epsilon(x-y) j_t^{R,+}(x) e^{\sqrt{C}\chi} j_t^{R,-}(y) \\ &= \frac{1}{2} j_t^{R,+}(x) \partial_y \left[e^{\sqrt{C}\chi(y)} \epsilon(x-y) \right], \\ \left\{ j_t^{R,-}(x), j_t^{R,2}(y) \right\}_{\text{P}} &= e^{\sqrt{C}\chi} j_t^{R,-}(x) \delta(x-y) \\ &= -\frac{1}{\sqrt{C}} \partial_x \left[e^{\sqrt{C}\chi(x)} \right] \delta(x-y). \end{aligned} \quad (4.8)$$

With (4.8) and the relations

$$\chi(\pm\infty) = \mp \frac{1}{2} Q^{R,-}, \quad (4.9)$$

the Poisson brackets of $Q^{R,a}$ are evaluated as

$$\begin{aligned} \{Q^{R,+}, Q^{R,-}\}_{\text{P}} &= -Q^{R,2}, \\ \{Q^{R,+}, Q^{R,2}\}_{\text{P}} &= -Q^{R,+} \cosh\left(\frac{\sqrt{C}}{2} Q^{R,-}\right), \\ \{Q^{R,-}, Q^{R,2}\}_{\text{P}} &= \frac{2}{\sqrt{C}} \sinh\left(\frac{\sqrt{C}}{2} Q^{R,-}\right). \end{aligned} \quad (4.10)$$

In the $C \rightarrow 0$ limit, this algebra becomes the $SL(2, \mathbb{R})$ algebra.

In order to get a familiar expression, let us rescale $Q^{R,+}$ as

$$Q^{R,+} \rightarrow \frac{\sqrt{C}}{2} Q^{R,+}. \quad (4.11)$$

Then the algebra is rewritten as

$$\{Q^{R,+}, Q^{R,-}\}_{\text{P}} = -\frac{\sqrt{C}}{2} Q^{R,2},$$

$$\begin{aligned}
\{Q^{R,+}, Q^{R,2}\}_{\text{P}} &= -Q^{R,+} \cosh\left(\frac{\sqrt{C}}{2}Q^{R,-}\right), \\
\{Q^{R,-}, Q^{R,2}\}_{\text{P}} &= \frac{2}{\sqrt{C}} \sinh\left(\frac{\sqrt{C}}{2}Q^{R,-}\right)
\end{aligned} \tag{4.12}$$

and this algebra is known as a q -deformed Poincaré algebra [24, 25]. A two-dimensional Poincaré algebra is reproduced from this expression in the $C \rightarrow 0$ limit.

4.2 Lax pair, monodromy matrix. r/s -matrices

Let us next consider a Lax pair in the right description. The following Lax pair,

$$\begin{aligned}
L_t^R(x; \lambda) &= \frac{1}{2} [L_+^R(x; \lambda) + L_-^R(x; \lambda)], \\
L_x^R(x; \lambda) &= \frac{1}{2} [L_+^R(x; \lambda) - L_-^R(x; \lambda)], \\
L_{\pm}^R(x; \lambda) &= -\frac{1}{1 \pm \lambda} \left\{ -T^+ J_{\pm}^- - T^- \left[J_{\pm}^+ \mp C \left(\lambda \pm \frac{\lambda^2}{2} \right) J_{\pm}^- \right] + T^2 J_{\pm}^2 \right\}, \\
J_{\pm} &= J_t \pm J_x
\end{aligned} \tag{4.13}$$

leads to the equations of motion (2.16). This Lax pair can be reproduced by taking an appropriate scaling limit of the Lax pair in the warped AdS₃ cases, as explained in detail in Appendix.

It is a simple practice to show the commutation relation

$$[\partial_t - L_t^R(\lambda), \partial_x - L_x^R(\lambda)] = 0 \tag{4.14}$$

leads to the equations of motion and the monodromy matrix defined as

$$U^R(\lambda) \equiv \text{P exp} \left[\int_{-\infty}^{\infty} dx L_x^R(x; \lambda) \right] \tag{4.15}$$

is conserved:

$$\frac{d}{dt} U^R(\lambda) = 0. \tag{4.16}$$

The Poisson brackets of the spatial components of Lax pair are given by

$$\begin{aligned}
\{L_x^{R,-}(x; \lambda), L_x^{R,+}(y; \mu)\}_{\text{P}} &= \frac{1}{\lambda - \mu} \left[\frac{\mu^2}{1 - \mu^2} L_x^{R,2}(x; \lambda) - \frac{\lambda^2}{1 - \lambda^2} L_x^{R,2}(x; \mu) \right] \delta(x - y) \\
&\quad + \frac{\lambda + \mu}{(1 - \lambda^2)(1 - \mu^2)} \partial_x \delta(x - y),
\end{aligned}$$

$$\begin{aligned}
\{L_x^{R,-}(x; \lambda), L_x^{R,2}(y; \mu)\}_P &= \frac{1}{\lambda - \mu} \left[\frac{\mu^2}{1 - \mu^2} L_x^{R,-}(x; \lambda) - \frac{\lambda^2}{1 - \lambda^2} L_x^{R,-}(x; \mu) \right] \delta(x - y), \\
\{L_x^{R,+}(x; \lambda), L_x^{R,2}(y; \mu)\}_P &= \frac{1}{\lambda - \mu} \left[-\frac{\mu^2}{1 - \mu^2} L_x^{R,+}(x; \lambda) + \frac{\lambda^2}{1 - \lambda^2} L_x^{R,+}(x; \mu) \right] \delta(x - y) \\
&\quad + \frac{C}{2} (\lambda - \mu) \frac{\lambda^2}{1 - \lambda^2} L_x^{R,-}(x; \mu) \delta(x - y), \\
\{L_x^{R,-}(x; \lambda), L_x^{R,-}(y; \mu)\}_P &= 0, \\
\{L_x^{R,+}(x; \lambda), L_x^{R,+}(y; \mu)\}_P &= \frac{C}{2} (\lambda - \mu) \left[\frac{\mu^2}{1 - \mu^2} L_x^{R,2}(x; \lambda) + \frac{\lambda^2}{1 - \lambda^2} L_x^{R,2}(x; \mu) \right] \delta(x - y) \\
&\quad - \frac{C}{2} \frac{(\lambda + \mu)(\lambda - \mu)^2}{(1 - \lambda^2)(1 - \mu^2)} \partial_x \delta(x - y), \\
\{L_x^{R,2}(x; \lambda), L_x^{R,2}(y; \mu)\}_P &= -\frac{\lambda + \mu}{(1 - \lambda^2)(1 - \mu^2)} \partial_x \delta(x - y).
\end{aligned}$$

With the tensor product notation, it is possible to rewrite the above brackets into a simple form,

$$\begin{aligned}
\{L_x^R(x; \lambda), \otimes L_x^R(y; \mu)\}_P &= [r^R(\lambda, \mu), L_x^R(x; \mu) \otimes 1 + 1 \otimes L_x^R(x; \mu)] \delta(x - y) \\
&\quad - [s^R(\lambda, \mu), L_x^R(x; \mu) \otimes 1 - 1 \otimes L_x^R(x; \mu)] \delta(x - y) \\
&\quad - 2s^R(\lambda, \mu) \partial_x \delta(x - y),
\end{aligned}$$

where we have introduced the r - and s -matrices defined as, respectively,

$$\begin{aligned}
r^R(\lambda, \mu) &= \frac{1}{2(\lambda - \mu)} \left(\frac{\mu^2}{1 - \mu^2} + \frac{\lambda^2}{1 - \lambda^2} \right) (-T^+ \otimes T^- - T^- \otimes T^+ + T^2 \otimes T^2) \\
&\quad + \frac{C}{4} (\lambda - \mu) \left(\frac{\mu^2}{1 - \mu^2} + \frac{\lambda^2}{1 - \lambda^2} \right) T^- \otimes T^-, \\
s^R(\lambda, \mu) &= \frac{\lambda + \mu}{2(1 - \lambda^2)(1 - \mu^2)} (-T^+ \otimes T^- - T^- \otimes T^+ + T^2 \otimes T^2) \\
&\quad + \frac{C(\lambda + \mu)(\lambda - \mu)^2}{4(1 - \lambda^2)(1 - \mu^2)} T^- \otimes T^-. \tag{4.17}
\end{aligned}$$

It is easy to show that the extended classical Yang-Baxter equation is satisfied,

$$\begin{aligned}
&[(r + s)_{13}^R(\lambda, \nu), (r - s)_{12}^R(\lambda, \mu)] + [(r + s)_{23}^R(\mu, \nu), (r + s)_{12}^R(\lambda, \mu)] \\
&\quad + [(r + s)_{23}^R(\mu, \nu), (r + s)_{13}^R(\lambda, \nu)] = 0. \tag{4.18}
\end{aligned}$$

Thus the classical integrability has been shown also in the right description.

5 Conclusion and Discussion

We have discussed the classical integrable structure of Schrödinger sigma models. Its classical dynamics can be described by the two descriptions, 1) the left description based on $SL(2, \mathbb{R})_L$ and 2) the right description based on enhanced $U(1)_R$.

The left description is based on the $SL(2, \mathbb{R})_L$ symmetry. The symmetry is enhanced to the Yangian symmetry. To construct the Yangian charges the flat and conserved $SL(2, \mathbb{R})_L$ current is used. By using the current, one can also construct the Lax pair. This Lax pair leads to the rational classical r/s -matrix algebra.

The right description is based on the enhanced $U(1)_R$ symmetry. We have shown that a non-local symmetry is realized and it enhances $U(1)_R$ to a q -deformed Poincaré symmetry. The Lax pair and monodromy matrix concerning the hidden symmetry have also been constructed by taking a scaling limit of the Lax pair in sigma models defined on warped AdS_3 geometries. The classical r/s -matrices explicitly depend on the value of C , but nevertheless those satisfy the classical Yang-Baxter equation.

The two descriptions are equivalent via a non-local map. In fact, as in the case of squashed S^3 and warped AdS_3 [23], one can figure out the map between the improved $SL(2, \mathbb{R})_L$ current and the non-local current concerning the enhanced $U(1)_R$ as follows:

$$\begin{aligned} j_\mu^{R,-} &= -2\text{Tr} (T^- g^{-1} j_\mu^L g) , & j_\mu^{R,2} &= -2e^{\sqrt{C}\chi} \text{Tr} (T^2 g^{-1} j_\mu^L g) , \\ j_\mu^{R,+} &= -2e^{\sqrt{C}\chi} \text{Tr} (T^+ g^{-1} j_\mu^L g) . \end{aligned} \tag{5.1}$$

Note that this is the map within the universality class of rational type, while there exists a map between the rational and the trigonometric in cases of squashed S^3 and warped AdS_3 .

One of the next steps is to construct and solve the corresponding lattice statistical model, which should be called “null-deformed spin chain models (XXN model)”. It seems non-diagonalizable and we are not sure whether it is well defined or not. It may be interesting to consider how the Bethe ansatz equations are modified in this system, for example, by taking a scaling limit. Indeed, quantum solutions for squashed spheres are already known [30, 31, 32] and so it would not be difficult to extend them to the warped AdS_3 cases. The q -deformed Poincaré symmetry should be realized in the “XXN model” and hence the S-matrix should get some constraint by the q -deformed Poincaré symmetry. It would also be nice to analyze the Schrödinger sigma models at quantum level following [33] as another direction.

It would be a challenging problem to try to extend the present argument to higher dimensional cases. The coset does not satisfy the reductive condition any more, hence

it would be difficult to follow the present analysis completely. However, since higher-dimensional Schrödinger algebras always contain $SL(2, \mathbb{R})_{\text{L}} \times U(1)_{\text{R}}$ as a subalgebra, we may expect to use the classical integrability discussed here to describe, at least, the motions restricted to a subspace described as a three-dimensional Schrödinger space-time.

It is also interesting to consider the relation of our result to the recent progress on the Kerr/CFT correspondence [17] from the view point of integrability.

Acknowledgments

We would like to thank Takashi Okada, Domenico Orlando and Akihiro Tsuchiya for illuminating discussions. The work of IK was supported by the Japan Society for the Promotion of Science (JSPS). The work of KY was supported by the scientific grants from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan (No.22740160). This work was also supported in part by the Grant-in-Aid for the Global COE Program “The Next Generation of Physics, Spun from Universality and Emergence” from MEXT, Japan.

Appendix

A Derivation of Lax pair in the right description

The derivation of the Lax pair in the right-description (4.13) is a bit complicated and hence it is explained in detail here.

We begin with the action of sigma models defined on space-like warped AdS_3 spaces,

$$S = - \iint dt dx \eta^{\mu\nu} \left[\text{Tr} (J_\mu J_\nu) - 2\tilde{C} \text{Tr} (T^1 J_\mu) \text{Tr} (T^1 J_\nu) \right]. \quad (\text{A.1})$$

The classical equations of motion are

$$\partial^\mu J_\mu^0 + \tilde{C} J_\mu^2 J^{1,\mu} = 0, \quad \partial^\mu J_\mu^1 = 0, \quad \partial^\mu J_\mu^2 + \tilde{C} J_\mu^0 J^{1,\mu} = 0. \quad (\text{A.2})$$

By performing a double Wick rotation to the Lax pair in the squashed S^3 case [34]⁴, the Lax pair in the warped AdS_3 case is obtained as

$$\begin{aligned} L_t^R(x; \lambda) &= \frac{1}{2} [L_+^R(x; \lambda) + L_-^R(x; \lambda)], & L_x^R(x; \lambda) &= \frac{1}{2} [L_+^R(x; \lambda) - L_-^R(x; \lambda)], \\ L_\pm^R(x; \lambda) &= -\frac{\sinh \alpha}{\sinh(\alpha \pm \lambda)} \left[-T^0 J_\pm^0 + T^2 J_\pm^2 + \frac{\cosh(\alpha \pm \lambda)}{\cosh \alpha} T^1 J_\pm^1 \right], \end{aligned} \quad (\text{A.3})$$

⁴We work on the notation used in [23]. The convention of left and right in [34] is opposite to us.

$$J_{\pm} = J_t \pm J_x, \quad \tilde{C} = \tanh^2 \alpha.$$

The following commutation relation

$$[\partial_t - L_t^R(x; \lambda), \partial_x - L_x^R(x; \lambda)] = 0 \quad (\text{A.4})$$

leads to the equations of motion (A.2).

The next task is to perform the scaling limit to the Lax pair (A.3). Let us first rewrite the Lax pair (A.3) by using T^{\pm} and J^{\pm} as

$$L_{\pm}^R(x; \lambda) = -\frac{\sinh \alpha}{\sinh(\alpha \pm \lambda)} \left[-\frac{1}{2} (T^+ + T^-) (J_{\pm}^+ + J_{\pm}^-) + T^2 J_{\pm}^2 \right. \\ \left. + \frac{\cosh(\alpha \pm \lambda)}{2 \cosh \alpha} (T^+ - T^-) (J_{\pm}^+ - J_{\pm}^-) \right]. \quad (\text{A.5})$$

Consider the redefinition (2.12). $J_{\mu}^{\pm} = 2\text{Tr}(T^{\pm} J_{\mu})$ is also transformed under the redefinition as:

$$J_{\mu}^{-} \rightarrow \sqrt{\frac{2C}{\tilde{C}}} J_{\mu}^{-}, \quad J_{\mu}^{+} \rightarrow \sqrt{\frac{\tilde{C}}{2C}} J_{\mu}^{+}. \quad (\text{A.6})$$

When rescaling as $\lambda \rightarrow \alpha\lambda$, the Lax pair has the following form,

$$L_{\pm}^R(x; \lambda) = -\frac{\sinh \alpha}{\sinh[\alpha(1 \pm \lambda)]} \\ \times \left(-T^+ \left[\frac{1}{2} \left(1 + \frac{\cosh[\alpha(1 \pm \lambda)]}{\cosh \alpha} \right) J_{\pm}^{-} + \frac{\tanh^2 \alpha}{4C} \left(1 - \frac{\cosh[\alpha(1 \pm \lambda)]}{\cosh \alpha} \right) J_{\pm}^{+} \right] \right. \\ \left. -T^- \left[\frac{1}{2} \left(1 + \frac{\cosh[\alpha(1 \pm \lambda)]}{\cosh \alpha} \right) J_{\pm}^{+} + \frac{C}{\tanh^2 \alpha} \left(1 - \frac{\cosh[\alpha(1 \pm \lambda)]}{\cosh \alpha} \right) J_{\pm}^{-} \right] \right. \\ \left. +T^2 J_{\pm}^2 \right). \quad (\text{A.7})$$

Taking a limit in which $\alpha \rightarrow 0$ with C and λ fixed, the Lax pair in the right description of Schrödinger sigma models is obtained as

$$L_{\pm}^R = -\frac{1}{1 \pm \lambda} \left(-T^+ J_{\pm}^{-} - T^- \left[J_{\pm}^{+} \mp C \left(\lambda \pm \frac{\lambda^2}{2} \right) J_{\pm}^{-} \right] + T^2 J_{\pm}^2 \right). \quad (\text{A.8})$$

Note that the $\alpha \rightarrow 0$ limit is the same as $\tilde{C} \rightarrow 0$ because $\tilde{C} = \tanh^2 \alpha$.

References

- [1] J. M. Maldacena, “The large N limit of superconformal field theories and supergravity,” *Adv. Theor. Math. Phys.* **2** (1998) 231 [*Int. J. Theor. Phys.* **38** (1999) 1113] [arXiv:hep-th/9711200].
- [2] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, “Gauge theory correlators from non-critical string theory,” *Phys. Lett. B* **428** (1998) 105 [arXiv:hep-th/9802109]; E. Witten, “Anti-de Sitter space and holography,” *Adv. Theor. Math. Phys.* **2** (1998) 253 [arXiv:hep-th/9802150]
- [3] N. Beisert *et al.*, “Review of AdS/CFT Integrability: An Overview,” arXiv:1012.3982 [hep-th].
- [4] G. Mandal, N. V. Suryanarayana and S. R. Wadia, “Aspects of semiclassical strings in AdS₅,” *Phys. Lett. B* **543** (2002) 81 [arXiv:hep-th/0206103].
- [5] I. Bena, J. Polchinski and R. Roiban, “Hidden symmetries of the AdS₅ × S⁵ superstring,” *Phys. Rev. D* **69** (2004) 046002 [arXiv:hep-th/0305116].
- [6] K. Zarembo, “Strings on semisymmetric superspaces,” *JHEP* **1005** (2010) 002 [arXiv:1003.0465 [hep-th]].
- [7] M. Lüscher and K. Pohlmeyer, “Scattering Of Massless Lumps And Nonlocal Charges In The Two-Dimensional Classical Nonlinear Sigma Model,” *Nucl. Phys. B* **137** (1978) 46.
- [8] V. G. Drinfel’d, “Hopf Algebras and the Quantum Yang-Baxter Equation,” *Sov. Math. Dokl.* **32** (1985) 254; “A New realization of Yangians and quantized affine algebras,” *Sov. Math. Dokl.* **36** (1988) 212.
- [9] D. Bernard, “Hidden Yangians in 2-D massive current algebras,” *Commun. Math. Phys.* **137** (1991) 191.
- [10] N. J. MacKay, “On the classical origins of Yangian symmetry in integrable field theory,” *Phys. Lett. B* **281** (1992) 90 [Erratum-ibid. B **308** (1993) 444].
- [11] N. J. MacKay, “Introduction to Yangian symmetry in integrable field theory,” *Int. J. Mod. Phys. A* **20** (2005) 7189. [arXiv:hep-th/0409183]
- [12] D. T. Son, “Toward an AdS/cold atoms correspondence: A Geometric realization of the Schrodinger symmetry,” *Phys. Rev. D* **78**, 046003 (2008) [arXiv:0804.3972 [hep-th]].
- [13] K. Balasubramanian and J. McGreevy, “Gravity duals for non-relativistic CFTs,” *Phys. Rev. Lett.* **101**, 061601 (2008) [arXiv:0804.4053 [hep-th]].
- [14] Y. Nishida and D. T. Son, “Nonrelativistic conformal field theories,” *Phys. Rev. D* **76** (2007) 086004 [arXiv:0706.3746 [hep-th]].
- [15] C. R. Hagen, “Scale and conformal transformations in galilean-covariant field theory,”

- Phys. Rev. D **5** (1972) 377.
- [16] U. Niederer, “The maximal kinematical invariance group of the free Schrodinger equation,” *Helv. Phys. Acta* **45** (1972) 802.
- [17] S. El-Showk and M. Guica, “Kerr/CFT, dipole theories and nonrelativistic CFTs,” arXiv:1108.6091 [hep-th].
- [18] D. Anninos, W. Li, M. Padi, W. Song and A. Strominger, “Warped AdS₃ black holes,” *JHEP* **0903** (2009) 130 [arXiv:0807.3040 [hep-th]].
- [19] M. Guica, T. Hartman, W. Song and A. Strominger, “The Kerr/CFT correspondence,” *Phys. Rev. D* **80** (2009) 124008 [arXiv:0809.4266 [hep-th]].
- [20] S. Schafer-Nameki, M. Yamazaki and K. Yoshida, “Coset Construction for Duals of Non-relativistic CFTs,” *JHEP* **0905**, 038 (2009) [arXiv:0903.4245 [hep-th]].
- [21] I. Kawaguchi and K. Yoshida, “Hidden Yangian symmetry in sigma model on squashed sphere,” *JHEP* **1011** (2010) 032 [arXiv:1008.0776 [hep-th]].
- [22] I. Kawaguchi, D. Orlando and K. Yoshida, “Yangian symmetry in deformed WZNW models on squashed spheres,” *Phys. Lett. B* **701** (2011) 475 [arXiv:1104.0738 [hep-th]].
- [23] I. Kawaguchi and K. Yoshida, “Hybrid classical integrability in squashed sigma models,” arXiv:1107.3662 [hep-th].
- [24] J. Lukierski, H. Ruegg, A. Nowicki and V. N. Tolstoi, “Q deformation of Poincare algebra,” *Phys. Lett. B* **264** (1991) 331.
- [25] Ch. Ohn, “A $*$ -product on SL(2) and the corresponding nonstandard quantum-U(sl(2)),” *Lett. Math. Phys.* **25** (1992) 85.
- [26] E. Brezin, C. Itzykson, J. Zinn-Justin and J. B. Zuber, “Remarks About The Existence Of Nonlocal Charges In Two-Dimensional Models,” *Phys. Lett. B* **82** (1979) 442.
- [27] J. M. Maillet, “New integrable canonical structures in two-dimensional models,” *Nucl. Phys. B* **269** (1986) 54.
- [28] D. Orlando, S. Reffert and L. I. Uruchurtu, “Classical integrability of the squashed three-sphere, warped AdS₃ and Schrödinger spacetime via T-Duality,” *J. Phys. A* **44** (2011) 115401 [arXiv:1011.1771 [hep-th]].
- [29] N. Seiberg and E. Witten, “String theory and noncommutative geometry,” *JHEP* **9909** (1999) 032 [arXiv:hep-th/9908142].
- [30] P. B. Wiegmann, “Exact solution of the O(3) nonlinear sigma model,” *Phys. Lett. B* **152** (1985) 209.
- [31] V. A. Fateev, “The sigma model (dual) representation for a two-parameter family of integrable quantum field theories,” *Nucl. Phys. B* **473** (1996) 509.

- [32] J. Balog and P. Forgacs, “Thermodynamical Bethe ansatz analysis in an $SU(2) \times U(1)$ symmetric sigma model,” Nucl. Phys. B **570** (2000) 655 [arXiv:hep-th/9906007].
- [33] D. Bernard and A. Leclair, “Quantum group symmetries and nonlocal currents in 2-D QFT,” Commun. Math. Phys. **142** (1991) 99.
- [34] L. D. Faddeev and N. Y. Reshetikhin, “Integrability of the principal chiral field model in (1+1)-dimension,” Annals Phys. **167** (1986) 227.