TITLE:

Response of ER β and aromatase expression in the monkey hippocampal formation to ovariectomy and menopause.

AUTHOR(S):

Higaki, Sayuri; Takumi, Ken; Itoh, Mariko;
Watanabe, Gen; Taya, Kazuyoshi; Shimizu, Keiko;
Hayashi, Motoharu; Oishi, Takao

CITATION:

Higaki, Sayuri ...[et al]. Response of ER β and aromatase expression in the monkey hippocampal formation to ovariectomy and menopause.. Neuroscience research 2012, 72(2): 148-154

ISSUE DATE:

2012-02

URL:

<http://hdl.handle.net/2433/153279>

RIGHT:

© 2011 Elsevier Ireland Ltd and the Japan Neuroscience Society.; この論文は著者最終稿です。内容が印刷版と異なることがありますので、引用の際には出版社版をご確認ご利用ください。This is the Accepted Author Manuscript. Please cite only the published version.

Title

Response of ER β and aromatase expression in the monkey hippocampal
formation to ovariectomy and menopause

Sayuri Higaki^a, Ken Takumi^{a, b}, Mariko Itoh^c, Gen Watanabe^{d, e}, Kazuyoshi

Taya^{d, e}, Keiko Shimizu^{a, f}, Motoharu Hayashi^{a, g, h}, Takao Oishi^{a*}

^a Department of Cellular and Molecular Biology, Primate Research

Institute, Kyoto University, Aichi 484-8506, Japan

^b Department of Anatomy and Neurobiology, Graduate School of Medicine,

Nippon Medical School, Tokyo 113-8602, Japan

^c Futuristic Environmental Simulation Center, Research Institute of

Environmental Medicine, Nagoya University, Aichi 464-8601, Japan

^d Laboratory of Veterinary Physiology, Department of Veterinary

Medicine, Faculty of Agriculture, Tokyo University of Agriculture and

Technology, Tokyo 183-8509, Japan

^e Department of Basic Veterinary Science, The United School of

Veterinary Sciences, Gifu University, Gifu 501-1193, Japan

^f Department of Zoology, Faculty of Science, Okayama University of

Science, Okayama 700-0005, Japan

^g Division of Applied System Neuroscience, Advanced Medical Science

Course, Nihon University School of Medicine, Tokyo 173-8610, Japan

^h Department of Physiology, Nippon Medical School, Tokyo 113-8602,

Japan

* Corresponding author at: Department of Cellular and Molecular Biology,

Primate Research Institute, Kyoto University, Aichi 484-8506, Japan

Tel.: +81-568-63-0573; Fax: +81-568-63-0416.

E-mail address: toishi@pri.kyoto-u.ac.jp (T. Oishi).

Abstract

Changes in the expression of estrogen-related substances in monkeys' brains at the menopausal transition, when estrogen deficit starts to occur, have not yet been examined thoroughly. In the present study, we immunohistochemically investigated the expression levels of estrogen receptor beta (ER β) and aromatase (local estrogen synthesizing enzyme) in the hippocampal formation of premenopausal, menopausal, and ovariectomized premenopausal monkeys. In all monkeys tested, ER β immunoreactivity was observed in interneurons located in the subiculum and the Ammon's horn, and most of these ER β -immunoreactive neurons coexpressed a GABAergic neuron marker, parvalbumin. In the menopausal monkeys who exhibited a decline in estrogen concentration, hippocampal ER β was highly upregulated, while aromatase expression was not markedly changed. By contrast, aromatase in the ovariectomized monkeys was significantly upregulated, while ER β expression was not changed. In the brains of ovariectomized and menopausal monkeys, depletion of ovary-derived estrogen brought about different reactions which may be attributed to the senescence of brain aging.

Keywords: estrogen receptor beta, aging, hippocampus, macaque

1. Introduction

Estrogen homeostatically keeps exerting influences on the brain, as well as on peripheral organs, at regular cycles over almost half of women's life until menopause. A relationship between estrogen and cognitive functions has been clinically documented indicating that menopausal women are at a higher risk of developing Alzheimer's disease (AD) and depression (Launer et al., 1999; Weissman et al., 1993) although there remains some controversy over the sex difference (Hebert et al., 2001). The aged hippocampus is particularly vulnerable to age-related neurodegenerative disorders presenting as atrophy (De Leon et al., 1997), increased levels of neurofilament protein (Vickers et al., 1994), senile plaques and neurofibrillary tangles (Hof, 1997; Rapp et al., 2006), and synapse loss (West et al., 1994). The neural circuits between the hippocampus and the neocortex are highly degenerated in AD patients whose episodic memory is often affected even at an early stage of AD (Morrison and Hof, 1997). Studies of aged monkeys have demonstrated that age-related cognitive impairment reflects the vulnerability of these circuits and that the synaptic alterations occurring in pyramidal cells can

be restored by estrogen administration (Hof, 1997). Although the estrogen replacement therapy (ERT) for menopausal women is expected to have protective effects on the prefrontal cortex and the hippocampus to protect against AD-related cognitive declines (Gibbs and Aggarwal, 1998), there is thought to be a critical window for the initiation of ERT (Henderson et al., 2003). The menopausal transition, a limited period following the loss of ovary function, is considered to be highly susceptible to hormone treatment (Gibbs and Gabor, 2003). Macaques, including Japanese monkeys and rhesus monkeys, enter menopause at around 25 years old, and have a postreproductive life span of 5 or more years (Pavelka and Fedigan, 1999; Walker, 1995). Aged monkeys can be an appropriate model in terms of sharing similar endocrinological and histological senescence of the ovary with humans (Gilardi et al., 1997; Nichols et al., 2005; Nozaki et al., 1995).

In broad areas of the brain, estradiol binds almost equally to two different subtypes of estrogen receptors, ER α and ER β (Kuiper et al., 1997). Previous studies have demonstrated that ER β mRNA (Gundlah et al., 2000) and protein (Blurton-Jones et al., 1999) were more abundant than those of ER α in the monkey hippocampus. Selective ER β agonists enhance

synaptic plasticity in the hippocampus and cognitive performance, which was confirmed by absence of the effects in ER β knockout mice (Liu et al., 2008; Walf et al., 2008). ER β is suggested to colocalize with GABAergic neuron marker, parvalbumin and indirectly upregulate brain-derived neurotrophic factor in the rat (Blurton-Jones and Tuszynski, 2002; 2006). However, the primate hippocampus has not been analyzed in terms of colocalization of ER β and GABAergic neuron markers, such as calcium binding proteins.

Estrogen is derived not only from endocrine glands, but also from the brain. Aromatase contributes to the conversion of androgens into estrogens, the last step of estrogen synthesis. Previous immunohistochemical studies have shown the presence of de novo synthesized neurosteroids in the monkey (Wehrenberg et al., 2001; Yague et al., 2008) and human (Webber et al., 2006; Yague et al., 2010) hippocampus. In order to study menopause-related changes in the brain, it is necessary to consider local estrogen synthesis in the brains of young and aged females, whose peripheral hormone levels are utterly different. In the present study, we attempted to elucidate the changes of ER β and aromatase expression in the

monkey hippocampus under the diminished blood levels of estrogen which
were caused by natural menopause and surgical ovariectomy.

2. Materials and methods

2.1 Animals

Twelve female Japanese monkeys (*Macaca fuscata*) used in this study were housed individually in indoor cages. Based on the age and daily menstruation records, nine monkeys were assigned to one of three categories: premenopausal (n=3; age 9 yr, 11 yr and 13 yr; regular monthly menstruation), perimenopausal (n=3; age 25, 26 and 26 yr; irregular menstruation within past 1 year), and postmenopausal (n=3; aged 29, 30 and 31 yr; no episodes of menstrual bleeding for at least 2 years). The cycling premenopausal monkeys were sacrificed on the second day (D2) (monkey ID, Mff1538 and Mff1564) or D5 (monkey ID, Mff1729) of menses (i.e. in the early follicular phase). The remaining three premenopausal monkeys were bilaterally ovariectomized (n=3; age 9, 13 and 18 yr; regular monthly menstruation) on D20 (monkey ID, Mff1512) or D24 (monkey ID, Mff1283 and Mff1736) of menses (i.e. in the luteal phase). Then, they were sacrificed two months after the operation, as a perimenopausal model in endocrinological aspects. To verify circulating hormone levels of all

monkeys, blood samples at the point of sacrifice were assayed for 17 β -estradiol, progesterone and luteinizing hormone (LH). Additionally, in several premenopausal, perimenopausal and ovariectomized monkeys, the hormonal rhythmicity was monitored by analyzing blood samples collected every 2 - 3 days during more than 120 days. All animals were pretreated with 10 mg/kg ketamine hydrochloride (Sankyo Yell Yakuhin Co., Ltd., Tokyo, Japan) and deeply anesthetized with 30 mg/kg pentobarbital sodium (Dainippon Sumitomo Pharma Co., Ltd., Osaka, Japan). Monkeys were perfused through the heart with the prewash solution, and then perfused with 4% paraformaldehyde. After perfusion, the brains were immediately removed and then immersed in 2% PFA and 5% sucrose in phosphate-buffered saline (PBS) for 24 hours, followed by successive immersions in 10%, 20% and 30% sucrose in PBS. The brain blocks cut into the coronal plane at 5-mm thickness were embedded in Tissue-Tek OCT compound (Miles Inc, Elkhart, IN, USA) followed by rapid freezing in a dry-ice acetone bath. All experimental procedures were planned and executed in accordance with The Guide for the Care and Use of Laboratory Primates (2002) established by the Primate Research Institute of Kyoto

University and NIH Guide for the Care and Use of Laboratory Animals
(1996).

2.2 Measurement of hormone levels by radioimmunoassay (RIA)

The blood samples were analyzed for LH levels using a heterologous RIA system described previously (Watanabe et al., 1990). The iodinated preparation was NIDDK-rat LH-I-5. The antiserum was anti-ovine LH (YM#18; Dr. Y. Mori, University of Tokyo, Tokyo, Japan). Concentrations of 17 β -estradiol and progesterone were measured using double-antibody RIA systems with ¹²⁵I-labeled radioligands, as previously described (Taya et al., 1985) using antiserum against 17 β -estradiol (GDN#244; Dr. G. D. Niswender, Animal Production and Biotechnology, Colorado State University, Fort Collins, CO, USA) and against progesterone (GDN#337; Dr. G. D. Niswender, Animal Production and Biotechnology, Colorado State University, Fort Collins, CO, USA). The intra- and interassay coefficients of variation were 4.4% and 16.1% for LH, 4.7% and 13.3% for 17 β -estradiol, and 5.5% and 16.8% for progesterone, respectively.

2.3 Immunohistochemistry

The coronal sections were cut at 40- μ m thickness with a cryostat (HM500-M, Micron, Waldorf, Germany) and stored in cryoprotectant at -20°C until the time of processing (Watson et al., 1986). Immunohistochemistry of monkey brains was performed as described elsewhere (Hayashi et al., 2001). The sections were washed in PBS, processed with additional antigen retrieval by boiling for 30 min in citrate buffer (pH 6.8) for anti-aromatase staining, and then soaked in 0.3% H₂O₂ in methanol for 30 min to inactivate endogenous peroxidase. They were preincubated with PBS containing 0.5% TritonX-100, 4% goat or donkey serum and 10% fish gelatin for 1 h at room temperature.

For the single-label staining, the sections were incubated with antibody against human ER β (N-19, 1:50, Santa Cruz Biotechnology Inc., Santa Cruz, CA, USA), or human placental aromatase (R-8-1, 1:1000, generously supplied by Dr. Y. Osawa) for 48 h at 4°C. The specificities of these antibodies were shown in the following studies: ER β (Kruijver et al., 2003) and aromatase (Ishimura et al., 1989) antibodies. After brief washes with PBS, sections were treated with a biotinylated second antibody for 24

h at 4°C, and then processed with the avidin-biotin complex peroxidase method (ABC elite kit, Vector Laboratories, Burlingame, CA, USA). The immunoreactive sites were visualized with a solution of 0.003% H₂O₂, 0.6 mg/ml 3, 3'-diaminobenzidine tetrahydrochloride (DAB-buffer tablets, Merck, Darmstadt, Germany). The adjacent sections were stained by the Nissl-staining method with cresyl violet (Merck, Darmstadt, Germany) for cytoarchitectonic reference.

For the double-label immunofluorescence to identify colocalization of ER β with one of the calcium binding proteins, parvalbumin (PV) , calretinin (CR) or calbindin (CB), ER β immunoreactivity was detected using a tyramide signal amplification kit (TSA Perkin Elmer, Boston, MA, USA) after endogenous peroxidase activity was reduced with 3% H₂O₂ in PBS for 30 min. Then, the sections were incubated with diluted antibody for PV (clone PARV-19, 1:2000, Sigma, Saint Louis, MO, USA), CR (clone 6B3, 1:2000, Swant, Bellinzona, Switzerland) or CB (clone 300, 1:2000, Swant, Bellinzona, Switzerland) for 48 h at 4°C and immunoreactivities were visualized with second antibody conjugated to AlexaFluor-546 (1:200, Molecular Probes, Eugene, OR, USA). Immunolabeled images were directly

captured by an Olympus BX-50 fluorescence microscope connected to a digital camera Olympus DP20 (Olympus Corporation, Tokyo, Japan). Confocal images were obtained with a LSM5 Pascal laser-scanning microscope (Carl Zeiss, Jena, Germany).

2.4 Semi-quantitative image analysis

The degree of immunoreactivity for ER β and aromatase in the subiculum was expressed as the integrated optical density (IOD; the product of optical density and area) by using computer-assisted image analysis (NIH ImageJ; <http://rsb.info.nih.gov/nih-image/>). We identified the subiculum from adjacent Nissl-stained images. The specified subicular region on the gray scale image stained for ER β was processed with thresholding filter (maximum entropy) to select ER β -immunostained pixels for IOD calculation. The number of ER β -immunoreactive cells was also counted manually in each subicular region. These analyses for ER β were performed in five sections of each animal. For the aromatase immunoreactivity, two different fields from a section, three sections from each animal, including pyramidal neurons homogeneously in the

subiculum were analyzed. Unlike ER β immunoreactivity, aromatase immunoreactivity showed uniform cellular and subcellular localization in all the four monkey groups. In order to precisely evaluate the intensity of aromatase immunostaining in individual cells, 30 pyramidal cells with relatively large cross-section (minor axis length of the soma > 9 μ m) were selected in a field and delineated manually. The mean IOD value of the selected cells in an image was calculated. ER β -immunoreactive neurons colocalizing with PV were counted manually in two different fields within the merged fluoroscopic subiculum area. The analysis for ER β /PV colocalization was performed in three sections of each animal. All results were statistically analyzed with one-way analysis of variance followed by the Games-Howell test for multiple comparisons. Differences were considered statistically significant if the p value was less than 0.05.

3. Results

3.1 Hormonal profile at each reproductive stage

The premenopausal monkeys exhibited regular menstruation and estrogen, progesterone and LH surges until they were sacrificed (Fig. 1). In ovariectomized monkeys, the mean circulating estrogen level dropped drastically to below one-sixth the pre-ovariectomy level within 4 days and the LH level became elevated to over six-fold within 8 days after the operation. Like the ovariectomized monkeys at postoperation, the perimenopausal monkeys exhibited low estrogen and high LH levels with no menstrual bleeding, in spite of having shown irregular menstruation during the previous year. Individual endocrine backgrounds at the time of sacrifice confirmed the reproductive stage of each monkey.

3.2 ER β and aromatase immunoreactivity localizations

We found conspicuous ER β immunoreactivity in the hippocampal formation of all monkey groups (Fig. 2a). Within the hippocampal formation, ER β was confined to the plasma membrane and perikaryon, but

not to the nucleus. Membranous ER β immunoreactivity was most intensely observed throughout the subiculum (Fig. 2c). These subicular ER β -immunoreactive neurons were found to consist of interneurons of fusiform or multipolar type that could be distinguished from typical pyramidal neurons. In addition, ER β -immunoreactive neurons were located in the stratum oriens of the Ammon's horn (Fig. 2e). To characterize these ER β -immunoreactive neurons, double immunofluorescence histochemistry was carried out for calcium-binding proteins: PV, CR and CB. A large population of ER β -immunoreactive neurons exhibited PV immunoreactivity in the hippocampal formation (Fig. 3a, b). Within the subiculum, approximately 70 to 85% of the total ER β -immunoreactive neurons were co-localized with PV in each of the premenopausal, ovariectomized, perimenopausal and postmenopausal monkey groups, and about 75 to 80% of the total PV-immunoreactive neurons were co-localized with ER β . Throughout the hippocampal formation, no ER β -immunoreactivity was found to be co-localized with CR or CB in a cell (Fig. 3c, d).

The immunoreactivity of aromatase, the enzyme responsible for local estrogen biosynthesis, was present homogeneously in pyramidal neurons and interneurons extending from Ammon's horn to the subiculum (Fig. 2b, d, f).

3.3 Menopause-related changes in ER β and aromatase immunoreactivities

In the ovariectomized monkeys, our quantitative analysis showed that the intensity of ER β immunoreactivity was similar to that in the premenopausal group (Fig.4a). In the premenopausal and ovariectomized monkeys, ER β immunoreactivity was localized primarily in somatic membrane sites and, to a lesser extent, in proximal dendrites. By contrast, the intensity of aromatase immunoreactivity in the subiculum was significantly higher in the ovariectomized monkeys than in the premenopausal monkeys ($p < 0.01$, Fig. 4b). Upregulation after ovariectomy and a lack of age-dependent change in aromatase expression were also observed in other subregions of the hippocampus, such as CA1.

In both the perimenopausal and postmenopausal monkeys, on the other hand, the ER β levels were highly elevated as compared to those in

the premenopausal monkeys ($p < 0.05$, Fig. 4a). The number of ER β -immunoreactive neurons, however, was not altered with menopause (the mean cell number $\pm$ standard error per mm² subicular area, premenopausal: 365.5 ± 19.0 , ovariectomized: 338.6 ± 18.6 , perimenopausal: 362.5 ± 21.9 , postmenopausal: 345.0 ± 13.9 ; one-way analysis of variance, $p = 0.708$). Upregulation of ER β immunoreactivity of subicular interneurons in perimenopausal and postmenopausal monkeys was observed in somatic membranes and dendrites (both proximal and distal) compared with that in premenopausal monkeys. Meanwhile, aromatase immunoreactivity in the subiculum of the perimenopausal and the postmenopausal monkeys was similar to that of the premenopausal monkeys (Fig. 4b). Size of aromatase immunoreactive soma in the postmenopausal monkeys was significantly smaller than that of premenopausal and ovariectomized monkeys (the mean soma sizes $\pm$ standard error (μm^2), premenopausal: 183.81 ± 4.59 , ovariectomized: 194.57 ± 4.24 , perimenopausal: 184.01 ± 5.53 , postmenopausal: 167.00 ± 3.13).

4. Discussion

Here we report that ER β expression is prominent in the subiculum of the hippocampal formation of menopausal monkeys. In the present study, ER β immunoreactivity in the hippocampus was identified at the plasma membrane of non-pyramidal neurons. The estrogen receptors are classically well known to exert genomic actions through translocation into the cell nucleus from the cytoplasm when they bind to 17 β -estradiol (McKenna and O'Malley, 2002). Both ER α and ER β also exert non-genomic rapid action to regulate cell signaling pathways, including calcium ion channels and kinase signaling pathways (Raz et al., 2008). ER β expression at the plasma membrane has been convincingly confirmed by electron microscopy and time lapse microscopy (Milner et al., 2005; Sheldahl et al., 2008). Rapid action of membranous ER β was confirmed with the study using membrane-impermeable estradiol conjugated to bovine serum albumin and selective ER β agonist (Wu et al., 2011). Such rapid action may be exerted in the monkey hippocampus.

Our finding that many of the ER β -immunoreactive neurons co-expressed PV in the subiculum of monkey hippocampal formation was

consistent with the study in the rat (Blurton-Jones and Tuszynski, 2002).

It has been well-known that estrogen affects the mRNA expression of a GABA transporter and a GABA-synthesizing enzyme, glutamic acid decarboxylase (Herbison et al., 1995; McCarthy et al., 1995). In the CA1 of hippocampus, estrogen reduces GABA synthesis in inhibitory interneurons that regulates pyramidal neurons (Murphy et al., 1998). Our study suggests that ER β is concerned with regulation of GABAergic systems in the subiculum, which sends excitatory output signals from the hippocampus to widespread areas.

Aromatase localized in hippocampal neurons synthesize 17 β -estradiol (Hojo et al., 2004). Aromatase expression in the hippocampus is confirmed immunohistochemically in both cycling and ovariectomized monkeys (Yague et al., 2008) and in menopausal humans (Ishunina et al., 2007). We confirmed aromatase expression in all groups examined.

Our study showed that ovariectomized monkeys retained an ER β level in the subiculum equivalent to that of premenopausal monkeys. Alternately, immunoreactivity for aromatase was upregulated in the subiculum of ovariectomized monkeys. In contrast to ovariectomized

monkeys, perimenopausal and postmenopausal monkeys exhibited obvious upregulation of ER β with unaltered aromatase expression in the subiculum of hippocampal formation. Though both ovariectomized and menopausal monkeys experienced peripheral estrogen depletion, reactive regulation in the expression of aromatase and ER β was quite different. Though difference in the senescence of the brain may cause these differences, molecular mechanism remains unclear.

In contrast to our results, aged rats showed significant downregulation of ER β in the hippocampus (Mehra et al., 2005). Middle aged rats exhibit the opposite endocrinological status (i.e. high estrogen level and low LH level) to perimenopausal macaques (low estrogen level and high LH level), though prolonged cycles and gradual cessation of estrus of middle aged rats are similar to characteristics in human or macaque menopausal transition (Chakraborty and Gore, 2004). In aged rats, high estrogen level from ovary may bring about the decrease of ER β expression in the hippocampus. Thus, in both aged monkeys and rats, ER β expression in the hippocampus seems to keep homeostasis in response to age-related changes in the level of ovary-derived estrogen.

It has been reported that postmortem brains developing AD showed apparent upregulation of ER β in the CA1-4 regions of the hippocampus compared to age-matched control patients (Savaskan et al., 2001). The upregulation of ER β in the hippocampus of AD patients and our menopausal monkeys may be induced to enhance the neuroprotective effect of estrogen. In the hippocampus of healthy menopausal women, aromatase immunoreactivity were upregulated (Ishunina et al., 2007), which is similar to our finding in the hippocampus of ovariectomized monkeys rather than that of menopausal monkeys. Similarity in the response of aromatase against estrogen depletion between healthy menopausal humans and ovariectomized premenopausal monkeys is consistent with a hypothesis that healthy aging of the brain is one of the features of brain evolution in hominids (Allen et al., 2005). We assume that adequate level of estrogen actively supplied with aromatase in the hippocampus may be crucial for the aged women to keep brains healthy for the longest life expectancy.

Acknowledgments

The authors acknowledge Dr. A. F. Parlow, the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK), CA, USA for providing RIA materials for LH, and Dr. Y. Mori, University of Tokyo, Tokyo, Japan for providing antiserum against ovine LH (YM#18), and Dr. G. D. Niswender, Colorado State University, Fort Collins, CO, USA for providing antiserum against 17 β -estradiol (GDN#244) and progesterone (GDN#337). The invaluable help of Dr. A. Yamashita and the technical assistance of T. Kunieda are greatly appreciated. We sincerely thank Professor M. Takada for helpful suggestions on this manuscript. This work was supported in part by a Grant-in-Aid for Scientific Research (B) (Grant No. 19370100), Global COE Program A06 of Kyoto University, and a grant for "Highly creative animal model development for brain sciences" carried out under the Strategic Research Program for Brain Sciences of the Ministry of Education, Culture, Sports, Science and Technology of Japan.

Figure Legends

Fig.1. Representative hormonal profiles of premenopausal (pre), ovariectomized (ovx), and perimenopausal (peri) monkeys from September to January. Solid vertical bars indicate the first day of menstruation, and an arrow indicates the day of ovariectomy. The hormonal profile of the ovx monkey after the operation was similar to that of the perimenopausal monkey. Filled circles: estradiol, open circles: progesterone, gray squares: luteinizing hormone (LH).

Fig. 2. ER β -immunoreactivity (ir; a, c, e) and Aromatase (Arom)-ir (b, d, f) in the hippocampal formation. Membranous ER β -ir was observed in fusiform and multipolar interneurons in the subiculum (Sub; c) and in the stratum oriens of CA1 (e) of the hippocampus.. Aromatase (Arom)-ir was also observed in the Sub (d) and CA1 (f). ER β -ir and Arom-ir were present in the hippocampus of all the examined groups. Bars = 500 μ m (a, b) and 50 μ m (c-f), respectively.

Fig. 3. Colocalization of ER β and calcium binding proteins in single cells are confirmed with z-projection of the confocal image stacks. Nearly all of ER β (green) was colocalized with parvalbumin (PV, red) in the subiculum (a, b). Double-immunofluorescence for ER β (green) and calretinin (CR, red in c) or calbindin (CB, red in d) in the subiculum failed to demonstrate colocalization. Bars = 20 μ m (a, c, d), 50 μ m (b)

Fig.4. Integrated optical density (IOD) for semi-quantified ER β (a) and Arom (b)-ir in the subiculum of ovariectomized (ovx), perimenopausal (peri) and postmenopausal (post) monkeys is represented as the ratio to that of premenopausal (pre) monkeys. Note the ER β -ir was prominently elevated in peri and post monkeys, and Arom-ir was elevated in ovx monkeys compared with pre monkeys. Each column indicates an index $\pm$ standard error. Bars with different superscripts differ significantly (a : b, $p < 0.05$). Photographs are representative clipped images used for the image analysis at each reproductive status. Bars = 50 μ m

Fig. 1

Fig. 2

Fig. 3

Fig. 4

binding specificity and transcript tissue distribution of estrogen receptors alpha and beta. *Endocrinology* 138, 863-870.

Launer, L. J., Andersen, K., Dewey, M. E., Letenneur, L., Ott, A., Amaducci, L. A., Brayne, C., Copeland, J. R., Dartigues, J. F., Kragh-Sorensen, P., Lobo, A., Martinez-Lage, J. M., Stijnen, T., Hofman, A., 1999. Rates and risk factors for dementia and Alzheimer's disease: results from EURODEM pooled analyses. *Neurology* 52, 78-84.

Liu, F., Day, M., Muniz, L. C., Bitran, D., Arias, R., Revilla-Sanchez, R., Grauer, S., Zhang, G., Kelley, C., Pulito, V., Sung, A., Mervis, R. F., Navarra, R., Hirst, W. D., Reinhart, P. H., Marquis, K. L., Moss, S. J., Pangalos, M. N., Brandon, N. J., 2008. Activation of estrogen receptor-beta regulates hippocampal synaptic plasticity and improves memory. *Nat. Neurosci.* 11, 334-343.

McCarthy, M. M., Kaufman, L. C., Brooks, P. J., Pfaff, D. W., Schwartz-Giblin, S., 1995. Estrogen modulation of mRNA levels for the two forms of glutamic acid decarboxylase (GAD) in female rat brain. *J. Comp. Neurol.* 360, 685-697.

References

- Allen, J. S., Bruss, J., Damasio, H., 2005. The aging brain: the cognitive reserve hypothesis and hominid evolution. *Am J Hum Biol* 17, 673-689.
- Blurton-Jones, M. M., Roberts, J. A., Tuszynski, M. H., 1999. Estrogen receptor immunoreactivity in the adult primate brain: neuronal distribution and association with p75, trkA, and choline acetyltransferase. *J. Comp. Neurol.* 405, 529-542
- Blurton-Jones, M., Tuszynski, M. H., 2002. Estrogen receptor-beta colocalizes extensively with parvalbumin-labeled inhibitory neurons in the cortex, amygdala, basal forebrain, and hippocampal formation of intact and ovariectomized adult rats. *J. Comp. Neurol.* 452, 276-287.
- Blurton-Jones, M., Tuszynski, M. H., 2006. Estradiol-induced modulation of estrogen receptor-beta and GABA within the adult neocortex: a potential transsynaptic mechanism for estrogen modulation of BDNF. *J. Comp. Neurol.* 499, 603-612.

Chakraborty, T. R., Gore, A. C., 2004. Aging-related changes in ovarian hormones, their receptors, and neuroendocrine function. *Exp Biol Med* (Maywood) 229, 977-987.

De Leon, M. J., George, A. E., Golomb, J., Tarshish, C., Convit, A., Kluger, A., De Santi, S., McRae, T., Ferris, S. H., Reisberg, B., Ince, C., Rusinek, H., Bobinski, M., Quinn, B., Miller, D. C., Wisniewski, H. M., 1997. Frequency of hippocampal formation atrophy in normal aging and Alzheimer's disease. *Neurobiol. Aging* 18, 1-11.

Gibbs, R. B., Aggarwal, P., 1998. Estrogen and basal forebrain cholinergic neurons: implications for brain aging and Alzheimer's disease-related cognitive decline. *Horm. Behav.* 34, 98-111.

Gibbs, R. B., Gabor, R., 2003. Estrogen and cognition: applying preclinical findings to clinical perspectives. *J. Neurosci. Res.* 74, 637-643.

Gilardi, K. V., Shideler, S. E., Valverde, C. R., Roberts, J. A., Lasley, B. L., 1997. Characterization of the onset of menopause in the rhesus macaque. *Biol. Reprod.* 57, 335-340.

Gundlah, C., Kohama, S. G., Mirkes, S. J., Garyfallou, V. T., Urbanski, H. F., Bethea, C. L., 2000. Distribution of estrogen receptor beta

(ERbeta) mRNA in hypothalamus, midbrain and temporal lobe of
spayed macaque: continued expression with hormone replacement.

Brain Res. Mol. Brain Res. 76, 191-204.

Hayashi, M., Mistunaga, F., Ohira, K., Shimizu, K., 2001. Changes in
BDNF-immunoreactive structures in the hippocampal formation of
the aged macaque monkey. Brain Res. 918, 191-196.

Hebert, L. E., Scherr, P. A., McCann, J. J., Beckett, L. A., Evans, D. A.,
2001. Is the risk of developing Alzheimer's disease greater for
women than for men? Am. J. Epidemiol. 153, 132-136.

Henderson, V. W., Guthrie, J. R., Dudley, E. C., Burger, H. G., Dennerstein,
L., 2003. Estrogen exposures and memory at midlife: a
population-based study of women. Neurology 60, 1369-1371.

Herbison, A. E., Augood, S. J., Simonian, S. X., Chapman, C., 1995.
Regulation of GABA transporter activity and mRNA expression by
estrogen in rat preoptic area. J. Neurosci. 15, 8302-8309.

Hof, P. R., 1997. Morphology and neurochemical characteristics of the
vulnerable neurons in brain aging and Alzheimer's disease. Eur.
Neurol. 37, 71-81.

- Hojo, Y., Hattori, T. A., Enami, T., Furukawa, A., Suzuki, K., Ishii, H. T., Mukai, H., Morrison, J. H., Janssen, W. G., Kominami, S., Harada, N., Kimoto, T., Kawato, S., 2004. Adult male rat hippocampus synthesizes estradiol from pregnenolone by cytochromes P45017alpha and P450 aromatase localized in neurons. *Proc. Natl. Acad. Sci. U. S. A.* 101, 865-870.
- Ishimura, K., Yoshinaga-Hirabayashi, T., Tsuru, H., Fujita, H., Osawa, Y., 1989. Further immunocytochemical study on the localization of aromatase in the ovary of rats and mice. *Histochemistry* 90, 413-416.
- Ishunina, T. A., Fischer, D. F., Swaab, D. F., 2007. Estrogen receptor alpha and its splice variants in the hippocampus in aging and Alzheimer's disease. *Neurobiol. Aging*. 28, 1670-1681
- Kruijver, F. P., Balesar, R., Espila, A. M., Unmehopa, U. A., Swaab, D. F., 2003. Estrogen-receptor-beta distribution in the human hypothalamus: similarities and differences with ER alpha distribution. *J. Comp. Neurol.* 466, 251-277.
- Kuiper, G. G., Carlsson, B., Grandien, K., Enmark, E., Haggblad, J., Nilsson, S., Gustafsson, J. A., 1997. Comparison of the ligand

- McKenna, N. J., O'Malley, B. W., 2002. Combinatorial control of gene expression by nuclear receptors and coregulators. *Cell* 108, 465-474.
- Mehra, R. D., Sharma, K., Nyakas, C., Vij, U., 2005. Estrogen receptor alpha and beta immunoreactive neurons in normal adult and aged female rat hippocampus: a qualitative and quantitative study. *Brain Res.* 1056, 22-35.
- Milner, T. A., Ayoola, K., Drake, C. T., Herrick, S. P., Tabori, N. E., McEwen, B. S., Warriar, S., Alves, S. E., 2005. Ultrastructural localization of estrogen receptor beta immunoreactivity in the rat hippocampal formation. *J. Comp. Neurol.* 491, 81-95.
- Morrison, J. H., Hof, P. R., 1997. Life and death of neurons in the aging brain. *Science* 278, 412-419.
- Murphy, D. D., Cole, N. B., Greenberger, V. and Segal, M., 1998. Estradiol increases dendritic spine density by reducing GABA neurotransmission in hippocampal neurons. *J. Neurosci.* 18, 2550-2559.

Nichols, S. M., Bavister, B. D., Brenner, C. A., Didier, P. J., Harrison, R. M.,

Kubisch, H. M., 2005. Ovarian senescence in the rhesus monkey

(*Macaca mulatta*). *Hum. Reprod.* 20, 79-83.

Nozaki, M., Mitsunaga, F., Shimizu, K., 1995. Reproductive senescence in

female Japanese monkeys (*Macaca fuscata*): age- and season-related

changes in hypothalamic-pituitary-ovarian functions and fecundity

rates. *Biol. Reprod.* 52, 1250-1257.

Pavelka, M. S., Fedigan, L. M., 1999. Reproductive termination in female

Japanese monkeys: A comparative life history perspective. *Am. J.*

Phys. Anthropol. 109, 455-464.

Rapp, M. A., Schnaider-Beeri, M., Grossman, H. T., Sano, M., Perl, D. P.,

Purohit, D. P., Gorman, J. M., Haroutunian, V., 2006. Increased

hippocampal plaques and tangles in patients with Alzheimer disease

with a lifetime history of major depression. *Arch. Gen. Psychiatry* 63,

161-167.

Raz, L., Khan, M. M., Mahesh, V. B., Vadlamudi, R. K., Brann, D. W., 2008.

Rapid estrogen signaling in the brain. *Neurosignals* 16, 140-153.

Savaskan, E., Olivieri, G., Meier, F., Ravid, R., Muller-Spahn, F., 2001.

Hippocampal estrogen beta-receptor immunoreactivity is increased
in Alzheimer's disease. *Brain Res.* 908, 113-119.

Sheldahl, L. C., Shapiro, R. A., Bryant, D. N., Koerner, I. P., Dorsa, D. M.,
2008. Estrogen induces rapid translocation of estrogen receptor beta,
but not estrogen receptor alpha, to the neuronal plasma membrane.
Neuroscience 153, 751-761.

Taya, K., Watanabe, G., Sasamoto, S., 1985. Radioimmunoassay for
progesterone, testosterone and estradiol-17beta using (125)
I-iodohistamine redioligands. *Japanese Journal of Animal
Reproduction (Japan)* 31, 186-197.

Vickers, J. C., Riederer, B. M., Marugg, R. A., Buee-Scherrer, V., Buee, L.,
Delacourte, A., Morrison, J. H., 1994. Alterations in neurofilament
protein immunoreactivity in human hippocampal neurons related to
normal aging and Alzheimer's disease. *Neuroscience* 62, 1-13.

Walf, A. A., Koonce, C. J., Frye, C. A., 2008. Estradiol or diarylpropionitrile
administration to wild type, but not estrogen receptor beta knockout,

mice enhances performance in the object recognition and object placement tasks. *Neurobiol. Learn. Mem.* 89, 513-521.

Walker, M. L., 1995. Menopause in Female Rhesus-Monkeys. *Am. J. Primatol.* 35, 59-71.

Watanabe, G., Nozaki, M., Taya, K., Katakai, Y., Sasamoto, S., 1990. Immunoreactive inhibin levels in peripheral blood during the breeding season in the female Japanese monkey. *Biol. Reprod.* 43, 196-201.

Watson, R. E., Jr., Wiegand, S. J., Clough, R. W., Hoffman, G. E., 1986. Use of cryoprotectant to maintain long-term peptide immunoreactivity and tissue morphology. *Peptides* 7, 155-159.

Webber, K. M., Stocco, D. M., Casadesus, G., Bowen, R. L., Atwood, C. S., Previll, L. A., Harris, P. L., Zhu, X., Perry, G., Smith, M. A., 2006. Steroidogenic Acute Regulatory Protein (StAR): Evidence of Gonadotropin-Induced Steroidogenesis in Alzheimer Disease. *Mol Neurodegener* 1, 14.

- Wehrenberg, U., Prange-Kiel, J., Rune, G. M., 2001. Steroidogenic factor-1 expression in marmoset and rat hippocampus: co-localization with StAR and aromatase. *J. Neurochem.* 76, 1879-1886.
- Weissman, M. M., Bland, R., Joyce, P. R., Newman, S., Wells, J. E., Wittchen, H. U., 1993. Sex differences in rates of depression: cross-national perspectives. *J. Affect. Disord.* 29, 77-84.
- West, M. J., Coleman, P. D., Flood, D. G., Troncoso, J. C., 1994. Differences in the pattern of hippocampal neuronal loss in normal ageing and Alzheimer's disease. *Lancet* 344, 769-772.
- Wu, T. W., Chen, S., Brinton, R. D., 2011. Membrane estrogen receptors mediate calcium signaling and MAP kinase activation in individual hippocampal neurons. *Brain Res.* 1379, 34-43.
- Yague, J. G., Azcoitia, I., Defelipe, J., Garcia-Segura, L. M., Munoz, A., 2010. Aromatase expression in the normal and epileptic human hippocampus. *Brain Res.* 1315, 41-52
- Yague, J. G., Wang, A. C., Janssen, W. G., Hof, P. R., Garcia-Segura, L. M., Azcoitia, I., Morrison, J. H., 2008. Aromatase distribution in the

monkey temporal neocortex and hippocampus. Brain Res. 1209,
115-127.