TITLE:

Sediment-complex-binding cellulose breakdown in wetlands of rivers

AUTHOR(S):

Liu, Wen; Toyohara, Haruhiko

CITATION:

Liu, Wen ...[et al]. Sediment-complex-binding cellulose breakdown in wetlands of rivers. Fisheries Science 2012, 78(3): 661-665

ISSUE DATE:

2012-05

URL:

<http://hdl.handle.net/2433/156153>

RIGHT:

The final publication is available at www.springerlink.com; This is not the published version. Please cite only the published version.; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。

和文要旨

河川底泥成分に吸着したセルロース分解活性

劉 文，豊原治彦（京大院農）

東海、近畿及び九州地方の5河川（筑後川，緑川，浜戸川，淀川，田中川）の河口域湿地帯から採集した底泥のセルラーゼ活性を測定した結果、活性レベルは河川により異なっていた。抗生物質で殺菌処理しても底泥は強いセルラーゼ活性を示したことから，セルラーゼの底泥成分への吸着が推測された。上記5河川の底泥のカビ由来セルラーゼの吸着能には違いがあったが，それは底泥中の植物残渣量の違いによるものと考えられた。本研究は、河川底泥においてセルラーゼが植物残渣等の底泥成分に吸着して分解機能を発現している可能性を示唆した。

キーワード：湿地帯、底泥、セルラーゼ、泥、植物残渣

1

2 1 **Sediment-complex-binding cellulose breakdown in wetlands of rivers**

3

4

5 2

6

7

8

9 3 Wen Liu¹ • Haruhiko Toyohara^{1*}

10

11

12 4

13

14

15

16 5 ¹Division of Applied Biosciences, Graduate School of Agriculture, Kyoto University,

17

18 6 Kyoto 606-8502, Japan

19

20

21 7

22

23

24

25 8 **Corresponding author**

26

27

28 9 Haruhiko Toyohara

29

30

31

32 10 Tel/Fax: 81-075-753-6446

33

34

35 11 Email: toyohara@kais.kyoto-u.ac.jp

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

Abstract We have been assessing the activity level of cellulase in wetland sediments to clarify the significance of cellulase for the turnover of plant cellulose in wetlands. In the present study, we investigated the cellulase degrading function of sediment in wetlands to clarify the biochemical breakdown mechanism of cellulose. Specifically, we measured cellulase activities of sediments collected from wetlands of the Chikugo River (Fukuoka Prefecture), Midori River (Kumamoto Prefecture), Hamado River (Kumamoto Prefecture), Yodo River (Osaka Prefecture), and Tanaka River (Mie Prefecture). The results revealed that the activity levels differed significantly among rivers. Additionally, the cellulase activities of the sediment were not completely suppressed in the presence of chloramphenicol. These findings suggested that a portion of the cellulase activities were derived from cellulases binding to the components of sediments. Actually, sediments also showed the ability to bind fungal cellulase. Comparison of the binding ability of clay and plant residues, the main components of sediments, revealed that plant residues had significantly higher abilities to bind cellulase. This finding was supported by the fact that there was a strong correlation between the organic matter content in the sediment and the cellulase binding ability ($R = 0.66$). Results of the present study show that sediment complexes harboring cellulases might be function as a bioreactor to degrade cellulose in wetlands.

Keywords Bioreactor · Cellulase · Clay · Plant residue · River · Sediment · Wetland

32 Introduction

33

34 Cellulose is the main component of the cell walls of plants. Cellulose that is transported
35 from forests to wetlands by rivers is assumed to be utilized by a variety of organisms as
36 a carbon source [1,2,3,4,5,6]; however, the details of the degradation process of
37 cellulose in wetlands remain unknown.

38 Cellulose is a high molecular weight polysaccharide comprised of glucose
39 bound by β -1, 4 linkage that is biochemically stable when compared with starch, in
40 which glucose is bound by α -1, 4 linkage and α -1, 6 linkage. Accordingly, a series of
41 enzymes such as endo- β -glucanase, cellobiohydrolase, and β -glucosidase, which are
42 collectively designated as cellulases, are required for the enzymatic breakdown of
43 cellulose [7].

44 Until recently, it has been assumed that herbivores digest cellulose using
45 cellulases derived from the symbiotic microorganisms [8]. Moreover, cellulases in
46 invertebrates, including insects, were long been assumed to originate from symbiotic
47 microbes before demonstration of the endogenous origin of termite cellulase [9]. During
48 the last decade, cellulase genes have been reported in the aquatic organisms such as
49 crayfish [1], mussel [2], abalone [3], bivalve [4], and sea urchin [5].

50 In addition to these macrobenthos, other organisms such as meiobenthos, fungi,
51 and bacteria are expected to be involved in the degradation of cellulose in the sediments

of wetlands [6,10,11]. In the present study, we attempted to identify organisms contributing to the degradation of cellulose in sediments. The results revealed that cellulases derived from organisms are bound to the components of sediments such as clay and plant residues. Overall, the results of the present study suggest that sediments in wetlands function as a bioreactor to degrade cellulose.

Materials and Methods

Collection of sediments

Sediments from the Tanaka River (Mie Prefecture) were collected on October 20, 2009 and September 8, 2010. Sediments from the Yodo River (Osaka Prefecture) were collected on September 23, 2009. Sediments from the Chikugo River (Fukuoka Prefecture), Hamado River (Kumamoto Prefecture), and Midori River (Kumamoto Prefecture) were collected on October 29, 2010. We selected one collecting site within 50 m from the river mouth and transported these samples at 4 °C back to the laboratory at Kyoto University. Sediment samples were stored at 4 °C until analyses. Macrobenthos such as bivalves and crustaceans were removed prior to collection of the sediments. Upon arrival at the laboratory, meiobenthos such as nematodes and oligochaetes were carefully removed using tweezers in conjunction with microscopic observation (OLYMPUS-SZX12, OLYMPUS, Tokyo, Japan). To remove the meiobenthos

73 completely, the sediments were further filtered through a 63 μm mesh and the
74 pass-through fraction was used for the subsequent experiments.
75
76 Measurement of cellulase activity of sediments
77
78 The cellulase activities of sediments collected from rivers were estimated by measuring
79 reducing sugar released from carboxymethyl cellulose (CMC, Sigma, St. Louis, MO,
80 USA) according to the method described by Somogyi-Nelson [12]. Briefly, 5.0 g of
81 sediment were mixed with 0.5 ml toluene, 10 ml of 0.2 M sodium acetate buffer (pH 5.9)
82 and 10 ml of 1% CMC solution. The CMC was dissolved in water to give a 1% solution.
83 For the blank, water was used instead of CMC solution. Following incubation at 30 $^{\circ}\text{C}$
84 for 24 h, the reaction mixture was centrifuged for 10 min at $2,500\times g$, and 1 ml of the
85 supernatant was then transferred to another tube and combined with 1 ml of Somogyi
86 solution (1.2% $\text{KNaC}_4\text{H}_4\text{O}_6 \cdot 4\text{H}_2\text{O}$, 2.4% Na_2CO_3 , 0.4% CuSO_4 , 1.6% NaHCO_3 , 18%
87 Na_2SO_4). The mixture was then incubated at 100 $^{\circ}\text{C}$ for 10 min. Next, the samples were
88 cooled with cold water, then 1 ml of Nelson solution (1.2% $(\text{NH}_4)_6\text{Mo}_7\text{O}_{24} \cdot 4\text{H}_2\text{O}$, 0.15%
89 $\text{Na}_2\text{HAsO}_4 \cdot 7\text{H}_2\text{O}$, 4.2% H_2SO_4) and 18 ml of water were added. The absorbance at 600
90 nm was then measured using a spectrophotometer (UV-mini-1240, Shimadzu, Kyoto,
91 Japan) and the activity was represented as the reducing sugar released by 1 g of
92 sediment for 1 h [12]. Unless otherwise specified, reagents of specific grades were
93 purchased from nacalai tesque (Kyoto, Japan).

94

95 Effect of chloramphenicol on cellulase activity of sediments

96 To determine if cellulase activity detected in the sediments was derived from
97 microorganisms, the effects of antibiotics on the cellulase activity of the sediments were
98 investigated. As a preliminary experiment, the effects of ampicillin, kanamycin,
99 tetracycline, and chloramphenicol on the growth of microorganisms in the sediment
100 were tested. Sediments collected from the Midori River were mixed with 400 mg, 40
101 mg, or 4 mg of ampicillin, kanamycin, tetracycline, or chloramphenicol per gram of wet
102 sediment, after which they were incubated at 37 °C for 24 h. The effects of antibiotics
103 were evaluated by plating the incubated sediments on LB (Luria Bertani medium) plates
104 and incubating for five days. Only chloramphenicol (400 mg and 40 mg per 1g
105 sediment) completely inhibited the growth of the microorganisms including bacteria and
106 fungi. Thus, we selected chloramphenicol to sterilize microorganisms in the sediments.

107 A solution of 2 ml of chloramphenicol dissolved in 50% ethanol to give 100
108 mg/ml was added to 5 g of per gram of wet sediment collected from the Tanaka and
109 Midori Rivers, which gave 40 mg chloramphenicol per gram of wet sediment in a final
110 concentration. These samples were then incubated for 24 h at 30 °C. As a control, 2 ml
111 of 50% ethanol in water was added. After incubation, the sediment was spread on a LB
112 plate to validate the sterilizing effect of chloramphenicol on microorganisms.

113 Meanwhile the sediment was spread on CMC agar plates containing 1% CMC,
114 0.15% Ca(NO₃)₂, 0.05% MgSO₄, 0.05% K₂HPO₄ and 1.5% agar and the samples were
115 then incubated at 30 °C for three days to detect the remaining cellulase activity of the

116 sediment (data not shown). Then, the remaining cellulase activities of the two rivers
117 were quantified by Somogyi Nelson method described above [12].

118

119 Evaluation of cellulase binding ability of sediments

120 Sediments collected from all rivers were autoclaved at 121 °C for 15 min to inactivate
121 sediment-bound cellulases after vigorous washing with water. After cooling, the
122 following procedures were conducted in the clean bench. Two grams (dry weight) of
123 autoclaved sediment and 6.25 mg of *Aspergillus niger* cellulase (MP Biomedicals,
124 California, USA) dissolved in 5 ml H₂O were mixed and shaken for 1 h at room
125 temperature to bind the cellulase to the sediments. After centrifugation at 2,500×g for
126 10 min, the pellet was vigorously washed twice with excess water to remove the
127 unbound cellulase. The cellulase activities of the sediments were measured as described
128 above to determine cellulase binding ability. The activity was expressed as the reducing
129 sugar released per dry weight of sediments.

130

131 Comparison of cellulase binding ability between clay and plant residues

132

133 Sediments collected from the Midori River were separated into clay and plant residues
134 using tweezers in conjunction with microscopic observation. To inactivate the originally
135 bound cellulase, separated clay and plant residues were autoclaved for 121 °C for 15 min.
136 Next, the commercially obtained fungal cellulase and autoclaved clay or plant residues

were mixed for 1 h at room temperature as described above. After washing vigorously with an excess amount of water, the cellulase activities bound to clay and plant residues were measured, respectively. The activity was expressed as the reducing sugar released per dry weight of clay or plant residues.

Correlation between organic matter content and cellulase binding ability

Sediments were heated at 600 °C for 3 h using a mantle heater (KCA-10A, Koito, Tokyo, Japan). The organic matter content was determined based on the loss on ignition values. The correlation between the cellulase binding ability determined in Table 3 and the organic matter content was then estimated.

Statistical analyses

All data were statistically analyzed by ANOVA.

Results

Cellulase activity of sediments from five rivers

Table 1

As shown in Table 1, sediments collected from all five rivers exhibited cellulase activities. However, the activity levels differed among rivers. Specifically, sediments

from the Yodo River showed significantly higher activity than those from the Hamado River, Midori River and Tanaka River, while those from the Chikugo River had higher values than those from the Midori River and Tanaka River. Sediments from the Hamado River showed significantly higher activity than those from the Tanaka River. However, the activity levels differed among rivers so far as compared by using the data obtained from the collecting sites of each river.

Effect of chloramphenicol on the cellulase activity of sediment

Table 2

As shown in Table 2, chloramphenicol exhibited no effect on the cellulase activity of sediment from the Midori River, while it showed a partially inhibiting effect on that of the Tanaka River. These findings suggest that part of the cellulase activity of the Tanaka River was derived from microorganisms sensitive to chloramphenicol. It should be stressed that a substantial amount of the activity of sediments from both rivers remained, even in the presence of chloramphenicol, suggesting that these activities were derived from cellulases extracellularly secreted from microorganisms and /or benthic animals.

Binding of fungal cellulase to the sediments

Table 3

The finding above suggested us that cellulases secreted from organisms would directly bind to sediments under natural condition. To validate this, we examined the binding ability of sediments to commercial available fungal cellulase. As shown in Table 3,

sediments collected from all five rivers showed fungal cellulase binding ability (Table 3). The sediments from Tanaka River showed significantly lower binding ability than those from Hamado River and Chikugo River, while those from Yodo River showed lower binding ability than those from Chikugo River.

The finding clearly suggested that sediment has the ability to bind cellulase. Sediments are mainly composed of clay and plant residues. Thus, we subsequently compared the cellulase binding ability between clay and plant residues using sediment collected from the Midori River. The result revealed that 11.7 ± 2.3 $\mu\text{g/g}$ of cellulase was bound to clay, while 419.5 ± 47.1 $\mu\text{g/g}$ of cellulase was bound to plant residues. These results suggest that plant residues have significantly higher ($P < 0.05$) cellulase binding ability than clay.

Correlation between organic matter content and cellulase binding ability

Fig.1

As shown in Fig. 1, the organic matter content varied among rivers. Organic matter in sediment was assumed to primarily consist of plant residues. For example, sediment from the Midori River included a large content of plant residues (14.2%), while that of the Tanaka River included a small amount of plant residues (1.3%). A strong correlation between cellulase binding ability and the organic matter content in the sediments was

195 also observed ($R = 0.66$), suggesting that plant residues can function as an efficient
196 binder of cellulase in the sediment.

197 Discussion

198 A significant amount of cellulase activities remained in the presence of
199 chloramphenicol (Table 2). When sediments incubated in the presence of
200 chloramphenicol were inoculated onto agar plates containing LB medium without
201 chloramphenicol, no colonies of microorganisms (fungi or bacteria) were observed
202 during incubation for five days (data not shown), suggesting that microorganisms in the
203 sediment were completely attenuated by chloramphenicol. Based on these findings, we
204 assumed that sediment-complex-binding cellulases secreted from microorganisms
205 and/or benthic animals function as a bioreactor independent of organisms under natural
206 conditions.

207 High cellulase activities of sediments in the Yodo River and Chikugo River
208 were detected, while low activity was observed for the Tanaka River (Table 1). This is
209 because sediments from the Tanaka River showed low cellulase binding ability,
210 possibly due to the low content of plant residues (Table 3 and Fig. 1). The cellulase
211 activity in sediment from the Midori River was significantly lower than that of the Yodo

212 River (Table 1), although the cellulase binding ability of the sediment from the Midori
213 River was not significantly different from Yodo River (Table 3). As described above,
214 these results suggest that a significant amount of cellulase was secreted extracellularly
215 from microorganisms and/or benthic animals. Accordingly, these findings strongly
216 suggest that the low cellulase activity in sediment of the Midori River could be ascribed
217 to the low level of cellulase supplied by microorganisms and/or benthic animals.

218 The results shown in Table 3 suggest that river sediments have cellulase binding
219 ability but that the level differed among rivers. As shown in Fig.1, the cellulase binding
220 ability of sediments depends on the content of plant residues. Indeed, there was a strong
221 correlation between the organic matter content in the sediments and cellulase binding
222 abilities ($R = 0.66$). Organic materials in sediments were considered to be almost
223 entirely a result of plant residues because no weight loss was observed before and after
224 heat treatment of sediments from which the plant residues were removed by tweezers as
225 possible as carefully (data not shown). The difference in the level of cellulase activity of
226 the sediments of various rivers could be ascribed to the content of plant residues as well
227 as the content of cellulase supplied by microorganisms and/or benthic animals.

228 To confirm that plant residues had a higher binding ability than clay, we
229 separated plant residues and clay from the sediment of the Midori River and compared
230 the cellulase binding abilities. Results indicate that the plant residues exhibited
231 approximately 36 times higher cellulase binding than clay. These findings strongly
232 suggest that plant residue functions as an efficient binder of cellulase in the sediment.

233 Overall, this is the first study to demonstrate that sediments composed of clay
234 and plant residues bind cellulase and act as a bioreactor independent of organisms. We
235 are now identifying the organisms that secrete cellulases and will report these results in
236 the near future.

238 Acknowledgement

239 This study was partly supported by a Grant-in-Aid for Scientific Research from the
240 Ministry of Education, Culture, Sports, Science and Technology of Japan (No.
241 21380131).

243 References

- 1
2 244 1. Byrne KA, Lehnert SA, Johnson SE, Moore SS (1999) Isolation of a cDNA
3
4 245 encoding a putative cellulase in the red claw crayfish *Cherax quadricarinatus*. Gene
5
6
7 246 239:317-324
8
9
10 247 2. Guo R, Ding M, Zhang SL, Xu GJ, Zhao FK (2008) Molecular cloning and
11
12 248 characterization of two novel cellulase genes from the mollusk *Ampullaria crosseana*.
13
14 249 J Comp Physiol B 178:209-215
15
16
17
18 250 3. Nikapitiya C, Oh C, Zoysa MD, Whang I, Kang DH, Lee SR, Kim SJ, Lee J (2010)
19
20 251 Characterization of beta-1,4-endoglucanase as a polysaccharide-degrading digestive
21
22 252 enzyme from disk abalone, *Haliotis discus discus*. Aquacult Int 18:1061-1078
23
24
25
26 253 4. Sakamoto K, Touhata K, Yamashita M, Kasai A, Toyohara H (2007) Cellulase
27
28 254 digestion by common Japanese freshwater clam *Corbicula japonica*. Fish Sci
29
30 255 73:675-683
31
32
33
34 256 5. Nishida Y, Suzuki K, Kumagai Y, Tanaka H, Inoue A, Ojima T (2007) Isolation and
35
36 257 primary structure of a cellulase from the Japanese sea urchin *Strongylocentrotus*
37
38 258 *nudus*. Biochimie 89:1002-1011
39
40
41
42 259 6. Hyde KD, Jones EBG, Leano E, Pointing SB, Poonyth AD, Vrijmoed LLP (1998)
43
44 260 Role of fungi in marine ecosystems. Biodivers Conserv 7:1147-1161
45
46
47
48 261 7. Watanabe H, Tokuda G (2010) Cellulolytic systems in insects. Annu Rev Entomol
49
50 262 55:609-632
51
52
53
54 263 8. Brezmal JA, Brune A (1994) Role of microorganisms in the digestion of
55
56 264 lignocelluloses by termites. Annu Rev Entomol 39:453-487
57
58
59
60
61
62
63
64
65

- 265 9. Watanabe H, Noda H, Tokuda G, Lo N (1998) A cellulase gene of termite origin.
266 Nature 394:330-331
- 267 10. Niiyama T, Toyohara H (2011) Widespread distribution of cellulase and
268 hemi-cellulase activities among aquatic invertebrates. Fish Sci 77:649-655
- 269 11. Toyohara H, Park Y, Tsuchiya K, Liu W (2011) Cellulase Activity in meiobenthos
270 in wetlands. Fish Sci in press
- 271 12. Nelson N (1944) A photometric adaptation of the Somogyi method for the
272 determination of glucose. J Biol Chem 153:375-380
273
- 274 Figure caption
- 275 Fig. 1 Correlation between the organic matter content and the cellulase binding ability
276 of the sediments from various rivers. A strong correlation was demonstrated between
277 them ($R = 0.66$).
278
279

Fig. 1

Table 1 Cellulase activity of the sediments collected from five rivers

Collecting sites	Chikugo	Midori	Hamado	Yodo	Tanaka
	River	River	River	River	River
Cellulase activity (nmol/gh)	$59.2 \pm 26.0^{a,d}$	$22.5 \pm 3.4^{b,c}$	$33.6 \pm 8.45^{a,c}$	77.6 ± 7.1^d	12.9 ± 8.2^b

Different letters indicate a significant difference ($P < 0.05$). Values are means $\pm$ SD ($n = 3$).

Table 2 Effect of Chloramphenicol on cellulase activities of the sediments from Tanaka River and Midori River

Collecting sites	Tanaka River		Midori River	
	Chloramphenicol		Chloramphenicol	
	+	-	+	-
Cellulase activity (nmol/gh)	27.4 ± 1.7 ^a	43.1 ± 4.7 ^b	18.8 ± 2.0 ^a	28.3 ± 7.7 ^a

Concentration of chloramphenicol was 40 mg/g sediment. The statistical analyses on both rivers were performed independently. Different letters indicate a significant difference ($P < 0.05$).

Table 3 Binding of fungal cellulase to the autoclaved sediments collected from five rivers

Collecting sites	Chikugo	Midori	Hamado	Yodo	Tanaka
	River	River	River	River	River
Cellulase binding ability ($\mu\text{g/g}$)	29.0 ± 5.4^a	19.7 ± 4.3^b	$29.2 \pm 9.13^{a,b}$	$16.6 \pm 10.6^{b,c}$	11.6 ± 4.0^c

Values are means $\pm$ SD ($n = 3$). Different letters indicate a significant difference ($P < 0.05$).

Reviewer #2: The reviewer admits that the manuscript has been improved from the original version; however, there are still many points that should be addressed by the authors.

Materials and Methods

- L86: $\text{KNaC}_4\text{H}_4\text{O}_6 \cdot \text{Na}_2\text{O}$ ---- Na_2O may be $4\text{H}_2\text{O}$.

-We have corrected it as the reviewer suggested.

- L89: $\text{Na}_2\text{HAsO}_4 \cdot \text{As}_2\text{O}$ ---- As_2O may be $7\text{H}_2\text{O}$

-We have corrected it as the reviewer suggested.

- L133-L134 --- *The reviewer cannot imagine how the authors distinguished clay and plant residues and how separated them using tweezers? Clay may be too small to pick up with tweezers.*

-As shown in the photo, clay and plant could be separated easily by tweezers, because the shapes of them completely different. The arrow indicates a plant residue.

- L136: *How many grams of clay and plant residues did the authors use for cellulase-binding assay?*

-We used approximately 5g (in wet condition) of clay and plant residues for the assay.

Results

- L165: *exhibited no effect on ... from the Midori River, --- The difference in the cellulase activity between plus and minus chloramphenicol for Midori River*

sample, i.e., 18.8 vs 28.3, indicates that the effect of CP is not negligible. The reviewer cannot agree the authors' explanation "exhibited no effect".

-Because of large SD of minus CP (28.3 ± 7.7) we could not recognize a significant difference between the values of plus CP and minus CP in the Midori River samples. Thus, we considered that CP exhibited no effect.

- *L178: The sediment from Tanaka River showed significantly lower... ---- The reviewer does not think that the value for Tanaka River is "significantly" low. The value for Tanaka River is comparable level with that for Yodo River.*

-We would like to show that there is a statistical difference between Tanaka River and Yodo River. We revised as follows to avoid misunderstanding.

Line178-180: "The sediments from Tanaka River showed statistically lower binding ability than those from Hamado River and Chikugo River."

Discussion

- *L203-204: secreted from ... and/or benthic animals ---- Do the benthic animals secrete cellulase to the sediment? Generally, the benthic animals are considered to ingest plant tissues and digest them in the digestive tract with cellulase secreted in the tract.*

-We found that cellulases are secreted in feces as active form. We will report this in the next paper.

- *L224: the plant residues were completely removed --- How did the authors confirmed the "complete" removal of plant tissues.*

-It is easy to remove plant residues by tweezers as described above from their shapes. However, as suggested by the referee, it is difficult to declare to be "complete". Thus we revised the sentence as follows in the new manuscript.

L222-225: Organic materials in sediments were considered to be almost entirely a result of plant residues because no weight loss was observed before and after heat treatment of sediments from which the plant residues were removed by tweezers as possible as carefully (data not shown).

- *Fig. 1. The values should be presented as the average values with SDs.*

-According to the suggestion, we added SDs in the figure.

- *Table 1-3. What do "a-d" mean?*

-These alphabet letters show the statistical difference. If the two values are attached by different letters such as “a” and “b”, it means both values are statistically different. This expression is usually used to demonstrate a statistical difference between values. For example, please refer Tables 3-8 in the paper published in Fisheries Science 77, 357-365 (2011).

- *Table 2. The activities for Tanaka River and Midori River in the absence of chloramphenicol, 43.1 and 28.3, are inconsistent with those in Table 1. How can the author explain?*

-We collected sediments from Tanaka River on October 20, 2009 used for the experiment in Table 1, and September 8, 2010 used for the experiment in Table 2. Therefore, the activities in Table 1 (12.9 ± 8.2) and Table 2 (43.1 ± 4.7) are inconsistent. This is possibly because the slight difference in the collecting sites, difference in the month and/or the difference in weather condition between 2009 and 2010.

As for Midori River, we used the same sediments collected on October 29, 2010. Thus, no statistical difference was observed; 22.5 ± 3.4 (Table 1) and 28.3 ± 7.7 (Table 2).

- *Table 3. The unit "nmol/gh" is not suitable for indication of "Cellulose binding ability".*

-According to the suggestion, we expressed “cellulase binding ability” as the ability to bind the amount of cellulase per one gram weight of dry sediment. To determine the amount of cellulase, we validated the relationship between cellulase activity (absorbance at 600 nm) and amount of fungal cellulase (μg) as shown below. The amount of cellulase binding to sediment was determined according to the formula inserted in the figure. Cellulase binding ability shown in Table 3 is expressed as the value which was determined by dividing the amount of cellulase by the dry weight of sediment.

Fig. Relationship between the cellulase activity (absorbance at 600 nm) and amount of fungal cellulase (μg).

By this change, the results of statistical analysis in Table.3 and Fig.1 were altered. In Table.3, altered portions are shown in red letters as described below.

Table 3 Binding of fungal cellulase to the autoclaved sediments collected from five rivers

Collecting sites	Chikugo	Midori	Hamado	Yodo	Tanaka
	River	River	River	River	River
Cellulase binding ability (μg/g)	29.0 ± 5.4 ^a	19.7 ± 4.3 ^b	29.2 ± 9.13 ^{a,b}	16.6 ± 10.6 ^{b,c}	11.6 ± 4.0 ^c

Values are means ±SD ($n = 3$). Different letters indicate a significant difference ($P < 0.05$).

In Fig.1, the slope of line was altered. Sentences in the manuscript corresponded were also altered as described below (altered portions are shown in red letters):

Line 26-27: This finding was supported by the fact that there was a strong correlation between the organic matter content in the sediment and the cellulase binding ability ($R = 0.66$).

Line 179-181: ... while those from Yodo River showed lower binding ability than those from Chikugo River.

Line 184-185: The result revealed that $11.7 \pm 2.3 \mu\text{g/g}$ of cellulase was bound to clay, while $419.5 \pm 47.1 \mu\text{g/g}$ of cellulase was bound to plant residues.

Line 193-195: A strong correlation between cellulase binding ability and the organic matter content in the sediments was also observed ($R = 0.66$),...

Line 210-213: The cellulase activity in sediment from the Midori River was significantly lower than that of the Yodo River (Table 1), although the cellulase binding ability of the sediment from the Midori River was not significantly different from Yodo River (Table 3).

Line 220-222: Indeed, there was a strong correlation between the organic matter content in the sediments and cellulase binding abilities ($R = 0.66$).

Line 230-231: Results indicate that the plant residues exhibited approximately 36 times higher cellulase binding than clay.

Line 274-277: Fig. 1 Correlation between the organic matter content and the cellulase binding ability of the sediments from various rivers. A strong correlation was demonstrated between them (**R=0.66**)