TITLE:

Cooperative functions of Hes/Hey genes in auditory hair cell and supporting cell development.

AUTHOR(S):

Tateya, Tomoko; Imayoshi, Itaru; Tateya, Ichiro; Ito, Juichi; Kageyama, Ryoichiro

CITATION:

Tateya, Tomoko ...[et al]. Cooperative functions of Hes/Hey genes in auditory hair cell and supporting cell development.. Developmental biology 2011, 352(2): 329-340

ISSUE DATE:

2011-04-15

URL:

<http://hdl.handle.net/2433/158370>

RIGHT:

© 2011 Elsevier Inc.; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。; This is not the published version. Please cite only the published version.

Cooperative functions of *Hes/Hey* genes in auditory hair cell and supporting cell development

Tomoko Tateya¹, Itaru Imayoshi^{1,2}, Ichiro Tateya³, Juichi Ito³, and Ryoichiro Kageyama^{1,4}

¹Institute for Virus Research, Kyoto University, Kyoto 606-8507, Japan;

²Japan Science and Technology Agency, PRESTO, Kyoto 606-8507, Japan;

³Department of Otolaryngology - Head and Neck Surgery, Kyoto University Graduate School of Medicine, Kyoto 606-8507, Japan;

⁴Japan Science and Technology Agency, CREST, Kyoto 606-8507, Japan

Corresponding author: Ryoichiro Kageyama

Institute for Virus Research

Kyoto University

Shogoin-Kawahara, Sakyo-ku

Kyoto 606-8507, Japan

Tel: 81-75-751-4011

Fax: 81-75-751-4807

E-mail: rkageyam@virus.kyoto-u.ac.jp

Running title: *Hes/Hey* genes in cochlear development

Abstract

Notch-mediated lateral inhibition has been reported to regulate auditory hair cell and supporting cell development from common precursors. While the Notch effector genes *Hes1*, *Hes5* and *Hey1* are expressed in the developing cochlea, inactivation of either of them causes only mild abnormality, suggesting their functional redundancy. To explore the roles of *Hes/Hey* genes in cochlear development, we examined compound heterozygous or homozygous mutant mice that lacked *Hes1*, *Hes5* and *Hey1* alleles. We found that a reduction in *Hes/Hey* gene dosage led to graded increase of hair cell formation. However, if at least one allele of *Hes1*, *Hes5* or *Hey1* was intact, excessive hair cells were accompanied by overproduction of supporting cells, suggesting that the hair cell increase does not occur at the expense of supporting cells, and that each *Hes/Hey* gene functions to induce supporting cells. By contrast, when all alleles of *Hes1*, *Hes5* and *Hey1* were inactivated, the number of hair cells increased more drastically, whereas that of supporting cells was unchanged compared with control, suggesting that supporting cell formation was balanced by their overproduction and fate conversion into hair cells. The increase of the cell numbers seemed to occur after the prosensory domain formation in the mutants because the proliferation state and the size of the prosensory domain were not affected. Thus, *Hes1*, *Hes5* and *Hey1* cooperatively inhibit hair cell formation, and one allele of either *Hes1*, *Hes5* or *Hey1* is sufficient for supporting cell production probably by lateral inhibition in the sensory epithelium. Strikingly, *Hes/Hey* mutations lead to disorganized cell alignment and polarity and to hearing loss despite hair cell overproduction. These results suggest that *Hes/Hey* gene dosage is essential not only for generation of appropriate numbers of hair cells and supporting cells by controlling cell proliferation and lateral inhibition but also for the hearing ability by regulating the cell alignment and polarity.

Key words: hair cell; supporting cell; *Hes1*; *Hes5*; *Hey1*; developing cochlea

Introduction

A specialized sensory epithelium of the cochlea of the mammalian inner ear, called the organ of Corti, contains sensory hair cells and non-sensory supporting cells. These cells are aligned in rows: one row of inner hair cells (IHC), three rows of outer hair cells (OHC), and supporting cells that surround each row of hair cells. During development, the prosensory domain is formed in the floor of the cochlear duct, and then hair cells and supporting cells differentiate from common precursors in the prosensory domain. The mechanism of the alternate hair cell and supporting cell formation has been extensively analyzed, and it has been shown that Notch-mediated lateral inhibition plays an important role in this process (Kelly, 2006). Notch-mediated lateral inhibition is known to generate two cell types: Notch ligands activate Notch signaling in neighboring cells, while activation of Notch signaling leads to expression of repressor genes such as *Hes1*, thereby inducing adoption of a fate different from Notch ligand-expressing cells (Kageyama et al., 2007).

Hair cells express the Notch ligands *Deltalike1* (*Dll1*), *Deltalike3* (*Dll3*) and *Jagged2* (*Jag2*) (Lanford et al., 1999; Morrison et al., 1999; Hartman et al., 2007), while supporting cells express the Notch effector genes *Hes1*, *Hes5*, *Hey1*, *Hey2* and *HeyL* (Hayashi et al. 2008; Li et al. 2008; Doetzlhofer et al., 2008), suggesting that hair cells activate Notch signaling in the neighboring supporting cells. Genetic ablation of Notch ligand or effector genes leads to overproduction of hair cells (Kiernan et al., 2005), and treatment of a γ -secretase inhibitor, which blocks Notch signaling, increases the number of hair cells at the expense of supporting cells in the cochlear explant culture (Takebayashi et al., 2007). Thus, it is likely that hair cells inhibit the neighboring cells from differentiating into the same cell type by activation of Notch signaling, and that the latter cells adopt the alternative fate, supporting cells. In agreement with this idea, hair cells can induce surrounding cells to develop as supporting cells (Woods et al., 2004). These results suggest that Notch-mediated lateral inhibition regulates hair cell versus supporting cell specification.

In spite of extensive studies, the role of Notch signaling in lateral inhibition is still obscure. In *Dll1*; *Jag2* double mutant cochlear, hair cells increased in number, but the concomitant decrease of the supporting cells was much milder (Kiernan et al., 2005). Furthermore, in compound mutant mice that lacked the Notch effectors *Hes* and *Hey* genes, excessive hair cell formation was accompanied by overproduction of supporting cells (Li et al., 2008). Thus, the fate change from supporting cells to hair cells does not account for overproduction of hair cells, and it has been suggested that Notch

signaling has an additional role in cell proliferation (Kiernan et al., 2005; Murata et al., 2009).

To investigate the pleiotropic roles of Notch signaling, we examined the cochlea of compound heterozygous or homozygous mutant mice that lacked *Hes1*, *Hes5* and *Hey1* alleles. We found that a reduction in *Hes/Hey* gene dosage led to graded increase in hair cell formation accompanied by overproduction of supporting cells, and that only when all of *Hes1*, *Hes5* and *Hey1* were inactivated, the number of hair cells increased at the expense of supporting cells. These results suggest that Notch signaling is essential for generation of appropriate numbers of hair cells and supporting cells by regulating cell proliferation and lateral inhibition. We also found that *Hes/Hey* mutations lead to hearing loss despite hair cell overproduction.

Materials and methods

Mice breeding

Hes1 floxed, *Hes3;Hes5* knockout (Hatakeyama et al., 2004; Imayoshi et al., 2008), *Hey1(Hesr1)* knockout (Kokubo et al., 2005) and *Emx2^{+Cre}* mice (Kimura et al., 2005) were used to produce various compound heterozygous or homozygous mutant mice. The loci of *Hes3* and *Hes5* are close to each other, and *Hes5* deficiency was accompanied by *Hes3* deficiency in this study. However, because there was no *Hes3* expression in the cochlea (see Supplemental Fig. S5C), *Hes3* was not mentioned in this study. *Hes1^{ff};Hes5^{-/-};Hey1^{-/-}* mice were crossed with *Emx2^{+Cre};Hes1^{+ff};Hes5^{+/-};Hey1^{+/-}* mice, *Emx2^{+Cre};Hes1^{+ff};Hes5^{+/-};Hey1^{-/-}* mice, *Emx2^{+Cre};Hes1^{+ff};Hes5^{-/-};Hey1^{+/-}* mice, or *Emx2^{+Cre};Hes1^{+ff};Hes5^{-/-};Hey1^{-/-}* mice. In the following, *Hes1^{+Δ}* and *Hes1^{ΔΔ}* are the abridgements of *Emx2^{+Cre};Hes1^{+ff}* and *Emx2^{+Cre};Hes1^{ff}*, respectively. For analysis of Cre recombinase activity, *Emx2^{+Cre}* mice were crossed with ROSA26-CFP mice (Srinivas et al., 2001). These mice were maintained on C57BL/6; ICR mixed background. Plug date was defined as embryonic day 0.5 (E0.5).

Histochemistry and in situ hybridization

Whole heads (E10.5–E14.5) or inner ears (E17.5–adult) were immediately fixed in 4% paraformaldehyde (PFA) in 0.1M phosphate-buffered saline pH7.4 (PBS), cryoprotected in 30% sucrose in PBS and embedded in OCT for cryostat sectioning.

Immunostaining of cochlear sections was performed, as described previously (Imayoshi et al., 2008). Primary antibodies used for this study were: anti-BrdU

(Oxford Biotechnology, rat monoclonal, 1:100 dilution), anti-p27Kip1 (BD Transduction Laboratories, mouse monoclonal, 1:200), anti-MyosinVI (Proteus Bio-Sciences, rabbit polyclonal, 1:200), and anti-Prox1 (Chemicon, rabbit polyclonal, 1:1000). Secondary antibodies were goat or donkey anti-species IgG conjugated with Alexa 405, Alexa 488 or Alexa 594. Nuclei were labeled by 4,6'-diamidino-2-phenylindole (DAPI). For p27^{Kip1}, Prox1 and BrdU staining, samples were heated in 10mM sodium citrate at 90°C for 10 min (p27^{Kip1} and Prox1) or 30 min (BrdU) prior to staining procedure. For staining of whole mount preparations of the cochlea, cochlear ducts were opened to expose the developing sensory epithelia prior to staining procedure (Yamamoto et al., 2009). Apoptotic cells were detected using Apoptag® Fluorescein In Situ Apoptosis Detection Kit (Millipore), following the manufacturer's instruction.

In situ hybridization was carried out using mouse *Hes1*, *Hes3*, *Hes5*, *Hey1*, *Hey2*, *HeyL* and *Hes3* and *Atoh1* probes, as described previously (Imayoshi et al., 2008). Immunolabeling of MyosinVI was performed after in situ hybridization as described above.

Quantification of hair cells

The number of hair cells in E18.5 mice was counted on cochlear surface preparations processed with MyosinVI immunohistochemistry, as described previously (Zine et al., 2001). We analyzed a region that covered a 1.6-mm length of the organ of Corti including the hook region, beginning at the basal end of cochlea and extending toward the middle turn.

Cell counts of supporting cell marker- and MyosinVI-immunolabeled sections

Every 2 sections of all mid-modiolar serial sections were labeled for DAPI, Prox1 and p27^{Kip1} or for DAPI and MyosinVI. Sections were analyzed with LSM510 confocal microscopy (Zeiss). Inner hair cells, outer hair cells, Prox1⁺ cells, and p27^{Kip1}-positive and Prox1-negative cells in the basal and middle turns of the organ of Corti were counted. Prox1⁺ cells include Pillar cells and Deiters' cells. p27^{Kip1}-positive and Prox1-negative cells in the greater epithelial ridge were regarded as inner phalangeal cells. The total number of each cell type per cochlea duct section was calculated. At least 25 sections per cochlea and at least 3 animals per genotype were used.

Bromodeoxyuridine (BrdU) administration

BrdU (Sigma) was dissolved in PBS. 50 µg/g body weight of BrdU was given to the

pregnant mice at E13.5, E14.5 or E17.5 by a single intraperitoneal injection at a concentration of 10 mg/ml. Every 2 sections of all mid-modiolar serial sections were labeled for DAPI, BrdU and MyosinVI, or Ki67, BrdU and MyosinVI. The number of BrdU⁺ cells and MyosinVI⁺ cells, or Ki67⁺ cells and BrdU⁺ cells were counted.

Quantitative reverse transcription PCR (qRT-PCR)

E14.5 cochlear epithelia were dissected and mesenchyme of cochlear tissue was removed using a thermolysin treatment as previously described (Yamamoto et al. 2009). Total RNA was reverse-transcribed by using Rever-Tra Ace (TOYOBO) and Random Primer (TOYOBO). Real-time PCR was done by using Applied Biosystems 7500 Real Time PCR System (Applied Biosystems, Foster City, CA) and THUNDERBIRD SYBR qPCR Mix (TOYOBO), according to the manufacturer's protocols. GAPDH was used as a control. The following primers were used for real-time PCR: Hes1 forward, tgaaggattccaaaaataaaattctctctggg; Hes1 reverse, cgcctctctctccatgataggctttgatgac; GAPDH forward, atcttcttgtgcagtgccagcctcgctcccg; and GAPDH reverse, agttgagggtcaatgaaggggtcgttgatgg.

Statistical analysis

Three or more embryos for each genotype group were analyzed in all experiments. A repeated-measures analysis of variance and a Student-Newman-Keuls test were used to detect differences among groups. Differences at $p < 0.05$ were regarded as statistically significant.

ABR recording

An Auditory Brainstem Response (ABR) recording was used to monitor the auditory function of the experimental animals. Under general anesthesia, ABR measurements were performed as previously described (Kada et al., 2008). Thresholds were determined for the frequencies of 10, 20, and 40 kHz from a set of responses at varying intensities with 5-dB Sound Pressure Level intervals.

Results

The expression of Hes1, Hes5 and Hey1 in the developing auditory sensory epithelium

We first examined expression of *Hes1*, *Hes5* and *Hey1* in the developing inner ear. *Hes1*

expression detected by in situ hybridization was faint, and therefore *Venus-Hes1* mice were used; Venus (a YFP variant) fragment was knocked in at the *Hes1* locus so that the Venus-Hes1 fusion protein was expressed under the control of the endogenous *Hes1* promoter (It will be reported elsewhere). At E11.5, *Venus-Hes1* was slightly expressed in the epithelium of presumptive cochlea (Fig. 1A arrow, and B), but more abundant in mesenchyme around the epithelium. *Hey1* was broadly expressed in the prosensory domain, while *Hes5* was not detected in the epithelium of otocyst (Fig. 1D-F,H-J). At E13.5, *Hey1* was expressed in the prosensory domain, whereas *Hes1* and *Hes5* expression were not seen in the cochlea duct (Fig. 1C,G,K). In agreement with the previous data (Hayashi et al, 2008; Li et al., 2008), *Hes1* and *Hey1* expression were seen in the E14.5 cochlea duct (Fig. 1R,S,U,V). *Hes5* was also expressed near hair cells (*Atoh1*⁺) (Fig. 1M,P). At E17.5, *Hes1* was expressed by Hensen cells, while *Hes5* was expressed by inner pharyngeal cells and Deiters' cells, and *Hey1* expression was seen in Kollikers organ, Deiters' cells and Hensen cells (Fig. 1Q,T,W), similar to the previous data (Zheng et al., 2000; Zine et al., 2001; Hayashi et al., 2008; Li et al., 2008; Doetzlhofer et al., 2008). Thus, *Hes/Hey* genes were continuously expressed both in the early presumptive cochlea epithelium and in the organ of Corti at a lateral inhibition phase.

Cre recombinase activity in the developing cochleae of *Emx2*^{+Cre} mice

To determine the Cre recombinase activity in the developing cochleae of *Emx2*^{+Cre} mice, we crossed these mice with ROSA26-CFP reporter mice. At E11.5, CFP expression was observed only in subsets of otocyst epithelial cells (Supplemental Fig. S1A-E). However, at E13.5, CFP expression occurred in most epithelial cells of the cochlea duct including the prosensory domain, which was labeled with anti-p27^{Kip1} antibody (Supplemental Fig. S1F-J, arrows), and of the organ of Corti at E18.5 and P1 (Supplemental Fig. S1K,L,P,Q,R). CFP expression also occurred in most cells of both the greater epithelial ledge and lesser epithelial ledge (Supplemental Fig. S1L) as well as in most hair cells (Supplemental Fig. S1Q) and supporting cells (Supplemental Fig. S1R). CFP expression was limited to the epithelial cells in the cochleae, and only several CFP⁺ cells were observed in the spiral ganglion (Supplemental Fig. S1K,P, arrowheads) and the maculae of saccule and utricle (Supplemental Fig. S1M,N). No CFP⁺ cells were seen in the crista ampullaris of the semicircular canals (Supplemental Fig. S1O). These reporter expression patterns were consistent with *Emx2* expression in the inner ears (Holley et al., 2010).

The above results indicated that the Cre recombinase became fully active by

E13.5 in the cochlea of *Emx2^{+Cre}* mice. In agreement with this idea, *Hes1* mRNA levels were reduced to 64% and 9% in the cochlea epithelium of *Emx2^{+Cre};Hes1^{+/f}* (designated *Hes1^{Δ/Δ}*) and *Emx2^{+Cre};Hes1^{flf}* (designated *Hes1^{Δ/Δ}*) mice, respectively, at E14.5 (Supplemental Fig. S1S), suggesting that *Hes1* expression was effectively lost by E14.5 in *Hes1* conditional knock out mice.

It was previously reported that extra inner hair cells and outer hair cells were found in *Emx2* heterozygous cochleae, although no quantification analysis was performed (Rhodes et al., 2003). It was also reported that *Emx2* knock out mice have shortened cochleae with effects on both proliferation and differentiation of hair cells whereas *Emx2* heterozygous mice have no phenotypes in the cochleae (Holley et al. 2010). We compared the inner and outer hair cell numbers of *Emx2^{+Cre}* cochleae with those of wild type animals, but there was no significant difference (Fig. 2I). Furthermore, we performed Auditory Brainstem Response (ABR) recording of adult *Emx2^{+Cre}* mice, and these mice were found to have no hearing impairment (data not shown). Thus, we decided to use *Emx2^{+Cre}* mice as a Cre driver to inactivate *Hes1* in the developing cochlea.

Graded increase in hair cell formation with a reduction in Hes/Hey gene dosage

The gross appearance of *Hes/Hey* compound mutant cochleae appeared to be normal (Fig. 2A,B), but a reduction in *Hes/Hey* gene dosage led to graded increase in hair cell formation (Fig. 2C-H). To compare the number of hair cells, we counted hair cell marker-positive cells over the length of 1.6 mm from the basal end of E18.5 cochlea, corresponding to approximately one quarter of the cochlea length. Wild type, *Emx2^{+Cre}* and *Hes1^{+/Δ};Hes5^{+/-};Hey1^{+/-}* mice were not significantly different from one another in both the inner and outer hair cell numbers (Fig. 2C,I). Mice that lacked either *Hes5* (*Hes5^{-/-};Hey1^{+/-}*) or *Hey1* (*Hes5^{+/-};Hey1^{-/-}*) or both (*Hes5^{-/-};Hey1^{-/-}*, Fig. 2D) but retained one *Hes1* allele (*Hes1^{+/Δ}*) exhibited mild increase of both inner and outer hair cells (Fig. 2D, red- and green-labeled, and Fig. 2I), although this increase was not significant (Supplemental Table S1). However, in mice that lacked *Hes1* (*Hes1^{Δ/Δ};Hes5^{+/-};Hey1^{+/-}*), the numbers of both inner and outer hair cells were significantly increased ($p < 0.01$) from the former groups (Fig. 2E,I, Supplemental Table S1). Additional deletion of *Hes5* (*Hes1^{Δ/Δ};Hes5^{-/-};Hey1^{+/-}*) or *Hey1* (*Hes1^{Δ/Δ};Hes5^{+/-};Hey1^{-/-}*) did not significantly change the inner and outer hair cell numbers compared with *Hes1^{Δ/Δ};Hes5^{+/-};Hey1^{+/-}* (Fig. 2F,G,I, Supplemental Table S1). When all genes were inactivated (*Hes1^{Δ/Δ};Hes5^{-/-};Hey1^{-/-}*), both inner and outer hair cells were further increased in number (Fig. 2H,I, and Supplemental Table S1). These

results indicate that *Hes1*, *Hes5* and *Hey1* cooperatively inhibit hair cell formation, while *Hes1* is the most effective.

The cochleae of *Hes1*^{ΔΔ};*Hes5*^{-/-};*Hey1*^{-/-} mice were shorter than those of the other genotypes ($p < 0.01$), but the difference was less than 10% (Supplemental Fig. S2), suggesting that not only the numbers per length but also the total numbers of both inner and outer hair cells were significantly increased in *Hes1*^{ΔΔ};*Hes5*^{-/-};*Hey1*^{-/-} mice. The average numbers of hair cell rows in the cochlear displayed 1.5-fold to 2-fold increase in *Hes/Hey* compound mutant cochlear (Figs. 2I, 3Q, 4J, and Supplemental Fig. S3), indicating that the total number of hair cells indeed increased in the absence of *Hes/Hey* genes. Hair cell overproduction occurred similarly in the regions of 25%, 50% and 75% from the basal end of the E18.5 mutant cochlea, suggesting that there is no apparent regional difference in the phenotypes (Supplemental Fig. S4).

Changes of supporting cells that surround outer hair cells in Hes/Hey mutants

We next examined how supporting cell formation is affected by hair cell increase in the mutant cochlear. We first analyzed supporting cells that surround outer hair cells. *Prox1* was expressed in the nuclei of specific subtypes of supporting cells, pillar cells (Fig. 3D, yellow arrows) and Deiters' cells (Fig. 3D, yellow arrowheads) at E18.5. *Hes1*^{+Δ};*Hes5*^{+/-};*Hey1*^{+/-} cochlea showed normal 5 rows of *Prox1*⁺ cells (Fig. 3D,G) and 3 rows of outer hair cells (Fig. 3A,H). In *Hes1*^{+Δ};*Hes5*^{-/-};*Hey1*^{-/-} cochlea, excessive hair cells were formed (Fig. 3J, arrows), and excessive *Prox1*⁺ cells were also present at the same place (Fig. 3I, arrowheads), indicating that an extra row of outer hair cells were accompanied by an extra row of Deiters' cells. *Hes1*^{ΔΔ};*Hes5*^{-/-};*Hey1*^{+/-} cochlea (Fig. 3B,L) and *Hes1*^{ΔΔ};*Hes5*^{+/-};*Hey1*^{-/-} cochlea (Fig. 3N) had more outer hair cells, and these excessive hair cells were accompanied by excessive *Prox1*⁺ cells, which mostly formed 6 rows (Fig. 3E,K,M). These results suggest that outer hair cells and supporting cells were overproduced in pairs in these compound mutant cochleae. However, in *Hes1*^{ΔΔ};*Hes5*^{-/-};*Hey1*^{-/-} cochlea, much more outer hair cells were formed (Fig. 3C,P), whereas *Prox1*⁺ cells were disarranged or missing at the same place (Fig. 3F,O, white asterisks). Thus, when all *Hes1*, *Hes5* and *Hey1* were inactivated, outer hair cell increase was accompanied by relative decrease of Deiters' cells (Fig. 3Q and Supplemental Fig. S3).

Quantification of outer hair cells and *Prox1*⁺ pillar and Deiters' cells in the basal and middle turn on serial sections of E18.5 cochleae showed that *Hes1*^{+Δ};*Hes5*^{+/-};*Hey1*^{+/-} cochlea had on average 3.1 outer hair cells and 5.2 *Prox1*⁺ pillar and Deiters' cells (Fig. 3Q). In *Hes1*^{ΔΔ};*Hes5*^{-/-};*Hey1*^{+/-} cochlea, both outer hair cells

and Prox1⁺ cells were significantly increased in number to 3.9 and 5.8, respectively (P<0.01, Supplemental Table S2). If the remaining *Hey* allele was further deleted (*Hes1*^{ΔΔ};*Hes5*^{-/-};*Hey1*^{-/-}), there were even more outer hair cells (4.7, P<0.01, Supplemental Table S2A), but less Prox1⁺ cells (5.1, P<0.01, Supplemental Table S2B) than *Hes1*^{ΔΔ};*Hes5*^{-/-};*Hey1*^{+/-} cochlea, although the number of Prox1⁺ cell rows varied between 4 to 6. When all *Hes1*, *Hes5* and *Hey1* were inactivated (*Hes1*^{ΔΔ};*Hes5*^{-/-};*Hey1*^{-/-}), excessive hair cell formation occurred in the same plane as decreased Deiters' cell formation (Fig. 3C,F), suggesting that excessive hair cells are formed at the expense of supporting cells, and that this fate conversion balanced supporting cell overproduction.

Changes of supporting cells that surround inner hair cells in Hes/Hey mutants

We also analyzed inner phalangeal cells, supporting cells that surround inner hair cells. Inner phalangeal cells are negative for pillar cell markers such as p75^{NTR} and Prox1 but express p27^{Kip1}, which can label all subtypes of supporting cells at E18.5 cochlea. In *Hes1*^{ΔΔ};*Hes5*^{-/-};*Hey1*^{+/-} cochlea, which had excessive inner hair cells, inner phalangeal cells increased in number (Fig. 4D,E, brackets, F, asterisks, J, and Supplemental Table S2B), compared with *Hes1*^{+Δ};*Hes5*^{+/-};*Hey1*^{+/-} and *Hes1*^{+Δ};*Hes5*^{-/-};*Hey1*^{-/-} cochlea (Fig. 4A,B, brackets, C, asterisks, J, and Supplemental Table S2B). These results indicated that excessive inner hair cell formation was accompanied by excessive supporting cell formation (Fig. 4J and Supplemental Fig. S3). However, when all *Hes1*, *Hes5* and *Hey1* were inactivated (*Hes1*^{ΔΔ};*Hes5*^{-/-};*Hey1*^{-/-}), inner hair cells increased further, whereas phalangeal cells relatively decreased in number to the level that was not significantly different from *Hes1*^{+Δ};*Hes5*^{+/-};*Hey1*^{+/-} cochleae (Fig. 4G,H, brackets, I, asterisks, J, Supplemental Fig. S3 and Supplemental Table S2B), suggesting that further increase of the inner hair cell number occurred at the expense of supporting cells, and that this fate conversion balanced supporting cell overproduction.

These results together suggest that the overproduction of hair cells by reduction of *Hes/Hey* gene dosage was accompanied by supporting cell overproduction if at least one allele of either *Hes1*, *Hes5* or *Hey1* was intact. However, if all alleles of *Hes1*, *Hes5* and *Hey1* were inactivated, it is likely that the number of hair cells increased at the expense of supporting cells. In *Hes1*^{ΔΔ};*Hes5*^{-/-};*Hey1*^{-/-} cochlea, however, there were still many supporting cells, suggesting that the fate conversion was not complete. This was probably due to compensation by *Hey2* and *HeyL*, which were expressed similarly in both the control and *Hes1*^{ΔΔ};*Hes5*^{-/-};*Hey1*^{-/-} cochlea (Supplemental Fig. S5).

Cell alignment and polarity were disturbed by *Hes/Hey* deficiency

In the developing cochleae, each hair cell gradually forms both stereocilia and kinocilium, which have a unique polarity: a bundle of stereocilia, visualized by phalloidin, forms a V shape directing outward, and a kinocilium (acetylated α -tubulin⁺) is located at the tip of the V shape. We next assessed the alignment and polarity of hair cells in *Hes/Hey* compound mutant cochlea by examining the stereocilia and kinocilium. In *Hes1^{+Δ};Hes5^{-/-};Hey1^{-/-}* cochlea, hair cells were properly aligned, and the polarity of stereocilia and kinocilium was mostly normal (Fig. 5A,B). In *Hes1^{ΔΔ};Hes5^{-/-};Hey1^{+/-}* cochlea, most outer hair cells were properly aligned, and the polarity of stereocilia and kinocilium was mostly normal (Fig. 5C,D). However, although excessive inner hair cells were aligned in two rows, the polarity of the stereocilia and kinocilium of inner hair cells was disorganized in *Hes1^{ΔΔ};Hes5^{-/-};Hey1^{+/-}* cochlea (Fig. 5C,D). When all *Hes1*, *Hes5* and *Hey1* were inactivated (*Hes1^{ΔΔ};Hes5^{-/-};Hey1^{-/-}*), the alignment of both inner and outer hair cells was severely disorganized and hair cell rows were wavy and disconnected (Fig. 5E,F). Furthermore, the polarity of stereocilia and kinocilium of inner and outer hair cells was abnormal, pointing rather randomly (Fig. 5E,F). These results indicated that the cell alignment and polarity were more severely disorganized as *Hes/Hey* gene dosage decreased.

Abnormal cell proliferation in the epithelium by *Hes/Hey* deficiency

If at least one allele of either *Hes1*, *Hes5* or *Hey1* was intact, excessive hair cells were accompanied by excessive formation of supporting cells, suggesting that the cell proliferation is enhanced by decrease of *Hes/Hey* gene dosage. We thus examined the cell proliferation of these mutant cochleae. At E18.5, the proliferation marker Ki67 was not detectable in hair cells or their neighboring supporting cells of *Hes1^{+Δ};Hes5^{+/-};Hey1^{+/-}* and *Hes1^{ΔΔ};Hes5^{-/-};Hey1^{+/-}* cochleae as well as in the control (Fig. 6A-D and data not shown) except for Kolliker's organ. In contrast, in *Hes1^{ΔΔ};Hes5^{-/-};Hey1^{-/-}* cochlea, Ki67 expression was observed in some cells next to hair cells (Fig. 6E, F, arrows), suggesting that cell proliferation was prolonged in the absence of *Hes/Hey* genes.

We also administered BrdU at E14.5 and examined BrdU uptake at E18.5. There were no BrdU⁺ hair cells or supporting cells in the control and *Hes1^{+Δ};Hes5^{+/-};Hey1^{+/-}* cochleae (Fig. 6G,H,M), suggesting that all hair cell/supporting cell precursors exited the cell cycle by E14.5. However, BrdU⁺ supporting cells were observed in *Hes1^{ΔΔ};Hes5^{-/-};Hey1^{+/-}* and *Hes1^{ΔΔ};Hes5^{-/-};Hey1^{-/-}* cochleae (Fig. 6I-L, arrows). Moreover, some hair cells incorporated BrdU in *Hes1^{ΔΔ};Hes5^{-/-};Hey1^{-/-}*

cochleae (Fig. 6K,L, arrows). The proportions of BrdU⁺ hair cells were calculated in the basal and middle turn on serial sections of E18.5 cochleae. No BrdU⁺ hair cells were found in *Hes1*^{+/Δ};*Hes5*^{+/-};*Hey1*^{+/-} cochleae (Fig. 6M). In contrast, the proportion of BrdU incorporation in hair cells of *Hes1*^{Δ/Δ};*Hes5*^{-/-};*Hey1*^{-/-} cochleae was about 2.7%, which was significantly higher than that of *Hes1*^{+/Δ};*Hes5*^{+/-};*Hey1*^{+/-} ($p < 0.01$), *Hes1*^{+/Δ};*Hes5*^{-/-};*Hey1*^{-/-} ($p < 0.05$), and *Hes1*^{Δ/Δ};*Hes5*^{-/-};*Hey1*^{+/-} cochleae ($p < 0.05$) (Fig. 6M). These proportions could be underestimated because BrdU was administered only once (at E14.5) to the mother mice in this experiment. These results together suggest that cell proliferation is prolonged by decrease of *Hes/Hey* gene dosage, and that particularly in *Hes1*^{Δ/Δ};*Hes5*^{-/-};*Hey1*^{-/-} cochlear, a substantial number of hair cell/supporting cell precursors are proliferating at E14.5, when normal precursors have already exited the cell cycle.

We further evaluated the increased cell proliferation of *Hes1*^{Δ/Δ};*Hes5*^{-/-};*Hey1*^{-/-} cochlear. BrdU was administered at E17.5 and the cell proliferation in cochlea sensory epithelium was analyzed at E18.5 (Supplemental Fig. S6). The proportion of Ki67⁺BrdU⁺ cells (Supplemental Fig. S6A,B, arrows) to Ki67⁺ cells was only 9.2%, suggesting that the majority of dividing cells (Ki67⁺) did not enter the S phase during the past day. Similarly, the proportion of Ki67⁺BrdU⁺ cells to BrdU⁺ cells was only 6.6%, suggesting that the majority of the cells that entered the S phase during the past day exited the cell cycle. Furthermore, only a few Ki67⁺BrdU⁺ and BrdU⁺ cells were found in pairs, suggesting that continuously dividing cells are rare during E17.5-18.5 in *Hes1*^{Δ/Δ};*Hes5*^{-/-};*Hey1*^{-/-} cochlear. BrdU⁺ or Ki67⁺ hair cells were occasionally detectable (Supplemental Fig. S6C,D), suggesting that some dividing cells can give rise to hair cells. Nevertheless, the number of dividing cells in the prosensory domain seemed to be too small to account for the overall increase of the cell number in the mutant cochlear (IHC:OHC = 1:3 in the wild type and 2:5 in *Hes1*^{Δ/Δ};*Hes5*^{-/-};*Hey1*^{-/-}). Because there were many Ki67⁺ and BrdU⁺ cells near the border of the prosensory domain (Fig. 6), some excessive hair and supporting cells could derive from the outside of the prosensory domain, although definitive evidence for this possibility remains to be demonstrated.

Prosensory domain appears to be unaffected in Hes/Hey mutants

We next examined whether increased cell proliferation is involved in expansion of the prosensory domain. In the control cochleae, p27^{Kip1} and Sox2 expression occurred at both E14.5 and E15.5 (Fig. 7C,D,K,L), and similar expression was observed in *Hes1*^{Δ/Δ};*Hes5*^{-/-};*Hey1*^{-/-} cochlea (Fig. 7G,H,O,P). The size of the p27^{Kip1} and Sox2

expression region was similar between the control and *Hes1^{ΔΔ};Hes5^{-/-};Hey1^{-/-}* cochlea, suggesting that the prosensory domain did not expand in *Hes/Hey* mutants. MyosinVI⁺ hair cells and Prox1⁺ supporting cells were mostly absent at E14.5 in the control and *Hes1^{ΔΔ};Hes5^{-/-};Hey1^{-/-}* cochlea (Fig. 7A,B,E,F). At E15.5, there were some MyosinVI⁺ outer hair cells and Prox1⁺ supporting cells in both the control and *Hes1^{ΔΔ};Hes5^{-/-};Hey1^{-/-}* cochlea (Fig. 7I,J,M,N). Thus, the onset of hair cell and supporting cell differentiation in the prosensory domain was not affected by *Hes/Hey* deficiency.

The cell proliferation feature of the prosensory domain of the control siblings and *Hes1^{ΔΔ};Hes5^{-/-};Hey1^{+/-}* mutants was further examined. BrdU was administered at E13.5 and the cell proliferation was analyzed at E14.5. The majority of BrdU⁺ cells were negative for Ki67 in both cochleae at E14.5, indicating that most proliferating prosensory cells at E13.5 became quiescent at E14.5 (Supplemental Fig. S7). These results suggest that *Hes/Hey* deficiency did not affect the timing of cell cycle exit.

Hearing loss by Hes/Hey deficiency

Some of the *Hes/Hey* compound mutant mice, which exhibited excessive hair cell and supporting cell formation, survived more than 2 months. We next recorded auditory brainstem responses of these mutant mice to determine whether excessive hair cells affect the hearing ability. Both control and mutant mice showed hearing loss at a high frequency (40kHz) probably due to the background containing C57BL/6. The thresholds of the lower frequencies were not significantly affected in *Hes5*-null (#566) and *Hey1*-null (#495) mice (Table 1), while *Hes1^{+Δ};Hes5^{-/-};Hey1^{-/-}* mice (#692) exhibited some mild hearing defect at 20kHz (Table 1). By contrast, in *Hes1^{ΔΔ};Hes5^{+/-};Hey1^{+/-}* (#562) and *Hes1^{ΔΔ};Hes5^{-/-};Hey1^{+/-}* mice (#923), the thresholds of the lower frequencies were significantly increased, indicating severe hearing loss (Table 1).

We next examined the histology of the cochlea of adult mutant mice. The cochlea of *Hes1^{+Δ};Hes5^{-/-};Hey1^{+/-}* mice had a mild increase in the hair cell number at E18.5 (Fig. 2I) but seemed to be normal at 9 months of age (Supplemental Fig. S8A), suggesting that excessive hair cells were eliminated after birth. *Hes1^{ΔΔ};Hes5^{-/-};Hey1^{+/-}* cochlea had nearly two rows of inner hair cells (Fig. 2F,I) and the polarity of the stereocilia and kinocilium of inner hair cells was disorganized at E18.5 (Fig. 5C,D). Notably, histological analysis of *Hes1^{ΔΔ};Hes5^{-/-};Hey1^{+/-}* cochlea at six months of age showed that inner hair cells formed into almost one row (Supplemental Fig. S8B,C), indicating that the number of excessive inner hair cells decreased after birth. There

were still excessive outer hair cells in the adult *Hes1^{Δ/Δ};Hes5^{-/-};Hey1^{+/-}* cochlea at six months of age (Supplemental Fig. S8B,C), but they seemed to be decreased compared with E18.5 cochlea. Indeed, many extra hair cells underwent apoptosis in *Hes/Hey* mutant cochlear at P3 (Supplemental Fig. S9). The polarity of outer hair cells was maintained almost normally, but the alignment of outer hair cells was disturbed (Supplemental Fig. S8B,C). Neurons of spiral ganglion were maintained in the adult *Hes1^{Δ/Δ};Hes5^{-/-};Hey1^{+/-}* cochlea, suggesting that the hearing loss was not due to the defect of spiral ganglion (Supplemental Fig. S8D,E). These data together suggest that excessive hair cells do not improve the hearing ability, and that the polarity and alignment of these cells are important for hearing. Moreover, many excessive hair cells died rather than became integrated into the hair cell rows to function as normal hair cells.

Discussion

Reduced Hes/Hey gene dosage leads to graded phenotypes in hair and supporting cell formation

It has been shown that Notch signaling inhibits hair cell formation, and that Notch-mediated lateral inhibition regulates hair cell versus supporting cell development from common precursors (Lanford et al., 1999; Kienan et al., 2005; Brooker et al., 2006; Takebayashi et al., 2007). Several Notch target genes, including *Hes1*, *Hes5*, *Hey1*, *Hey2* and *HeyL*, have been reported to be expressed in supporting cells (Lanford et al., 2000; Zheng et al., 2000; Zine et al., 2001; Murata et al., 2006; Hayashi et al., 2008; Li et al., 2008; Hartman et al., 2009; Doetzlhofer et al., 2009). The expression of *Hes1*, *Hes5* and *Hey1* occurred in different but overlapping subtypes of supporting cells. *Hey1* mRNA was expressed broadly and continuously from E11.5 to E17.5, whereas *Hes1* expression was faint at all ages. *Hes5* expression was limited to differentiating cells after E14.5. These expression patterns do not necessarily reflect the activities of endogenous Hes/Hey proteins. It has been reported that *Hes1* deficiency leads to predominant increase of the inner hair cell number, whereas *Hes5* deficiency causes predominant increase of the outer hair cell number (Zheng et al., 2000; Zine et al., 2001). Such tendency was not observed in *Hes1*, *Hes5* and *Hey1* compound hetero- or homozygous mutant mice. Mice that lacked either *Hes5* or *Hey1* or both in *Hes1^{+/-}* background exhibited mild increase of inner and outer hair cell formation, while mice that lacked *Hes1* in *Hes5^{+/-};Hey1^{+/-}* background showed more significant increase of

both inner and outer hair cell formation. When all *Hes1*, *Hes5* and *Hey1* were inactivated, both inner and outer hair cells were further increased in number. These results indicate that all *Hes1*, *Hes5* and *Hey1* are involved in inhibition of both inner and outer hair cell formation, although *Hes1* seems to be the most effective. In addition, hair cell overproduction was always accompanied by supporting cell overproduction if at least one allele of either *Hes1*, *Hes5* or *Hey1* was intact. Thus, all *Hes1*, *Hes5* and *Hey1* have a similar activity to promote supporting cell versus hair cell specification.

When all *Hes1*, *Hes5* and *Hey1* genes were inactivated, even more hair cells were produced without excessive formation of supporting cells. This defect is similar to the one caused by treatment with a γ -secretase inhibitor (Takebayashi et al., 2007) and to the one of *Dll1*; *Jag2* double mutant cochlea (Kiernan et al., 2005). Furthermore, death of supporting cells was not increased in *Hes1* ^{$\Delta\Delta$} ;*Hes5*^{-/-};*Hey1*^{-/-} cochlea at E14.5 (data not shown), suggesting that hair cell overproduction without excessive formation of supporting cells is due to fate conversion of supporting cells to hair cells in addition to increased supporting cell formation. Therefore, *Hes/Hey* genes may cooperatively hold the supporting cell property and prevent the fate conversion from supporting cells into hair cells. Our results suggest that *Hes1*, *Hes5* and *Hey1* are the main Notch effectors in the developing cochleae, cooperatively determine the strength of Notch signaling, and maintain the supporting cell property.

***Hes/Hey* genes function to induce supporting cells**

It has been reported that ectopic hair cells were surrounded by ectopic supporting cells (Woods et al., 2004; Driver et al., 2008; Li et al., 2008). Hair cells are thought to use specific inductive mechanisms to recruit surrounding cells to develop as supporting cells, although the precise mechanisms for this induction are unknown (Driver et al., 2009). Our results suggest that *Hes/Hey* genes are needed for induction of supporting cells.

We observed that supporting cells still remained in *Hes1* ^{$\Delta\Delta$} ;*Hes5*^{-/-};*Hey1*^{-/-} cochlea, and that *Hey2* and *HeyL* were expressed in these supporting cells, suggesting that *Hey2* and *HeyL* might compensate the deficiency of *Hes1*, *Hes5* and *Hey1*. It is known that *Hey2* is controlled by the Fgf pathway, and that this pathway specifically regulates pillar cell development (Doetzlhofer et al. 2009), suggesting that *Hey2* is involved in supporting cell formation. *HeyL* may also have an activity to induce supporting cells, although the specific function of *HeyL* in cochlea remains to be determined.

What contributes excessive hair and supporting cell production

We found that if at least one allele of *Hes1*, *Hes5*, or *Hey1* was intact, both hair cells and supporting cells were formed excessively in the mutant cochlea. The total numbers of hair cells and supporting cells increased in *Hes1*^{ΔΔ};*Hes5*^{-/-};*Hey1*^{+/-}, *Hes1*^{ΔΔ};*Hes5*^{+/-};*Hey1*^{-/-} and *Hes1*^{ΔΔ};*Hes5*^{-/-};*Hey1*^{-/-} cochleae compared with *Hes1*^{+/-};*Hes5*^{+/-};*Hey1*^{+/-} cochleae. Among the former three genotypes, the total cell number seemed to be unchanged. The precise mechanism of the increase of the total cell number in these mutants remained to be determined, but our results suggest that abnormal Ki67⁺ cells and BrdU-incorporating cells remained at a later stage in the sensory epithelium of *Hes1*^{ΔΔ};*Hes5*^{-/-};*Hey1*^{-/-} cochleae, indicating that prolonged proliferation contributes to excessive hair cell and supporting cell formation. The cell types that are inhibited from proliferation by *Hes/Hey* genes are not known, but these cells could be common precursors for hair cells and supporting cells because excessive hair cells and supporting cells are often found in pairs in the mutant cochleae (Figs. 3 and 4). However, there were much fewer Ki67⁺ or BrdU⁺ hair cells in *Hes1*^{ΔΔ};*Hes5*^{-/-};*Hey1*^{+/-} and *Hes1*^{ΔΔ};*Hes5*^{+/-};*Hey1*^{-/-} cochleae than in *Hes1*^{ΔΔ};*Hes5*^{-/-};*Hey1*^{-/-} cochleae (Fig. 6M), although the total cell numbers were similar in these mutants (Supplementary Fig. S3), suggesting that prolonged proliferation alone cannot explain the defects.

Alternatively, excessive cells could be provided by reserved cells in the prosensory domain that are not supposed to differentiate into hair cells or adjacent supporting cells if *Hes/Hey* genes properly function. The total number of hair and supporting cells seemed to reach the plateau in *Hes1*^{ΔΔ};*Hes5*^{-/-};*Hey1*^{+/-}, *Hes1*^{ΔΔ};*Hes5*^{+/-};*Hey1*^{-/-} and *Hes1*^{ΔΔ};*Hes5*^{-/-};*Hey1*^{-/-} cochleae (Supplementary Fig. S3), and it is possible that such reserved cells had been consumed in these mutants. Proliferating cells near the border of the prosensory domain also could contribute to hair and supporting cell increase.

The mechanism of Notch-mediated regulation of cell proliferation

The mechanism of how Notch signaling inhibits cell proliferation in the developing cochlea is not known. Previous studies showed that sustained *Hes1* expression leads to G1 arrest of neural progenitor cells by repressing the expression of cell cycle regulators such as cyclin D1, whereas oscillating *Hes1* expression promotes cell cycle progression (Baek et al., 2006; Shimojo et al., 2008). *Hes1* expression could be sustained in supporting cells, and without sustained *Hes1* expression, these cells could

undergo extra cell divisions, although the dynamics of *Hes1* expression remains to be determined.

It was reported that the CDK inhibitor p27^{Kip1} regulates the cell cycle exit, and that deletion of p27^{Kip1} results in extension of cell proliferation in the auditory sensory epithelium (Chen et al., 1999; Löwenheim et al., 1999., Lee et al., 2006). In the developing cochlear epithelium of *Hes1*-null mice, p27^{Kip1} expression occurred prematurely, and BrdU-incorporating dividing cells were reduced in number, suggesting that the Notch-*Hes1* pathway may promote proliferation of sensory precursor cells via down-regulation of p27^{Kip1} expression (Murata et al., 2009). This phenotype is opposite to our present results of *Hes1* conditional knock out cochlea. This difference could be due to the different timing of *Hes1* inactivation. In the cochlea of *Emx2*^{+Cre} mice, Cre activity is weak in the prosensory domain at E11.5 but fully active by E13.5, suggesting that Cre-induced *Hes1* inactivation mostly occurred after specification of the prosensory domain. By contrast, in conventional *Hes1*-null mice, *Hes1* is inactive before the prosensory domain formation. The mechanism of how Notch signaling differentially regulates cell proliferation depending on developmental stages remains to be determined.

Notch signaling regulates the cell alignment and polarity

Both inner and outer hair cells are aligned in rows (one and three rows, respectively), and each hair cell has a unique polarity of stereociliary bundles. If at least one allele of *Hes1*, *Hes5*, or *Hey1* was intact, the hair cell rows including extra ones were relatively well maintained in the *Hes/Hey* compound mutant cochlea. However, when all three genes were inactivated, hair cell rows became wavy and sometimes disconnected. In the former case, excessive hair cells were accompanied by excessive supporting cells, whereas in the latter case, hair cells were overproduced at the expense of supporting cells. These results suggest that proper ratios of hair cells and supporting cells may be important for organization of the cell alignment. In E18.5 *Hes1*^{ΔΔ};*Hes5*^{-/-};*Hey1*^{-/-} cochlea, the polarity of stereocilia and kinocilium was also severely disorganized, suggesting that supporting cells may help hair cells keep the normal polarity. An increase in cell numbers in the organ of Corti could also affect the mechanical properties of the tissue, thereby affecting the cell alignment. Further studies will be required to understand the role of supporting cells in maintenance of the alignment and polarity of hair cells.

Some of the compound mutant mice survived more than two months, and we were able to assess their hearing ability. Both *Hes1*^{+Δ};*Hes5*^{-/-};*Hey1*^{+/-} and

Hes1^{+/-};Hes5^{+/-};Hey1^{-/-} mice exhibited normal hearing ability. In contrast, *Hes1^{Δ/Δ};Hes5^{+/-};Hey1^{+/-}* and *Hes1^{Δ/Δ};Hes5^{-/-};Hey1^{+/-}* mice exhibited severe hearing impairment. Histological analysis showed that *Hes1^{Δ/Δ};Hes5^{-/-};Hey1^{+/-}* cochlea had less excessive inner hair cells in the adult than at E18.5. It suggests that excessive inner hair cells died after birth probably because they could be non-functional. There were still excessive outer hair cells in the adult *Hes1^{Δ/Δ};Hes5^{-/-};Hey1^{+/-}* cochlea but notably, these excessive hair cells were not helpful to the hearing ability. The polarity of outer hair cells was maintained almost normally, but the alignment of outer hair cells seemed to be disturbed. It is not clear whether the disturbance of outer hair cell alignment can be the only reason for hearing loss or not, and further analysis will be needed to know the effects of *Hes/Hey* deficiency on hearing function. These results imply that excessive hair cells are not useful for hearing, and that the polarity and alignment may be important for the survival and the function of hair cells. Thus, for the purpose of regenerative medicine, not only the number but also the polarity and alignment of hair cells might be important for recovery of the hearing ability.

Conclusion

Genetic inactivation of *Hes1*, *Hes5* and *Hey1* was achieved in the cochlea epithelium after prosensory domain formation. These results suggest that *Hes1*, *Hes5* and *Hey1* cooperatively inhibit hair cell formation and abnormal proliferation, and that at least one allele of either *Hes1*, *Hes5* or *Hey1* is sufficient for supporting cell production and inhibition of fate conversion from supporting cells to hair cells. Abnormal hair cell and supporting cell formation leads to disorganized cell alignment and polarity, leading to hearing loss despite hair cell overproduction.

Acknowledgements

The authors thank Dr. Frank Costantini and Dr. Shinichi Aizawa for ROSA26-CFP mice and *Emx2^{+/-Cre}* mice, respectively. T.T. was supported by Restart Postdoctoral Fellowship of the Japan Society for the Promotion of Science.

References

Baek JH, Hatakeyama J, Sakamoto S, Ohtsuka T, Kageyama R (2006) Persistent and high levels of

- Hes1 expression regulate boundary formation in the developing central nervous system. *Development* 133:2467-76.
- Brooker R, Hozumi K, Lewis J (2006) Notch ligands with contrasting functions: Jagged1 and Delta1 in the mouse inner ear. *Development* 133:1277-86.
- Chen P, Segil N (1999) p27 (Kip1) links cell proliferation to morphogenesis in the developing organ of Corti. *Development* 126:1581-90.
- Doetzlhofer A, Basch ML, Ohyama T, Gessler M, Groves AK, Segil N (2009) Hey2 regulation by FGF provides a Notch-independent mechanism for maintaining pillar cell fate in the organ of Corti. *Dev Cell* 16:58-69.
- Driver EC, Pryor SP, Hill P, Turner J, Rüther U, Biesecker LG, Griffith AJ, Kelley MW (2008) Hedgehog signaling regulates sensory cell formation and auditory function in mice and humans. *J Neurosci* 28:7350-8.
- Driver EC, Kelley MW (2009) Specification of cell fate in the mammalian cochlea. *Birth Defects Res C Embryo Today* 87:212-21.
- Hartman BH, Hayashi T, Nelson BR, Bermingham-McDonogh O, Reh TA (2007) Dll3 is expressed in developing hair cells in the mammalian cochlea. *Dev Dyn* 236:2875-83.
- Hartman BH, Basak O, Nelson BR, Taylor V, Bermingham-McDonogh O, Reh TA (2009) Hes5 expression in the postnatal and adult mouse inner ear and the drug-damaged cochlea. *J Assoc Res Otolaryngol* 10:321-40.
- Hatakeyama J, Bessho Y, Katoh K, Ookawara S, Fujioka M, Guillemot F, Kageyama, R (2004) Hes genes regulate size, shape and histogenesis of the nervous system by control of the timing of neural stem cell differentiation. *Development* 131:5539-5545.
- Hayashi T, Kokubo H, Hartman BH, Ray CA, Reh TA, Bermingham-McDonogh O (2008) Hesr1 and Hesr2 may act as early effectors of Notch signaling in the developing cochlea. *Dev Biol* 316:87-99.
- Holley M, Rhodes C, Kneebone A, Herde MK, Fleming M, Steel KP (2010) Emx2 and early hair cell development in the mouse inner ear. *Dev Biol* 340:547-56.
- Imayoshi I, Shimogori T, Ohtsuka T, Kageyama R (2008) Hes genes and neurogenin regulate non-neural versus neural fate specification in the dorsal telencephalic midline. *Development* 135:2531-41.
- Kada S, Nakagawa T, Ito J (2008) A mouse model for degeneration of the spiral ligament. *J Assoc Res Otolaryngol* 10:161-72.
- Kageyama R, Ohtsuka T, Kobayashi T (2007) The Hes gene family: repressors and oscillators that orchestrate embryogenesis. *Development* 134:1243-1251.
- Kelley MW (2006) Regulation of cell fate in the sensory epithelia of the inner ear. *Nat Rev Neurosci* 7:837-849.

- Kiernan AE, Cordes R, Kopan R, Gossler A, Gridley T (2005) The Notch Ligands DLL1 and JAG2 act synergistically to regulate hair cell development in the mammalian inner ear. *Development* 132:4353–4362.
- Kimura J, Suda Y, Kurokawa D, Hossain ZM, Nakamura M, Takahashi M, Hara A, Aizawa S (2005) *Emx2* and *Pax6* function in cooperation with *Otx2* and *Otx1* to develop caudal forebrain primordium that includes future archipallium. *J Neurosci* 25:5097-108.
- Kokubo H, Miyagawa-Tomita S, Nakazawa M, Saga Y, Johnson RL (2005) Mouse *hesr1* and *hesr2* genes are redundantly required to mediate Notch signaling in the developing cardiovascular system. *Dev Biol* 278: 301-309.
- Lanford PJ, Lan Y, Jiang R, Lindsell C, Weinmaster G, Gridley T, Kelley MW (1999) Notch signaling pathway mediates hair cell development in mammalian cochlea. *Nat Genet* 21:289–292.
- Lanford PJ, Shailam R, Norton CR, Gridley T, Kelley MW (2000) Expression of *Math1* and *HES5* in the cochleae of wildtype and *Jag2* mutant mice. *J Assoc Res Otolaryngol*. 1:161-71.
- Lee YS, Liu F, Segil N (2006) A morphogenetic wave of p27Kip1 transcription directs cell cycle exit during organ of Corti development. *Development* 133:2817-26.
- Li S, Mark S, Radde-Gallwitz K, Schlisner R, Chin MT, Chen P (2008) *Hey2* functions in parallel with *Hes1* and *Hes5* for mammalian auditory sensory organ development. *BMC Dev Biol* 26: 8-20.
- Löwenheim H, Furness DN, Kil J, Zinn C, Gültig K, Fero ML, Frost D, Gummer AW, Roberts JM, Rubel EW, Hackney CM, Zenner HP (1999) Gene disruption of p27(Kip1) allows cell proliferation in the postnatal and adult organ of corti. *Proc Natl Acad Sci U S A* 96:4084-8.
- Morrison A, Hodgetts C, Gossler A, Hrabé de Angelis M, Lewis J (1999) Expression of *Delta1* and *Serrate1* (*Jagged1*) in the mouse inner ear. *Mech Dev* 84; 169-172.
- Murata J, Tokunaga A, Okano H, Kubo T (2006) Mapping of notch activation during cochlear development in mice: implications for determination of prosensory domain and cell fate diversification. *J Comp Neurol* 497:502-18.
- Murata J, Ohtsuka T, Tokunaga A, Nishiike S, Inohara H, Okano H, Kageyama R (2009) Notch-Hes1 pathway contributes to the cochlear prosensory formation potentially through the transcriptional down-regulation of p27Kip1. *J Neurosci Res* 87:3521-34.
- Rhodes CR, Parkinson N, Tsai H, Brooker D, Mansell S, Spurr N, Hunter AJ, Steel KP, Brown SD (2003) The homeobox gene *Emx2* underlies middle ear and inner ear defects in the deaf mouse mutant *pardon*. *J Neurocytol* 32:1143-54.
- Shimojo H, Ohtsuka T, Kageyama R (2008) Oscillations in notch signaling regulate maintenance of neural progenitors. *Neuron* 58:52-64.
- Srinivas S, Watanabe T, Lin CS, William CM, Tanabe Y, Jessell TM, Costantini F (2001) Cre

reporter strains produced by targeted insertion of EYFP and ECFP into the ROSA26 locus. BMC Dev. Biol. 1: 4.

Takebayashi S, Yamamoto N, Yabe D, Fukuda D, Kojima K, Ito J, Honjo T (2007) Multiple roles of notch signaling in cochlear development. Dev Biol 307:165–178.

Woods C, Montcouquiol M, Kelley MW (2004) Math1 regulates development of the sensory epithelium in the mammalian cochlea. Nat Neurosci 7: 1310–1318.

Yamamoto N, Okano T, Ma X, Adelstein RS, Kelley MW (2009) Myosin II regulates extension, growth and patterning in the mammalian cochlear duct. Development 136:1977-86.

Zheng JL, Shou J, Guillemot F, Kageyama R, Gao WQ (2000) Hes1 is a negative regulator of inner ear hair cell differentiation. Development 127:4551-60.

Zine A, Aubert A, Qiu J, Therianos S, Guillemot F, Kageyama R, de Ribaupierre F (2001) Hes1 and Hes5 activities are required for the normal development of the hair cells in the mammalian inner ear. J Neurosci 21:4712–4720.

Figure legends

Figure 1. Expression of *Hes1*, *Hes5* and *Hey1* in the developing otic vesicle and cochlea. **A-B**, *Venus-Hes1* was detected by anti-GFP antibody at E11.5 (green, arrow). Venus (a YFP variant) fragment was knocked in at the *Hes1* locus so that the Venus-Hes1 fusion protein was expressed under the control of the endogenous *Hes1* promoter. *Hes1* was slightly expressed in the epithelium of presumptive cochlea at E11.5. Co-labeling is a prosensory marker, Sox2 (red), and DAPI (blue). **C-W**, in-situ hybridization of *Hes1* (C, R-T), *Hes5* (D-G,O-Q) and *Hey1* (H-K,U-W). At E11.5, *Hey1* was broadly expressed in the prosensory domain, while *Hes5* was not detected in the epithelium of otocyst (D-F, H-J, arrows indicate the presumptive cochlea). At E13.5, *Hey1* was expressed in the prosensory domain, whereas *Hes1* and *Hes5* expression were not seen in the cochlea duct (C,G,K, asterisks indicate the prosensory domain). At E14.5, *Hes1* expression was faint, but occurred at a higher level in the greater epithelial ledge (S). *Hes5* was expressed around hair cells. *Hey1* expression was broad in the apical turn (U) and more limited to the prosensory domain in the mid-basal turn (V). At E17.5, *Hes1* was expressed in Hensen cells (T), while *Hes5* was expressed in inner pharyngeal cells and Deiters's cells (Q). *Hey1* expression occurred in Kollikers organ, Deiters' cells and Hensen cells (W). *Atoh1* expression was also shown for the landmark of inner (i) and outer (o) hair cells (L-N). Scale bars: A, 500µm; B, 100µm; C (for C-K), 200µm; L (for L,M,O,P,R,S,U,V), 100µm; N (for N,Q,T,W), 50µm.

Figure 2. Graded increase in hair cell formation with a reduction in *Hes/Hey* gene dosage. **A,B**, E18.5 cochleae of *Emx2^{+Cre}* and *Hes1^{Δ/Δ};Hes5^{-/-};Hey1^{-/-}* mice. The gross appearance of the cochlea epithelia in *Hes/Hey* mutants seemed to be normal. Otic capsules and stria vascularis were removed. **C-H**, Whole mount immunolabeling of MyosinVI. A reduction in *Hes/Hey* gene dosage led to graded increase in hair cell formation. Red and green asterisks indicate excessive inner and outer hair cells, respectively. *Hes1* conditional knockout cochlea (E-H) had more inner and outer hair cells than *Hes1* heterozygous mutants (C,D). **I**, The inner (IHC) and outer hair cell (OHC) numbers per 100μm were shown. Hair cells were counted over the length of 1600μm from the basal end of E18.5 cochlea, corresponding to approximately one quarter of the cochlea length. Wild type (WT), *Emx2^{+Cre}* and *Hes1^{+Δ};Hes5^{+/-};Hey1^{+/-}* mice were not significantly different from one another in both inner and outer hair cell numbers. In *Hes1* conditional knockout mice, the numbers of both inner and outer hair cells were significantly increased ($p<0.01$) from the other groups. The effect of *Hes5* and *Hey1* was limited compared with *Hes1*, but *Hes5* and *Hey1* deficiency strengthened the phenotype of *Hes1* conditional knockout. Error bars indicate standard deviation. Scale bars: A (for A, B), 100μm; C (for C-H), 100μm

Figure 3. Defects of supporting cells surrounding outer hair cells in *Hes/Hey* mutants. **A-F**, Representatives of MyosinVI⁺ hair cells and Prox1⁺ supporting cells in E18.5 cochlea sections. At this stage, Prox1 was expressed in the nuclei of pillar cells, supporting cells between inner and outer hair cells (D, yellow arrows), and Deiters' cells, supporting cells surrounding outer hair cells (D, yellow arrowheads). *Hes1^{+Δ};Hes5^{+/-};Hey1^{+/-}* cochlea sections showed a normal pattern; 1 inner hair cell, 3 outer hair cells and 5 Prox1⁺ cells (A,D). In *Hes1^{Δ/Δ};Hes5^{-/-};Hey1^{+/-}* cochlea sections, there were 2 inner hair cells, 4 outer hair cells and 6 Prox1⁺ cells (B,E), suggesting that the outer hair cell increase was accompanied by Deiters' cell increase (see yellow asterisks in K). In *Hes1^{Δ/Δ};Hes5^{-/-};Hey1^{-/-}* cochlea sections, there were 2 inner hair cells, 5 outer hair cells and 4-6 Prox1⁺ cells in most regions (C,F). There were 4 Prox1⁺ cells in F. **G-P**, Whole mount staining of *Hes/Hey* mutant cochleae. Hair cells and supporting cells were visualized by Phalloidin and anti-Prox1 antibody, respectively. In *Hes1^{+Δ};Hes5^{+/-};Hey1^{+/-}* cochlea, there were normal 1 inner hair cell, 3 outer hair cells and 5 rows of Prox1⁺ cells (G,H). In *Hes1^{+Δ};Hes5^{-/-};Hey1^{-/-}* cochlea, excessive hair cells (J, white arrows) and excessive Prox1⁺ cells (I, white arrowheads) were present at the same site, indicating that the fourth row of outer hair cells were surrounded by the

fourth row of Deiters' cells. More hair cells were seen in *Hes1^{Δ/Δ};Hes5^{-/-};Hey1^{+/-}* cochlea and *Hes1^{Δ/Δ};Hes5^{+/-};Hey1^{-/-}* cochlea, and Prox1⁺ cells also increased in number at the same place, forming mostly 6 rows (K-N). In *Hes1^{Δ/Δ};Hes5^{-/-};Hey1^{-/-}* cochlea, even more hair cells were formed (P), and at the same site, Prox1⁺ cells were disarranged or missing (O, asterisks). **Q**, Quantification of outer hair cells (OHC) and Prox1⁺ pillar and Deiters' cells (PC + DC) in the basal and middle turn on serial sections of E18.5 cochleae. The total number of each cell type per cochlea duct section was calculated. Error bars indicate standard deviation. Scale bars: C (for A-F), 20μm; H (for G-P), 20μm

Figure 4. Defects of supporting cells surrounding inner hair cells in *Hes/Hey* mutants. **A-I**, Whole mount immunostaining of p27^{Kip1} and p75^{NTR} (A,B,D,E,G,H) and section immunohistochemistry of p27^{Kip1} and Prox1 (C,F,I). Brackets indicate inner phalangeal cells (A,D,G) and hair cells (B,E,H, shadow of p27^{Kip1}). Inner phalangeal cells and inner hair cells are indicated by asterisks and arrowheads, respectively (C,F,I). In *Hes1^{+Δ};Hes5^{-/-};Hey1^{-/-}* cochlea, an inner hair cell was surrounded by two inner phalangeal cells (A-C). In *Hes1^{Δ/Δ};Hes5^{-/-};Hey1^{+/-}* cochlea, two inner hair cells were surrounded by three inner phalangeal cells (D-F). By contrast, in *Hes1^{Δ/Δ};Hes5^{-/-};Hey1^{-/-}* cochlea, two inner hair cells were surrounded by two inner phalangeal cells (G-I). Thus, overproduction of hair cells by reduction of *Hes/Hey* gene dosage was accompanied by supporting cell overproduction in *Hes1^{Δ/Δ};Hes5^{-/-};Hey1^{+/-}* cochlea, but when all of *Hes1*, *Hes5* and *Hey1* were inactivated, the number of hair cells increased without excess supporting cells. **J**, Quantification of inner hair cells (IHC) and inner phalangeal cells (IPhC) in the basal and middle turn on serial sections of E18.5 cochleae. p27^{Kip1}-positive and Prox1-negative cells were inner phalangeal cells in the greater epithelial ridge. The total number of each cell type per cochlea duct section was calculated. Error bars indicate standard deviation. Scale bar: 10μm.

Figure 5. Disturbed cell alignment and polarity by *Hes/Hey* deficiency. Hair cell stereocilia, kinocilium and pillar cells were labeled with phalloidin (red), anti-acetylated α-tubulin antibody (green), and anti-p75^{NTR} antibody (blue), respectively. In the control cochlea, a bundle of stereocilia (phalloidin⁺) forms a V shape directing outward, and a kinocilium is located at the tip of the V shape (anti-acetylated α-tubulin⁺). The cell alignment and polarity were mostly normal in *Hes1^{+Δ};Hes5^{-/-};Hey1^{-/-}* (A,B). In *Hes1^{Δ/Δ};Hes5^{-/-};Hey1^{+/-}* cochlea, most outer hair cells were properly aligned, and their polarity of stereocilia and kinocilium was mostly normal (C,D). However, although

excessive inner hair cells were aligned in two rows, the polarity of the stereocilia and kinocilium of inner hair cells was somewhat disorganized in *Hes1^{Δ/Δ};Hes5^{-/-};Hey1^{+/-}* cochlea (C,D). In *Hes1^{Δ/Δ};Hes5^{-/-};Hey1^{-/-}* cochlea, the alignment of outer hair cells was severely disorganized, while excessive inner hair cells were also aligned in a disorganized manner (E,F). Furthermore, the polarity of stereocilia and kinocilium of outer and inner hair cells was abnormal, pointing rather randomly (E,F). Scale bar: 10μm.

Figure 6. Abnormal proliferation in the epithelium by *Hes/Hey* deficiency. E18.5 cochlea sections of *Hes/Hey* mutants were examined. Hair cells were labeled by anti-MyosinVI antibody. Sections from apical to middle turn (A,C,E,G,I,K) and from mid-basal to basal turn (B,D,F,H,J,L) of cochleae were analyzed. **A-F**, Ki67⁺ proliferating cells. Proliferating cells were occasionally observed next to hair cells in *Hes1^{Δ/Δ};Hes5^{-/-};Hey1^{-/-}* cochleae (E,F, arrows). **G-L**, BrdU⁺ cells. 50 μg/g body weight of BrdU was given to pregnant mice at E14.5 by a single i.p. injection. BrdU⁺ hair cells were occasionally found in *Hes1^{Δ/Δ};Hes5^{-/-};Hey1^{-/-}* cochleae (K,L, arrows). Scale bars: B (for A-F), 20μm; H (for G-L), 20μm. **M**, Proportion of BrdU⁺ MyosinVI⁺ cells to total MyosinVI⁺ cells. Error bars indicate standard errors.

Figure 7. The prosensory domain appeared to be unaffected in *Hes/Hey* mutants. Immunohistochemistry of E14.5 and E15.5 cochlea sections. **A-H**, At E14.5, MyosinVI⁺ hair cells and Prox1⁺ supporting cells did not appear in both the control and *Hes1^{Δ/Δ};Hes5^{-/-};Hey1^{-/-}* cochlea (A,B,E,F). p27^{Kip1} and Sox2 were expressed similarly in the control and *Hes1^{Δ/Δ};Hes5^{-/-};Hey1^{-/-}* cochlea at this stage (C,D,G,H). **I-P**, At E15.5, there were some MyosinVI⁺ hair cells and Prox1⁺ supporting cells in both the control and *Hes1^{Δ/Δ};Hes5^{-/-};Hey1^{-/-}* cochlea (I,J,M,N). Thus, the onset of hair cell and supporting cell differentiation in the prosensory domain was not affected by *Hes/Hey* deficiency. p27^{Kip1} and Sox2 were expressed similarly in the control and *Hes1^{Δ/Δ};Hes5^{-/-};Hey1^{-/-}* cochlea (K,L,O,P). Scale bars: A (for A-H), 20μm; I (for I-P), 20μm.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

E18.5

Ki67 MyosinVI

Hes1^{+/-}*Δ*;*Hes5*^{+/-};*Hey1*^{+/-}

Hes1^{Δ/Δ};*Hes5*^{+/-};*Hey1*^{+/-}

Hes1^{Δ/Δ};*Hes5*^{+/-};*Hey1*^{-/-}

BrdU MyosinVI

Hes1^{+/-}*Δ*;*Hes5*^{+/-};*Hey1*^{+/-}

Hes1^{Δ/Δ};*Hes5*^{+/-};*Hey1*^{+/-}

Hes1^{Δ/Δ};*Hes5*^{+/-};*Hey1*^{-/-}

M

Figure 7

E14.5

Cre(-) control

Hes1^{Δ/Δ};Hes5^{-/-};Hey1^{-/-}

E15.5

Cre(-) control

Hes1^{Δ/Δ};Hes5^{-/-};Hey1^{-/-}

Mouse No.	Sex	Age (M)	BW (g)	Genotype			ABR thresholds (dB)		
				Hes1	Hes5	Hey1	10kHz	20kHz	40kHz
302	F	6	25.8	+/ Δ	+/-	+/-	25	35	SO
304	F	6	27.7	+/ Δ	+/-	+/-	15	40	95
313	F	6	24.5	+/ Δ	+/-	+/-	15	25	95
495	F	4	31.5	+/ Δ	+/-	-/-	20	20	100
566	F	9	29.5	+/ Δ	-/-	+/-	15	10	90
692	F	2	24.0	+/ Δ	-/-	-/-	20	55	SO
562	F	3	19.1	Δ/Δ	+/-	+/-	90	95	SO
923	M	6	43.3	Δ/Δ	-/-	+/-	85	85	SO

Table 1. Auditory brainstem responses of *Hes/Hey* mutant mice

Auditory brainstem responses (ABR) were recorded with *Hes/Hey* mutant mice and their siblings. All mice showed hearing loss at a high frequency (40kHz), and this was probably due to the background containing C57BL/6. The thresholds of the lower frequencies were increased when both alleles of *Hes1* were lost. SO: scale out.

Supplementary Information

Figure S1. Cre recombinase activity was examined in the developing cochlea of *Emx2^{+/Cre};ROSA26-CFP* mice. **A-E**, E11.5 otic vesicle, sectioned from rostral to caudal. The prosensory domain was labeled with Sox2 (red). The cochlea develops from a tubular outpocketing at the ventral-medial region of the otic vesicle (asterisks in D and E). CFP expression was sparsely seen in the prosensory domain (Sox2⁺) of the presumptive cochlea. **F-J**, E13.5 otic vesicle, sectioned from rostral to cordal. The cochlea appears from G to J (arrows). The prosensory domain was labeled with p27^{Kip1} (red). CFP expression was seen in most epithelial cells of the cochlea duct and some cells of the vestibular system. **K-O**, E18.5 inner ear sections. CFP expression was seen in most epithelial cells of the cochlea duct and some cells of the spiral ganglion (K, L). **P-R**, P1 inner ear sections. CFP expression was seen in most epithelial cells of the cochlea duct and some cells of the spiral ganglion (P, Q). **S**, Bar graph showing the relative expression of *Hes1* mRNA at E14.5 for Cre(-) control, *Hes1*^{+/Δ}, and *Hes1*^{Δ/Δ} genotypes.

arrowheads). Both hair cells and supporting cells were CFP⁺ in the organ of Corti (L). Approximately a half of hair cells in the maculae of the vestibular system were CFP⁺ (M, N). There were no CFP⁺ cells in the ampullae of semicircular canals (O). **P-R**, Whole-mount immunostaining of P1 cochleae. CFP expression was seen throughout the cochlea epithelium (P) and hardly found in the spiral ganglion (P, arrowheads). Almost all hair cells (Q) and Prox1⁺ cells (supporting cells surrounding hair cells, R) expressed CFP. **S**, *Hes1* mRNA level in E14.5 cochlea epithelia was reduced to 64% and 9% in *Emx2^{+Cre};Hes1^{+lf} (Hes1^{+Δ})* and *Emx2^{+Cre};Hes1^{flf} (Hes1^{Δ/Δ})* mice, respectively, compared to the control. Thus, *Hes1* expression was effectively lost by E14.5 in the cochlear of *Hes1* conditional knock out mice. The average mRNA level of the control, Cre(-), was taken as 1. Scale bars: E (for A-E), 100μm; J (for F-J), 100μm; K, 100μm; L, 20μm; M (for M-O), 100μm; P, 200μm; Q (for Q, R), 20μm.

Figure S2. E18.5 cochlea length. The cochleae of *Hes1^{Δ/Δ};Hes5^{-/-};Hey1^{-/-}* mice were significantly shorter than the other genotypes (* $p < 0.01$), but the difference was less than 10%. The cochlea lengths of other genotypes were not significantly different from the wild type (WT).

Figure S3. The average numbers of hair and supporting cells per cochlear duct section. IHC, inner hair cells; IPhC, inner phalangeal cells; OHC, outer hair cells; PC+DC, pillar cells and Deiters' cells.

Figure S4. E18.5 cochlea surface at length of 25%, 50% and 75% from the apex. Excessive hair cells were similarly observed in any part of the mutant cochlea ducts.

Figure S5. *Hey2*, *HeyL* and *Hey3* expression in E18.5 cochleae. In situ hybridization of *Hey2*, *HeyL* and *Hey3* was performed with E18.5 cochlea sections of *Hes1*^{+Δ};*Hes5*^{+/-};*Hey1*^{+/-} and *Hes1*^{Δ/Δ};*Hes5*^{-/-};*Hey1*^{-/-} mice. Immunostaining of MyosinVI was shown in yellow. *Hey2* was expressed at a similar level in the pillar cells of *Hes1*^{+Δ};*Hes5*^{+/-};*Hey1*^{+/-} and *Hes1*^{Δ/Δ};*Hes5*^{-/-};*Hey1*^{-/-} cochleae (A and D, arrows). *Hey2* expression occurred in Hensen cells of *Hes1*^{Δ/Δ};*Hes5*^{-/-};*Hey1*^{-/-} cochlea (D, arrowhead) at a higher level than those of *Hes1*^{+Δ};*Hes5*^{+/-};*Hey1*^{+/-} cochlea. *HeyL* was expressed at a similar level in Kollikers organ of *Hes1*^{+Δ};*Hes5*^{+/-};*Hey1*^{+/-} and *Hes1*^{Δ/Δ};*Hes5*^{-/-};*Hey1*^{-/-} cochleae (B and E, asterisks). *Hes3* was not expressed in *Hes1*^{+Δ};*Hes5*^{+/-};*Hey1*^{+/-} cochlea (C).

Figure S6. E18.5 cochlea sections of *Hes1*^{ΔΔ};*Hes5*^{-/-};*Heyl*^{-/-} mutants. BrdU was administered at E17.5, and immunohistochemistry of Ki67 (red), BrdU (green) and MyosinVI (blue) was performed at E18.5. **A,B**, Arrows indicate Ki67⁺BrdU⁺ cells. Ki67⁺BrdU⁺ cells were rather rare in the organ of Corti and did not accompany adjacent BrdU⁺ cells. **C,D**, There were occasionally BrdU⁺ hair cells (C, arrow) and Ki67⁺ hair cells (D, arrow).

Figure S7. E14.5 cochlea sections of *Hes/Hey* mutants. 50 $\mu\text{g/g}$ body weight of BrdU was administered at E13.5, and immunohistochemistry of Ki67 (red), BrdU (green) and p27Kip1 (blue) was performed at E14.5. p27Kip1⁺BrdU⁺ cells were seen in the prosensory domain. There were more in the basal turn (B,D) than in the apical turn (A,C), but BrdU⁺Ki67⁺ positive cells were not seen in the prosensory domain of control (A,B) and *Hes1* ^{Δ/Δ} ;*Hes5*^{-/-};*Hey1*^{+/-} cochleae (C,D). This result suggests that proliferating prosensory cells at E13.5 became quiescent at E14.5, and that *Hes/Hey* deficiency did not affect the timing of cell cycle exit.

Figure S8. Adult *Hes/Hey* mutant cochleae. **A-C**, Immunostaining of whole-mount cochleae of *Hes1*^{+/Δ};*Hes5*^{-/-};*Hey1*^{+/-} (A, #566 in Table 1) and *Hes1*^{Δ/Δ};*Hes5*^{-/-};*Hey1*^{+/-} (B,C, #923 in Table 1). Hair cells of *Hes1*^{+/Δ};*Hes5*^{-/-};*Hey1*^{+/-} appeared normal (A). In *Hes1*^{Δ/Δ};*Hes5*^{-/-};*Hey1*^{+/-} cochlea, inner hair cells formed into almost one row at 6 months of age (B,C), although the mutant cochlea had two rows of inner hair cells at E18.5. Therefore, excessive inner hair cells decreased in number. Arrow indicates paired inner hair cells (B,C). There were still excessive outer hair cells in the adult *Hes1*^{Δ/Δ};*Hes5*^{-/-};*Hey1*^{+/-} cochlea (B). The polarity and alignment of outer hair cells seemed to be disturbed (B,C). **D,E**, Neurons in spiral ganglion were normally maintained in the cochleae of *Hes1*^{+/Δ};*Hes5*^{-/-};*Hey1*^{+/-} (D) and *Hes1*^{Δ/Δ};*Hes5*^{-/-};*Hey1*^{+/-} (E). Scale bars: A (for A,B,D,E), 100μm; C, 10μm.

Figure S9. Postnatal day 3 *Hes/Hey* mutant cochleae. **A,B**, Immunohistochemistry of MyosinVI (red), TUNEL (green) and DAPI (blue) was performed. **A',B'**, TUNEL (gray scale). Hair cell loss was not observed in *Hes1*^{+/ Δ} ;*Hes5*^{+/-};*Hey1*^{-/-} cochlea (**A** and **A'**), whereas outer hair cells were TUNEL⁺ in *Hes1* ^{Δ/Δ} ;*Hes5*^{+/-};*Hey1*^{-/-} cochlea (**B** and **B'**, arrows), suggesting that extra hair cells died rather than became integrated into the normal network.

	IHC	Hes1	+/ Δ	+/ Δ	+/ Δ	+/ Δ	Δ/Δ	Δ/Δ	Δ/Δ	Δ/Δ
OHC		Hes3&5	+/-	-/-	+/-	-/-	+/-	-/-	+/-	-/-
Hes1	Hes3&5	Hey1	+/-	+/-	-/-	-/-	+/-	+/-	-/-	-/-
+/ Δ	+/-	+/-		NS	NS	NS	**	**	**	**
+/ Δ	-/-	+/-	NS		NS	NS	**	**	**	**
+/ Δ	+/-	-/-	NS	NS		NS	**	**	**	**
+/ Δ	-/-	-/-	NS	NS	NS		**	**	**	**
Δ/Δ	+/-	+/-	**	**	**	**		NS	NS	NS
Δ/Δ	-/-	+/-	**	**	**	**	NS		NS	NS
Δ/Δ	+/-	-/-	**	**	**	**	NS	NS		*
Δ/Δ	-/-	-/-	**	**	**	**	**	**	*	

Table S1. Statistical analysis of the result shown in Fig. 2I.

IHC, inner hair cells (pink); OHC, outer hair cells (light blue). *: $p < 0.05$, **: $p < 0.01$, NS, not significant.

A		IHC	Hes1	+/ Δ	+/ Δ	Δ/Δ	Δ/Δ	Δ/Δ
	OHC		Hes3&5	+/-	-/-	-/-	+/-	-/-
	Hes1	Hes3&5	Hey1	+/-	-/-	+/-	-/-	-/-
	+/ Δ	+/-	+/-		NS	**	**	**
	+/ Δ	-/-	-/-	NS		**	**	**
	Δ/Δ	-/-	+/-	**	**		NS	**
B		IPhC	Hes1	+/ Δ	+/ Δ	Δ/Δ	Δ/Δ	Δ/Δ
	PC+DC		Hes3&5	+/-	-/-	-/-	+/-	-/-
	Hes1	Hes3&5	Hey1	+/-	-/-	+/-	-/-	-/-
	+/ Δ	+/-	+/-		NS	*	*	NS
	+/ Δ	-/-	-/-	NS		NS	NS	NS
	Δ/Δ	-/-	+/-	**	**		NS	NS
				+/ Δ	+/ Δ	Δ/Δ	Δ/Δ	Δ/Δ
				+/-	-/-	-/-	+/-	-/-
				+/-	-/-	+/-	-/-	-/-
				+/ Δ	+/-	-/-		
				+/ Δ	-/-	-/-		
				Δ/Δ	-/-	-/-		

Table S2. Statistical analysis of the results shown in Figs. 3Q and 4J. **A**, Inner hair cells (IHC, pink) and outer hair cells (OHC, light blue). **B**, Inner phalangeal cells (IPhC, yellow) and Prox1-positive pillar cells and Deiters' cells (PC+DC, light green). *, $p < 0.05$; **, $p < 0.01$; NS, not significant.