TITLE:

Inflammation-associated cancer development in digestive organs: mechanisms and roles for genetic and epigenetic modulation.

AUTHOR(S):

Chiba, Tsutomu; Marusawa, Hiroyuki; Ushijima, Toshikazu

CITATION:

Chiba, Tsutomu ...[et al]. Inflammation-associated cancer development in digestive organs: mechanisms and roles for genetic and epigenetic modulation.. Gastroenterology 2012, 143(3): 550-563

ISSUE DATE:

2012-09

URL:

<http://hdl.handle.net/2433/160134>

RIGHT:

© 2012 AGA Institute. Published by Elsevier Inc.; This is not the published version. Please cite only the published version.; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。

Inflammation-Associated Cancer Development in Digestive Organs: Mechanisms and Roles for Genetic and Epigenetic Modulation-

Tsutomu Chiba*, Hiroyuki Marusawa* and Toshikazu Ushijima‡

*Department of Gastroenterology and Hepatology, Graduate School of Medicine, Kyoto University, Kyoto, and ‡Division of Epigenomics, National Cancer Center Research Institute, Tokyo, Japan

Abstract

Chronic inflammation, regardless of infectious agents, plays important roles in the development of various cancers particularly in digestive organs, including *Helicobacter pylori* (*H. pylori*)-associated gastric cancer, hepatitis C virus (HCV)-positive hepatocellular carcinoma, and colitis-associated colon cancers. Cancer development is characterized by stepwise accumulation of genetic and epigenetic alterations of various proto-oncogenes and tumor suppressor genes. During chronic inflammation, infectious agents such as *H. pylori* and HCV as well as intrinsic mediators of inflammatory responses, including proinflammatory cytokines and reactive oxygen and nitrogen species, can induce genetic and epigenetic changes, including point mutations, deletions, duplications, recombinations, and methylation of various tumor-related genes through various mechanisms. Furthermore, inflammation also modulates the expressions of microRNAs that influence the production of several tumor-related mRNAs or proteins. These molecular events induced by chronic inflammation work in concert to alter important pathways involved in normal cellular function, and hence accelerate inflammation-associated cancer development. Among these, recent studies highlighted an important role of activation-induced cytidine deaminase, a nucleotide-editing enzyme essential for somatic hypermutation and class-switch recombination of the immunoglobulin gene, as a genomic modulator in inflammation-associated cancer development.

Keywords: Inflammation; Cancer; *H. pylori*; HCV; Mutation Induction; Epigenetics; DNA methylation; microRNA; Activation-Induced Cytidine Deaminase (AID).

Abbreviations used in this paper: 5-mC, 5-methyl cytosine; 8-OHdG, 8-hydroxydeoxyguanosine; A, adenine; AID, activation-induced cytidine deaminase; APC, adenomatous polyposis coli; AUC, area under the curve; Basp1, brain abundant, membrane attached signal protein 1; BCR, breakpoint cluster region; C, cytosine; *cagPAI*, cytotoxin-associated gene pathogenicity island; CD40, cluster of differentiation 40- TNF receptor superfamily member 5; CDH1, cadherin-1; CDKN, cyclin-dependent kinase inhibitor; COX-2, cyclooxygenase 2; Cxcl2, chemokine (C-X-C motif) ligand 2; DNMT, DNA methyltransferase; DSS, dextran sulfate sodium; EGF, epidermal growth factor; ERK, extracellular signal-regulated kinase; EZH2, enhancer of zeste homolog 2; FIH-1, regulating factor inhibiting hypoxia inducible factor 1; FLNc, filamin C γ ; G, guanine; H, histone; HBV, hepatitis B virus; HCC, hepatocellular carcinoma; HCV, hepatitis C virus; HIF-1, hypoxia-inducible factor-1; IBD, inflammatory bowel disease; IGH, immunoglobulin H; IKK, I- κ B kinase; iNOS, inducible nitric oxide synthase; IL-1 β , interleukin-1 β ; IFN, interferon; JNK, c-Jun N-terminal kinase; Let-7, lethal-7; Lphn2, latrophilin 2; LPS, lipopolysaccharide; Mafg, musculoaponeurotic fibrosarcoma oncogene homolog; MALT, mucosa-associated lymphatic tissue; miRNA, microRNA; MAPK, mitogen activated protein kinase; MBD4, methyl-CpG binding domain protein 4; *miR-7*, microRNA-7; MLH1, mutL homolog 1; MSH2, mutS homolog 2; MSH6, mutS homolog 6; NF- κ B, activation of transcription factor nuclear factor κ B; NLRs, nod-like receptors; PDCD4, programmed cell death 4; PSC, primary sclerosing cholangitis; PTEN, phosphatase and tensin homolog; RASSF1A, RAS-association domain family 1, isoform A; RB, retinoblastoma protein; RNS, reactive nitrogen species; ROC, Receiver-operating characteristic; ROS, reactive oxygen species; SCID, severe combined immunodeficiency; STAT3, signal transducer and activator of transcription 3; SIRT1, sirtuin 1; STAT6, signal transducer and activator of transcription 6; SVR, sustained virological response; T, thymine; Th2, T helper 2 cell; THDB, thrombomodulin; TET1, ten-eleven translocation 1; TLRs, toll-like receptors; TNF- α , tumor necrosis factor- α ; U, uracil; UNG, uracil DNA glycosylase; WEE1, mitosis inhibitor protein kinase

Introduction

Nearly 150 years ago, Rudolf Virchow noted that inflammatory cells are present in tumor tissues and that tumors develop at sites of chronic inflammation; he suggested that chronic inflammation plays important roles in cancer development. Since then, many clinical and epidemiological studies have confirmed a strong association between inflammation and cancer^{1, 2}. For instance, epidemiological studies have shown that approximately 10-15% of cancers were related to chronic infections with viruses, bacteria or parasites³⁻⁷, and moreover, that up to 25% of all cancers were associated with chronic inflammation irrespective of the presence or absence of infection⁵⁻⁷.

In inflammation-associated cancer development, in addition to infectious agents such as *Helicobacter pylori* (*H. pylori*) and hepatitis C virus (HCV), many intrinsic mediators of inflammation including proinflammatory cytokines, eicosanoids, growth factors, and reactive oxygen species (ROS) and reactive nitrogen species (RNS) exert important effects in cancer development through various mechanisms. These include enhancement of cell growth and mobility, induction of angiogenesis, and inhibition of apoptosis. However, a hallmark of cancer development is the stepwise accumulation of various genetic and epigenetic alterations of the genome. Indeed, recent genome-wide analysis of human cancer tissues revealed that a single cancer cell generally possesses approximately 100 mutations in coding regions, 10-20 of which are known as “driver genes” that contribute to cancer development⁸⁻¹⁰, and moreover, that there are many somatic gene rearrangements, including duplications, deletions and inversions in human cancer genomes^{11, 12}. In addition to genetic alterations, recent studies have also shown that chronic inflammation enhances epigenetic changes as represented by DNA methylation¹³. It is estimated that several hundreds to thousands of genes are methylated in a cancer cell¹⁴, and that aberrant DNA methylation is present even in normal-appearing tissues, being involved in field cancerization^{13, 15, 16}.

Digestive organs are inhabited by many microorganisms and are infiltrated by many immune cells in physiological and pathological conditions, and thus they are more or less accompanied by certain levels of inflammation. Here, we review mechanisms of how inflammation is involved in cancer development in digestive organs, particularly focusing on the role of chronic inflammation in inducing genetic and epigenetic changes.

Cancers in Digestive Organs Associated with Inflammation

Many cancers arise in digestive organs. Indeed, gastric cancer remains the third leading cause of cancer death in men and the fifth in women, and colorectal cancer is the third most commonly diagnosed cancer in men and the second most in women world-wide¹⁷. In addition, hepatocellular carcinoma (HCC) is one of the most frequent malignancies and its incidence is increasing not only in an endemic area for the hepatitis virus but also in the United States and other western countries¹⁸. Digestive organs cover a large part of the body surface in contact with the outer environment. Accordingly, they are not only inhabited by many microorganisms but also exposed to ingested food or chemical agents, and therefore infiltrated by many immune cells in pathological as well as normal conditions, supporting the perpetuation of chronic inflammation. Therefore, it is reasonable that many cancers in digestive organs are associated with inflammation.

The best examples of inflammation-associated cancer in humans are gastric cancer and HCC. Since the discovery of *H. pylori* by Warren and Marshall in 1982¹⁹, it has been well established that *H. pylori*-positive patients with chronic gastritis have a significantly higher risk for gastric cancer than *H. pylori*-negative subjects²⁰, and moreover, careful investigations have shown more than 95% positivity for *H. pylori* infection in gastric cancer patients²¹. On the other hand, hepatitis B virus (HBV) and HCV infections account for approximately 60% and 33% of the total HCC cases in developing countries and 23% and 20% in developed countries, respectively^{6, 22}, and the majority of HCCs develop in patients who have chronic hepatitis or liver cirrhosis. Other inflammation-associated cancers in digestive organs are colitic cancers developed in patients with inflammatory bowel disease (IBD) or celiac disease²³⁻²⁵, primary sclerosing cholangitis (PSC)-associated cholangiocarcinoma²⁶, primary biliary cirrhosis-associated HCC²⁷, and Barrett's cancer developed in patients with reflux esophagitis²⁸. In addition, the incidence of pancreatic cancer in patients with chronic pancreatitis is reported to be 4-8 times higher than in the general population²⁹, and more strikingly, the incidence of pancreatic cancer in patients with hereditary pancreatitis is 53 times higher than in the normal population³⁰, indicating that chronic pancreatitis is a risk for pancreatic cancer.

In addition to cancers, inflammation is also a risk for developing various lymphomas in digestive organs. These include *H. pylori*-induced mucosa-associated lymphatic tissue (MALT) lymphoma or plasmacytoma^{31, 32}, HCV-related lymphoma³³, and lymphoma related to celiac disease³⁴.

Mechanisms for Inflammation-Associated Cancer Development

The inflammatory response is coordinated by a large range of mediators, which are released from immune cells, mesenchymal cells and epithelial cells; these mediators exert various functions in maintaining or resolving inflammation, and at the same time are involved in cancer development. Among the mediators, cytokines play central roles in diversifying the inflammatory process, and interleukin (IL)-1 β , tumor necrosis factor (TNF)- α , and IL-6 are known to be the major cytokines important for inflammation and cancer development³⁵⁻³⁷.

IL1- β and TNF- α act directly on epithelial cells to induce activation of transcription factor nuclear factor- κ B (NF- κ B), a key transcription factor mediating inflammation and cancer development^{36, 37}. NF- κ B activation not only promotes growth or suppresses apoptosis of epithelial cells but also stimulates the production of growth factors and cytokines such as epidermal growth factor (EGF) and IL-6, enhances cyclooxygenase (COX)2 induction, and increases ROS production³⁸. The induced COX-2 subsequently has many functions, including enhancement of cell growth and angiogenesis³⁹. ROS modifies protein function⁴⁰. IL-6 activates signal transducer and activator of transcription 3 (STAT3) and thereby enhances cell growth, and stimulates growth factor production, including the Reg protein⁴¹. Interestingly, TNF- α and IL-6 often create a positive-feedback loop during cancer development⁴².

At the same time, these cytokines also activate mitogen-activated protein kinase (MAPK) cascades. For instance, TNF- α and IL-6 have been shown to activate the extracellular signal-regulated kinase (ERK)/MAPK cascade, an important signaling pathway involved in many processes in carcinogenesis including cell proliferation, migration, and angiogenesis^{43, 44}. Similarly, IL1- β , TNF- α , and IL-6 all activate c-Jun N-terminal kinase (JNK). Although JNKs are primarily attributed to proapoptotic cell death or tumor suppression in response to inflammation or various stressors⁴⁵, JNK activation, particularly JNK1, by proinflammatory cytokines has been reported to contribute to inflammation-associated cancer development through cell death-induced “compensatory proliferation”⁴⁵⁻⁴⁸. In this regard, an interesting thing to note is that *H. pylori* directly activates ERK/MAPK and JNK in human gastric cells via a type IV secretion system-dependent mechanism^{49, 50}.

Thus, these mediators of inflammation form a complex of regulatory networks, and appear to work in concert to enhance cancer development. However, for normal cells to be eventually transformed and become cancer cells with clonal expansion, inflammation has to damage cellular DNA, either genetically or epigenetically, leading to permanent alteration within the genome.

Inflammation and Genetic Modulation

Cancer is a genetic disease resulting from stepwise accumulation of genetic and epigenetic alterations that drives the progressive transformation of normal cells into malignant derivatives⁵¹. Inactivation of tumor-suppressor genes and/or activation of oncogenes caused by somatic mutations, DNA copy number changes, or chromosomal aberrations are widely detectable in human cancer cells. Among them, the tumor suppressor *TP53* gene is one of the most frequent targets for genetic alterations in many human cancers⁵². An important point to note is that *TP53* mutations are frequently present also in non-cancerous tissues with chronic inflammation before cancer development. Indeed, multiple genetic changes in the *TP53* gene have been detected in various inflammatory tissues such as IBD^{53, 54}, Barrett's esophagus⁵⁵ and HCV-associated chronic hepatitis⁵⁶. For example, by analyzing the individual crypt mutation burden across plaques of the dysplasia, it was shown that mutations in *TP53* genes could be identified in the majority of inflamed crypts of patients with ulcerative colitis (UC)⁵⁷. Moreover, *TP53* mutations are detectable at the frequencies of 4-15 nucleotides out of 10^4 nucleotides in the hepatocytes of the patients with chronic HCV infection⁵⁶. Normal mutation rates cannot account for such abundant genetic changes that accumulate in inflamed epithelial cells, suggesting that certain molecular mechanisms underlie such a large number of genetic alterations. Therefore, to understand the mechanisms of inflammation-associated tumorigenesis, several possible intrinsic mutagens responsible for genetic aberrations in the inflammatory condition have been proposed. Among them, free radicals and intrinsic DNA mutator enzymes appear to be important candidates in the setting of chronic inflammation (Figure 1).

Free radicals refer to any molecular species with one or more unpaired electron(s), including ROS and RNS³⁸. Interestingly, increases in *TP53* gene mutations at codons 247 and 248 are paralleled by an enhanced expression of nitric oxide synthase (iNOS) in the inflamed lesions of the colonic tissues of patients with ulcerative colitis⁵⁴. HCV infection also induces iNOS mRNA expression, thereby enhancing nitric oxide (NO) production, which in turn results in DNA breaks and enhanced mutation frequencies⁵⁸. Moreover, an increased level of NO accelerated spontaneous tumor development, mostly lymphomas, in *Trp53*-deficient mouse model infected with *Cryptosporidium parvum*⁵⁹.

In the inflammatory condition, cellular ROS levels are substantially elevated, and nucleic acids exposed to ROS generate various modified bases such as oxidatively altered purines and pyrimidines⁶⁰. These modified nucleic acids could induce the

putative DNA damage, including single- or double-stranded DNA breaks, DNA intrastrand adducts, and DNA protein crosslinks⁶¹. In addition, ROS alters the mismatch repair function and allows mutations to accumulate in microsatellite sequences⁶². It has been well recognized that oncogene activation is capable of inducing genomic instability in precancerous lesions as well as cancer cells⁶³. In this regard, ROS is also a putative mediators that links excessive activity of oncogene products and DNA damage. For example, oncogene *c-MYC* overexpression results in DNA damage prior to the S phase in association with the ROS induction in normal human fibroblasts⁶⁴. These findings suggested that the cumulative situation of ROS production, a condition of so-called oxidative stress, is involved in both the initiation and progression of inflammation-associated cancers through the induction of genetic instability.

Importantly, the typical mutation pattern induced by oxidative stress cannot account for a mutation signature observed in many human cancer tissues, particularly in inflammation-associated cancers. Among the oxidized nucleosides, one of the common products of free radical attack on DNA is 8-hydroxydeoxyguanine (8-OHdG), which is considered to be a biomarker of oxidative stress⁶⁵. The typical pattern of nucleotide alterations induced by 8-OHdG is guanine (G)/cytosine (C) to thymine (T)/adenine (A) transversions, which have been observed in the *RAS* oncogene and *TP53* tumor suppressor gene in lung and liver cancers^{66, 67}. However, recent genome wide analyses clearly demonstrated that G/C to T/A transversions account for a minor proportion of the total mutations identified in human cancer cells, and instead C/G to T/A transitions are the most prevalent mutation pattern in various cancer tissues, including inflammation-associated cancers⁶⁸. Thus, it appears reasonable to assume that there is an alternative mechanism that accounts for the most frequent mutational pattern, C/G to T/A transitions, detected in many human cancer tissues.

Recently, several human enzymes that are capable of inducing nucleotide alterations have been identified, providing a new avenue for understanding mutagenesis mechanisms⁶⁹. Among them, activation-induced cytidine deaminase (AID) is a well defined molecule involved in DNA mutations in the human genome. Through its enzymatic activity, AID can deaminate C on target DNA to produce a uracil (U), and therefore turns a DNA C:G pair into a U:G mismatch. When DNA replication starts before recognition by the repair system, a U:G mismatch gives rise to C/G to T/A transition. Alternatively, recognition of a U:G mismatch by uracil-DNA-glycosylase (UNG) or mutS homolog 2 (MSH2)/mutS homolog 6 (MSH6) heterodimer induces mutations in the U:G mismatch or at the nearby A:T site (Figure 2). As a result, AID can induce any type of mutations⁷⁰. Under physiological conditions, AID contributes to

generating antibody gene diversification in activated B lymphocytes by inducing somatic hypermutation and class switch recombination of immunoglobulin gene⁷¹. In sharp contrast to the favorable function of AID in the immune system, the role of AID in tumorigenesis through induction of genetic instability was first suggested in hematopoietic malignancies. A number of studies have demonstrated that increased AID expression in various neoplasms of the B lymphocytic lineage was associated with unfavorable mutations and chromosomal translocations^{72, 73}. For instance, AID has been shown to be responsible for the chromosomal breaks in *c-MYC* leading to a *c-MYC/immunoglobulin H (IGH)* translocation in B cell lymphoma⁷⁴. Moreover, AID induces *breakpoint cluster region (BCR)-Abelson murine leukemia viral oncogene homolog 1 (ABLI)* mutations leading to Imatinib resistance in chronic myeloid leukemia cells⁷⁵. Since the target of AID-mediated genotoxic effects was not restricted to immunoglobulin genes and a variety of other genes also received the AID-mediated mutations in B cells⁷⁰, it was not surprising that aberrant upregulation of AID induced genetic alterations in various tumor-related genes, leading to the transformation of hematopoietic cells.

As described, activation of NF- κ B is induced in response to various inflammatory stimulations, and is deeply involved in multiple processes of cancer initiation and progression³⁶. Interestingly, NF- κ B is a major transcription factor for AID in B cells that is activated through cluster of differentiation 40-TNF receptor superfamily member 5 (CD40) ligation by T cells⁷⁶, suggesting that AID might link NF- κ B activation and genetic instability in non-lymphoid cells in the setting of inflammation. In agreement with this hypothesis, AID expression is induced in response to proinflammatory cytokine stimulation via the NF- κ B-dependent pathway in various epithelial cells (Figure 3). In hepatocytes, AID expression is induced by TNF- α through the I- κ B kinase (IKK)-dependent NF- κ B signaling pathway⁷⁷. Consistent with a previous finding that the HCV core protein triggers the activation of NF- κ B in hepatocytes⁷⁸, the HCV core protein itself also up-regulates endogenous AID in cultured hepatocytes⁷⁷. NF- κ B-mediated induction of AID expression is not limited to hepatocytes. In human gastric epithelial cells, AID expression is induced by TNF- α stimulation via activation of NF- κ B, but not detected in non-stimulated cells⁷⁹. More interestingly, aberrant AID expression is induced by the infection of a pathogenic *H. pylori* strain, the cytotoxin-associated gene pathogenicity island (*CagPAI*)-positive strain that is capable of introducing bacterial virulence factors into the host cells through a type-IV secretion system and activating NF- κ B, indicating that both bacterial factors introduced into epithelial cells and the inflammatory mediators such as TNF- α and IL-1 β induced by

H. pylori infection cooperatively promote aberrant AID expression in *H. pylori*-infected gastric mucosal cells. Similar to hepatocytes and gastric mucosal cells, TNF- α stimulation resulted in upregulation of endogenous AID in human colonic cells via the IKK-dependent NF- κ B signaling pathway⁸⁰. In addition, IL-4 and IL-13, which are involved in Th2 type immune response in IBD, induced aberrant AID expression in a signal transducer and activator of transcription 6 (STAT6)-dependent manner in human colonic epithelial cells⁸⁰. Of note, IL-4 is known to induce AID also in B cells⁷¹.

Consistent with the in vitro analyses, aberrant AID expression is widely detectable in not only various inflammation-associated cancer tissues but also in a variety of inflamed epithelial tissues where tumorigenic risk is high, including chronic hepatitis and cirrhosis caused by HCV infection⁵⁶, chronic gastritis caused by *H. pylori* infection⁷⁹, IBD⁸⁰, PSC⁸¹, and the columnar cell-lined Barrett's esophagus⁸².

The impact of AID expression in non-lymphoid epithelial cells was clarified using both in vivo and in vitro systems with aberrant AID expression. Constitutive and ubiquitous AID expression in transgenic mice induced lymphoma development via the accumulation of somatic mutations in various non-immunoglobulin genes, including the proto-oncogene *c-Myc*⁸³. More importantly, further phenotypic analyses revealed that AID transgenic mice also develop neoplasia in epithelial tissues, including lung, liver and stomach accompanied by the emergence of *Trp53* mutations, indicating that aberrant AID expression in epithelial cells can induce genetic instability leading to cancer development^{83, 84}. It is widely recognized that the frequently mutated tumor-related genes differ among different cancers. For instance, nucleotide alterations in the *K-RAS* are detectable in almost all human pancreatic cancers⁸⁵, while it is relatively low in other human tumors. Similarly, the *c-MYC* is a frequent target for genetic alterations in human lung cancers, while its nucleotide alterations are rare in hepatocellular carcinoma⁸⁶. However, the mechanisms underlying the accumulation of organ-specific genomic changes in oncogenic pathways are not well known. Interestingly, organ-specific changes in mutational profiles were observed in the epithelial tissues of the AID transgenic mice. Indeed, the *c-Myc* gene was frequently mutated in non-cancerous tissue of the lung, while *K-ras* gene mutations were frequently detectable in gastric cancer developed in AID transgenic mice⁸⁴. Thus, the organ-specific differences in the mutational profiles in AID transgenic mice suggest the possibility that the target preference of AID-induced mutagenesis in different tissues might contribute to the diversity of tissue-specific oncogenic pathways in various epithelial organs.

In vitro analyses using human cultured cells with constitutive AID expression

revealed that *TP53* mutations were frequently induced by AID genotoxic activity in hepatocytes, and gastric, colonic, and bile duct epithelial cells^{77, 79-81}. Similar to the *TP53* gene, the cyclin-dependent kinase inhibitor (*CDKN*)-2*B*-*CDKN2A* locus was identified as a target for AID-mediated genotoxic activity. The *CDKN2B-CDKN2A* locus encodes the potent suppressor proteins, p16^{INK4a}, p15^{INK4b}, and p14^{ARF}, that regulate the activities of the retinoblastoma protein (RB) and the TP53 transcription factor. Aberrant AID expression preferentially induces somatic mutations at the *CDKN2B-CDKN2A* locus in gastric epithelial cells and biliary cells^{81, 87}. Moreover, comparative genomic hybridization analysis clearly demonstrated that constitutive AID activation in cultured gastric epithelial cells caused submicroscopic deletions as represented by copy number losses of various chromosomal loci, especially at the *CDKN2B-CDKN2A* locus at 9p21. Copy number reduction of *Cdkn2b-Cdkn2a* was also seen in the gastric mucosa of AID transgenic mice⁸⁷. In agreement with the preferential deletions at the *CDKN2B-CDKN2A* locus in gastric epithelial cells by AID introduction, AID expression was required for inducing DNA single-strand breaks in the *CDKN2B* gene in leukemia cells⁸⁸, and furthermore, the deletion of the *CDKN2B-CDKN2A* locus is frequently detectable in AID-expressing lymphoid blast crisis leukemia cells⁷⁵. These findings suggest that AID can induce both mutations and deletions at the same gene locus, and moreover, that the representative tumor-suppressor genes, *TP53* and *CDKN2B-CDKN2A* may be common targets for AID-mediated genotoxic effects in various human tissues in the setting of inflammation.

Finally, a recent finding that a deficiency of endogenous AID reduced the incidence of both accumulation of somatic mutations in the *Trp53* gene and the development of colitis-associated colorectal cancers further supports the critical role of AID in inflammation-associated cancer development via its ability to induce genetic alterations in tumor-related genes⁸⁹.

Inflammation and Epigenetic Modulation

Epigenetic modifications are DNA-associated modifications that are inherited upon somatic cell replication, which include DNA methylation and histone modifications⁹⁰. Coordinated changes of epigenetic modifications control development and tissue differentiation, and erasure of epigenetic modifications is involved in reprogramming. In somatic cells, DNA methylation is present in repetitive elements, CpG-sparse regions, and in a very limited number of CpG islands^{91, 92}. DNA methylation of a CpG island in a promoter region causes silencing of its downstream gene, whether it is a protein-coding

gene or a miRNA gene, by forming nucleosomes and thus possibly blocking access of RNA polymerase II to the promoter^{93, 94}. In contrast, DNA methylation of a gene body is often associated with increased gene expression^{91, 95}.

Histone modifications denote chemical modifications, such as acetylation, methylation, and ubiquitination of lysine and arginine residues of histones, mainly H3 and H4 but also H2A and H2B⁹³. Specific histone modifications, such as acetylation of histones H3 and H4 (H3Ac and H4Ac) and trimethylation of lysine 4 of histone H3 (H3K4 me3) are associated with active gene transcription. In contrast, di- and trimethylation of H3 lysine 9 (H3K9 me2 and H3K9 me3) and trimethylation of H3 lysine 27 (H3K27 me3) are associated with gene repression. H3K9 me2 represses gene transcription in concert with DNA methylation, while H3K27 me3 works independently of DNA methylation⁹⁶. Trimethylation of H3 lysine 36 (H3K36 me3) is considered to mark exonic regions of active genes. However, the mechanisms of how histone modifications are inherited upon somatic cell replication remains unclear⁹⁷.

In cancer cells, the presence of regional hypermethylation and global hypomethylation has been described^{98, 99}. Regional hypermethylation refers to aberrant DNA methylation of promoter CpG islands physiologically kept unmethylated^{95, 100}. If aberrant methylation is induced in a promoter CpG island, it consistently induces silencing of its downstream gene⁹⁰. Many tumor-suppressor genes that have promoter CpG islands, such as *CDKN2A*, mutL homolog 1 (*MLH1*), cadherin-1 (*CDH1*), and RAS-association domain family 1, isoform A (*RASSF1A*), can be permanently inactivated by aberrant DNA methylation as drivers, that have significant roles in cancer development. At the same time, most of the aberrant DNA methylation of promoter CpG islands are considered to be passengers, that play no role in carcinogenesis¹⁴. Several hundreds to thousands of promoter CpG islands are aberrantly methylated in a cancer, and the number is too large for all of them to be drivers. Moreover, most of the genes methylated in cancers are not expressed in normal tissues^{101, 102}, and such genes are considered to be not involved in carcinogenesis. Global hypomethylation was shown to be causally involved in carcinogenesis by inducing genomic instability¹⁰³. In addition, induction of H3K27 me3 is considered to be an alternative mechanism to induce gene silencing⁹⁶, and aberrant H3K27 me3 was observed in promoter regions consisting of 200-600 genes^{96, 104}. Again, the number is very large, and most are expected to be passengers.

As inducers of aberrant DNA methylation, aging was first indicated¹⁰⁵, and chronic inflammation was then suggested by the presence of aberrant DNA methylation of specific tumor-suppressor genes in non-cancerous colonic mucosae of patients with

IBD^{106, 107}. Aberrant DNA methylation was present more frequently in liver tissues of patients with HCC than in those with metastatic liver tumors¹⁰⁸. By measuring methylation levels of passenger genes in gastric mucosae of *H. pylori*-infected individuals, a very close association between *H. pylori* infection and high methylation levels in gastric mucosa was demonstrated¹⁵. Aberrant DNA methylation is particularly prominent in chronic inflammation-associated cancers, such as gastric cancer, HCCs, colitic cancer, cholangiocarcinoma, Barrett's cancer, and pancreatic cancer¹³. These findings strongly indicated that the major inducer of aberrant DNA methylation is chronic inflammation.

Levels of aberrant DNA methylation accumulated in normal-appearing tissues correlate with the risk of gastric, colon, breast, and renal cancers^{15, 109-112}. Such accumulation mainly involves passenger and driver genes to some extent, and is considered to form an epigenetic field for cancerization (epigenetic field defect) (Figure 4)¹¹³. Chronic inflammation-associated cancers are known to show multiple events, which can be explained by the presence of a field defect in normal-appearing tissues. Along with the accumulation of genetic alterations, an epigenetic field defect is deeply involved in the development of inflammation-associated cancers. The degree of epigenetic field defect can be easily measured using methylation levels of marker genes¹¹⁴, which are passenger genes in most cases and show relatively high methylation levels in predisposed tissues¹¹³.

Mechanistic studies, including cause and effect of accumulated aberrant DNA methylation and chronic inflammation, were conducted using animal models. When *H. pylori*-induced inflammation was suppressed by cyclosporine A in Mongolian gerbils, induction of aberrant DNA methylation was markedly suppressed, while the number of *H. pylori* in gastric mucosae was unaffected¹⁶. This indicated that inflammation, not *H. pylori* itself, is critical for induction of aberrant DNA methylation. Expression analysis of inflammation-related genes showed that expression levels of *Il1b*, *Nos*, *Tnf*, and chemokine (C-X-C motif) ligand 2 (*Cxcl2*) correlated with methylation levels in gastric mucosae. *H. pylori*-induced inflammation was capable of inducing aberrant DNA methylation, but not repeated induction of acute inflammation by ethanol or a high sodium concentration¹¹⁵. *Il1 β* , *Nos2* and *Tnf* were specifically upregulated by the *H. pylori*-induced inflammation. Notably, in humans, a polymorphism of the *IL1B* promoter was associated with not only gastric cancer susceptibility³⁵, but also the presence of the CpG island methylation phenotype in gastric cancers¹¹⁶.

Another animal model for methylation induction by chronic inflammation is mouse colitis induced by administration of dextran sulfate sodium (DSS)¹¹⁷. Aberrant DNA

methylation of multiple genes occurred in DSS-induced colitis mucosae before induction of colon tumors, showing an epigenetic field¹¹⁸. The induction of aberrant DNA methylation was unaffected even in severe combined immunodeficiency (SCID) mice that lacked T and B cells, suggesting that infiltrated macrophages might be critical for methylation induction. Gene expression analysis in colonic mucosae in wild-type and SCID mice showed that expression levels of *Illb*, *Nos*, and *Ifng* were associated with methylation induction in colonic mucosae. Taken together with the finding in the *H. pylori*-infected gerbils, infiltration of macrophages and resulting secretion of IL-1 β and TNF- α as well as production of active oxygen species are believed to be involved in induction of aberrant DNA methylation in epithelial cells (Figure 1).

Several in vitro studies have been conducted to examine inflammatory signals that lead to methylation induction in target cells. Treatment of insulinoma or blood cells with IL-1 β or a NO donor induced methylation of endogenous genes by increasing activity of DNA methyltransferase(s) (DNMTs)¹¹⁹. IL-6 induces DNMT1 transcription by increasing its promoter activity and suppressing miR-148a and miR-152, both of which target DNMT1^{120,121}. Although some studies suggested that DNA methylation is induced by IL-1 β or IL-6, the changes were marginal possibly because identification of appropriate target CpG islands was difficult and the levels of increase were too small to be detected by ordinary methods. Prostaglandin E2 treatment of cancer cell lines increased DNMT1 and DNMT3B expression, and induced DNA methylation of specific genes, which was also observed in vivo¹²².

In contrast to in vitro studies, mRNA expression levels of Dnmt1, Dnmt3A, and Dnmt3B were not increased in vivo, such as colonic mucosae with DSS-induced colitis¹⁶, and human gastric tissues with *H. pylori* infection¹²³. In line with these *in vivo* findings, O'Hagan et al. recently showed in vitro that oxidative damage recruits complexes containing DNMTs, a histone deacetylase (sirtuin 1, SIRT1), and histone methyltransferase (enhancer of zeste homolog 2, EZH2) to damaged chromatin, and induces DNA methylation¹²⁴. They also showed that, in *Apc*^{Min} mice infected with an inflammation-inducing bacterium, Dnmt1 and Ezh2 are recruited to promoter CpG islands of untranscribed or minimally transcribed genes. Promoter CpG islands with H3K27 me3 and without RNA polymerase II are susceptible to DNA methylation induction^{101, 102}.

Taken together, we can hypothesize a model for aberrant DNA methylation induction in vivo (Figure 5). Inflammatory signals mainly from macrophages, such as IL-1 β , TNF- α , and IL-6, and oxidative stress, possibly produced by NO synthase, are likely to recruit a complex with DNMT1 and EZH2 to promoter CpG islands with

H3K27 me3 flag and without protection by RNA polymerase II. Since DNA methylation is harmful to a gene, aberrant DNA methylation is likely to be induced only rarely and at scattered CpG sites within a CpG island (seeds of methylation)¹²³. Most "seeds of methylation" are erased during cell replication, but can lead to dense methylation of a CpG island at very low frequencies^{125, 126}. If such dense methylation is induced in a promoter CpG island of a tumor-suppressor gene, the tissue becomes predisposed to carcinogenesis, and forms an epigenetic field defect.

In addition to aberrant DNA methylation of promoter CpG islands, cancer cells are characterized by global DNA hypomethylation as well as aberrant hypomethylation of oncogenes^{99, 127}. Gastric mucosa infected by *H. pylori* displays global hypomethylation¹²⁸. In this regard, it is interesting to note that AID has been recently shown to be involved in active DNA demethylation during fetal development¹²⁹. Mechanistically, AID deaminates 5-methyl cytosine (5-mC) to yield T. This T would be subsequently removed by either of the T:G mismatch-specific glycosylases, thymidine DNA glycosylase or methyl-CpG binding domain protein 4 (MBD4). The resulting abasic site would then be replaced by an unmethylated C via base excision repair processes, resulting in DNA demethylation. Notably, AID participates in active demethylation by 5-mC hydroxylase, ten-eleven translocation 1 (TET1), and subsequent gene expression in the dentate gyrus of adult mouse brain¹³⁰. Thus, whether AID is involved in DNA demethylation during cancer development is an interesting topic for future studies¹³¹. The fact that AID targets the chromatin marked by H3K4 me3 histone modification¹³², in contrast to preferential DNA methylation at promoter CpG islands with H3K27 me3 histone modification^{101, 102}, might suggest opposing mechanisms for induction of DNA methylation and demethylation.

Inflammation and MicroRNA Modulation

MicroRNAs (miRNAs) are short noncoding RNAs that regulate the expression of many target genes post-transcriptionally, and are thus involved in a variety of cellular functions. Recent studies have revealed that miRNAs have important roles in cancer development as either oncogenes or tumor suppressor genes by regulating various cancer-related proteins or mRNA expressions^{133, 134}. Indeed, cancer cells are associated with dysregulation of many miRNA expressions, which occurs through a variety of mechanisms, such as genetic changes, epigenetic regulation, or altered expression of transcription factors¹³⁵. On the other hand, miRNA expression is also altered in inflammatory conditions, and such alterations in miRNA expression appear to play roles

not only in controlling chronic inflammation, but also in promoting cancer development^{136, 137}. Many of the changes in miRNA expressions observed in inflammatory tissues are derived from immune cells that may participate in hematopoietic tumorigenesis¹³⁸. However, recent reports have shown that inflammation also induces changes in cancer-related miRNAs in epithelial cells, suggesting a direct link between alteration of miRNA expressions and inflammation-associated cancer development^{139, 140}.

miRNA expressions in epithelial cells can be altered during inflammation through various mechanisms such as NF- κ B activation by toll-like receptors (TLRs) or cytokine stimulation and STAT3 phosphorylation by IL-6 or other cytokines¹³⁹⁻¹⁴³. Among those, several miRNAs are identified as tumor suppressor miRNAs. *miR-7* targets not only *Egfr* but also latrophilin (*Lphn2*), brain abundant, membrane attached signal protein 1 (*Baspl1*) and musculoaponeurotic fibrosarcoma oncogene homolog (*Mafg*), and thus is considered to be a tumor suppressor miRNA¹⁴². In a mouse model of inflammation-associated cancer development, expression of *miR-7* has been shown to be inhibited by activated macrophages in *Helicobacter*-infected gastritis mucosa, being involved in gastric cancer development, whereas it was increased in germ-free conditions¹⁴². Lethal-7 (*Let-7*), consisting of 12 members, targets the *RAS* family and *c-MYC*^{144, 145}, and genomic locations of *let-7* family members are frequently deleted in colon cancers and other solid cancers¹⁴⁶. NF- κ B activation enhances *Lin28B* transcription that causes posttranscriptional inhibition of *let7* family member expression, and *let-7* directly inhibits IL-6 expression, a cytokine often produced in cancer cells. Thus, reduction of *let-7* expression by NF- κ B activation appears to play a role in a positive feedback loop for NF- κ B activation through an increase of IL-6 in cancer cells¹⁴⁷.

miR-155, a possible oncogenic miRNA, is involved in blood cell maturation, immune responses and autoimmune disorders, and high expression of *miR-155* is associated with the development of myeloproliferative disorders¹⁴⁸. Recent studies have revealed a direct link between elevation of *miR-155* and tumor formation and development in gastric and colon cancers^{148, 149}. *miR-155* expression is induced by NF- κ B, IFN- β and TLR stimulation¹⁵⁰, and thus enhanced by *H. pylori* and lipopolysaccharide (LPS) treatment¹⁵¹. Recently, Croce et al.¹⁴³ reported that TNF- α /LPS stimulation enhances *miR-155* expression in association with an increased mutation rate. They also showed that *miR-155* targets mitosis inhibitor protein kinase 1 (WEE1), which blocks cell-cycle progression, and therefore reasoned that reduction of WEE1 by *miR-155* allowed cell division to continue even in the presence of DNA

damage, leading to enhanced mutation induction. In another study, they also demonstrated that *miR-155* promotes gene mutations by down-regulating the core mismatch repair proteins, hMSH2, hMSH6 and hMLH1¹⁵². Of particular interest are the recent reports showing that *miR-155* negatively regulates AID in B cells. Teng et al.¹⁵³ demonstrated that *miR-155* is upregulated in B cells undergoing class-switch recombination, and regulates the germinal center reaction by modulating AID. Moreover, *miR-155* has been suggested to inhibit *MYC-IGH* translocation by reducing AID mRNA and protein in B cells¹⁵⁴. Thus, although an inhibitory effect of *miR-155* on AID has not been examined in non-B cells, *miR-155* may also have a tumor suppressor function in epithelial cells by inhibiting AID production.

A miRNA expression pattern distinct from normal colonic mucosa has been found in the colonic mucosa and colitic tumor of patients with IBD as well as mice with colonic inflammation, including upregulation of *miR-21* and *miR-3*¹⁵⁵. *miR-21* is one of the most highly expressed miRNAs in colonic tissues of patients with ulcerative colitis¹⁵⁵, and its expression is enhanced by LPS and IL-6 through STAT3 activation, targeting key regulators of cell proliferation and apoptosis such as phosphatase and tensin homolog (PTEN) and programmed cell death 4 (PDCD4)¹⁵⁶. Olaru et al.¹⁵⁷ recently demonstrated that in colitic cancer development *miR-31* expression increases in a stepwise fashion from IBD to cancer, and that *miR-31* directly targets regulating factor inhibiting hypoxia inducible factor 1 (FIH-1), decreasing its repressor activity for hypoxia-inducible factor 1 (HIF-1).

It is now evident that miRNAs exert various functions in inflammation-associated cancer development. However, alterations of miRNA expression observed in inflammatory tissues occur in both immune cells and epithelial cells. Accordingly, it is important to dissect miRNA changes in the two cell types, as the patterns of the miRNA changes are different between immune cells and epithelial cells. Further elucidation of the changes of miRNA expression, particularly in epithelial cells, will facilitate our understanding of the role of tumor-related miRNAs in inflammation-associated cancer development.

Application to Cancer Prevention, Diagnostics, and Therapeutics

In order to prevent inflammation-associated cancer development, it is crucial to cure or control inflammation. Indeed, it has been repeatedly demonstrated that long-term therapy with anti-inflammatory drugs resulted in fewer appearances of tumors¹⁵⁸. The best way to control chronic inflammation is, of course, to eliminate

causative infections. In other cases unrelated to infection such as IBD and PSC, one approach is to block the action of key regulators of inflammation. In this regard, NF- κ B or STAT3, and their activators TNF- α or IL-6, respectively, may be good targets for suppressing the inflammatory response. However, since treatment usually needs to be continued for long periods to control chronic inflammation, agents without serious side effects with lower costs should be developed. For this purpose, many natural agents derived from vegetables, fruits, spices, and their components have been tested. Among them, curcumin, derived from yellow spice turmeric (*Curcuma longa*) has been used for centuries, and has been shown to suppress NF- κ B- as well as STAT3-regulated inflammation¹⁵⁹, and thus can be administered safely over the long-term¹⁶⁰. Indeed, a recent study showed that curcumin reduced TNF- α expression, prevented cancer-associated weight loss, and induced apoptosis of tumors in patients with colorectal cancer¹⁶¹. Resveratrol, a natural polyphenolic, non-flavonoid antioxidant found in grapes and other berries has been shown to have generalized inhibitory effects on inflammation-related molecules such as NF- κ B, COX2 and tyrosine kinases¹⁶². Recently, Resveratrol was found to alter the expression of many tumor-related miRNAs¹⁶³. Similar types of agents may have the potential to both prevent and treat cancers¹⁶⁴.

In contrast to controlling inflammatory mediators, blocking genetic modulation appears to be difficult. One might consider inhibiting AID. However, because AID plays a critical role in immunoglobulin maturation in B cells, specific targeting for AID in the epithelial cells without affecting AID in B cells is critical. Control of epigenetic modulation can be considered from two aspects: suppression of methylation induction and reversal of induced methylation. Since induction of methylation is not essential in adult somatic cells, control of this process is a promising approach to prevent chronic inflammation-associated cancers. On the other hand, reversal of aberrant DNA methylation is an attractive idea to repair an epigenetic field defect, but targeting only aberrant DNA methylation without affecting physiological DNA methylation is currently very difficult.

H. pylori eradication ameliorates chronic inflammation, and reduces the risk for gastric cancer. However, it is apparent that eradication cannot completely resolve chronic inflammation, as some patients develop gastric cancer even after successful eradication¹⁶⁵. Likewise, some patients with chronic hepatitis or liver cirrhosis due to HCV infection also develop HCC after obtaining sustained virological response (SVR)¹⁶⁶. As such, when inflammation is not appropriately controlled or even when inflammation is resolved after long-standing inflammation, accurate prediction for the

risk of developing cancers in the inflammatory tissues becomes important. As was discussed, carcinogenesis is characterized by a stepwise accumulation of both genetic and epigenetic changes. Importantly, previous data suggested that the extent of those genetic and epigenetic modulations is paralleled with duration or severity of inflammation^{15, 167}, and the degree of epigenetic field defect can be measured relatively easily and accurately. Thus, both qualitative and quantitative detection of these genetic and epigenetic changes in inflammatory tissues or tissues previously exposed to inflammation may provide a good risk marker for inflammation-associated cancer development. Indeed, epigenetic risk markers that can differentiate gastric mucosae of cancer patients from those of healthy individuals with odds ratios between 12.7-36.0 have been isolated^{168, 169}, and a prospective study is now being conducted.

Conclusion

Many cancers in digestive organs develop in the background of chronic inflammation. During chronic inflammation, a variety of mediators for inflammation such as cytokines, growth factors, eicosanoids, ROS and NOS form complex networks for not only maintaining or reducing inflammation but also promoting cell growth, angiogenesis and inhibiting apoptosis. These events eventually merge into and result in both genetic and epigenetic changes of the cellular genome, leading to inflammation-associated cancer development. In particular, AID plays a crucial role in inducing not only mutations, but also chromosomal aberrations during inflammation. Moreover, signals from macrophages with resulting mislocalization of DNMTs appear to be involved in the induction of epigenetic alterations.

Interestingly, epigenetic inactivation of *MLH1* leads to accumulation of genetic alterations¹⁷⁰. At the same time, recent studies have demonstrated that AID induces DNA demethylation through its deaminating activity on methylated cytosines¹³¹. Thus, genetic and epigenetic events are mutually related and work in concert in the development of inflammation-associated cancers.

References

1. Coussens LM, Werb Z. Inflammation and cancer. *Nature* 2002;420:860-867.
2. Walczak H. TNF and ubiquitin at the crossroads of gene activation, cell death, inflammation, and cancer. *Immunol Rev* 2011;244:9-28.
3. Yeh JM, Goldie SJ, Kuntz KM, et al. Effects of *Helicobacter pylori* infection and smoking on gastric cancer incidence in China: a population-level analysis of trends and projections. *Cancer Causes Control* 2009;20:2021-2029.
4. Vennervald BJ, Polman K. Helminths and malignancy. *Parasite Immunol* 2009;31:686-696.
5. Balkwill F, Mantovani A. Inflammation and cancer: back to Virchow? *Lancet* 2001;357:539-545.
6. Hussain SP, Harris CC. Inflammation and cancer: an ancient link with novel potentials. *Int J Cancer* 2007;121:2373-2380.
7. Mantovani A, Allavena P, Sica A, et al. Cancer-related inflammation. *Nature* 2008;454:436-444.
8. Stephens PJ, McBride DJ, Lin ML, et al. Complex landscapes of somatic rearrangement in human breast cancer genomes. *Nature* 2009;462:1005-1010.
9. Pleasance ED, Stephens PJ, O'Meara S, et al. A small-cell lung cancer genome with complex signatures of tobacco exposure. *Nature* 2010;463:184-190.
10. Sjoblom T, Jones S, Wood LD, et al. The consensus coding sequences of human breast and colorectal cancers. *Science* 2006;314:268-274.
11. Pleasance ED, Cheetham RK, Stephens PJ, et al. A comprehensive catalogue of somatic mutations from a human cancer genome. *Nature* 2010;463:191-196.
12. Kumar-Sinha C, Tomlins SA, Chinnaiyan AM. Recurrent gene fusions in prostate cancer. *Nat Rev Cancer* 2008;8:497-511.
13. Ushijima T, Hattori N. Molecular Pathways: Involvement of *Helicobacter pylori*-Triggered Inflammation in the Formation of an Epigenetic Field Defect, and Its Usefulness as Cancer Risk and Exposure Markers. *Clin Cancer Res* 2012;18:923-929.
14. Ushijima T, Asada K. Aberrant DNA methylation in contrast with mutations. *Cancer Sci* 2010;101:300-305.
15. Maekita T, Nakazawa K, Mihara M, et al. High levels of aberrant DNA methylation in *Helicobacter pylori*-infected gastric mucosae and its possible association with gastric cancer risk. *Clin Cancer Res* 2006;12:989-995.
16. Niwa T, Tsukamoto T, Toyoda T, et al. Inflammatory processes triggered by

- Helicobacter pylori* infection cause aberrant DNA methylation in gastric epithelial cells. *Cancer Res* 2010;70:1430-1440.
17. Jemal A, Bray F, Center MM, et al. Global cancer statistics. *CA Cancer J Clin* 2011;61:69-90.
 18. Altekruse SF, McGlynn KA, Reichman ME. Hepatocellular carcinoma incidence, mortality, and survival trends in the United States from 1975 to 2005. *J Clin Oncol* 2009;27:1485-1491.
 19. Warren JR, Marshall B. Unidentified curved bacilli on gastric epithelium in active chronic gastritis. *Lancet* 1983;1:1273-1275.
 20. Uemura N, Okamoto S, Yamamoto S, et al. *Helicobacter pylori* infection and the development of gastric cancer. *N Engl J Med* 2001;345:784-789.
 21. Chiba T, Marusawa H, Seno H, et al. Mechanism for gastric cancer development by *Helicobacter pylori* infection. *J Gastroenterol Hepatol* 2008;23:1175-1181.
 22. Parkin DM. The global health burden of infection-associated cancers in the year 2002. *Int J Cancer* 2006;118:3030-3044.
 23. Bernstein CN, Blanchard JF, Kliever E, et al. Cancer risk in patients with inflammatory bowel disease: a population-based study. *Cancer* 2001;91:854-862.
 24. Askling J, Linet M, Gridley G, et al. Cancer incidence in a population-based cohort of individuals hospitalized with celiac disease or dermatitis herpetiformis. *Gastroenterology* 2002;123:1428-1435.
 25. Elfstrom P, Granath F, Ye W, et al. Low risk of gastrointestinal cancer among patients with celiac disease, inflammation, or latent celiac disease. *Clin Gastroenterol Hepatol* 2012;10:30-36.
 26. Patel T. Cholangiocarcinoma. *Nat Clin Pract Gastroenterol Hepatol* 2006;3:33-42.
 27. Imam MH, Silveira MG, Sinakos E, et al. Long-term Outcomes of Patients With Primary Biliary Cirrhosis and Hepatocellular Carcinoma. *Clin Gastroenterol Hepatol* 2012;10:182-185.
 28. Shaheen NJ, Richter JE. Barrett's oesophagus. *Lancet* 2009;373:850-861.
 29. DiMagno EP, Reber HA, Tempero MA. AGA technical review on the epidemiology, diagnosis, and treatment of pancreatic ductal adenocarcinoma. American Gastroenterological Association. *Gastroenterology* 1999; 117: 1464-1484.
 30. Whitcomb DC, Applebaum S, Martin SP. Hereditary pancreatitis and pancreatic carcinoma. *Ann N Y Acad Sci* 1999;880:201-209.
 31. Wotherspoon AC, Doglioni C, Diss TC, et al. Regression of primary low-grade B

- cell gastric lymphoma of mucosa-associated lymphoid tissue type after eradication of *Helicobacter pylori*. *Lancet* 1993;342:575-577.
32. Kodama Y, Kawabata K, Yoshida S, et al. Malt lymphoma simulating an extramedullary plasmacytoma of the stomach. *Am J Med* 1999;107:530-532.
 33. Hartridge-Lambert SK, Stein EM, Markowitz AJ, et al. Hepatitis C and non-hodgkin lymphoma: The clinical perspective. *Hepatology* 2012;55:634-641.
 34. Smedby KE, Akerman M, Hildebrand H, et al. Malignant lymphomas in coeliac disease: evidence of increased risks for lymphoma types other than enteropathy-type T cell lymphoma. *Gut* 2005;54:54-59.
 35. El-Omar EM, Carrington M, Chow WH, et al. Interleukin-1 polymorphisms associated with increased risk of gastric cancer. *Nature* 2000;404:398-402.
 36. Ben-Neriah Y, Karin M. Inflammation meets cancer, with NF-kappaB as the matchmaker. *Nat Immunol* 2011;12:715-723.
 37. Kuraishy A, Karin M, Grivennikov SI. Tumor promotion via injury- and death-induced inflammation. *Immunity* 2011;35:467-477.
 38. Hussain SP, Hofseth LJ, Harris CC. Radical causes of cancer. *Nat Rev Cancer* 2003;3:276-285.
 39. Wang D, DuBois RN. Eicosanoids and cancer. *Nat Rev Cancer* 2010;10:181-193.
 40. Ziech D, Franco R, Pappa A, et al. Reactive oxygen species (ROS)--induced genetic and epigenetic alterations in human carcinogenesis. *Mutat Res* 2011;711:167-173.
 41. Sekikawa A, Fukui H, Fujii S, et al. REG Ialpha protein mediates an anti-apoptotic effect of STAT3 signaling in gastric cancer cells. *Carcinogenesis* 2008;29:76-83.
 42. Kanda K, Komekado H, Sawabu T, et al. Nardilysin and ADAM proteases promote gastric cancer cell growth by activating intrinsic cytokine signalling via enhanced ectodomain shedding of TNF-alpha. *EMBO Mol Med* 2012;4:396-411.
 43. Schievella AR, Chen JH, Graham JR, et al. MADD, a novel death domain protein that interacts with the type I tumor necrosis factor receptor and activates mitogen-activated protein kinase. *J Biol Chem* 1997;272:12069-12075.
 44. Kamimura D, Ishihara K, Hirano T. IL6-signal transduction and its physiological roles: the signal orchestration model. *Rev Physiol Biochem Pharmacol* 2003;149:1-38.
 45. Chen F. JNK-induced apoptosis, compensatory growth, and cancer stem cells.

- Cancer Res 2012;72:379-386.
46. Liu J, Yan J, Jiang S, et al. Site-specific ubiquitination is required for relieving the transcription factor Miz1-mediated suppression on TNF α -induced JNK activation and inflammation. *Proc Natl Acad Sci* 2012;109:191-196.
 47. Inokuchi S, Aoyama T, Miura K, et al. Disruption of TAK1 in hepatocytes causes hepatic injury, inflammation, fibrosis and carcinogenesis. *Proc Natl Acad Sci USA* 2010;107:844-849.
 48. Chang Q, Zhang Y, Beezhold KJ, et al. Sustained JNK1 activation is associated with altered histone H3 methylations in human liver cancer. *J Hepatol* 2009; 50: 323-333.
 49. Higashi H, Tsutsumi R, Muto S, et al. SHP-2 tyrosine phosphatase as an intracellular target of *Helicobacter pylori* CagA protein. *Science* 2002;295:683-686.
 50. Snider JL, Allison C, Bellaire BH, et al. The β 1 integrin activates JNK independent of CagA, and JNK activation is required for *Helicobacter pylori* CagA+-induced motility of gastric cancer cells. *J Biol Chem* 2008; 283: 13952-13963.
 51. Hanahan D, Weinberg RA. The hallmarks of cancer. *Cell* 2000;100:57-70.
 52. Joerger AC, Fersht AR. Structure-function-rescue: the diverse nature of common p53 cancer mutants. *Oncogene* 2007;26:2226-2242.
 53. Brentnall TA, Haggitt RC, Rabinovitch PS, et al. Risk and natural history of colonic neoplasia in patients with primary sclerosing cholangitis and ulcerative colitis. *Gastroenterology* 1996;110:331-338.
 54. Hussain SP, Amstad P, Raja K, et al. Increased p53 mutation load in noncancerous colon tissue from ulcerative colitis: a cancer-prone chronic inflammatory disease. *Cancer Res* 2000;60:3333-3337.
 55. Barrett MT, Sanchez CA, Prevo LJ, et al. Evolution of neoplastic cell lineages in Barrett oesophagus. *Nat Genet* 1999;22:106-109.
 56. Kou T, Marusawa H, Kinoshita K, et al. Expression of activation-induced cytidine deaminase in human hepatocytes during hepatocarcinogenesis. *Int J Cancer* 2007;120:469-476.
 57. Leedham SJ, Graham TA, Oukrif D, et al. Clonality, founder mutations, and field cancerization in human ulcerative colitis-associated neoplasia. *Gastroenterology* 2009; 136:542-550.
 58. Machida K, Cheng KT, Sung VM, et al. Hepatitis C virus infection activates the immunologic (type II) isoform of nitric oxide synthase and thereby enhances

- DNA damage and mutations of cellular genes. *J Virol* 2004;78:8835-8843.
59. Hussain SP, He P, Subleski J, et al. Nitric oxide is a key component in inflammation-accelerated tumorigenesis. *Cancer Res* 2008;68:7130-7136.
 60. Demple B, Harrison L. Repair of oxidative damage to DNA: enzymology and biology. *Annu Rev Biochem* 1994;63:915-948.
 61. Federico A, Morgillo F, Tuccillo C, et al. Chronic inflammation and oxidative stress in human carcinogenesis. *Int J Cancer* 2007;121:2381-2386.
 62. Gasche C, Chang CL, Rhees J, et al. Oxidative stress increases frameshift mutations in human colorectal cancer cells. *Cancer Res* 2001;61:7444-7448.
 63. Halazonetis TD, Gorgoulis VG, Bartek J. An oncogene-induced DNA damage model for cancer development. *Science* 2008;319:1352-1355.
 64. Vafa O, Wade M, Kern S, et al. c-Myc can induce DNA damage, increase reactive oxygen species, and mitigate p53 function: a mechanism for oncogene-induced genetic instability. *Mol Cell* 2002;9:1031-1044.
 65. Evans MD, Dizdaroglu M, Cooke MS. Oxidative DNA damage and disease: induction, repair and significance. *Mutat Res* 2004;567:1-61.
 66. Takahashi T, Nau MM, Chiba I, et al. p53: a frequent target for genetic abnormalities in lung cancer. *Science* 1989;246:491-494.
 67. Hsu IC, Metcalf RA, Sun T, et al. Mutational hotspot in the p53 gene in human hepatocellular carcinomas. *Nature* 1991;350:427-428.
 68. Greenman C, Stephens P, Smith R, et al. Patterns of somatic mutation in human cancer genomes. *Nature* 2007;446:153-158.
 69. Conticello SG. The AID/APOBEC family of nucleic acid mutators. *Genome Biol* 2008;9:229 (1-10).
 70. Liu M, Duke JL, Richter DJ, et al. Two levels of protection for the B cell genome during somatic hypermutation. *Nature* 2008;451:841-845.
 71. Honjo T, Kinoshita K, Muramatsu M. Molecular mechanism of class switch recombination: linkage with somatic hypermutation. *Annu Rev Immunol* 2002;20:165-196.
 72. Greeve J, Philipsen A, Krause K, et al. Expression of activation-induced cytidine deaminase in human B-cell non-Hodgkin lymphomas. *Blood* 2003;101:3574-3580.
 73. Pasqualucci L, Guglielmino R, Houldsworth J, et al. Expression of the AID protein in normal and neoplastic B cells. *Blood* 2004;104:3318-3325.
 74. Robbiani DF, Bothmer A, Callen E, et al. AID is required for the chromosomal breaks in *c-myc* that lead to *c-myc/IgH* translocations. *Cell* 2008;135:1028-1038.

75. Klemm L, Duy C, Iacobucci I, et al. The B cell mutator AID promotes B lymphoid blast crisis and drug resistance in chronic myeloid leukemia. *Cancer Cell* 2009;16:232-245.
76. Nagaoka H, Tran TH, Kobayashi M, et al. Preventing AID, a physiological mutator, from deleterious activation: regulation of the genomic instability that is associated with antibody diversity. *Int Immunol* 2010;22:227-235.
77. Endo Y, Marusawa H, Kinoshita K, et al. Expression of activation-induced cytidine deaminase in human hepatocytes via NF-kappaB signaling. *Oncogene* 2007;26:5587-5595.
78. Marusawa H, Hijikata M, Chiba T, et al. Hepatitis C virus core protein inhibits Fas- and tumor necrosis factor alpha-mediated apoptosis via NF-kappaB activation. *J Virol* 1999;73:4713-4720.
79. Matsumoto Y, Marusawa H, Kinoshita K, et al. Helicobacter pylori infection triggers aberrant expression of activation-induced cytidine deaminase in gastric epithelium. *Nat Med* 2007;13:470-476.
80. Endo Y, Marusawa H, Kou T, et al. Activation-induced cytidine deaminase links between inflammation and the development of colitis-associated colorectal cancers. *Gastroenterology* 2008;135:889-898.
81. Komori J, Marusawa H, Machimoto T, et al. Activation-induced cytidine deaminase links bile duct inflammation to human cholangiocarcinoma. *Hepatology* 2008;47:888-896.
82. Morita S, Matsumoto Y, Okuyama S, et al. Bile acid-induced expression of activation-induced cytidine deaminase during the development of Barrett's oesophageal adenocarcinoma. *Carcinogenesis* 2011;32:1706-1712.
83. Okazaki IM, Hiai H, Kakazu N, et al. Constitutive expression of AID leads to tumorigenesis. *J Exp Med* 2003;197:1173-1181.
84. Morisawa T, Marusawa H, Ueda Y, et al. Organ-specific profiles of genetic changes in cancers caused by activation-induced cytidine deaminase expression. *Int J Cancer* 2008;123:2735-2740.
85. Almoguera C, Shibata D, Forrester K, et al. Most human carcinomas of the exocrine pancreas contain mutant c-K-ras genes. *Cell* 1988;53:549-554.
86. Thorgeirsson SS, Grisham JW. Molecular pathogenesis of human hepatocellular carcinoma. *Nat Genet* 2002;31:339-346.
87. Matsumoto Y, Marusawa H, Kinoshita K, et al. Up-regulation of activation-induced cytidine deaminase causes genetic aberrations at the *CDKN2b-CDKN2a* in gastric cancer. *Gastroenterology* 2010;139:1984-1994.

88. Feldhahn N, Henke N, Melchior K, et al. Activation-induced cytidine deaminase acts as a mutator in BCR-ABL1-transformed acute lymphoblastic leukemia cells. *J Exp Med* 2007;204:1157-1166.
89. Takai A, Marusawa H, Minaki Y, et al. Targeting activation-induced cytidine deaminase prevents colon cancer development despite persistent colonic inflammation. *Oncogene* 2012;31:1733-1742.
90. Jones PA, Baylin SB. The epigenomics of cancer. *Cell* 2007;128:683-692.
91. Rauch TA, Wu X, Zhong X, et al. A human B cell methylome at 100-base pair resolution. *Proc Natl Acad Sci USA* 2009;106:671-678.
92. Lister R, Pelizzola M, Dowen RH, et al. Human DNA methylomes at base resolution show widespread epigenomic differences. *Nature* 2009;462:315-322.
93. Li B, Carey M, Workman JL. The role of chromatin during transcription. *Cell* 2007;128:707-719.
94. Lin JC, Jeong S, Liang G, et al. Role of nucleosomal occupancy in the epigenetic silencing of the MLH1 CpG island. *Cancer Cell* 2007;12:432-444.
95. Yamashita S, Hosoya K, Gyobu K, et al. Development of a novel output value for quantitative assessment in methylated DNA immunoprecipitation-CpG island microarray analysis. *DNA Res* 2009;16:275-286.
96. Kondo Y, Shen L, Cheng AS, et al. Gene silencing in cancer by histone H3 lysine 27 trimethylation independent of promoter DNA methylation. *Nat Genet* 2008;40:741-750.
97. Margueron R, Reinberg D. Chromatin structure and the inheritance of epigenetic information. *Nat Rev Genet* 2010;11:285-296.
98. Feinberg AP, Tycko B. The history of cancer epigenetics. *Nat Rev Cancer* 2004;4:143-153.
99. Yoshida T, Yamashita S, Takamura-Enya T, et al. Alu and Sata hypomethylation in *Helicobacter pylori*-infected gastric mucosae. *Int J Cancer* 2011;128:33-39.
100. Rauch TA, Zhong X, Wu X, et al. High-resolution mapping of DNA hypermethylation and hypomethylation in lung cancer. *Proc Natl Acad Sci USA* 2008;105:252-257.
101. Takeshima H, Ushijima T. Methylation destiny: Moira takes account of histones and RNA polymerase II. *Epigenetics* 2010;5:89-95.
102. Takeshima H, Yamashita S, Shimazu T, et al. The presence of RNA polymerase II, active or stalled, predicts epigenetic fate of promoter CpG islands. *Genome Res* 2009;19:1974-1982.
103. Chen RZ, Pettersson U, Beard C, et al. DNA hypomethylation leads to elevated

- mutation rates. *Nature* 1998;395:89-93.
104. Enroth S, Rada-Iglesias A, Andersson R, et al. Cancer associated epigenetic transitions identified by genome-wide histone methylation binding profiles in human colorectal cancer samples and paired normal mucosa. *BMC cancer* 2011;11:450.
 105. Issa JP, Ottaviano YL, Celano P, et al. Methylation of the oestrogen receptor CpG island links ageing and neoplasia in human colon. *Nat Genet* 1994;7:536-540.
 106. Hsieh CJ, Klump B, Holzmann K, et al. Hypermethylation of the p16INK4a promoter in colectomy specimens of patients with long-standing and extensive ulcerative colitis. *Cancer Res* 1998;58:3942-3945.
 107. Issa JP, Ahuja N, Toyota M, et al. Accelerated age-related CpG island methylation in ulcerative colitis. *Cancer Res* 2001;61:3573-3577.
 108. Kondo Y, Kanai Y, Sakamoto M, et al. Genetic instability and aberrant DNA methylation in chronic hepatitis and cirrhosis--A comprehensive study of loss of heterozygosity and microsatellite instability at 39 loci and DNA hypermethylation on 8 CpG islands in microdissected specimens from patients with hepatocellular carcinoma. *Hepatology* 2000;32:970-979.
 109. Nakajima T, Maekita T, Oda I, et al. Higher methylation levels in gastric mucosae significantly correlate with higher risk of gastric cancers. *Cancer Epidemiol Biomarkers Prev* 2006;15:2317-2321.
 110. Shen L, Kondo Y, Rosner GL, et al. MGMT promoter methylation and field defect in sporadic colorectal cancer. *J Natl Cancer Inst* 2005;97:1330-1338.
 111. Yan PS, Venkataramu C, Ibrahim A, et al. Mapping geographic zones of cancer risk with epigenetic biomarkers in normal breast tissue. *Clin Cancer Res* 2006;12:6626-6636.
 112. Arai E, Kanai Y, Ushijima S, et al. Regional DNA hypermethylation and DNA methyltransferase (DNMT) 1 protein overexpression in both renal tumors and corresponding nontumorous renal tissues. *Int J Cancer* 2006;119:288-296.
 113. Ushijima T. Epigenetic field for cancerization. *J Biochem Mol Biol* 2007;40:142-150.
 114. Shin CM, Kim N, Park JH, et al. Prediction of the risk for gastric cancer using candidate methylation markers in the non-neoplastic gastric mucosae. *J Pathol* 2012;226:654-665.
 115. Hur K, Niwa T, Toyoda T, et al. Insufficient role of cell proliferation in aberrant DNA methylation induction and involvement of specific types of inflammation.

- Carcinogenesis 2011;32:35-41.
116. Yoo EJ, Park SY, Cho NY, et al. Influence of IL1B polymorphism on CpG island hypermethylation in Helicobacter pylori-infected gastric cancer. *Virchows Arch* 2010;456:647-652.
 117. Rosenberg DW, Giardina C, Tanaka T. Mouse models for the study of colon carcinogenesis. *Carcinogenesis* 2009;30:183-196.
 118. Katsurano M, Niwa T, Yasui Y, et al. Early-stage formation of an epigenetic field defect in a mouse colitis model, and non-essential roles of T- and B-cells in DNA methylation induction. *Oncogene* 2012;31:342-351.
 119. Hmadcha A, Bedoya FJ, Sobrino F, et al. Methylation-dependent gene silencing induced by interleukin 1beta via nitric oxide production. *J Exp Med* 1999;190:1595-1604.
 120. Hodge DR, Xiao W, Clausen PA, et al. Interleukin-6 regulation of the human DNA methyltransferase (HDNMT) gene in human erythroleukemia cells. *J Mol Biol* 2001;276:39508-39511.
 121. Braconi C, Huang N, Patel T. MicroRNA-dependent regulation of DNA methyltransferase-1 and tumor suppressor gene expression by interleukin-6 in human malignant cholangiocytes. *Hepatology* 2010;51:881-890.
 122. Xia D, Wang D, Kim SH, et al. Prostaglandin E(2) promotes intestinal tumor growth via DNA methylation. *Nat Med* 2012;18:224-226.
 123. Nakajima T, Yamashita S, Maekita T, et al. The presence of a methylation fingerprint of Helicobacter pylori infection in human gastric mucosae. *Int J Cancer* 2009;124:905-910.
 124. O'Hagan HM, Wang W, Sen S, et al. Oxidative damage targets complexes containing DNA methyltransferases, SIRT1, and polycomb members to promoter CpG Islands. *Cancer Cell* 2011;20:606-619.
 125. Stirzaker C, Song JZ, Davidson B, et al. Transcriptional gene silencing promotes DNA hypermethylation through a sequential change in chromatin modifications in cancer cells. *Cancer Res* 2004;64:3871-3877.
 126. Ushijima T, Watanabe N, Shimizu K, et al. Decreased fidelity in replicating CpG methylation patterns in cancer cells. *Cancer Res* 2005;65:11-17.
 127. Ehrlich M. DNA hypomethylation in cancer cells. *Epigenomics* 2009;1:239-259.
 128. Bae JM, Shin SH, Kwon HJ, et al. ALU and LINE-1 hypomethylations in multistep gastric carcinogenesis and their prognostic implications. *Int J Cancer* 2012 (in press)
 129. Cortellino S, Xu J, Sannai M, et al. Thymine DNA glycosylase is essential for

- active DNA demethylation by linked deamination-base excision repair. *Cell* 2011;146:67-79.
130. Guo JU, Su Y, Zhong C, et al. Hydroxylation of 5-methylcytosine by TET1 promotes active DNA demethylation in the adult brain. *Cell* 2011;145:423-434.
 131. Fritz EL, Papavasiliou N. Cytidine deaminases: AIDing DNA demethylation? *Genes Dev* 2010;24:2107-2114.
 132. Kato L, Begum NA, Burroughs AM, et al. Nonimmunoglobulin target loci of activation-induced cytidine deaminase (AID) share unique features with immunoglobulin genes. *Proc Natl Acad Sci USA* 2012;109:2479-2484.
 133. Di Leva G, Croce CM. Roles of small RNAs in tumor formation. *Trends Mol Med* 2010;16:257-267.
 134. Croce CM. Causes and consequences of microRNA dysregulation in cancer. *Nat Rev Genet* 2009;10:704-714.
 135. Lu J, Getz G, Miska EA, et al. MicroRNA expression profiles classify human cancers. *Nature* 2005;435:834-838.
 136. Sonkoly E, Pivarcsi A. MicroRNAs in inflammation and response to injuries induced by environmental pollution. *Mutat Res* 2011;717:46-53.
 137. O'Connell RM, Rao DS, Baltimore D. MicroRNA Regulation of Inflammatory Responses. *Annu Rev Immunol* 2012;30:295-312.
 138. Zhao JL, Rao DS, Boldin MP, et al. NF-kappaB dysregulation in microRNA-146a-deficient mice drives the development of myeloid malignancies. *Proc Natl Acad Sci USA* 2011;108:9184-9189.
 139. Padgett KA, Lan RY, Leung PC, et al. Primary biliary cirrhosis is associated with altered hepatic microRNA expression. *J Autoimmun* 2009;32:246-253.
 140. Wu F, Zikusoka M, Trindade A, et al. MicroRNAs are differentially expressed in ulcerative colitis and alter expression of macrophage inflammatory peptide-2 alpha. *Gastroenterology* 2008;135:1624-1635.
 141. Schetter AJ, Heegaard NH, Harris CC. Inflammation and cancer: interweaving microRNA, free radical, cytokine and p53 pathways. *Carcinogenesis* 2010;31:37-49.
 142. Kong D, Piao YS, Yamashita S, et al. Inflammation-induced repression of tumor suppressor miR-7 in gastric tumor cells. *Oncogene* 2011 Dec 5 [Epub ahead of print].
 143. Tili E, Michaille JJ, Wernicke D, et al. Mutator activity induced by microRNA-155 (miR-155) links inflammation and cancer. *Proc Natl Acad Sci USA* 2011;108:4908-4913.

144. Johnson SM, Grosshans H, Shingara J, et al. RAS is regulated by the let-7 microRNA family. *Cell* 2005;120:635-647.
145. Akao Y, Nakagawa Y, Naoe T. let-7 microRNA functions as a potential growth suppressor in human colon cancer cells. *Biol Pharm Bull* 2006;29:903-906.
146. Calin GA, Sevignani C, Dumitru CD, et al. Human microRNA genes are frequently located at fragile sites and genomic regions involved in cancers. *Proc Natl Acad Sci USA* 2004;101:2999-3004.
147. Iliopoulos D, Hirsch HA, Struhl K. An epigenetic switch involving NF-kappaB, Lin28, Let-7 MicroRNA, and IL6 links inflammation to cell transformation. *Cell* 2009;139:693-706.
148. Tili E, Croce CM, Michaille JJ. miR-155: on the crosstalk between inflammation and cancer. *Int Rev Immunol* 2009;28:264-284.
149. Volinia S, Calin GA, Liu CG, et al. A microRNA expression signature of human solid tumors defines cancer gene targets. *Proc Natl Acad Sci USA* 2006;103:2257-2261.
150. O'Connell RM, Taganov KD, Boldin MP, et al. MicroRNA-155 is induced during the macrophage inflammatory response. *Proc Natl Acad Sci USA* 2007;104:1604-1609.
151. Xiao B, Liu Z, Li BS, et al. Induction of microRNA-155 during *Helicobacter pylori* infection and its negative regulatory role in the inflammatory response. *J Infect Dis* 2009;200:916-925.
152. Valeri N, Gasparini P, Fabbri M, et al. Modulation of mismatch repair and genomic stability by miR-155. *Proc Natl Acad Sci USA* 2010;107:6982-6987.
153. Teng G, Hakimpour P, Landgraf P, et al. MicroRNA-155 is a negative regulator of activation-induced cytidine deaminase. *Immunity* 2008;28:621-629.
154. Dorsett Y, McBride KM, Jankovic M, et al. MicroRNA-155 suppresses activation-induced cytidine deaminase-mediated Myc-Igh translocation. *Immunity* 2008;28:630-638.
155. Pekow JR, Kwon JH. MicroRNAs in inflammatory bowel disease. *Inflamm Bowel Dis* 2012;18:187-193.
156. Iliopoulos D, Jaeger SA, Hirsch HA, et al. STAT3 activation of miR-21 and miR-181b-1 via PTEN and CYLD are part of the epigenetic switch linking inflammation to cancer. *Mol Cell* 2010;39:493-506.
157. Olaru AV, Selaru FM, Mori Y, et al. Dynamic changes in the expression of MicroRNA-31 during inflammatory bowel disease-associated neoplastic transformation. *Inflamm Bowel Dis* 2011;17:221-231.

158. Rothwell PM, Fowkes FG, Belch JF, et al. Effect of daily aspirin on long-term risk of death due to cancer: analysis of individual patient data from randomised trials. *Lancet* 2011;377:31-41.
159. Aggarwal BB, Sung B. Pharmacological basis for the role of curcumin in chronic diseases: an age-old spice with modern targets. *Trends Pharmacol Sci* 2009;30:85-94.
160. Kanai M, Imaizumi A, Otsuka Y, et al. Dose-escalation and pharmacokinetic study of nanoparticle curcumin, a potential anticancer agent with improved bioavailability, in healthy human volunteers. *Cancer Chemother Pharmacol* 2012;69:65-70.
161. He ZY, Shi CB, Wen H, et al. Upregulation of p53 expression in patients with colorectal cancer by administration of curcumin. *Cancer Invest* 2011;29:208-213.
162. Delmas D, Lancon A, Colin D, et al. Resveratrol as a chemopreventive agent: a promising molecule for fighting cancer. *Current Drug Targets* 2006;7:423-442.
163. Tili E, Michaille JJ, Adair B, et al. Resveratrol decreases the levels of miR-155 by upregulating miR-663, a microRNA targeting JunB and JunD. *Carcinogenesis* 2010;31:1561-1566.
164. Aggarwal BB, Van Kuiken ME, Iyer LH, et al. Molecular targets of nutraceuticals derived from dietary spices: potential role in suppression of inflammation and tumorigenesis. *Exp Biol Med* 2009;234:825-849.
165. Fukase K, Kato M, Kikuchi S, et al. Effect of eradication of *Helicobacter pylori* on incidence of metachronous gastric carcinoma after endoscopic resection of early gastric cancer: an open-label, randomised controlled trial. *Lancet* 2008;372:392-397.
166. Ikeda K, Marusawa H, Osaki Y, et al. Antibody to hepatitis B core antigen and risk for hepatitis C-related hepatocellular carcinoma: a prospective study. *Ann Int Med* 2007;146:649-656.
167. Sato Y, Takahashi S, Kinouchi Y, et al. IL-10 deficiency leads to somatic mutations in a model of IBD. *Carcinogenesis* 2006;27:1068-1073.
168. Ando T, Yoshida T, Enomoto S, et al. DNA methylation of microRNA genes in gastric mucosae of gastric cancer patients: its possible involvement in the formation of epigenetic field defect. *Int J Cancer* 2009;124:2367-2374.
169. Nanjo S, Asada K, Yamashita S, et al. Identification of gastric cancer risk markers that are informative in individuals with past *H. pylori* infection. *Gastric Cancer* 2012 (in press).

170. Toyota M, Ahuja N, Ohe-Toyota M, et al. CpG island methylator phenotype in colorectal cancer. *Proc Natl Acad Sci USA* 1999;96:8681-8686.

Table 1. Inflammation-associated cancers in digestive organs

Inflammation-associated cancer	Underling inflammation
Barrett's cancer	Reflux esophagitis
Gastric cancer	<i>H. pylori</i> -induced chronic gastritis
Colitic cancer	Inflammatory bowel disease
	Celiac disease
HCC	HCV and HBV chronic hepatitis
	Primary biliary cirrhosis
Cholangiocarcinoma	Primary sclerosing cholangitis
Pancreatic cancer	Chronic pancreatitis
	Hereditary pancreatitis
Lymphoma	<i>H. pylori</i> -induced MALT lymphoma
	HCV-associated lymphoma
	Celiac disease-associated lymphoma
HCC; hepatocellular carcinoma	

Figure Legends

Figure 1. Molecular link between inflammation, genetic and epigenetic alterations, and carcinogenesis. Inflammation contributes to ROS production and transcriptional upregulation of the DNA mutator enzyme, AID. These two factors were capable of inducing somatic mutations and chromosomal aberrations in tumor-related genes. On the other hand, inflammation results in mislocalization of DNMTs, inducing aberrant DNA methylation. The resulting genetic and epigenetic changes, including the activation of oncogenes, inactivation of tumor-suppressor genes, and dysregulation of DNA repair genes, could further enhance genetic instability, finally leading to carcinogenesis.

Figure 2. Mechanism of mutation induction by AID activity. AID deaminates cytosine (C), resulting in the generation of a uracil (U) and therefore can transform a DNA C:G pair into a U:G mismatch. The AID-generated U:G mismatch can be recognized by UNG or MSH2/MSH6 heterodimer and repaired correctly (a). If DNA replication starts before recognition by the repair system, a U:G mismatch gives rise to C/G to T/A transition (b). Alternatively, generation of an abasic site by UNG (c) or recognition of the U:G mismatch by the MSH2/MSH6 heterodimer (d) induces any mutations in the AID-generated U:G mismatch or at a nearby A:T site, respectively, in an error-prone manner (indicated as M).

Figure 3. AID exerts both favorable and unfavorable effects. AID is a molecule that is indispensable for the diversification of immunoglobulin genes by inducing both somatic hypermutation and class-switch recombination in activated B lymphocytes. The genotoxic activity of AID, however, can be aimed to trigger the genetic alterations at both the nucleotide and chromosomal levels in not only B lymphocytes but also epithelial cells in the inflammatory conditions.

Figure 4. The degree of epigenetic field defects can be assessed using methylation levels of appropriate marker CpG islands, mostly passengers. Receiver-operating characteristic (ROC) curves were drawn to distinguish gastric mucosae of gastric cancer patients and those of healthy individuals with past infections by *H. pylori*. The ROC curves of newly isolated methylation risk markers, CpG islands #3 and #7, had a much larger area under the curve (AUC) values than those of two previously isolated markers, filamin C γ (FLNc) and thrombomodulin (THBD), reaching 0.78–0.84. Modified from

Nanjo et al¹⁶⁹.

Figure 5. Current model of aberrant DNA methylation induction by chronic inflammation. Cytokines, such as IL-1 β and TNF- α from macrophages, and oxidative stress, such as NO, are likely to recruit EZH2 and DNMT1 to a promoter CpG island of a gene with a flag of H3K27 me3 and without protection by RNA polymerase II. Sparsely induced methylation leads to dense methylation with a low probability.

A

B

▲ H3K27me3

○ Unmethylated CpG site

● Methylated CpG site

● Nucleosome