TITLE:

Characterization and solvent engineering of wheat β -amylase for enhancing its activity and stability.

AUTHOR(S):

Daba, Tadessa; Kojima, Kenji; Inouye, Kuniyo

CITATION:

Daba, Tadessa ...[et al]. Characterization and solvent engineering of wheat β -amylase for enhancing its activity and stability.. *Enzyme and Microbial Technology* 2012, 51(5): 245-251

ISSUE DATE:

2012-10-10

URL:

<http://hdl.handle.net/2433/160221>

RIGHT:

© 2012 Elsevier Inc.; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。; This is not the published version. Please cite only the published version.

1 Enzyme and Microbial Technology

2

3 **Characterization and solvent engineering of wheat β -amylase for enhancing its**
4 **activity and stability**

5

6

7 Tadessa Daba, Kenji Kojima, and Kuniyo Inouye*

8

9

10 *Division of Food Science and Biotechnology, Graduate School of Agriculture, Kyoto*

11 *University, Sakyo-ku, Kyoto 606-8502, Japan*

12

13

14 *corresponding author. Tel.: +81-75-753-6266; fax: +81-75-753-6265.

15 E-mail address: inouye@kais.kyoto-u.ac.jp (K. Inouye)

16

17

18

ABSTRACT

The kinetic and thermodynamic parameters of wheat β -amylase (WBA) were characterized and various additives were evaluated for enhancing its activity and thermostability. WBA activity was examined by neocuproine method using soluble starch as substrate. The Michaelis constant (K_m) and molecular activity (k_{cat}) were determined to be 1.0 ± 0.1 % (w/v) and 94 ± 3 s⁻¹, respectively, at pH 5.4 and at 25°C. The optimum reaction temperature (T_{opt}) for WBA activity was 55°C and the temperature (T_{50}) at which it loses half of the activity after 30-min incubation was 50 ± 1 °C. Modifications of the solvent with 182 mM glycine and 0.18% (w/v) gelatin have increased the T_{50} by 5°C. Glycerol, ethylene glycol, dimethylformamide (DMF) and dimethyl sulfoxide have also slightly enhanced the thermostability plausibly through weakening the water structure and decreasing the water shell around the WBA protein. Ethanol and DMF activated WBA by up to 24% at 25°C probably by inducing favorable conformation for the active site or changing the substrate structure by weakening the hydrogen bonding. Its half-life in the inactivation at 55°C was improved from 23 to 48 min by 182 mM glycine. The thermodynamic parameters indicate that WBA is thermo-labile and sufficient stabilization was achieved through solvent modification

37 with additives and that the heat inactivation of WBA is entropic-driven. It is suggested
38 that WBA could be applied more widely in starch-saccharification industries with
39 employing suitable additives.

40

41 *Keywords:* activation; additives; β -amylase; thermostablility; solvent engineering, wheat

42

43

44

45

46

47

48

49

50

51

52

53

54

1. Introduction

β -Amylase hydrolyses the α -1, 4-glucan bonds in amylosaccharide chains from the non-reducing ends and generates maltose. It has considerable application in the production of high maltose syrups together with starch debranching enzymes and α -amylases. β -Amylase has been well characterized in higher plants [1-3] and micro-organisms [4-7]. However, the well-characterized β -amylases are neither active nor stable at higher temperatures $> 65^{\circ}\text{C}$ [8]. In many findings, α -amylase is more stable compared to β -amylase of the same origin [9-11]. Enzymes may easily be denatured by slight change of the environmental conditions such as temperature, pressure, pH, and ionic strength [9]. Nevertheless, stabilization of the enzymes could be achieved in several ways: screening for more stable ones (favorably from thermophiles and extremophiles), chemical modification, site-directed mutagenesis, immobilization and solvent engineering or modifying the enzyme reaction conditions with stabilizing additives [12-17]. In solvent engineering, selection of appropriate additives is dependent on the nature of the enzyme and there are no established rules to select effective additives for improving enzyme functions [18]. The thermostability of β -amylase has been substantially enhanced by modifying the solvent with additives [18, 19].

73 Different amylases give oligosaccharides with specific lengths of end products.
74 For this reason, amylases with unique properties need to be studied for various
75 applications in starch-saccharification for production of food and bio-ethanol [20].
76 Unlike soybean, barely and sweet potato, wheat was not the common source of
77 β -amylase for starch-saccharification so far. However, the sources of β -amylase supply
78 has drastically changed due to the escalating prices of the major sources like soybean.
79 Wheat β -amylase (WBA) is prepared from wheat bran, which is an industrial
80 by-product in the production of wheat starch and gluten. It is a cheaper alternative
81 source of β -amylase for industries. Nevertheless, it is lower in thermostability as
82 compared with β -amylases of other crops and microbes. For instance, the optimum
83 temperature (T_{opt}) of *Clostridium thermosuiphurogenes* β -amylase is 75°C [8]; the
84 temperature at which it loses half of its activity after 30-min incubation (T_{50}) of barely
85 β -amylase is 56.8°C and that of soybean β -amylase is 63°C [21] while the T_{opt} and T_{50}
86 after 30-min incubation of WBA are 55 and 50°C, respectively (Data obtained in this
87 study).

88 Therefore, enhancing the activity and thermostability of WBA has an excellent
89 prospect for starch-saccharification industries. In this study, we used a
90 commercially-available WBA preparation, Himaltosin, without further purification

because it is already purified from other protein contaminants and utilized industrially.

In this paper, we describe the kinetic and thermodynamic properties of WBA and

improvement in its activity and thermostability via solvent engineering using various

additives. This suggests that WBA would be likely applicable to a wide range of

starch-saccharification industries.

2. Materials and methods

2.1. Materials

Himaltosin GS (Lot 2S24A), a commercial preparation of WBA, was

purchased from HBI Enzymes (Osaka, Japan). This preparation was filtered with a

Millipore membrane filter (Type HA; pore size: 0.45 μ m) and used without further

purification. According to the manufacturer, the Himaltosin preparation contains 90%

starch as a stabilizer, and almost all of the protein is β -amylase and α -amylase was not

detected. Himaltosin was suspended to 20 mM sodium acetate buffer (pH 5.4) at 25°C

to be 0.3 mg/ml. In this paper, this buffer was hereinafter referred to as buffer A. The

WBA protein content was expected to be 0.03 mg/ml in the suspension and it was

109 followed by filtration with the Millipore membrane filter. However, the protein
110 concentration in the filtrate was less than 10% of the expected content, suggesting that >
111 90% of the WBA protein was remained with starch on the filter. The WBA
112 concentration was determined spectrophotometrically in buffer A using the absorptivity
113 value (*A*) of 1.40 ± 0.02 at 281 nm with a 1.0-cm light-path for the WBA solution at the
114 concentration of 1.0 mg/ml [22]. The molecular mass of 57.5 kDa for WBA was used to
115 determine the molar concentration of WBA (see sections 2.3 and 3.1). Under the
116 standard condition in this study, the concentration of WBA in the enzyme-reaction
117 solution was set to 15.0-30.0 nM. The starch concentration due to the stabilizer starch
118 (0.027%, w/v) was completely removed by filtration with the Millipore membrane filter.
119 Soluble starch (Lot M7H1482) as substrate and maltose (Lot M1F7568) as standard for
120 the activity assay were obtained from Nacalai Tesque (Kyoto, Japan). The substrate has
121 a weight-average molecular weight of 1.0×10^6 according to the manufacturer, and thus
122 the average degree of polymerization of the glucose unit is estimated to be 6,000.
123 Neocuproine-HCl (2, 9-dimethyl-1, 10-phenanthroline, Lot 032K2533) as coloring
124 reagent B in the neocuproine method was from Sigma (St. Louis, MO, USA). Coloring
125 reagent A (0.38 M Na_2CO_3 , 1.8 mM CuSO_4 , and 0.2 M glycine) in the neocuproine
126 method and all other chemicals were purchased from Nacalai Tesque. All enzyme

reactions were carried out in buffer A, pH 5.4.

2.2. Sodium dodecylsulfate-polyacrylamide gel electrophoresis (SDS-PAGE)

SDS-PAGE was performed in a 10% polyacrylamide gel under reducing conditions [23]. The Himaltosin preparation was suspended in buffer A to the concentrations of 1, 2, and 5% (w/v). The solution was filtered with a Millipore membrane filter (Type HA) and reduced by treatment with 10 mM dithiothreitol (DTT). The solution was applied to SDS-PAGE with a constant voltage of 150 V for 80 min. Proteins were stained with Coomassie Brilliant Blue R-250. The molecular-mass marker kit consisting of rabbit muscle phosphorylase *b* (97.2 kDa), bovine serum albumin (66.4 kDa), hen egg albumin (44.3 kDa), and hen egg white lysozyme (14.4 kDa) was a product of Takara Bio (Otsu, Japan).

2.3. Measurement of enzyme activity

Various initial concentrations of the soluble starch substrate [0.00, 0.09, 0.45, 0.90, 1.13, 1.35, 1.80, 2.02, 2.25, 2.70, 2.93, and 3.15% (w/v)] in the reaction solution

were prepared in buffer A at 25°C. The WBA solution in the same buffer was filtered with a Millipore membrane filter (Type HA) and kept in ice water for immediate use. The various concentrations of starch were hydrolyzed by WBA (30.0 nM) for 0, 2.5, 5.0, 7.5, and 10.0 min at 25°C. The reaction was stopped by adding 300 μ l of 0.1 M NaOH into 100 μ l of the enzyme-reaction solution. The amount of the reducing sugar in the enzyme-reaction solution was determined by the neocuploine method as follows [24]. Reagent A and reagent B, 250 μ l each, were mixed with 50 μ l of the enzyme-reaction solution, boiled for 8 min, and diluted with 550 μ l of water after cooling in ice water. The activity was measured at 450 nm using a Beckman-Coulter DU 800 spectrophotometer (Batavia, IL, USA) [25, 26]. The enzyme activity was determined by measuring the velocity (v) of reducing sugar production, and the reaction velocity was analyzed by the Michaelis-Menten kinetics. The maximum velocity ($V_{\max}$) and Michaelis constant (K_m) were obtained from the v vs. the substrate concentration ($[S]$) plots using KaleidaGraph 4.1 (Synergy Software, Reading, PA, USA). The molecular weight of WBA is 57,500 [27] and was used for the evaluation of the molecular activity (k_{cat}) (see section 3.1).

2.4. The optimum temperature

163

164 Starch solution (900 μ l) in buffer A was mixed with the WBA solution (0.150
165 μ M; 100 μ l) in the same buffer for 0, 2.5, 5.0, 7.5, and 10.0 min at 25°C after incubation
166 at 15, 25, 35, 45, 55, 65, and 75°C in a water bath for 10 min. The initial concentrations
167 of starch and WBA in the reaction solution were 1.80% and 15.0 nM, respectively. The
168 reaction was stopped by adding 300 μ l of 0.1 M NaOH into the enzyme-reaction
169 solution (100 μ l). Then the enzyme activity was determined by the neocuproine method
170 (see above). The optimal reaction temperature at which WBA exhibited the highest
171 activity was referred to as the optimal temperature (T_{opt}).

172

173 2.5. Thermal inactivation of WBA

174

175 The substrate and WBA solutions were prepared in buffer A. The enzyme
176 solution was incubated at 25, 35, 45, 55, and 65°C for 10, 20, and 30 min and cooled at
177 25°C for 3 min in a water bath. The substrate solution (2.00%, w/v; 900 μ l) was mixed
178 with the heat-treated WBA solution (0.300 μ M; 100 μ l) at 25°C and incubated for 0, 2.5,
179 5.0, 7.5, and 10.0 min. The initial concentrations of substrate and enzyme in the reaction
180 solution were 1.80% and 30.0 nM, respectively. The activity was assayed as

181 aforementioned by the neocuproine method and the first-order rate constant k of the
 182 thermal inactivation was determined assuming pseudo-first order kinetics by plotting $\ln$
 183 (v/v_o) against the heat-treatment time (t) (Eq. 1), where v is the initial reaction velocity
 184 of the enzyme with heat treatment at each incubation temperature and v_o is that obtained
 185 without heat treatment and at 25°C. The activation energy E_a of the thermal inactivation
 186 was obtained by the Arrhenius plot (Eq. 2), and the standard Gibbs energy difference of
 187 activation for thermal inactivation ($\Delta G^{o\ddagger}$), the standard enthalpy difference of activation
 188 ($\Delta H^{o\ddagger}$), and the standard entropy difference of activation ($\Delta S^{o\ddagger}$) were obtained from the
 189 Eyring plot according to Eqs. 3 and 4 [24, 28].

$$\ln (v/v_o) = k t \quad (1)$$

$$\ln k = - (E_a/R)(1/T) \quad (2)$$

$$\Delta G^{o\ddagger} = - RT[\ln hk/k_B T] \quad (3)$$

$$\ln (hk/k_B T) = (\Delta H^{o\ddagger}/RT) + (\Delta S^{o\ddagger}/R) \quad (4)$$

194 where k_B , h , and R are the Boltzmann, Plank, and gas constants, respectively. T is
 195 temperature in Kelvin.

2.6. Activation and thermostablization of WBA using additives

The WBA solution in buffer A was mixed and incubated with equal volume of various additives in the same buffer at 25, 45, 55, and 65°C in a water bath for 30 min before hydrolyzing soluble starch. The initial concentrations of WBA and substrate in the reaction solution were 30.0 nM and 1.80%, respectively. The additive concentrations in the reaction solution were: 45.5, 91, 182, and 364 mM glucose, NaCl, and glycine; 45.5 and 91 mM L-arginine; 0.9, 1.8, 3.6, and 45.5 mM of L-aspartate; 45.5 and 91 mM of L-cysteine and glutathione (GSH); 0.18 and 0.45% (w/w) gelatin; 0.91, 1.82, and 5.5% (w/w) ethanol and 2-methyl-2-butanol (2M2B); 0.45 and 2.7% (w/w) dimethyl sulfoxide (DMSO) and dimethylformamide (DMF); and 0.91 and 5.5% glycerol, ethylene glycol (EG), and β -mercaptoethanol (β ME). Their effects on activation and thermal stabilization of WBA were examined. The enzyme-additive mixture solution (0.20 ml) was diluted with water (0.55 ml), and the enzyme activity was measured by the neocuproine method.

3. Results

3.1. Kinetic parameters of WBA

The WBA preparation, Himaltosin, shows a single band in SDS-PAGE with molecular mass of 57.5 kDa (Fig. 1). This value is in good agreement with those (54.0-64.2 kDa) so far reported [22, 27]. In this paper, we used 57.5 kDa for the molecular mass of WBA. This indicates that the Himaltosin preparation is composed of solely β -amylase as a protein component, although it contains 90% (w/w) starch as a stabilizer. Substantially, there was no starch carried over from the stabilizer starch into the reaction solution as examined by the starch-iodine reaction. Therefore, the enzyme preparation was treated as WBA without further purification.

(Fig. 1)

The initial velocity (v) of the WBA-catalyzed hydrolysis of different concentrations of soluble starch was examined in buffer A at 25°C at the enzyme concentration of 30.0 nM. The dependence of v on substrate concentration exhibited the Michaelis-Menten profile (Fig. 2). The K_m , V_{max} , and k_{cat} were determined to be 1.0 ± 0.1 % (w/v); $2.8 \pm 0.1 \mu\text{M s}^{-1}$, and $94 \pm 3 \text{ s}^{-1}$, respectively, by fitting the experimental data to the Michaelis-Menten equation.

(Fig. 2)

3.2. Thermal inactivation of WBA

WBA at the concentration of 0.300 μM was treated thermally as described in the Material and methods section, and the WBA concentration in the enzyme-reaction solution was fixed to 30.0 nM. The enzyme activity was decreased with the progress of the heat treatment at every temperature examined (Fig. 3). The enzyme activity in the hydrolysis of soluble starch was evaluated by measuring the initial velocity (v) in the same buffer at 25°C. The v value measured in the 0-min incubation at each temperature was designated as v_0 . The relative activity (v/v_0) observed after incubation at various temperature decreased progressively with increasing the incubation time. The semi-log plots of v/v_0 against incubation time showed linear relationship at the respective incubation temperatures (Fig. 3), indicating that the thermal inactivation process of WBA follows the first-order kinetics. The first-order rate constant (k) at the indicated incubation temperature was evaluated from the slope of the plot.

(Fig. 3)

253

254 The activation energy E_a value of the thermal inactivation of WBA in buffer A was $36 \pm$
255 1 kJ mol^{-1} from the slope of the Arrhenius plot (Fig. 4). The $\Delta G^{0\ddagger}$, $\Delta H^{0\ddagger}$, and $T\Delta S^{0\ddagger}$
256 values for the thermal inactivation were found to be 90 ± 1 , 33 ± 1 , and $-59 \pm 1 \text{ kJ mol}^{-1}$
257 respectively, at 25°C from the slope of the Eyring plot (Fig. 5). As E_a is defined
258 theoretically as $\Delta H^{0\ddagger} + RT$, the $\Delta H^{0\ddagger}$ value is calculated to be $34 \pm 1 \text{ kJ mol}^{-1}$, which is
259 in good agreement with the value obtained from the Eyring plot.

260

261 (Fig. 4)

262 (Fig. 5)

263

264 3.3. The optimum temperature of WBA

265

266 WBA activity was measured at various reaction temperatures (Fig. 6). The
267 maximal activity in starch hydrolysis was obtained at 55°C , being the optimal
268 temperature (T_{opt}) of this enzyme in buffer A, pH 5.4.

269

270 (Fig. 6)

271

272 3.4. *The T_{50} of WBA*

273

274 The heat inactivation of WBA was studied by incubating the enzyme at various
275 temperatures for 30 min before hydrolyzing the substrate. The enzyme activity was
276 observed to decline with heat treatment even at moderate temperatures (45 and 55°C).
277 The T_{50} of WBA, which is the temperature at which the enzyme loses half of its activity
278 with thermal treatment for 30 min was determined to be $50 \pm 1^\circ\text{C}$ (Fig. 7) by plotting
279 the residual activities (%) against temperature of incubation. WBA has lost 30% of its
280 activity at 45°C, 76% at 55°C, and 95% at 65°C after 30 min of incubation.

281

282 (Fig. 7)

283

284 3.5. *Effect of additives on WBA activity*

285

286 The catalytic activity of WBA in starch hydrolysis was examined in the
287 presence of various additives at 25°C without heat treatment. Majority of the evaluated
288 additives did not enhance the enzyme activity (Fig. 8). However, it is noted that ethanol

at the concentration of 0.91-5.5%, w/w (or 0.20-1.20 M) enhanced the activity by 24% and lower concentration (1.8%) of DMF by 11%. On the other hand, the activity was not much affected with the addition of NaCl and glycine up to 364 mM; and DMF, DMSO, and EG up to 5.5%. The addition of cysteine up to 91 mM and aspartate up to 45.5 mM showed no substantial effect on the activity, and gelatin (0.18-0.45%, w/w) had no effect either. The other additives inhibited the activity to varying degrees. Especially, the activity was reduced to almost 10% with the addition of 0.91-5.5%, w/w (or 0.11-0.71 M) β ME; and to zero with 45.5-91 mM arginine. The activity decreased with increasing the glucose and GSH concentrations, and 50% of the activity was lost with 182 mM glucose; and 75% was with 91 mM GSH.

(Fig. 8)

3.6. Effect of additives on the thermostability of WBA

The rate of starch hydrolysis by heat treated WBA was examined in modified solvent by various additives. Glycine was found to be the best in improving the thermal stability of WBA followed by gelatin. The T_{50} of WBA was improved by 6°C with 182

mM glycine and by 4°C with 0.18% (w/w) gelatin (Fig. 9). The half-life times of the enzyme were enhanced by 25 min and 17 min with glycine and gelatin, respectively, at 55°C (Table 1). It should be noted that the enzyme activities remained after the thermal treatment at 45°C was 97% and 92% in the presence of 0.18% (w/w) gelatin and 182 mM glycine although it was only 75% in the absence of additives. Aspartate (45.5 mM) and DMSO (5.5%, w/w) had no effect on the stability of WBA. β ME (0.91%); ethanol (0.91%); NaCl (182 mM); and arginine (45.5 mM) slightly decreased the stability of WBA, and the T_{50} values were 46-47°C. Glucose (182 mM), cysteine (46 mM), and GSH (91 mM) considerably decreased the stability with the T_{50} values of 35-45°C.

(Fig. 9)

(Table 1)

4. Discussion

In general, depending on the nature and concentration of the additives, they affect the protein conformation in: (a) screening effect, where the electrostatic repulsion between similarly charged groups of proteins is reduced by cosolvent ions; (b)

325 solvophobic effect, where ion pair formation occurs, favoring protein folding; and (c)
326 modification of water structure leading to hydrophobic interactions in proteins [10, 29].
327 The catalytic activity and thermostability of various enzymes were considerably
328 enhanced through solvent engineering [17-19, 30, 31].

329 The Michaelis constant ($K_m = 1.0 \pm 0.1$ %, w/v) and molecular activity ($k_{cat} =$
330 94 ± 3 s⁻¹) of WBA was evaluated at pH 5.4 and at 25°C using soluble starch as
331 substrate (Fig. 2). Similarly, the K_m of glutenin-adsorbed WBA was reported to be
332 0.15% (w/v) using soluble starch substrate [32], indicating that WBA (which is free in
333 the reaction solution) used in our study has lower affinity to soluble starch as compared
334 with WBA adsorbed on glutenin. The other available reports on the kinetic parameters
335 of WBA were using maltotriose, maltoheptaose, amyloextrin, and amylopectin
336 substrates under various reaction conditions [27, 33]. The K_m values of plant β -amylases
337 have been reported to be in the range of 0.2-0.7 % (w/v) for soluble starch. The value
338 obtained in this study is apparently higher than those. However, the kinetic parameters
339 for WBA and other plant β -amylases have not yet examined under the same reaction
340 conditions using the same soluble starch preparation. Therefore, it is not suitable to
341 discuss the enzyme functions by comparing the kinetic parameters so far reported. In
342 this paper, we have reported the kinetic parameters (K_m and k_{cat}) of WBA obtained under

the fixed conditions with the WBA and soluble starch preparations in the same lot. It should be noted that the activity and stability of WBA are affected easily by additives added in the reaction solution. This means that the kinetic parameters of WBA must be evaluated with the data collected from the experiments conducted carefully under the same conditions.

Thermodynamic parameters for the heat inactivation of WBA showed that the $T\Delta S^{\circ\ddagger}$ value ($59 \pm 1 \text{ kJ mol}^{-1}$) was greater than the $\Delta H^{\circ\ddagger}$ value ($33 \pm 1 \text{ kJ mol}^{-1}$), and thus it can be inferred that the heat inactivation of the enzyme is entropic-driven. The optimum reaction temperature (T_{opt}) of WBA is 55°C (Fig. 6). It has been reported that the T_{opt} of *Sorghum bicolor* cv β -amylase is 50°C [11, 20, 34], while α -amylase from the same cereal crop has T_{opt} of 70°C corresponding with the fact that β -amylases in most of the cases are lower in thermostability than their respective α -amylases of the same origin [10, 11]. The T_{50} of WBA was determined to be $50 \pm 1^{\circ}\text{C}$ (Fig. 6), which is lower than those of soybean β -amylase, 63.2°C , and of barely β -amylase, 56.8°C [21].

In the present study, we have examined the effects of additives on the kinetic parameters and thermostability of WBA. The additives are supposed to have effects on the structures of WBA, soluble starch substrate, and bulk water. For example, when we observed decrease in activity by the addition of an additive, there might be some

reasons considered such as inhibition of the enzyme by the additive, conformational changes of the enzyme and/or starch substrate by the additive, etc. Therefore, strict interpretation of the molecular effects of the additives on the activity and stability of WBA seems to be difficult, and further studies must be needed. Thus in this paper we tried to present the effects of the additives without describing the molecular-mechanistic insights of the cause for the effects, although some possible comments have been made with references to other enzymes. The molecular-mechanistic study for the effects of the representative additives is underway.

Ethanol and low concentration of DMF have enhanced the activity of WBA by up to 24% (Fig. 8). The activation of acetylcholinesterase (AChE) by ethanol was reported and explained as that it non-competitively alters the hydrophobic-interaction site and subsequently induces favorable conformation to the active center [35]. However, higher concentration of ethanol had a destabilization effect on WBA (Fig. 9B). This agrees with a finding that a high concentration of ethanol (> 800 mM) can perturb the structure of water around hydrophilic area of AChE causing instability to the conformation of the enzyme [35]. This might be the case for the effects of ethanol on WBA. These effects should be considered also from the viewpoint of solvent polarity (see below). The effects of various alcohols on the enzyme structure and activity have

been extensively studied with thermolysin, a thermophilic and halophilic metalloproteinase produced by *Bacillus thermoproteolyticus*. Thermolysin is remarkably activated and stabilized by neutral salts such as NaCl, and is inhibited by increasing concentration of alcohols and the degree of inhibition is dependent on the size of alcohols [36]. It is interesting to note that alcohol such as 2-methyl-1-propanol (2MP) which binds tightly to the active site inhibits thermolysin strongly, whereas the alcohols with the sizes larger or smaller than 2MP bind weakly to the active site and inhibit thermolysin weakly [37]. This suggests that the enzyme activity could be controlled intentionally using additives with suitable sizes. These lines of evidence have provided information for the optimal conditions for thermolysin-catalyzed synthesis of a precursor of the sweetener, aspartame [14]. As shown with thermolysin to find the optimal conditions for its inhibition by alcohols, it should be possible to find the optimal conditions for industrial application of WBA by examining the effects of alcohols on the activity and stability of WBA by changing systematically the size of alcohols.

Contrary to the effect of ethanol, the T_{50} of WBA is slightly improved by polyols, indicating that polyols are WBA stabilizer. Our finding agrees with reports that solvent modification by sugars and polyols had marked thermal stabilization in various enzymes [9, 30]. Polyols were explained to preferentially be excluded from the surface

layer of the protein and form a water shell around the protein, so that the protein is preserved and the conformation becomes more rigid or stable [9, 30]. Similar mechanism of stabilization by polyols was also reported for organic polar solvents (DMSO and DMF) [31], and actually polar organic solvents have conferred a slight thermostabilization to WBA. However, sugars, which are classified also as polyols, interestingly had no stabilizing effect on WBA in our study.

The activation and thermostabilization of WBA were tested using different additives with various values of empirical parameters of solvent polarity, $E_T(30)$ [38]. In particular, ethanol ($E_T(30) = 217 \text{ kJ mol}^{-1}$), DMSO (189 kJ mol^{-1}), DMF (183 kJ mol^{-1}), glycerol (238 kJ mol^{-1}), EG (235 kJ mol^{-1}), β ME (224 kJ mol^{-1}), and 2M2B (172 kJ mol^{-1}) were examined at different temperatures. According to the results, the half-life time of WBA was enhanced at 55°C with ethanol. It is suggested that thermostabilization by additives could be induced by reducing the degree of water-solvation or deformation of a water shell around the protein. However, a solvent 2M2B with a low $E_T(30)$ value decreased the activity and stability of WBA (Figs. 8B and 9B), suggesting that the effects of additives on the enzyme activity and stability are complicated and that the effect on the water-solvation might not be the main one [39].

Glycine was the most favorable among the evaluated additives in stabilizing

WBA followed by gelatin (Table 1) and the stabilization effects of various additive concentrations were presented in Fig. 9. It should be noted that the thermostability of WBA in the presence of favorable additives such as glycine and gelatin is almost comparable with that of soybean β -amylase widely used in starch-saccharification industry. This suggests that WBA could be applied more widely in food and bio-ethanol industries with employing suitable additives. A good empirical correlation between the relative stabilizing effects of glycine with the change in solvent-accessible hydrophobic surface area of the folded protein was reported [40]. The possible protective effect of gelatin on penicillinase was stated that it combines with enzyme and form a thermostable complex [41]. The molecular mechanism of glycine and gelatin on stabilizing WBA should be examined in the next step from these lines.

Arginine was found to destabilize WBA in our study. Similarly, chloroperoxidase was confirmed to be inactivated by arginine mainly by the binding of a guanidinium group with the catalytic site [42]. This inactivation effect might be the same as that given by the denaturant, guanidine hydrochloride; namely, arginine might cleave the hydrogen bonds in the protein structure and increase the solubility of hydrophobic residues of the protein. On the other hand, arginine is known to work effectively in recovering human matrix metalloproteinase 7 from inclusion bodies [43].

433 This suggests that arginine promotes unfolding the misfolded protein structure to lead it
434 into the correctly folded form. While arginine seems to show multiple effects on protein
435 stability depending on target proteins, it could be a good tool to increase the stability of
436 WBA by selecting suitable conditions.

437 We tried to modify thiol groups of WBA using reducing agents like β ME and
438 GSH, whereas they destabilized WBA. This is because they attack disulfide bonds and
439 expose proteins to heat denaturation and, hence reduces their thermostability [44].
440 Plausible involvement of thiol groups in the catalytic activity of amylases has been
441 suggested, although it is not known currently with WBA. The effects of β ME and GSH
442 should be considered from this point in the next step.

443 Various concentrations of NaCl exhibited destabilizing effect on WBA in our
444 study. Three possible reasons were suggested for salt-induced inactivation or
445 destabilization effect on enzymes: (a) break weak hydrogen-bonds and disrupt the
446 protein conformation; (b) attract water molecules and the enzyme coagulates by
447 protein-protein hydrophobic interaction; and (c) high concentration of salt makes the
448 enzyme more likely to bind with the salt ions instead of the substrate by electrostatic
449 interaction [45]. However, high concentration of neutral salts remarkably improves
450 thermolysin activity [14, 15]. The catalytic activity of thermolysin is enhanced by 4

451 times with the substitution of the active-site zinc with cobalt and further exponentially
452 by NaCl up to 13-15 times [29].

453 It is well known that enzyme activity is controlled by the factors of enzyme
454 structure and reaction environment. The structural factors and environmental factors are
455 sometimes independent and sometimes closely connected. With their optimal
456 combination, the optimal catalytic activity should be realized. Protein engineering
457 (namely, site-directed mutagenesis and chemical modification) is a tool for changing the
458 enzyme structure in a predictable and precise manner to effect a change on the catalytic
459 process. Since the enzyme is even improved in only one side of a reaction, any changes
460 in the rest of the reaction may also alter the catalytic process [46]. Solvent engineering
461 is a powerful tool in rational control of enzyme activity. In reality, both approaches are
462 still somewhat difficult to confirm their effects, whereas they have been used
463 successfully to alter the protein properties [47]. At last, it should be reminded that the
464 protein-engineered enzymes are strictly limited to use for food processing in many
465 countries in order to avoid unpredictable harms. Thus solvent engineering must be an
466 inevitable alternative to find the optimal conditions from the viewpoints of enhancing
467 enzyme catalytic efficiency, guaranteeing safeness of the products, reducing the costs
468 for production, etc. When the enzyme activity in its industrial application is increased

by solvent engineering, the cost of the additive should be considered in the total cost, although it is generally much cheaper than that of the enzyme, and the improved enzyme activity and stability by the additive should decrease the enzyme amount needed and thus shortened the reaction time, which results in decreasing the running costs, utility, labor, etc.

5. Conclusion

From the thermodynamic parameters evaluated in this study, it is eminent that WBA is not stable at temperatures higher than 55°C. The thermal stability of WBA was improved by the addition of glycine and gelatin explicably through convening conformation of the enzyme and reducing the interaction of the protein with the solvent. Polyols and organic polar solvents (DMSO and DMF) also conferred slight stability to the enzyme while some evaluated additives have exhibited destabilizing effect. The thermodynamic parameters indicate that WBA is thermo-labile and sufficient stabilization was achieved by solvent engineering with additives and that the heat inactivation of WBA is entropic-driven. On the other hand, it was shown that WBA activity was enhanced by the addition of ethanol and DMF persuasively by altering the

hydrophobic interaction and inducing favorable conformation to its active center. It is suggested that WBA would be applicable to a wide range of saccharification industries such as food and bio-ethanol production with employing suitable additives.

References

- [1] Higashihara M, Okada S. Studies on β -amylase of *Bacillus megaterium* Strain No. 32. *Agric Biol Chem* 1974;38:1023-9.
- [2] Murao S, Ohyama K, Arai M. β -Amylase from *Bacillus polymyxa* No. 72. *Agric Biol Chem* 1979;43:719-26.
- [3] Robyt J, French D. Purification and action pattern of an amylase from *Bacillus polymyxa*. *Arch Biochem Biophys* 1964;104:338-42.
- [4] Shinke R, Kunimi Y, Nishira H. Production and some properties of β -amylases of *Bacillus* sp. BQ 10. *J Ferment Technol* 1975;53:693-7.
- [5] Takasaki Y. Purification and enzymatic properties of β -amylase pullulanase from *Bacillus cereus* var. *mycoides*. *Agric Biol Chem* 1976;40:1523-30.
- [6] Thomas M, Priest G, Stark JR. Characterization of an extracellular β -amylase from

- 505 Bacillus megaterium sensu stricto. J Gen Microbiol 1980;118:67-72.
- 506 [7] Yamashiro K, Yokobori S-I, Koikeda S, Yamagishi A. Improvement of Bacillus
- 507 circulans β -amylase activity attained using the ancestral mutation method. Protein
- 508 Eng Des Sel 2010;23:519-28.
- 509 [8] Shen G-J, Saha BC, Bhatnagar Y-E, Zeikus J. Purification and characterization of a
- 510 novel thermostable β -amylase from Clostridium thermosulphurogenes. Biochem J
- 511 1988;254:835-40.
- 512 [9] Matsumoto M, Kida K, Kondo K. Effects of polyols and organic solvents on
- 513 thermostability of lipase. J Chem Technol Biotechnol 1997;70:188-92.
- 514 [10] Muralikrishna G, Nirmala M. Cereal α -amylases: an overview. Carbohydr Polym
- 515 2005;60:163-73.
- 516 [11] El Nour MEM, Yagoub S. Partial purification and characterization of α and
- 517 β -amylases isolated from Sorghum bicolor cv. (Feterita) malt. J Appl Sci
- 518 2010;10:1314-9.
- 519 [12] Costa SA, Tzanov T, Paar A, Gudelj M, Gübitz GM, Cavaco-Paulo A.
- 520 Immobilization of catalases from Bacillus SF on alumina for the treatment of
- 521 textile bleaching effluents. Enzyme Microb Technol 2001;28:815-9.
- 522 [13] Rocha JMS, Gil MH, Garcia FAP. Effects of additives on the activity of a

- 523 covalently immobilized lipase in organic media. J Biotechnol 1998;66:61-7.
- 524 [14] Inouye K. Effects of salts on thermolysin: activation of hydrolysis and synthesis of
- 525 N-carbobenzoxy-L-aspartyl-L-phenylalanine methyl ester and a unique change in
- 526 the absorption spectrum of thermolysin. J Biochem 1992;112:335-40.
- 527 [15] Inouye K, Kuzuya K, Tonomura B. Sodium chloride enhances markedly the
- 528 thermal stability of thermolysin as well as its catalytic activity. Biochim Biophys
- 529 Acta 1998;1388:209-14.
- 530 [16] Hashida Y, Inouye K. Molecular mechanism of the inhibitory effect of cobalt ion
- 531 on thermolysin activity and the suppressive effect of calcium ion on the cobalt
- 532 ion-dependent inactivation of thermolysin. J Biochem 2007;14:879-88.
- 533 [17] Sekiguchi S, Hashida Y, Yasukawa K, Inouye K. Effects of amines and
- 534 aminoalcohols on bovine intestine alkaline phosphatase activity. Enzyme Microb
- 535 Technol 2011;49:171-6.
- 536 [18] Costa SA, Tzanov T, Carneiroa AF, Paar A, Gübitz GM, Cavaco-Paulo A. Studies
- 537 of stabilization of native catalase using additives. Enzyme Microb Technol
- 538 2002;30:387-91.
- 539 [19] Devi NA, Rao AGA. Effect of additives on kinetic thermal stability of
- 540 polygalacturonase II from *Aspergillus carbonarius*: mechanism of stabilization by

- 541 sucrose. J Agric Food Chem 1998;46:3540-5.
- 542 [20] Wong DWS. Recent advances in enzyme development. In: Whitaker JR, Voragen
543 AGJ, Wong DWS, editors. Handbook of food enzymology. Marcel Dekker, New
544 York; 2003. p. 379-88.
- 545 [21] Yoshigi N, Okada Y, Maeba H, Sahara H, Tamaki T. Construction of a plasmid used
546 for the expression of a sevenfold-mutant barley β -amylase with increased
547 thermostability in Escherichia coli and properties of the sevenfold-mutant β -
548 amylase. J Biochem 1995;118:562-7.
- 549 [22] Tkachuk R, Tipples KH. Wheat beta-amylases. II. Characterization. Cereal Chem.
550 1966;43:62-79.
- 551 [23] Laemmli UK. Cleavage of structural proteins during the assembly of the head of
552 bacteriophage T4. Nature 1970;227:680-5.
- 553 [24] Oneda H, Lee S, Inouye K. Inhibitory effect of 0.19 α -amylase inhibitor from
554 wheat kernel on the activity of porcine pancreas α -amylase and its thermal stability.
555 J Biochem 2004; 135:421-7.
- 556 [25] Brown ME. Ultra-micro sugar determinations using 2, 9-dimethyl-1,10-
557 phenanthroline hydrochloride (neocuproine). Diabetes 1961;10:60-3.
- 558 [26] Baker WL, Panow A. Estimation of cellulose activity using a glucose-oxidase-Cu

- 559 (II) reducing assay for glucose. J Biochem Biophys Methods 1991;23:265-73.
- 560 [27] Kato M, Hiromi K, Morita Y. Purification and kinetic studies of wheat bran
- 561 β -amylase. Evaluation of subsite affinities. J Biochem 1974;75:563-76.
- 562 [28] Segel IH. Enzyme Kinetics: behavior and analysis of rapid equilibrium and steady
- 563 state enzyme systems. John Wiley and Sons, New York; 1975. p. 926-42.
- 564 [29] Kuzuya K, Inouye K. Effects of cobalt-substitution of the active zinc ion in
- 565 thermolysin on its activity and active-site microenvironment. J Biochem
- 566 2001;130:783-8.
- 567 [30] De Cordt S, Hendrickx M, Maesmans G, Tobback P. The influence of polyalcohols
- 568 and carbohydrates on the thermostability of *Bacillus licheniformis* α -Amylase.
- 569 Biotechnol Bioeng 1994;43:107-14.
- 570 [31] Rajendran S, Radha C, Prakash V. Mechanism of solvent-induced thermal
- 571 stabilization of α -amylase from *Bacillus amyloliquefaciens*. Int J Pept Protein Res
- 572 1995;45:122-8.
- 573 [32] Rothfus JA, Kennel SJ. Properties of wheat beta-amylase adsorbed on glutenin.
- 574 Cereal Chem 1970;47:140-6.
- 575 [33] Wong DWS, Robertson GH: β -Amylases. In: Whitaker JR, Voragen AGJ, Wong
- 576 DWS, editors. Handbook of food enzymology. Marcel Dekker; 2003. p. 719-26.

- 577 [34] Kumar RSS, Singh SA, Rao AGA. Conformational stability of α -amylase from
578 malted sorghum (*Sorghum bicolor*): reversible unfolding by denaturants.
579 *Biochimie* 2009;91:548-57.
- 580 [35] Shin S, Wu P, Chen CH. Biochemical studies of the actions of ethanol on
581 acetylcholinesterase activity: Ethanol-enzyme-solvent interaction. *Int J Biochem*
582 1991;23:169-74.
- 583 [36] Inouye K, Lee S-B, Nambu K, Tonomura B. Effects of pH, temperature, and
584 alcohols on the remarkable activation of thermolysin by salts. *J Biochem*
585 1997;122:358-64.
- 586 [37] Muta Y, Inouye K. Inhibitory effects of alcohols on thermolysin activity as
587 examined using a fluorescent substrate. *J Biochem* 2002;132:945-51.
- 588 [38] Reichardt C. Solvatochromic dyes as solvent polarity indicators. *Chem Rev* 1994;
589 94: 2319-58.
- 590 [39] Oneda H, Inouye K. Effects of dimethyl sulfoxide, temperature, and sodium
591 chloride on the activity of human matrix metalloproteinase 7 (matrilysin). *J*
592 *Biochem* 2000;128:785-91.
- 593 [40] Serrano L, Neira JL, Sancho J, Fersht AR. Effect of alanine versus glycine in
594 α -helices on protein stability. *Nature* 1992;356:453-5.

- 595 [41] Manson EED, Pollock MR. The thermostability of penicillinase. J Gen Microbiol
596 1953;8:163-7.
- 597 [42] Bai C, Bo H, Jiang Y, Hu M, Li S, Zhai Q. Inactivation of chloroperoxidase by
598 arginine. Process Biochem 2010;45:312-6.
- 599 [43] Oneda H, Inouye K. Refolding and recovery of recombinant human matrix
600 metalloproteinase 7 (matrilysin) from the inclusion bodies expressed by
601 Escherichia coli. J Biochem 1999;126:905-11.
- 602 [44] Kim S, Paik WK. Effect of glutathione on ribonuclease. Biochem J
603 1968;106:707-10.
- 604 [45] Caussette M, Planche H, Delepine S, Monsan P, Gaunand A, Lindet B. The self
605 catalytic enzyme inactivation induced by solvent stirring: a new example of
606 protein conformational change induction. Protein Eng 1997;10:1235-40.
- 607 [46] Michael MM, Lanyi JK. Threonine deaminase from extremely halophilic bacteria.
608 Cooperative substrate kinetics and salt dependence. Biochemistry 1972;10:211-6.
- 609 [47] Russell AJ, Vierheller C. Protein engineering. In: Bronzino JD, editor. The
610 biomedical engineering handbook. 2nd ed. CRC Press; 2000. p. 101-4.
- 611

Figure legends

Fig. 1. SDS-PAGE of the Himaltosin preparation. Molecular marker proteins (lane 1); Himaltosin preparation: 1% (w/v) (lane 2), 2% (w/v) (lane 3), and 5% (w/v) (lane 4). The experimental conditions were given in the text (Materials and methods).

Fig. 2. Dependence of the initial reaction velocity (v) in the hydrolysis of soluble starch catalyzed by wheat β -amylase (WBA) on substrate concentration. The hydrolysis was carried out in 20 mM sodium acetate buffer (buffer A), pH 5.4, at 25°C. The WBA concentration in the reaction solution was 30.0 nM.

Fig. 3. Progress of the decrease in WBA activity with the time of thermal treatment. WBA (0.300 μ M) was treated at 25, 35, 45, 55, and 65°C for 0-30 min in buffer A, pH 5.4. WBA activity in the hydrolysis of soluble starch was evaluated by measuring the initial velocity (v) in the same buffer at 25°C (see Materials and methods). The WBA concentration in the enzyme-reaction solution was 30.0 nM. The v value measured in the 0-min incubation at each temperature was designated as v_o . The logarithm of the relative activity [$\log (v/v_o)$] was plotted against the thermal-treatment time. The markers

are (temperature in °C): 25, ○; 35, ◇; 45, △; 55, □; and 65, ●. From these semi-log plots, the first-order rate constant for the thermal inactivation of WBA was evaluated at the specified temperature of the thermal treatment.

Fig. 4. Arrhenius plot of WBA for the first-order rate constant (k) of thermal inactivation. The rate constants were evaluated from the semi-log plots as shown in Fig. 3 obtained by thermal treatment of the enzyme at various temperatures. The reaction conditions were given in the text (Materials and methods).

Fig. 5. Eyring plot of WBA for the first-order rate constant (k) of thermal inactivation. The rate constants were evaluated as described in the legend of Fig. 4.

Fig. 6. Effect of temperature on the initial velocity (v) of WBA in hydrolyzing soluble starch. The enzyme reaction was carried out at the temperature indicated. The initial concentrations of starch and WBA in the reaction solution were 1.80% and 15.0 nM, respectively. The optimal temperature T_{opt} was determined to be 55°C.

Fig. 7. Effect of thermal inactivation of WBA on the enzyme activity. The activity was

assayed in buffer A (pH 5.4) in hydrolyzing starch after incubation at various temperatures indicated for 30 min. The reaction conditions are given in the text (Materials and methods). The T_{50} was determined to be $50 \pm 1^\circ\text{C}$.

Fig. 8. The effects of various concentrations of additives on the activity of WBA at 25°C . Symbols for additives (mM): (A) Gly, ○; NaCl, ◇; Asp, □; Cys, △; glucose, ●; GSH, ◆; and Arg, ▲. Symbols for additives (% w/w): (B) ethanol, ○; DMF, ◇; DMSO, △; EG, □; gelatin, ●; glycerol, ◆; BME, ▲; and 2M2B, ▽. The relative activity (%) of the enzyme obtained without additive was set as 100%.

Fig. 9. Effects various additives on the thermostabilization of WBA after incubation with the additives at various temperatures for 30 min. Symbols for concentrations (mM) of additives: (A) buffer, ○; 182 glycine, ◇; 45.5 Arg, □; 45.5 Asp, △; 182 NaCl, ●; 182 glucose, ◆; 45.5 Cys, ▲; and 91 GSH, ■. Symbols for concentrations (% w/w) of additives: (B) buffer, ○; 5.5 EG, ◇; 0.18 gelatin, △; 5.5 glycerol, □; 0.91 βME, ●; 2.7 DMF, ◆; 0.91 ethanol, ▲; 5.5 DMSO, ■; and 5.5 2M2B, ▽. Low concentrations of some additives were evaluated because of their solubility limit in buffer A (pH 5.4) at 25°C . The residual activity of WBA without additive was depicted in broken lines.

666

667 Table 1. Effects of selected additive concentrations on thermostablization of wheat

668 β -amylase (WBA) after incubation at 55°C with the additives for 30 min prior

669 to starch hydrolysis

Additives	T_{50} (°C)	k (min ⁻¹)	$t_{1/2}$ at 55°C (min)	$\Delta G^{o\ddagger}$ (kJ mol ⁻¹)
none	50 ± 1	0.030 ± 0.004	23 ± 2	90 ± 1
0.18% gelatin	54 ± 1	0.017 ± 0.003	40 ± 1	92 ± 1
0.45% gelatin	54 ± 1	0.018 ± 0.001	39 ± 2	92 ± 1
91 mM glycine	54 ± 1	0.017 ± 0.003	41 ± 1	92 ± 1
182 mM glycine	56 ± 2	0.014 ± 0.006	48 ± 3	92 ± 1

670 T_{50} is the temperature at which WBA loses half of it activity after 30 min of incubation;

671 k is the first-order rate constant of the thermal inactivation; $t_{1/2}$ (at 55°C) is the time at

672 which WBA loses half of its activity at 55°C; and $\Delta G^{o\ddagger}$ is the standard Gibbs energy

673 difference of activation. Each value is a mean of triplicate analysis ± standard deviation.

674 The initial concentrations of starch and WBA in the reaction solution were 1.80% and

675 30.0 nM, respectively.

kDa

97.2

66.4

44.3

14.3

1

2

3

4

← WBA

Top

Fig. 1

Top

Fig. 2

Top
↑

Fig. 3

Top
↑

Fig. 4

Top
↑

Fig. 5

Top

Fig. 6

Top

Fig. 7

Top

Fig. 8A

Top

Fig. 8B

Top
↑
Fig. 9A