

**Replication mechanisms of plant RNA viruses**

**-Current understanding and perspective-**

**Tetsuro Okuno**

T. Okuno

Graduate School of Agriculture, Kyoto University, Kyoto 606-8502, Japan

okuno@kais.kyoto-u.ac.jp

This article is an abstract of the Presidential Address presented at the 2012 Annual Meeting of the Phytopathological Society of Japan in Fukuoka.

20    **Introduction**

21

22    Viruses cause numerous diseases in economically important plants and animals,  
23    including humans. However, there are no efficient measures to counter viral infection,  
24    with very few exceptions for animal viral diseases. Although virus resistance genes  
25    have been used to minimize the loss of crop production in several important plant  
26    species, the number of genes that have been used to confer virus resistance is very  
27    limited. The delay in the development of efficient measures for the control of viral  
28    diseases might be caused in part by the lack of critical information on the mechanisms  
29    underlying viral replication, especially regarding the host proteins used by viruses.

30       Positive-strand RNA viruses with a genome composed of messenger-sense  
31    single-stranded RNAs, represent the largest group among the seven genetic classes of  
32    all viruses. Upon entry into host cells, the genomic RNAs of these viruses serve as  
33    mRNAs, and viral replication proteins are translated by exploiting the host translational  
34    machinery via diverse strategies, which include cap- and poly(A)-independent  
35    translation mechanisms. Subsequently, viral genomes are synthesized on intracellular  
36    membranes in infected cells by RNA replicase complexes that consist of viral  
37    RNA-dependent RNA polymerase (RdRp), viral auxiliary proteins, host-encoded  
38    proteins, and viral RNAs (Ahlquist et al. 2003; den Boon et al. 2010; Nagy and Pogany

2012). Increasing evidence has demonstrated that viral replication proteins are multifunctional and play critical roles in recruiting the viral genomic RNAs to, and rearranging the specific cellular membranes, which are the sites of replication of positive-strand RNA viruses. This paper presents a brief review of viral RNA translation and replication mechanisms and refers to recent data obtained in the study of the *Red clover necrotic mosaic virus* (RCNMV), which is used by our group as a model virus.

#### ***Red clover necrotic mosaic virus***

RCNMV is a positive-strand RNA plant virus and a member of the genus *Dianthovirus* in the family *Tombusviridae*. This genus includes the *Carnation ring spot virus* (CRSV) as the type member and the *Sweet clover necrotic mosaic virus* (SCNMV) (Hiruki 1987). The genome of RCNMV consists of two RNAs (RNA1 and RNA2). The bipartite genome is unique among viruses of the family *Tombusviridae*, the genome of which is monopartite. RNA1 encodes putative RNA replicase components, an auxiliary 27 kDa protein (p27), and an 88 kDa protein (p88) with an RdRP motif. RNA1 also encodes a 37 kDa coat protein (CP) that is expressed from a subgenomic RNA (CPsgRNA). Transcription of the CPsgRNA requires an intermolecular interaction

between RNA1 and RNA2 (Sit et al. 1998; Tatsuta et al. 2005). RNA2 is a monocistronic RNA that encodes a movement protein (MP), which is required for viral cell-to-cell movement in plants (Kaido et al. 2011; Xiong et al. 1993).

## **Translation of RCNMV replicase proteins**

Many plant RNA viruses, including members of the *Tombusviridae* and the *Luteoviridae* families, lack both a 5' cap and a 3' poly(A) tail. Instead, they have cap-independent translation elements (CITEs) in the 3' untranslated region (UTR) of their genomic RNAs (Nicholson and White 2011). In some of these viruses, 3'CITEs bind the eukaryotic translation initiation factor complex eIF4F or eIFiso4F (Iwakawa et al. 2012; Nicholson and White 2011).

RCNMV RNA1 and RNA2 possess neither a cap structure at the 5' end nor a poly(A) tail at the 3' end (Mizumoto et al. 2003). Therefore, RCNMV proteins must be translated via cap-independent translation mechanisms. Two RNA elements play essential roles in the cap-independent translation of the replicase proteins encoded in RNA1. One element is the 3'CITE that consists of five stem-loop structures in the 3' UTR of RNA1 (Mizumoto et al. 2003). Another element is an A-rich sequence (ARS) located upstream of the 3'CITE (Iwakawa et al. 2012). A search for cellular

factors that bind the 3' UTR of RNA1 using RNA aptamer-based one-step affinity chromatography and mass spectrometry analysis led to the identification of the ARS as an RNA element that binds to poly(A)-binding protein (PABP) (Iwakawa et al. 2012). Mutagenesis and a tethering assay revealed that the direct interaction between PABP and ARS stimulates the 3'CITE-mediated translation of RCNMV RNA1, and that the PABP–ARS interaction is required for the recruitment of eukaryotic translation initiation factors (eIFs), such as eIF4s or eIF(iso)4Fs, to the 3' UTR and of the 40S ribosomal subunit to the viral mRNA (Iwakawa et al. 2012). Dianthoviruses might have evolved the ARS and 3'CITE as substitutes for the 3' poly(A) tail and the 5' cap of eukaryotic mRNAs for the efficient recruitment of eIFs, PABP, and ribosomes to the uncapped/nonpolyadenylated viral mRNA.

p88, which overlaps N-terminally with p27, is produced by programmed –1 ribosomal frameshifting (–1 PRF) (Kim and Lommel 1994). In addition, the production of p88 requires the long-distance base pairing between a bulge sequence in the stem-loop structure predicted just downstream of the –1 PRF site and the loop sequence of a small stem-loop structure predicted between the ARS and the 3'CITE (Tajima et al. 2011). Because p88, but not p27, is required in *cis* for the replication of RNA1 (Okamoto et al. 2008), the existence of RNA elements responsible for –1 PRF and 3'CITE-mediated translation in the 3' UTR of RNA1 might be important for switching

translation to replication of RNA1 and for regulating the ratio of p88 to p27.

## **Roles of viral replication proteins in RNA replication**

Viral RdRP is a key enzyme in the synthesis of viral RNAs. In addition to the RdRP, many positive-strand RNA viruses encode an auxiliary replication protein. Accumulating evidence indicates that the viral auxiliary replication proteins play crucial roles in recognizing viral RNA templates, targeting them to RNA replication sites, and assembling the viral replicase complexes. Thus, the auxiliary replication proteins play multiple roles in the accomplishment of viral RNA replication.

For example, the auxiliary replicase protein p33 of *Tombusvirus* binds directly to an internal replication element present in the viral replicon RNA (Pogany et al. 2005), interacts with p92 RdRP (p92<sup>pol</sup>) via a protein–protein interaction, and recruits the replicon RNA and p92<sup>pol</sup> to peroxisomal membranes (Panavas et al. 2005). These protein–RNA and protein–protein interactions are required for the assembly of the *Tombusvirus* replicase complex. In the Brome mosaic virus (BMV), which has a tripartite genome, the 1a auxiliary replication protein recognizes RNA elements that are present at the 5' end of RNA2 and in the intergenic region of RNA3 (Chen et al. 2001; Schwartz et al. 2002). The 1a protein, 2a RdRP (2a<sup>pol</sup>), and viral nucleotide sequences

are required for the assembly of the functional BMV replicase complex. The 1a protein recruits replication templates and 2a<sup>pol</sup> to the endoplasmic reticulum (ER) membrane, which is the site of BMV RNA replication (Liu et al. 2009; Schwartz et al., 2002).

In RCNMV, the p27 auxiliary replication protein binds specifically and directly to a Y-shaped RNA element (YRE) located in the 3' UTR of RNA2. The replicase–YRE interaction is critical for recruiting RNA2 to the membrane fraction (Hyodo et al. 2011; Iwakawa et al. 2011). The domains in p27 that are required for YRE binding were mapped to its central and C-terminal regions, and the critical amino acids of these regions were identified (Hyodo et al. 2011). The C-terminal half of p27 is also involved in the interaction with p88 (Mine et al. 2010b) and localizes at the ER membrane, together with p88 (Turner et al. 2004). The p27–p88 interaction is required for the formation of RCNMV replication complexes, as described below.

#### **Viral RNA replication complexes and host proteins involved in RNA replication**

One of the useful approaches to identify host genes affecting virus RNA replication is a systematic genome-wide screen using yeast single-gene deletion or essential-gene libraries, when the viruses of interest can replicate in yeast (Nagy 2008). Using the yeast systems, about 130 and 100 genes that affect the replication of tomato bushy stunt

virus (TBSV) and BMV, respectively, have been identified (Nagy 2008). Functional analyses in several of these genes demonstrated that the encoded proteins include RNA-binding proteins, cellular chaperons, membrane-shaping proteins, proteins associated with membrane remodeling and lipid synthesis, and others (Nagy and Pogany 2012). Screens of Arabidopsis mutant plants also led to the identification of several genes, such as TOM1 and TOM3, that encode membrane proteins and affect tomato mosaic virus replication (Yamanaka et al. 2000).

Host proteins required for plant RNA virus replication have also been identified by purifying membrane-associated viral RNA replication complexes using biochemical and immunological methods, followed by mass spectrometry and other methods. This approach led to the identification of several proteins, including a yeast homologue of mammalian glyceraldehyde 3-phosphate dehydrogenase (GAPDH) for TBSV, the eukaryotic translation elongation factor 1A for TBSV and *Turnip mosaic virus*, heat shock protein 70 (Hsp70) for TBSV (Nagy and Pogany 2012), and Hsp70, Hsp90, and ADP-ribosylation factor 1 (ARF1) for RCNMV, as described below.

RCNMV RNA replication complexes were isolated from virus-infected plant tissues using immunoprecipitation and the membrane-associated replication complexes were analyzed using blue native polyacrylamide gel electrophoresis (BN-PAGE) (Mine et al. 2010a). The purified fractions contained viral and host proteins and retained two types



of RdRP activities. One was an endogenous template-bound RdRP activity that synthesizes virus-related RNAs without adding RNA templates. The other was an exogenous template-dependent RdRP activity capable of *de novo* initiation of complementary RNA synthesis from selected RNA templates. The RCNMV replication complexes are bound tightly with membranes and their apparent molecular weight is 480 kDa. The complex with a template-dependent RdRP synthesizes RNA fragments by specifically recognizing the 3' terminal core promoter sequences of RCNMV RNAs.

Using mass spectrometry, we identified possible host proteins present in the affinity-purified RCNMV RNA-dependent RNA polymerase fraction, or in the 480 kDa complex fraction that was purified further via BN-PAGE. The former fraction contained Hsp70, Hsp90, ubiquitin, actin, hin1-like protein, several ribosomal proteins, ARF1, and other proteins in addition to viral replication proteins and CP (Mine et al. 2010a; A. Mine and T. Okuno, unpublished data). The latter fraction contained ubiquitin, ATP synthase subunit  $\alpha$ , hin1-like protein, and other proteins, in addition to viral replicase proteins (Mine et al. 2010a). Among these candidate proteins, we analyzed Hsp70, Hsp90, and ARF1 for their contribution to RCNMV RNA replication. Hsp70 and Hsp90 are well-known protein chaperones and ARF1 is a highly conserved, ubiquitous, small GTPase that is implicated in the formation of the COPI vesicles on Golgi membranes. Our recent experimental data suggest that these three proteins

172 interact directly and colocalize with p27 within the virus-induced large punctate  
173 structures of ER membranes. Downregulation of these proteins by virus-induced gene  
174 silencing decreased RCNMV RNA accumulation in plants. Furthermore, specific  
175 inhibitors of Hsp70, Hsp90, and ARF1, such as 2-phenylethynesulfonamide (PES),  
176 geldanamycin, and brefeldin A, respectively, inhibited the formation of the 480 kDa  
177 replication complex, and the synthesis of RCNMV RNA in an *in vitro* cell-free viral  
178 translation/replication system and in protoplast experiments (A. Mine, K. Hyodo and T.  
179 Okuno, unpublished data). These data suggest that Hsp70, Hsp90, and ARF1 play  
180 important roles in the formation of replication complexes and viral RNA synthesis.  
181 Confocal microscopy using GFP-fused Arf1 suggested that p27 induces the  
182 redistribution of Arf1 to large aggregate structures from small punctate structures that  
183 could be the Golgi apparatus (K. Hyodo and T. Okuno, unpublished data). Interestingly,  
184 inhibition of the nucleotide exchange activity of ARF1 by brefeldin A disrupts the  
185 ER-localization of p27 in RCNMV-infected cells. Although the precise role of Arf1 in  
186 the replication of RCNMV RNA is unclear at present, it is tempting to speculate that  
187 p27 functions as the recruiter of Arf1 to the replication site and acts in viral RNA  
188 replication via membrane modification and the formation of RNA replication  
189 complexes at the membrane. It is noteworthy that enteroviral 3A proteins bind and  
190 modulate Arf1 and its guanine nucleotide exchange factor GBF1 to enhance the

preferential recruitment of phosphatidylinositol-4-kinase IIIb (PI4KIIIb) to membranes, yielding a phosphatidylinositol 4-phosphate (PI4P) lipid-enriched microenvironment that differs from that found in uninfected cells (Hsu et al. 2010). Thus, plant and animal viruses might use common host factors to create a favorable environment for viral replication. Conversely, many host proteins used by viruses seem to differ among viruses. TBSV is affected by a set of host factors that is vastly different from that observed for BMV (Li et al. 2009). Down regulation of GAPDH, which inhibits TBSV accumulation, does not affect TMV accumulation in GAPDH-silenced plants (Wang and Nagy 2008).

## **Perspective**

Over the past 10 years, a large amount of information has been accumulated on *cis*-acting RNA elements and host proteins that play important roles in the replication of positive-strand RNA viruses. The cumulative evidence supports the idea that viral genomic RNAs that encode replication proteins can be conductors that orchestrate multiple processes using viral and host proteins and viral RNA elements. However, our current understanding of viral RNA replication and translation mechanisms is far from complete. Viral RNAs appear to change their structure regarding either local or global

structures in a spatially and temporally regulated manner, probably via the concerted action of viral and host proteins.

Screenings of host proteins that bind viral RNAs may represent a way to study the regulatory mechanisms of viral gene expression. One of the approaches is the use of RNA aptamer-based one-step affinity chromatography and mass spectrometry analysis to identify host proteins that bind specifically to viral RNAs. Several host proteins have been identified for RCNMV (Iwakawa et al. 2012). Another approach is a high-throughput method using yeast protein microarrays. This method has led to the identification of as many as 50 viral RNA-binding host proteins in the studies of BMV and TBSV; (Li et al. 2009; Zhu et al. 2007). Further studies using proteomics and genome wide-screens of co-opted host factors will also expand this field of research. Functional analyses of the candidate host proteins using biochemical methods and single-molecule techniques in cell-free systems in combination with analyses in living cells will be needed to test the roles of candidate proteins in virus replication processes. Confocal and electron microscopy experiments will also be useful to study the manner via which the candidate proteins function spatially and temporally in virus replication processes. Our understanding of the roles of viral and host proteins and viral RNA elements in viral replication may lead to the development of novel strategies aimed at controlling viral diseases.

229

230 **Acknowledgments**

231

232 The author thanks the staff and students of the Laboratory of Plant Pathology, Kyoto  
233 University, and all collaborators who supported our research. The author also thanks the  
234 researchers who provided materials and experimental protocols used in the research.  
235 Research in the author's laboratory was supported in part by a Grant-in-Aid for  
236 Scientific Research from the Japan Society for the Promotion of Science.

237

238 **References**

239

240 Ahlquist P, Noueiry AO, Lee WM, Kushner DB, Dye BT (2003) Host factors in  
241 positive-strand RNA virus genome replication. *J Virol* 77: 8181-8186  
242 den Boon JA, Diaz A, Ahlquist P (2010) Cytoplasmic Viral Replication Complexes.  
243 *Cell Host Microbe* 8: 77-85  
244 Chen J, Noueiry A, Ahlquist P (2001) *Brome mosaic virus* protein 1a recruits viral  
245 RNA2 to RNA replication through a 5' proximal RNA2 signal. *J Virol*  
246 75:3207-3219  
247 Hiruki C (1987) The dianthoviruses: A distinct group of isometric plant viruses with

- 248 bipartite genome. *Adv Virus Res* 33:257-300
- 249 Hsu, NY, Ilnytska O, Belov G, Santiana M, Chen YH, Takvorian PM, Pau C, van der  
250 Schaar H, Kaushik-Basu N, Balla T, Cameron CE, Ehrenfeld E, van Kuppeveld  
251 FJM, Altan-Bonnet N (2010) Viral reorganization of the secretory pathway  
252 generates distinct organelles for RNA replication. *Cell* 141:799–811
- 253 Hyodo K, Mine A, Iwakawa H-O, Kaido M, Mise K, Okuno T (2011) Identification of  
254 amino acids in auxiliary replicase protein p27 critical for its RNA-binding activity  
255 and the assembly of the replicase complex in *Red clover necrotic mosaic virus*.  
256 *Virology* 413:300–309
- 257 Iwakawa H-O, Mine A, Hyodo K, An M, Kaido M, Mise K, Okuno T (2011) Template  
258 recognition mechanisms by replicase proteins differ between bipartite  
259 positive-strand genomic RNAs of a plant virus. *J Virol* 85:497–509
- 260 Iwakawa H-O, Tajima Y, Taniguchi T, Kaido M, Mise K, Taniguchi H, Okuno T  
261 (2012) Poly(A)-binding protein facilitates translation of an  
262 uncapped/nonpolyadenylated viral RNA by binding to the 3' untranslated region. *J*  
263 *Virol* 86 (in press)
- 264 Kaido M, Funatsu N, Tsuno Y, Mise K, Okuno T (2011) Viral cell-to-cell movement  
265 requires formation of cortical punctate structures containing Red clover necrotic  
266 mosaic virus movement protein. *Virology* 413:205–215

- 267 Kim KH, Lommel SA (1994) Identification and analysis of the site of -1 ribosomal  
268 frameshifting in red clover necrotic mosaic virus. *Virology* 200:574–582
- 269 Li Z, Pogany J, Panavas T, Xu K, Esposito AM, Kinzy TG, Nagy PD (2009)  
270 Translation elongation factor 1A is a component of the tombusvirus replicase  
271 complex and affects the stability of the p33 replication co-factor.  
272 *Virology*:385:245-60
- 273 Liu L, Westler WM, den Boon JA, Wang X, Diaz A, Steinberg HA, Ahlquist P (2009)  
274 An amphipathic  $\alpha$ -helix controls multiple roles of Brome mosaic virus protein 1a  
275 in RNA replication complex assembly and function. *PLoS Pathog* 5(3): e1000351
- 276 Mine A, Takeda A, Taniguchi T, Taniguchi H, Kaido M, Mise K, Okuno T. (2010a)  
277 Identification and characterization of the 480-kilodalton template-specific  
278 RNA-dependent RNA polymerase complex of *Red clover necrotic mosaic virus*. *J*  
279 *Virol* 84:6070-6081
- 280 Mine A, Hyodo K, Takeda A, Kaido M, Mise K, Okuno T (2010b) Interactions between  
281 p27 and p88 replicase proteins of *Red clover necrotic mosaic virus* play an  
282 essential role in viral RNA replication and suppression of RNA silencing via the  
283 480-kDa viral replicase complex assembly. *Virology* 407:213-224
- 284 Mizumoto H, Tatsuta M, Kaido M, Mise K and Okuno T (2003) Cap-independent  
285 translational enhancement by the 3' untranslated region of *Red clover necrotic*

- 286        *mosaic virus* RNA1. J Virol 77:12113–12121
- 287    Nagy PD (2008) Yeast as a model host to explore plant virus-host interactions. Ann  
288        Rev Phytopathol 46:217-242.
- 289    Nagy PD, Pogany J (2012) The dependence of viral RNA replication on co-opted host  
290        factors. Nat Rev Microbiol 10:137-149
- 291    Nicholson BL, White KA (2011) 3'cap-independent translation enhancers of  
292        positive-strand RNA plant viruses. Curr Opin Virol 1:373-380
- 293    Okamoto K, Nagano H, Iwakawa H, Mizumoto H, Takeda A, Kaido M, Mise K, Okuno  
294        T (2008) *cis*-Preferential requirement of a-1 frameshift product p88 for the  
295        replication of *Red clover necrotic mosaic virus* RNA1. Virology 375:205-212
- 296    Panavas T, Hawkins CM, Panaviene Z, Nagy PD (2005) The role of the p33:p33/p92  
297        interaction domain in RNA replication and intracellular localization of p33 and  
298        p92 proteins of *Cucumber necrosis* tombusvirus. Virology 338:81-95
- 299    Pogany J, White KA, Nagy PD (2005) Specific binding of tombusvirus replication  
300        protein p33 to an internal replication element in the viral RNA is essential for  
301        replication. J Virol 79:4859-4869
- 302    Sit TL, Vaewhongs AA, Lommel SA (1998) RNA-mediated trans-activation of  
303        transcription from a viral RNA. Science 281:829–832



- 304 Schwartz M, Chen J, Janda M, Sullivan M, den Boon J, Ahlquist P (2002) A  
305 positive-strand RNA virus replication complex parallels form and function of  
306 retrovirus capsids. *Mol Cell* 9:505–514
- 307 Tajima Y, Iwakawa H, Kaido M, Mise K, Okuno T (2011) A long-distance RNA-RNA  
308 interaction plays an important role in programmed-1 ribosomal frameshifting in  
309 the translation of p88 replicase protein of *Red clover necrotic mosaic virus*.  
310 *Virology* 417:169-178
- 311 Turner KA, Sit TL, Callaway AS., Allen NS, Lommel SA (2004) Red clover necrotic  
312 mosaic virus replication proteins accumulate at the endoplasmic reticulum.  
313 *Virology* 320:276-290
- 314 Tatsuta M, Mizumoto H, Kaido M, Mise K, Okuno T (2005) The *Red clover necrotic*  
315 *mosaic virus* RNA2 trans-activator is also a cis-acting RNA2 replication element.  
316 *J Virol* 79:978–986
- 317 Wang RY-L, Nagy PD (2008) *Tomato bushy stunt virus* co-opts the RNA-binding  
318 function of a host metabolic enzyme for viral genomic RNA synthesis. *Cell Host*  
319 *Microbes* 3:178–187
- 320 Xiong Z, Kim KH, Giesman-Cookmeyer D, Lommel SA (1993) The roles of the red  
321 clover necrotic mosaic virus capsid and cell-to-cell movement proteins in  
322 systemic infection. *Virology* 192:27–32

- 323 Yamanaka T, Ohta T, Takahashi M, Meshi T, Schmidt R, Dean C, Naito S, Ishikawa M  
324 (2000) *TOM1*, an *Arabidopsis* gene required for efficient multiplication of a  
325 tobamovirus, encodes a putative transmembrane protein. Proc Natl Acad Sci USA  
326 97:10107–12
- 327 Zhu J, Gopinath K, Murali A, Yi G, Hayward SD, Zhu H, Kao C (2007) RNA-binding  
328 proteins that inhibit RNA virus infection. Proc Natl Acad Sci USA 104:3129–34