TITLE:

The first assessment of cyanobacterial and diazotrophic diversities in the Japan Sea

AUTHOR(S):

Hashimoto, Ryoya; Yoshida, Takashi; Kuno, Sotaro;
Nishikawa, Tetsuya; Sako, Yoshihiko

CITATION:

Hashimoto, Ryoya ...[et al]. The first assessment of cyanobacterial and diazotrophic diversities in the Japan Sea. Fisheries Science 2012, 78(6): 1293-1300

ISSUE DATE:

2012-11

URL:

<http://hdl.handle.net/2433/163407>

RIGHT:

The final publication is available at www.springerlink.com

1 **Title**

2 The first assessment of cyanobacterial and diazotrophic diversities in the Japan Sea

3

4 **Authors**5 Ryoya Hashimoto¹, Takashi Yoshida¹, Sotaro Kuno¹, Tetsuya Nishikawa² and Yoshihiko Sako¹.

6

7 **Affiliation**8 ¹ Graduate School of Agriculture, Kyoto University, Kyoto 606-8502, Japan, ² Tajima Fisheries

9 Technology Institute, Hyogo Prefectural Technology Center for Agriculture, Forestry and Fisheries,

10 Kasumi, Kami, Mikata, Hyogo 669-6541, Japan

11

12 Corresponding author. Postal address: Faculty of Agriculture, Kitashirakawa-Oiwake, Sakyo-ku, Kyoto

13 606-8502, Japan. Phone: (+81) 75-753-6218. Fax: (+81) 75-753-6226. E-mail:

14 yoshiten@kais.kyoto-u.ac.jp

15

Abstract

The diversity of cyanobacteria and dizaotrophs in the Japan Sea was investigated by analyzing sequences of cyanobacterial 16S rRNA genes and nitrogen fixation genes (*nifH*) from seawater sampled at a depth ranging from surface to 100 m at two stations. Of the 107 cyanobacterial 16S rRNA gene sequences obtained, 97 sequences and 3 sequences were assigned to *Synechococcus* sub-cluster 5.1 and *Prochlorococcus* HL (II), respectively. Unlike other oceanic regions, compositional ratio of the sequences assignable to *Synechococcus* sub-cluster 5.3 was relatively high (8%). No sequences of diazotrophic cyanobacteria were found in the cyanobacterial 16S rRNA genes. In *nifH* clone library (36 sequences), 10 sequences were identified as a UCYN-A group of diazotrophic cyanobacteria. Residual 26 sequences (72%) were assigned to proteobacteria. These results suggest heterotrophic bacteria including UCYN-A dominated in diazotrophic community in the Japan Sea. Our study revealed the dominance of *Synechococcus* in cyanobacterial community and (photo)heterotrophic diazotrophs in diazotrophic community in the Japan Sea, suggesting its unique characteristics.

Key words: cyanobacteria, diazotroph, Japan Sea, *Synechococcus* sub-cluster 5.1, *Synechococcus* sub-cluster 5.3, UCYN-A, *nifH*

33 Introduction

34 Marine cyanobacteria, especially two genera, *Prochlorococcus* and *Synechococcus*, significantly
35 contribute to 32-80% of the primary productivity in various oceanic regions [1–3]. Some other
36 cyanobacteria retain ability for N₂ fixation which is the process of reducing N₂ gas to biologically
37 available ammonium (referred to as diazotrophic cyanobacteria). The biological N₂ fixation is carried out
38 by several species of bacteria and archaea, but mainly by diazotrophic cyanobacteria in the ocean [4].
39 Therefore, marine cyanobacteria play an important role in carbon cycle as well as in nitrogen cycle.

40 Among marine diazotrophic cyanobacteria, filamentous nonheterocystous *Trichodesmium* spp. are the
41 most abundant diazotrophs in the ocean. *Trichodesmium* spp. are observed throughout the tropical and
42 subtropical Atlantic, Pacific, Indian oceans, and East and South China seas [5,6]. In North Atlantic Ocean,
43 their contribution to nitrogen inputs can be as great or greater than nitrate advection from deep sea water
44 [7]. In addition, heterocystous cyanobacterial diatom symbionts such as *Richelia* spp. and *Calothrix* spp.
45 have also been considered to make a significant contribution to N₂ fixation in North Atlantic Ocean and
46 North Pacific Ocean [8,9]. They are detected in tropical and subtropical regions in North Pacific Ocean
47 and North Atlantic Ocean [10,11]. Recently, some unicellular cyanobacteria have been recognized as
48 important diazotrophs in the ocean [12–14]. The unicellular diazotrophic cyanobacteria (UCYN) are
49 phylogenetically sub-divided into three groups, UCYN-A, B and C based on amino acid sequences of
50 NifH (dinitrogenase reductase), a subunit of the nitrogenase complex enzyme. UCYN-A is an
51 uncultivated cyanobacterium lacking oxygenic photosystem II, ribulose biphosphate carboxylase
52 (Rubisco), and tricarboxylic acid cycle, therefore UCYN-A is considered to be photoheterotroph [15].

UCYN-A is shown to be more abundant than other two groups with broad latitudinal range in the North and South Pacific Oceans[10,14]. Unlike UCYN-A, UCYN-B including *Crocospaera watsonii* WH8501 shows a patchy distribution in the ocean, and is detected in South Pacific, Eastern Atlantic, and North Pacific [10,14,16]. Compared to the other two unicellular diazotrophic cyanobacterial groups, UCYN-C including *Cyanothece* TW3 has less abundance in the open ocean [17,18].

The Japan Sea is a semi-enclosed marginal sea in the northwest Pacific surrounded by the Japanese islands and Asian continent. The Japan Sea is often referred to as “the miniature ocean” because the situation, which the course of the western branch of the Tsushima Current and the cold water generated in north and the warm water in east, is similar to those of the global ocean [19]. Its water exchange is limited in the upper layer because the Japan Sea is connected to adjacent seas with the shallow straits [19]. Although many studies have examined the diversity of cyanobacteria in various oceanic regions, less is known about the diversity of cyanobacteria, especially diazotrophic cyanobacteria in the Japan Sea.

In this study, we examined the diversity of cyanobacteria and diazotrophic cyanobacteria vertically in the Japan Sea using polymerase chain reaction (PCR) approach targeting cyanobacterial 16S rRNA genes and *nifH* genes.

Materials and Methods

Seawater collection

Sampling was conducted during Augaust-2011 on board the R/V Tajima at Station 2 (35°50.2N,

134°19.8E, 275 m depth) and Station 7 (37°20.2N, 134°19.8E, 2591 m depth) in the Japan Sea. Seawater was collected in 2l Niskin bottles mounted on a CTD (SBE-9 plus, Sea Bird Electronics) rosette from 0, 10, 30, 75, 100m, and deep chlorophyll maximum (DCM, near 50m). We measured profiles of temperature, salinity, and concentrations of chlorophyll a using the CTD.

DNA extraction

Each 300 ml sample was filtered through 0.2 µm pore-size polycarbonate filters (ADVANTEC, Tokyo, Japan), and then filters were wrapped in aluminum foil and were stored at 4 °C until stored at -80 °C in laboratory. Genomic DNA was extracted from the 0.2 µm filters using ISOIL kit (Nippon Gene, Tokyo, Japan).

PCR amplification

The cyanobacterial 16S rRNA gene sequences were amplified by PCR using primers described by Nübel et al. [20] (Table 1). The genomic DNA sample (1 µl) recovered from each station was added to each reaction mixture. The reaction mixtures were made with 5 µl of 10x PCR Buffer for Blend Taq (TOYOBO, Osaka, Japan), 5 µl of dNTPs (2 mM each), 0.4 µl of forward primer, 0.2 µl of each reverse primer, 0.5 µl of Blend Taq (2.5 U µl⁻¹; TOYOBO), and 37.7 µl of sterile Milli-Q water. After the initial denaturation step (2 min at 94°C), the samples were amplified for 30 cycles (30 s at 94°C, 30 s at 60°C, and 1min at 72°C). To amplify *nifH* genes, we performed nested PCR using the degenerate oligonucleotide primer pair described by Zehr and Turner [21] (Table 1). The reaction mixtures were made with 5 µl of 10x PCR Buffer for Blend Taq (TOYOBO), 5 µl of dNTPs (2 mM each), 2 µl of nifH3, 2 µl of nifH4, 0.5 µl of Blend Taq (2.5 U µl⁻¹; TOYOBO), and 34.5 µl of sterile Milli-Q water. The first

round of PCR consisted of an initial 5 min denaturation at 95°C, then 30 cycles of 1 min at 94°C, 1 min at 50°C and 1 min at 72°C, and final elongation of 7 min at 72°C. Of this reaction product, 1 µl was transferred into a second 50 µl reaction mixture containing the same reagent mixture, and primers were replaced with nifH1 and nifH2. The second round PCR consisted of an initial 5 min denaturation at 95°C, then 30 cycles of 1 min at 94°C, 1 min at 57°C and 1min at 72°C, and final elongation of 7 min at 72°C.

Cloning, Sequencing, and Phylogenetic Analysis

The amplified fragments were purified and cloned into pTac-1 vector (TOYOBO). We sequenced a total of 107 and 36 clones containing cyanobacterial 16S rRNA gene and *nifH* inserts, respectively. Basic Local Alignment Search Tool (BLAST) comparisons were used to find the sequence closely related to those amplified from our samples with the National Center for Biotechnology Information (NCBI) database [22]. The NifH phylogenetic tree based on the deduced amino acid sequences was constructed by the neighbour-joining method with Poisson correction using MEGA version 5 [23].

Nucleotide sequence accession numbers

The 16S rRNA gene sequences and *nifH* sequences reported in this paper have been deposited in the DDBJ under accession numbers AB727363-AB727505.

Results

The vertical thermal stratification was observed in surface water (0-30 m) of both Stn. 2 and Stn. 7 (Fig. 1). The water temperature in surface was relatively constant (25~26 °C) in Stn. 2 and Stn. 7, and water

temperature below 30 m depth declined gradually to 8.2 and 4.5 °C at 200 m depth, respectively (Fig. 1).

Salinity of surface water in Stn. 2 was around 32.7 psu, which was a bit lower than that in Stn. 7 (33.3 psu) (Fig. 1). This lower salinity of Stn. 2 was considered to be derived from freshwater input from the terrestrial area. In each station, the concentrations of chlorophyll *a*, the maximum of which was observed around 50 m depth, were extremely low ($< 0.2 \mu\text{M}$) compared with other oceanic regions [24].

We obtained a total of 107 clones of cyanobacterial 16S rRNA gene sequences from 12 samples in the Japan Sea (Table 2). Of these, 96 sequences (90%) and 3 sequences (3%) were assignable (~99 % nucleotide similarity) to *Synechococcus* sub-cluster 5.1, and *Prochlorococcus* HL-(high-light) (II), respectively. In each station and depth, *Synechococcus* sub-cluster 5.1 dominated the cyanobacterial community. Of the 96 sequences assignable to *Synechococcus* sub-cluster 5.1, 6 sequences showed a 100% identity with *Synechococcus* KORDI-19 which was isolated from the Japan Sea [25]. Residual 8 sequences (8%) were assignable to *Synechococcus* sub-cluster 5.3. This type of sequences was recovered from samples ranging from 10-75 m depth in each station. In general, *Prochlorococcus* has two distinct ecotypes, the high-light(HL)- and low-light(LL)-adapted ecotypes[26]. The sequences similar to a subgroup of the HL ecotype, *Prochlorococcus* HL (II) were obtained only from surface water (0 m), which is comparable with the previous reports that the HL-adapted ecotype generally distributes the upper regions of the euphotic zone [26–28]. No sequences assignable to the LL-*Prochlorococcus* were obtained, suggesting low abundance of LL-*Prochlorococcus* in the subsurface areas (Table 2).

We performed the *nifH*-targeting PCR amplification against the same 12 samples in the Japan Sea (Table 3). We obtained a total of 36 clones of *nifH*. However, no PCR products were detected from DNA

samples of Stn. 2 at DCM, 75 m, 100 m and Stn. 7 at 100 m, suggesting low abundance of diazotrophs in these areas. On the basis of phylogenetic analysis, the *nifH* sequences were divided into five groups (Fig. 2). Sequences assigned to cyanobacteria showed a similarity to the sequences within UCYN-A, and accounted for 28% of the *nifH* clone library. All sequences assignable to UCYN-A were recovered from Stn. 7 at 30m, DCM, and 75m. This result is consistent with the observations that the peak abundances of UCYN-A was observed around 50 m depth in the Pacific Ocean [14]. Other sequences (72%) were grouped within α -, β -, and γ - Proteobacteria. Sequences assigned as α -Proteobacteria (Group 4) showed 100% amino acid identity with sequences recovered from Baltic Sea and Chesapeake Bay (Fig. 2). In β -Proteobacteria, the sequences (Group 3) were recovered from Stn. 2 at 10m and Stn. 7 at 10m with 100% amino acid identity to *Burkholderia vietnamensis* G4 and sequences recovered from North Pacific Ocean. *Burkholderia vietnamensis* were also observed in reservoirs and agricultural settings [29]. Of sequences assigned to γ - Proteobacteria, 12 sequences were highly similar to protein sequences recovered from Atlantic Ocean, South China Sea and Mediterranean Sea.

Discussion

Marine *Synechococcus* is phylogenetically subdivided into three major subclusters, 5.1, 5.2, and 5.3. *Synechococcus* sub-cluster 5.1 dominates *Synechococcus* community in the open ocean [30]. In general, *Synechococcus* are more abundant in eutrophic coastal and mesotrophic open ocean waters with mixing conditions in water column, while *Prochlorococcus* is abundant in oligotrophic waters with stratified

conditions [2,31–33]. Although surface water of the Japan Sea is extremely nutrient-limited environment (eg. under-detection limit for PO_4 , NO_3 , NO_2) [34], our data showed the dominance of *Synechococcus* in the cyanobacterial community. This discrepancy suggests unknown factors controlling their distribution in the Japan Sea, which will be the subject of future work.

Synechococcus sub-cluster 5.3 accounted for 8% of the clone library with vertical distribution in each station [30, 35, 36] (Table 2). Huang et al. [30] report that their distribution may be restricted to certain oceanic regions because few sequences assignable to *Synechococcus* sub-cluster 5.3 were detected in the GOS database. However, their distribution and characteristics is not well understood.

No sequences belonging to diazotrophic cyanobacteria were detected in the cyanobacterial 16S rRNA gene clone library. It is not surprising because the abundance of diazotrophic cyanobacteria is two to three orders of magnitude lower than that of *Synechococcus* or *Prochlorococcus* in most oceanic regions [10,18,37,38]. In *nifH* clone library, sequences belonging to UCYN-A were obtained from Stn. 7 at 30m, DCM, 75m depths. Due to the elevated phytoplankton biomass around DCM, the carbon source may be more readily available for UCYN-A lacking the carbon fixation enzyme Rubisco. Sequences related to other unicellular cyanobacteria (i.e. UCYN-B and C) were not detected in our *nifH* clone library. Moisander et al. [37] suggest the possibility of limiting the growth of UCYN-A and B by differential grazing pressure due to the smaller cell size of UCYN-A compared to UCYN-B. However, the factors of limiting the distribution of the unicellular diazotrophic cyanobacterial groups have not unraveled yet well.

Sequences of filamentous cyanobacteria were also not detected in our clone libraries (Table 2). Diazotrophic cyanobacteria require a lot of iron as cofactor of nitrogenase and phosphorus for energy

generation during diazotrophic growth [6]. As unicellular cyanobacteria have high surface/volume ratio comparing to filamentous cyanobacteria, they may efficiently grow in oligotrophic environment because phosphate uptake efficiency is dependent upon cell size [39]. Therefore, the presence of UCYN-A and no detection of filamentous cyanobacteria may be due to the extremely nutrient poor condition in the Japan Sea with iron repletion [34].

About two-thirds of *nifH* clones were derived from proteobacteria, and especially sequences most related to γ -Proteobacteria were abundant. Although the sequences belonging to γ -Proteobacteria are detected in various oceans such as North Pacific Ocean and North Atlantic Ocean with accounting for a substantial fraction of the DNA/RNA *nifH* clone libraries, it is still unknown what extent these heterotrophic bacteria contribute to the net nitrogen input in the ocean [40,41]. However, despite the low N_2 fixation rates, heterotrophic bacteria dominated the diazotrophic community in the South Pacific with significant nitrogen input into the ocean [42]. Therefore, given the absence of other diazotrophic cyanobacteria, heterotrophic bacteria may account for substantial fraction of nitrogen input through N_2 fixation in the Japan Sea.

To our knowledge, this is the first study describing molecular cyanobacterial 16S rRNA genes and *nifH* diversity in the Japan Sea. Our data showed wide distribution of *Synechococcus* sub-cluster 5.3 in the Japan Sea vertically. Considering patchy distribution of *Synechococcus* sub-cluster 5.3 in other oceanic regions [30], the Japan Sea may be one of the suitable environments for them to propagate. The dominance of *Synechococcus* with oligotrophic well-stratified condition environment was contradict the general distribution of cyanobacteria in the ocean, therefore these results may reflect the unique

characteristics in the Japan Sea. We also showed that heterotrophic bacteria dominate the diazotrophic community. Further studies are required for determining what factors affect the vertical and horizontal distributions of different groups of these microorganisms in the Japan Sea.

Acknowledgments

This study was supported by Challenging Exploratory Research (No. 23658160).

References

1. Veldhuis MJW, Kraay GW, VanBleijswijk JDL, Baars MA (1997) Seasonal and spatial variability in phytoplankton biomass, productivity and growth in the northwestern Indian Ocean: the southwest and northeast monsoon. 1992-1993. *Deep Sea Res Part I* 44: 425–449.
2. Liu HB, Nolla HA, Campbell L (1997) *Prochlorococcus* growth rate and contribution to primary production in the equatorial and subtropical North Pacific Ocean. *Aquat Microb Ecol* 12: 39–47.
3. Li WKW (1994) Primary production of prochlorophytes, cyanobacteria, and eucaryotic ultraphytoplankton: Measurements from flow cytometric sorting. *Limnol Oceanogr* 39: 169–175.
4. Zehr JP (2011) Nitrogen fixation by marine cyanobacteria. *Trends Microbiol* 19: 162–173.
5. Capone DG, Zehr JP, Paerl HW, Bergman B, Carpenter EJ (1997) *Trichodesmium*, a globally significant marine cyanobacterium. *Science* 276: 1221–1229.
6. Chen Y-ling L, Chen H-yung, Tuo S-how, Ohki K (2008) Seasonal dynamics of new production from *Trichodesmium* N₂ fixation and nitrate uptake in the upstream Kuroshio and South China Sea basin. *Limnol Oceanogr* 53: 1705–1721.

- 216 7. Capone DG Burns JA, Michaels AF, Montoya JP, Subramaniam, A, Gunderson T, Carpenter EJ
217 (2005) Nitrogen fixation by *Trichodesmium* spp.: An important source of new nitrogen to the
218 tropical and subtropical North Atlantic Ocean. *Glob Biogeochem Cycles* 19: 786–787.

- 219 8. Mague TH, Weare NM, Holm-Hansen O (1974) Nitrogen fixation in the north Pacific Ocean.
220 *Mar Biol* 24: 109–119.

- 221 9. Foster RA, Kuypers MMM, Vagner T, Paerl RW, Musat N, Zehr JP (2011) Nitrogen fixation and
222 transfer in open ocean diatom-cyanobacterial symbioses. *ISME J* 5: 1484–1493.

- 223 10. Church MJ, Björkman KM, Karl DM, Saito MA, Zehr JP (2008) Regional distributions of
224 nitrogen-fixing bacteria in the Pacific Ocean. *Limnol Oceanogr* 53: 63–77.

- 225 11. Foster RA, Subramaniam A, Mahaffey C, Carpenter EJ, Capone DG, Zehr JP (2007) Influence of
226 the Amazon River plume on distributions of free-living and symbiotic cyanobacteria in the
227 western tropical north Atlantic Ocean. *Limnol Oceanogr* 52: 517–532.

- 228 12. Church MJ, Mahaffey C, Letelier RM, Lukas R, Zehr JP, Karl DM (2009) Physical forcing of
229 nitrogen fixation and diazotroph community structure in the North Pacific subtropical gyre. *Glob*
230 *Biogeochem Cycles* 23. GB2020, doi:10.1029/2008GB003418.

- 231 13. Montoya JP, Holl CM, Zehr JP, Hansen A, Villareal TA, Capone DG (2004) High rates of N₂
232 fixation by unicellular diazotrophs in the oligotrophic Pacific Ocean. *Nature* 430: 1027–1031.

- 233 14. Moisaner PH, Beinart RA, Hewson I, White AE, Johnson KS, Carlson CA, Montoya JP, and
234 Zehr JP (2010) Unicellular cyanobacterial distributions broaden the oceanic N₂ fixation domain.
235 *Science* 327: 1512–1514.

- 236 15. Zehr JP, Bench SR, Carter BJ, Hewson I, Niazi F, Shi T, Tripp J, Affourtit JP (2008) Globally
237 Distributed Uncultivated Oceanic N₂-Fixing Cyanobacteria Lack Oxygenic Photosystem II.
238 *Science* 322: 1110–1112.

- 239 16. Staal M, Hekkert ST, Brummer GJ, Veldhuis M, Sikkens C, Persijn S, Stal LJ (2007) Nitrogen
240 fixation along a north-south transect in the eastern Atlantic Ocean. *Limnol Oceanogr* 52:
241 1305–1316.

- 242 17. Needoba JA, Foster RA, Sakamoto C, Zehr JP, Johnson KS (2007) Nitrogen fixation by
243 unicellular diazotrophic cyanobacteria in the temperate oligotrophic North Pacific Ocean. *Limnol*
244 *Oceanogr* 52: 1317 – 1327.

- 245 18. Goebel NL, Turk KA, Achilles KM, Paerl R, Hewson I, Morrison AE, Montoya JP, Edwards CA,
246 Zehr JP (2010) Abundance and distribution of major groups of diazotrophic cyanobacteria and
247 their potential contribution to N₂ fixation in the tropical Atlantic Ocean. *Environ Microbiol* 12:
248 3272–3289.

- 249 19. Ichiye T (1984) Some problems of circulation and hydrography of the Japan Sea and the
250 Tsushima Current. *Ocean hydrodynamics of the Japan and East China Seas*. Elsevier, Netherlands,
251 pp. 15–54.

- 252 20. Nübel U, Garcia-Pichel F, Muyzer G (1997) PCR primers to amplify 16S rRNA genes from
253 cyanobacteria. *Appl Environ Microbiol* 63:3327–3332.

- 254 21. Zehr JP, Turner PJ (2001) Nitrogen Fixation: Nitrogenase Genes and Gene Expression. *Methods*
255 *Microbiol* 30: 271–286.

- 256 22. Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ (1997) Gapped
257 BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids*
258 *Res* 25: 3389–3402.

- 259 23. Tamura K, Peterson D, Peterson N, Stecher G, Nei M and KS (2011) MEGA5: Molecular
260 Evolutionary Genetics Analysis using Maximum Likelihood, Evolutionary Distance, and
261 Maximum Parsimony Methods. *Mol Biol Evol* doi:10.1093/molbev/msr121.

- 262 24. Dasgupta S, Singh RP, Kafatos M (2009) Comparison of global chlorophyll concentrations using
263 MODIS data. *Adv Space Res* 43: 1090–1100.

- 264 25. Choi DH, Noh JH (2009) Phylogenetic diversity of *Synechococcus* strains isolated from the East
265 China Sea and the East Sea. *FEMS Microbiol Ecol* 69: 439–448.

- 266 26. Rocap G, Larimer FW, Lamerdin J, Malfatti S, Chain P, Ahlgren NA, Arellano A, Coleman M,
267 Hauser L, Hess WR, Johnson ZI, Land M, Lindell D, Post AF, Regala W, Shah M, Shaw SL,
268 Steglich C, Sullivan MB, Ting CS, Tolonen A, Webb EA, Zinser ER, Chisholm SW. (2003)
269 Genome divergence in two *Prochlorococcus* ecotypes reflects oceanic niche differentiation.
270 *Nature* 424: 1042–1047.

- 271 27. Ahlgren NA, Rocap G, Chisholm SW (2006) Measurement of *Prochlorococcus* ecotypes using
272 real-time polymerase chain reaction reveals different abundances of genotypes with similar light
273 physiologies. *Environ Microbiol* 8: 441–454.

- 274 28. Zinser ER, Coe A, Johnson ZI, Martiny AC, Fuller NJ, Scanlan DJ, Chisholm SW. (2006)
275 *Prochlorococcus* ecotype abundances in the north atlantic ocean as revealed by an improved
276 quantitative PCR method. Appl Environ Microbiol 72: 723–732.
- 277 29. Menard A, Monnez C, Santos PEDL, Segonds C, Caballero-Mellado J, LiPuma, JJ, Chabanon G,
278 Cournoyer B. (2007) Selection of nitrogen-fixing deficient *Burkholderia vietnamiensis* strains by
279 cystic fibrosis patients: involvement of *nif* gene deletions and auxotrophic mutations. Environ
280 Microbiol 9: 1176–1185.
- 281 30. Huang S, Wilhelm SW, Harvey HR, Taylor K, Jiao N, Chen F. (2011) Novel lineages of
282 *Prochlorococcus* and *Synechococcus* in the global oceans. ISME J 6: 285–297.
- 283 31. Johnson ZI, Zinser ER, Coe A, McNulty NP, Woodward EM, Chisholm SW. (2006) Niche
284 partitioning among *Prochlorococcus* ecotypes along ocean-scale environmental gradients.
285 Science 311: 1737–1740.
- 286 32. Olson RJ, Chisholm SW, Zettler ER, Altabet MA, Dusenberry JA. (1990) Spatial and temporal
287 distributions of prochlorophyte picoplankton in the North Atlantic Ocean. Deep-Sea Res Part A
288 37: 1033–1051.
- 289 33. Bouman HA, Ulloa O, Scanlan DJ, Zwirgmaier K, Li WKW, Platt T, Stuart V, Barlow R, Leth O,
290 Clementson L, Lutz V, Fukasawa M, Watanabe S, Sathyendranath S. (2006) Oceanographic basis
291 of the global surface distribution of *Prochlorococcus* ecotypes. Science 312: 918–921.
- 292 34. Takata H, Kuma K, Iwade S, Isoda Y, Kuroda H, Senjyu T (2005) Comparative vertical
293 distributions of iron in the Japan Sea, the Bering Sea, and the western North Pacific Ocean. J
294 Geophys Res Lett 33, L02613, doi:10.1029/2005GL024538.
- 295 35. Dufresne A, Ostrowski M, Scanlan DJ, Garczarek L, Mazard S, Palenik BP, Paulsen IT, Tandeau
296 de Marsac N, Wincker P, Dossat C, Ferriera S, Johnson J, Post AF, Hess WR, Partensky F.
297 (2008) Unraveling the genomic mosaic of a ubiquitous genus of marine cyanobacteria. Genome
298 Biol 9:R90.
- 299 36. Scanlan DJ, Ostrowski M, Mazard S, Dufresne A, Garczarek L, Hess WR, Post AF, Hagemann M,
300 Paulsen I, Partensky F. (2009) Ecological genomics of marine picocyanobacteria. Microbiol
301 Molecul Biol Rev 73: 249–299.
- 302 37. Moisaner PH, Zhang R, Boyle EA, Hewson I, Montoya JP, Zehr JP. (2012) Analogous nutrient
303 limitations in unicellular diazotrophs and *Prochlorococcus* in the South Pacific Ocean. ISME J 6:
304 733–744.

- 305 38. Zwirgmaier K, Jardillier L, Ostrowski M, Mazard S, Garczarek L, Vaultot D, Not F, Massana R,
306 Ulloa O, Scanlan DJ. (2008) Global phylogeography of marine *Synechococcus* and
307 *Prochlorococcus* reveals a distinct partitioning of lineages among oceanic biomes. Environ
308 Microbiol 10: 147–161.
- 309 39. Smith REH, Kalff J (1982) Size-dependent phosphorus uptake kinetics and cell quota in
310 phytoplankton. J Phycol 18: 275–284.
- 311 40. Langlois RJ, Hümmel D, LaRoche J (2008) Abundances and distributions of the dominant *nifH*
312 phylotypes in the Northern Atlantic Ocean. Appl Environ Microbiol 74: 1922–1931.
- 313 41. Moisander PH, Beinart RA, Voss M, Zehr JP (2008) Diversity and abundance of diazotrophic
314 microorganisms in the South China Sea during intermonsoon. ISME J 2: 954–967.
- 315 42. Halm H, Lam P, Ferdelman TG, Lavik G, Dittmar T, LaRoche J, D'Hondt S, Kuypers MMM.
316 (2011) Heterotrophic organisms dominate nitrogen fixation in the South Pacific Gyre. ISME J
317 6:1238-1249.
- 318
- 319
- 320
- 321
- 322
- 323
- 324
- 325
- 326
- 327
- 328
- 329

Figure Legends

Figure 1

Vertical distributions (0-200 m) of temperature, salinity, chlorophyll *a* concentrations (mg m⁻³) at Stn. 2 (solid line) and Stn. 7 (dotted line) in Japan Sea on August 2011. The asterisk shows that fluorescent intensity was below zero.

Figure 2

Phylogenetic trees for *nifH*-deduced amino acid sequences from each sample constructed using the neighbor-joining method. Bootstrap values greater than 50% for 1000 replicates are indicated at respective nodes. Database entries for sequences are as follows: *Azotobacter chroococcum* strain NRRL B-14637 (AAR11505); *Pseudomonas stutzeri* A1501 (YP_001171863); *Vibrio diazotrophicus* clone CC1104A1 (AAP48983); *Sinorhizobium meliloti* bv (ABC67302); *Azospirillum brasilense* (P17303); *Rhodopseudomonas* sp. 99D (BAC07290); *Rhodobacter sphaeroides* 2.4.1 (YP_353614); *Burkholderia vietnamiensis* G4 (YP_001115195); *Rhodferax antarcticus* strain ANT.BR. clone 685K1_sp6 (ABI14697); *Sphingomonas azotifigens* (BAE71134); *Bradyrhizobium japonicum* USDA 110 (BAC47034); *Xanthobacter autotrophicus* Py2 (YP_001415004); *Crocospaera watsonii* WH 8501 (AAP48976); *Anabaena cylindrica* clone CC1085A1 (AAP48968); *Lyngbya lagerheimii* UTEX 1930 (AAA19029); *Cyanothece* sp. ATCC51142 (AAB61408); *Gloeotheca* sp. SK40 (BAF51659); *Gloeotheca* sp. KO68DGA (BAF51663); *Cyanothece* sp. WH 8902 (AAV28035); *Cyanothece*

352 sp. TW3 (BAJ21365); Unidentified marine bacterial clone HT1902 (AAK83129); Uncultured
353 marine bacterium clone TAP14405 (AAY60051); Cyanobacterium UCYN-A (ACJ53724);
354 *Oscillatoria sancta* clone CC1091A1 (AAP48970); *Trichodesmium thiebautii* (AAA77023);
355 *Trichodesmium erythraeum* IMS101 (AAA77022); *Methanosarcina acetivorans* C2A (NP_616144)

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372 Hashimoto et al., Figure 1

373

374

375

376

377

378

379

380

381

382

383

384

385

386

387 Hashimoto et al., Figure 2

Table 1. Cyanobacterial 16S rRNA and *nifH* primers used in this study.

Primer	Target site	Sequence(5'-3')	Reference
CYA359F	16S rRNA of cyanobacteria	GGG GAA TYT TCC GCA ATG GG	Nübel et al. 1997
CYA781R(a)	16S rRNA of cyanobacteria	GAC TAC TGG GGT ATC TAA TCC CAT T	Nübel et al. 1997
CYA781(b)	16S rRNA of cyanobacteria	GAC TAC AGG GGT ATC TAA TCC CTT T	Nübel et al. 1997
nifH1	<i>nifH</i> of prokaryotes	TGY GAY CCN AAR GCN GA	Zehr and McReynolds 1989
nifH2	<i>nifH</i> of prokaryotes	AND GCC ATC ATY TCN CC	Zehr and McReynolds 1989
nifH3	<i>nifH</i> of prokaryotes	ATR TTR TTN GCN GCR TA	Zani et al. 2000
nifH4	<i>nifH</i> of prokaryotes	TTY TAY GGN AAR GGN GG	Zani et al. 2000

408

409

410

411

412

413

414

415

416

417

418

419

420

421

422

423 Table 2. Summary of cyanobacterial 16S rRNA gene clones obtained from each microbial group.

Station	Depth	Number of Clones			Total
		<i>Synechococcus</i> sub-cluster 5.1	<i>Synechococcus</i> sub-cluster 5.3	<i>Prochlorococcus</i> HL(II)	
2	0m	7		2	9
	10m	9	1		10
	30m	9	1		10
	DCM	8	2		10
	75m	8			8
	100m	9	1		10
7	0m	9		1	10
	10m	10			10
	30m	9	1		10
	DCM	9	1		10
	75m	9	1		10
	100m				-

424

425

426

427

428

429

430

431

432

433

434

435 Table 3. Summary of *nifH* clones obtained from each microbial group.

Station	Depth	Number of Clones			Total
		Cyanobacterium	α -, β -	γ -	
		UCYN-A	Proteobacteira	Proteobacteria	
2	0m			5	5
	10m		2	2	4
	30m			5	5
	DCM				-
	75m				-
	100m				-
7	0m			4	4
	10m		2	1	3
	30m	4			4
	DCM	2	5		7
	75m	4			4
	100m				-

436

437