

Title	<Note> Comparison of the Longevity of Chimpanzee Beds between Two Areas in the Mahale Mountains National Park, Tanzania
Author(s)	Zamma, Koichiro; Makelele, Masayuke
Citation	Pan Africa News (2012), 19(2): 25-28
Issue Date	2012-12
URL	http://hdl.handle.net/2433/168176
Right	Copyright © Pan Africa News.
Type	Article
Textversion	publisher

ACKNOWLEDGEMENTS

I thank the Tanzania Commission for Science and Technology, the Tanzania Wildlife Research Institute, and the Tanzania National Parks for permission to work at Mahale, the Mahale Mountains Wildlife Research Centre and Mahale Mountains National Park for logistic support, and Dr. H. Hayaki and Dr. M. Nakamura for their valuable comments on this manuscript. This study was financially supported by MEXT Grant in-Aid for Scientific Research (#12375003 and #16255007 to T. Nishida)

REFERENCES

1. Nishida T, Wallauer W 2003. Leaf-pile pulling: An unusual play pattern in wild chimpanzees. *Am J Primatol* **60**:167–173.
2. Nishida T, Matsusaka T, McGrew WC 2009. Emergence, propagation or disappearance of novel behavioral patterns in the habituated chimpanzees of Mahale: A review. *Primates* **50**:23–36.
3. Nishida T, Zamma K, Matsusaka T, Inaba A, McGrew WC 2010. *Chimpanzee Behavior in the Wild: An Audio-Visual Encyclopedia*. Springer, Tokyo.
4. Arcadi AC, Robert D, Boesch C 1998. Buttress drumming by wild chimpanzees: Temporal patterning, phrase integration into loud calls, and preliminary evidence for individual distinctiveness. *Primates* **39**:505–518.
5. Nishida T 2003. Individuality and flexibility of cultural behavior patterns in chimpanzees. In: *Animal Social Complexity*. de Waal FBM, Tyack PL (eds), Harvard University Press, Cambridge, MA, pp. 392–413.
6. Wallin NL, Merker B, Brown S (eds) 2000. *The Origins of Music*. A Bradford Book, Cambridge, MA.
7. Fitch WT 2006. The biology and evolution of music: A comparative perspective. *Cognition* **100**:173–215.

<NOTE>

Comparison of the Longevity of Chimpanzee Beds between Two Areas in the Mahale Mountains National Park, Tanzania

Koichiro Zamma^{1,2} & Masayuke Makelele²

¹ Great Ape Research Institute, Hayashibara, Japan

² Mahale Mountains Chimpanzee Research Project
(E-mail: zamma@gari.jp)

INTRODUCTION

Chimpanzee (*Pan troglodytes*) beds have been used as an indicator for the presence of chimpanzees, and the longevity of beds is one of the important variables for estimating chimpanzee population sizes^{1,2}. The longevity of chimpanzee beds can differ due to location, vegetation, and season^{3–6}. For example, beds in forested sites decay faster than those in woodlands⁵, and beds in the dry season decay faster than those in the rainy season^{3,5}.

A previous study⁶ on the longevity of chimpanzee beds was conducted in the Mahale Mountains National Park, Tanzania. Because the study⁶ was conducted only in the forested area in the dry season, we studied chimpanzee beds both in forested and woodland sites during the rainy season.

METHODS

This study was conducted in Mahale Mountains National Park, Tanzania. This site has a rainy season from early October to mid-May⁷. We focused our research in two areas in the park, Kasoje and Miyako (Figure 1). The Kasoje area is composed of gallery forest^{8,9}, and is the center of the range of the M group of chimpanzees. The Miyako area consists of *Brachystegia* woodland with gallery forest along riverside area^{8,9}, and is occupied by chimpanzees of the Y group¹⁰.

Across both sites, we found 102 fresh beds from 3 October 2006 to 16 February 2007, and classified them by their date of construction (Table 1). Of these, we selected 70 beds to estimate bed longevity, and 32 beds to study bed construction. For the longevity study, we classified beds into two categories based on their time of construction. Beds constructed during Period 1 (3 to 10 October) were monitored until 24 May 2007. Beds constructed during Period 2 (11 October to 10 November) were monitored for a minimum of 100 days. Beds constructed after Period 2 were not included in analyses.

Beds were monitored approximately once per week (mean interval = 8.7 days, range = 6–28 days). KZ monitored the beds from 3 October 2006 to 16 February 2007, and MM monitored them from 17 February to 24 May 2007. We defined bed leaves as decayed when all of leaves had fallen from the branches in the bed^{3,5}, and defined beds as decayed when they were no longer clearly discernible as chimpanzee beds^{1,2,4–6}.

We monitored 9 beds in the Kasoje area and 7 beds in the Miyako area during Period 1. During Period 2, we monitored 38 beds in Kasoje area. The minimum distance between bed sites of the two areas was about 2 km (Figure 1).




Figure 1. Locations of chimpanzee beds in Kasoje (closed circle) and Miyako (open circle) during period 1.

Table 1. Tree species used for chimpanzee bed.

Area	Local name (Kitongwe)	Scientific name	Family name	Number of beds			
				Total	Period 1	Period 2	After Period 2
Kasoje	Lulumasha*	<i>Pycnanthus angolensis</i> #1	Myristicaceae	19	7	10	
	Mkibu gwesimbwa*	<i>Cordia millenii</i>	Boraginaceae	18		12	3
	Buhono*	<i>Pseudospondias microcarpa</i> #2	Anacardiaceae	13	1	6	
	Mjohoro	<i>Senna spectabilis</i> #3	Fabaceae	6			6
	Ibonobono*	<i>Croton sylvaticus</i>	Euphorbiaceae	5			1
	Msabasaba	<i>Syzygium</i> sp.	Myrtaceae	5		5	
	Mnyoni	<i>Bridelia micrantha</i> #4	Euphorbiaceae	3		3	
	Kafulujege*	<i>Lecaniodiscus fraxinifolius</i>	Sapindaceae	3			2
	Kahwibili	<i>Xylopia parviflora</i> #5	Annonaceae	3			
	Kamilaninga*	<i>Bridelia atroviridis</i>	Euphorbiaceae	2			
	Mjohoro (dead)§	#3		2			2
	Mlama*	<i>Combretum molle</i> #6	Combretaceae	2			1
	Mlyansekesi*	<i>Synsepalum brevipes</i>	Sapotaceae	2	1		
	Bulonje*	<i>Dracaena usambarensis</i> #7	Dracaenaceae	1			
	Ikubila*	<i>Ficus sur</i>	Moraceae	1			
	Mtelele*	<i>Stereospermum kunthianum</i>	Bignoniaceae	1			1
	Muhoko	<i>Aporrhiza paniculata</i>	Sapindaceae	1			
	Buhono* & Kahwibili†	#2 & #5		2		1	
	Buhono* & Lulumasha*†	#2 & #1		1			
	Buhono* & Mlama*†	#2 & #6		1			
Buhono* & Bulonje*†	#2 & #7		1				
Kahwibili & Mnyoni†	#5 & #4		1		1		
Kahwibili & Lulumasha*†	#5 & #1		1				
Kahwibili & Mgongogongo*†	#5 & <i>Anthocleista schweinfurthii</i>	Loganiaceae	1				
Miyako	Mlonje*	<i>Mimusops bagshawei</i>	Sapotaceae	4	4		
	Kashindabilangulube	<i>Blighia unijugata</i>	Sapindaceae	1	1		
	Lisalike	?	?	1	1		
	Mkusi	<i>Croton megalocarpus</i>	Euphorbiaceae	1	1		
Total				102	16	38	16

*: Fruits, leaves, and/or piths of this plant were eaten by chimpanzees.

†: Two species of tree were used to make a bed.

§: Two beds were constructed at the base of dead trees but were built from living vines including *Saba comorensis* (Apocynaceae), *Keetia venosa* (Rubiaceae), and *Tetracera potatoria* (Dilleniaceae).

RESULTS

We recorded 21 tree species that were used for constructing chimpanzee beds (Table 1). The most frequently used species was *Pycnanthus angolensis*, followed by *Cordia millenii* and *Pseudospondias microcarpa*. Many of the trees in which beds were constructed (78/102) are known food trees of Mahale chimpanzees. Eight beds were also found in the invasive tree, *Senna spectabilis*, and two of them were constructed at the base of dead trees.

The mean longevity of chimpanzee beds during Period 1 was 87 days ($n = 9$, range = 21–231) in the Kasoje area and 164 days ($n = 7$, range = 13 to > 233) in the Miyako area (Figure 2). Three beds in the Miyako area were not decayed by the end of the survey period. Bed longevity was significantly longer in Miyako than in Kasoje (Mann-Whitney U test, $n_1 = 9$, $n_2 = 7$, $U = 12$, $p < 0.05$).

During Period 2, we monitored 38 chimpanzee beds in Kasoje. We set a 100 day limit for monitoring these beds and more than half of the beds (22/38) were decayed

by the end of this period (Figure 3). This was not statistically different from the decay rate for Period 1 beds in Kasoje over 100 days (6/9; Fisher's exact test $p = 0.72$), but was significantly greater than the decay rate for beds in Miyako (1/7; Fisher's exact test, $p < 0.05$).

The mean time for leaf decomposition in beds was 76 days in Kasoje ($n = 9$, range = 14–158) and 126 days in Miyako ($n = 7$, range = 13–157). These numbers were not statistically different according to a Mann-Whitney U test (Figure 4, $n_1 = 9$, $n_2 = 7$, $U = 15$, $p = 0.088$).

DISCUSSION

There are two methods for measuring the longevity of chimpanzee beds: the time until leaves decays and the time until the bed structure decays^{1–6}. Bed leaves decayed faster in Kasoje in the dry season (49 days, $n = 41$, range = 18–110)⁶ than in the rainy season (76 days) in this study. In tree species of bed no remarkable differences could be seen between this and previous study, and *Pycnanthus angolensis* was the most frequently used species in both studies. Leaves in the dry season are thought to have




Figure 2. Rate of decay of chimpanzee beds in period 1.
*: Three beds were remained undecayed.




Figure 3. Rate of decay of chimpanzee beds in Kasoje in Period 2.




Figure 4. Rate of decay of bed leaves in Period 1.

lower moisture content and, thus, decay faster³. The result in this study is consistent with previous studies^{3,5}.

In contrast, the longevity of bed structures in Kasoje was longer in the dry season (131 days, $n = 41$, range = 30–189)⁶ than in the rainy season (87 days) in this study. This indicates that different mechanisms are relevant for the decay of leaves in beds than for decay of the bed structure. Over the period of this study (3 October 2006–24 May 2007) 1,562 mm of precipitation fell and almost two-thirds of the total precipitation (1,001 mm) fell during the first 3 months (Mahale Mountains Chimpanzee Research Project data). We propose a hypothesis: because dead leaves often remain in branch structure in beds and absorb rainfall during the rainy season, the increased weight of the wet branches may contribute to the deterioration of the bed structure.

The longevity of chimpanzee beds in Miyako was longer than that in Kasoje, but this difference cannot be explained by rainfall because the amount of precipitation did not differ between the two areas¹¹. The longevity of chimpanzee beds in woodlands in the Ugalla area, Tanzania is long (more than 358 days (Yoshikawa M, personal communication); 432 days⁵), and beds in Miyako area also

constructed in woodland. The strength of branches of tree species in woodland may be a key factor contributing to the longevity of chimpanzee beds in these areas.

The present study and previous studies (e.g. ref. 1–6) indicate that the longevity of chimpanzee beds differs according to season and vegetation type. These studies indicate the need for broader range of bed longevity or for appropriate estimates of bed longevity in each site to accurately estimate chimpanzee population sizes.

ACKNOWLEDGEMENTS

We thank TAWIRI, TANAPA and COSTECH for permission to do the field research, and MMNP and MMWRC for logistic support. We also thank T. Sakamaki for providing information about the beds of Y group chimpanzees, H. Bunengwa, M. Mwami, H. Ihobe, N. Itoh, H. Ogawa, and M. Yoshikawa for their help and advices, the members of MMCRP for providing precipitation data, and M. Kiyono and C. Tokimatsu for data handling. This study was financially supported by the Global Environment Research Fund of the Ministry of Environment, Japan (F-061 to T. Nishida).

REFERENCES

1. Ghiglieri MP 1984. *The Chimpanzees of Kibale Forest*. Columbia University Press, New York.


2. Tutin CEG, Fernandez M 1984. Nationwide census of gorilla (*Gorilla g. gorilla*) and chimpanzee (*Pan t. troglodytes*) populations in Gabon. *Am J Primatol* 6:313–336.
3. Plumptre AJ, Reynolds V 1996. Censusing chimpanzees in the Budongo Forest, Uganda. *Int J Primatol* 17:85–99.
4. Ogawa H, Idani G, Moore J, Pintea L, Hernandez-Aguilar A 2007. Sleeping parties and nest distribution of chimpanzees in the savanna woodland, Ugalla, Tanzania. *Int J Primatol* 28:1397–1412.
5. Stewart FA, Piel AK, McGrew WC 2011. Living archaeology: Artifacts of specific nest site fidelity in wild chimpanzees. *J Hum Evol* 61:388–395.
6. Ihobe H 2005. Life span of chimpanzee beds at the Mahale Mountains National Park, Tanzania. *Pan Afr News* 12:10–12.
7. Itoh N, Nakamura M, Ihobe H, Uehara S, Zamma K, Pintea L, Seimon A, Nishida T 2012. Long-term changes in the social and natural environments surrounding the chimpanzees of the Mahale Mountains National Park. In: *The Ecological Impact of Long-Term Changes in Africa's Rift Valley*. Plumptre AJ (ed), NOVA Science, New York, pp. 249–277.
8. Nishida T 1972. A note on the ecology of the red-colobus monkeys (*Colobus badius tephrosceles*) living in the Mahali Mountains. *Primates* 13:57–64.
9. Nishida T, Uehara S 1981. Kitongwe name of plants: A preliminary listing. *Afr Stud Monogr* 1:109–131.
10. Sakamaki T, Nakamura M, Nishida T 2007. Evidence of cultural differences in diet between two neighboring unit groups of chimpanzees in Mahale Mountains National Park, Tanzania. *Pan Afr News* 14:3–5.
11. Takasaki H, Nishida T, Uehara S, Norikoshi K, Kawanaka K, Takahata Y, Hiraiwa-Hasegawa M, Hasegawa T, Hayaki H, Masui K, Huffman MA 1990. Summary of meteorological data at Mahale research camps, 1973–1988 In: *The Chimpanzees of the Mahale Mountains: Sexual and Life History Strategies*. Nishida T (ed), University of Tokyo Press, Tokyo, pp. 291–300.

<BOOK INFO>

Wild Cultures: A Comparison between Chimpanzee and Human Cultures

By Christophe Boesch

How do chimpanzees say, “I want to have sex with you?” By clipping a leaf or knocking on a tree trunk? How do they eat live aggressive ants? By using a short stick with one hand or long stick with both? Ivorian and Tanzanian chimpanzees answer these questions differently, as would humans from France and China if asked how they eat rice. Christophe Boesch takes readers into the lives of chimpanzees from different African regions, highlighting the debate about culture. His ethnography reveals how simple techniques have evolved into complex ones, how teaching styles differ, how material culture widens access to new food sources and how youngsters learn culture. This journey reveals many parallels between humans and chimpanzees and points to striking differences. Written in a vivid and accessible style, *Wild Cultures* places the reader in social and ecological contexts that shed light on our twin cultures.



296 pages
Cambridge University Press
Publication date: September 2012
Hardcover \$95.00
ISBN: 9781107025370

Contents

- Introduction
- 1. Studying culture in the wild
- 2. From human culture to wild culture
- 3. Shaping nature into home: about material culture
- 4. One for all and all for one: about social culture
- 5. I want to have sex with you: about symbolic culture
- 6. Learning culture: from pupils to teachers
- 7. Dead or alive? Towards a notion of death and empathy
- 8. Wild culture – wild intelligence
- 9. Uniquely chimpanzee – uniquely human
- Epilogue: will we have the time to study chimpanzee culture?
- References
- Index

Pan Africa News, Vol. 19, No.2
Published in December, 2012
Address: c/o Human Evolution Studies,
Dept. of Zoology, Faculty of Science,
Kyoto Univ., Kyoto, 606-8502, JAPAN
TEL: (+81)75-753-4093
FAX: (+81)75-753-4115
E-mail: pan.editor@gmail.com
URL: <http://mahale.main.jp/PAN/>
ISSN: 1884-751X (Print), 1884-7528 (Online)