

# On Algebraic Knots I

— Computability of Their Jones Polynomials —

Yuanan Diao<sup>†1</sup>, Claus Ernst<sup>‡2</sup> and Uta Ziegler<sup>§3</sup>

<sup>†</sup> Department of Mathematics and Statistics  
University of North Carolina at Charlotte  
Charlotte, NC 28223, USA

<sup>‡</sup> Department of Mathematics

<sup>§</sup> Department of Computer Science  
Western Kentucky University  
Bowling Green, KY 42101, USA

**Abstract:** We prove that the Jones polynomial of any Conway algebraic link diagram with  $n$  crossings can be computed in  $O(n^2)$  time. In particular, the Jones polynomial of any Montesinos link and two-bridge knot or link with minimum crossing number  $n$  can be computed in  $O(n^2)$  time.

## 1 Conway algebraic knots

A knot  $\mathcal{K}$  is called a *Conway algebraic knot* (or just algebraic knot for short) if it admits a diagram  $D$  in  $S^2$  such that there exists a set of disjoint simple closed curves  $C_1, C_2, \dots, C_n$  such that each  $C_j$  intersects  $D$  transversely in exactly four non-crossing points of  $D$  and the regions bounded by each  $C_j$  are of the two types shown in Figure 1. Each such simple closed curve  $C_j$  is referred to as a Conway circle. Notice that the type B regions bounded by the Conway circles shown in the figure do not contain any crossings of  $D$ .




Figure 1: A. A Conway region containing a single Conway circle with a single crossing from the knot diagram in it. B. A Conway region bounded by three Conway circles without any crossings from the knot diagram.

In general, it is not known which algebraic knots admit a minimal crossing diagram that can be decomposed by Conway circles as described above. There exist algebraic knots whose minimal diagrams are not algebraic diagrams. The class of algebraic knots is very large: it contains all two-bridge knots and all Montesinos knots, as well as many other knots. For a more detailed discussion on algebraic knots, one may refer to [2].

<sup>1</sup>E-mail: ydiao@uncc.edu

<sup>2</sup>E-mail: claus.ernst@wku.edu

<sup>3</sup>E-mail: uta.ziegler@wku.edu

## 2 Algebraic diagrams and tangle replacement operations

Given a simple link diagram  $D_0$  and a 2-string tangle  $\Gamma_1$ , a *tangle replacement operation* on  $D_0$  using  $\Gamma_1$  is to replace a crossing in  $D_0$  by  $\Gamma_1$ . See Figure 2 for an example. This results in a diagram  $D_1$ . The tangle replacement operation may be repeated on  $D_1$  using a different 2-string tangle  $\Gamma_2$  to obtain yet another new knot diagram  $D_2$ . If this process is repeated  $m$  times using tangles  $\Gamma_1, \Gamma_2, \dots, \Gamma_m$ , we obtain a knot diagram  $D_m$ . Let  $|\Gamma_j|$  be the number of crossings in  $\Gamma_j$  and let  $d = \max\{|\Gamma_1|, |\Gamma_2|, \dots, |\Gamma_m|, |D_0|\}$ . We say that  $D_m$  is obtained by  $m$  tangle replacement operations (using the tangles  $\Gamma_1, \Gamma_2, \dots, \Gamma_m$ ) with *tangle depth*  $d$ . Of course, the tangle depth of the knot diagram  $D_m$  depends on the particular choice of the tangle replacement operation sequence which may not be unique.



Figure 2: Tangle replacement operation: replacing a single crossing by a tangle.

By induction on the number of crossings in a diagram one can prove the following theorem.

**Theorem 2.1** *If  $D$  is an algebraic knot diagram with  $n$  crossings, then it can be obtained by a sequence of tangle replacement operations with tangle depth 2.*

## 3 Face graphs of algebraic knot diagrams

Figure 3 shows an example of how to obtain a (signed) face graph of a knot projection: shade the regions of  $D$  in a checkerboard fashion and each shaded region corresponds to a vertex in the face graph  $F_D$  and a crossing between two shaded regions becomes an edge in  $F_D$  connecting the two vertices corresponding to these regions. The signs of the edges in  $F_D$  are determined by the relation of the shaded regions at the crossing in  $D$  as shown in Figure 3.




Figure 3: A projection of the knot  $9_{45}$  and its corresponding face graph.

Replacing a crossing in  $D$  by tangle  $\Gamma$  corresponds to replacing an edge in  $F_D$  by a plane graph  $N$ , see Figure 4.  $N$  is obtained from a facegraph generated by shading a knot or link diagram  $D'$  that contains  $\Gamma$  and an additional crossing  $c$ . The extra crossing  $c$  corresponds to an edge with label  $e$  and  $N$  equals to the face graph of  $F_{D'} - e$ . The operation that replaces an edge  $e$  in  $F_D$  by a plane graph  $N$  as shown in the figure is called a (single edge) *tensor product* of  $F_D$  and  $N$  and is denoted by  $F_D \otimes_e N$  (or simply  $F_D \otimes N$ ). Using Theorem 2.1, we can then prove the following theorem.

**Theorem 3.1** *Let  $D$  be an algebraic knot diagram with  $n$  crossings, then there exists a sequence of (connected) plane graphs  $D_0, N_1, \dots, N_{n-2}$  such that each of these graphs contains exactly two edges and the face graph  $F_D = (\dots((D_0 \otimes N_1) \otimes N_2) \otimes \dots) \otimes N_{n-2}$ .*




Figure 4: A knot diagram before and after a tangle replacement and the corresponding face graphs.

### 4 Tutte polynomials of colored graphs

A colored graph is a graph  $G$  in which each edge is assigned a color from a color set  $\Lambda$ . If  $\Lambda = \{+, -\}$ , then  $G$  is called a signed graph. Four variables  $x_\lambda, y_\lambda, X_\lambda$  and  $Y_\lambda$  are associated with each color  $\lambda \in \Lambda$ . Let  $e_\lambda$  be an edge in  $G$  with color  $\lambda$ , then the (colored) Tutte polynomial of  $G$  can be defined recursively by

$$T(G) = \begin{cases} Y_\lambda T(G \setminus e), & \text{if } e \text{ is a loop} \\ X_\lambda T(G/e), & \text{if } e \text{ is a bridge,} \\ y_\lambda T(G \setminus e) + x_\lambda T(G/e), & \text{otherwise.} \end{cases}$$

It is a well known result of Bollobás and Riordan [1] that  $T(G)$  is well-defined (that is, it is independent of the order of the edges used in the recursions) as long as the color variables satisfy the following conditions:

$$\det \begin{pmatrix} X_\lambda & y_\lambda \\ X_\mu & y_\mu \end{pmatrix} = \det \begin{pmatrix} x_\lambda & Y_\lambda \\ x_\mu & Y_\mu \end{pmatrix} \text{ and } \det \begin{pmatrix} x_\lambda & Y_\lambda \\ x_\mu & Y_\mu \end{pmatrix} = \det \begin{pmatrix} x_\lambda & y_\lambda \\ x_\mu & y_\mu \end{pmatrix}.$$

For the sake of convenience, sometimes we just use the name of an edge as its color.

**Theorem 4.1** [5] *Let  $M$  and  $N$  be two colored graphs such that the color variables involved satisfy the above conditions. Furthermore, assume that  $N$  is a connected graph with exactly two edges (with colors  $c_1$  and  $c_2$ ), then  $T(M \otimes_e N)$  can be obtained from  $T(M)$  by the following color variable assignments to  $X_e, Y_e, x_e$  and  $y_e$ :*

Case 1.  $N$  consists of a two edge path:

$$X_{c_1} X_{c_2} \mapsto X_e \quad x_{c_1} x_{c_2} \mapsto x_e \quad y_{c_1} X_{c_2} + x_{c_1} Y_{c_2} \mapsto Y_e \quad y_{c_1} X_{c_2} + x_{c_1} y_{c_2} \mapsto y_e.$$

Case 2.  $N$  is a cycle consisting of two vertices and two edges:

$$y_{c_1} X_{c_2} + x_{c_1} Y_{c_2} \mapsto X_e \quad y_{c_1} x_{c_2} + x_{c_1} Y_{c_2} \mapsto x_e \quad Y_{c_1} Y_{c_2} \mapsto Y_e \quad y_{c_1} y_{c_2} \mapsto y_e.$$

### 5 The Jones polynomials of algebraic knots

For knots with a large crossing number, the computation of their knot invariants can be very difficult. For example, the computation of the Jones polynomial of a link is known to be NP-hard [6]. This prevents the computation of the Jones polynomial of knots with large crossing numbers in general. However, special classes of knots may allow a fast computation of some of their invariants. For example, Murakami et al. [9] recently proved that the Jones polynomial of any 2-bridge knot or link of crossing number  $n$  can be computed in  $O(n^2 \ln n)$  time.

The authors greatly extended the above result to the following theorem.

**Theorem 5.1** [4] *The Jones polynomial of any Conway algebraic link diagram with  $n$  crossings can be computed in  $O(n^2)$  time. Consequently, the Jones polynomial of any Montesinos link or any two-bridge knot or link of crossing number  $n$  can be computed in  $O(n^2)$  time.*

It is well known that the Jones polynomial  $J(t)$  of a knot  $\mathcal{K}$  can be obtained from a signed version of the Tutte polynomial of the face graph obtained from a regular projection  $D$  of  $\mathcal{K}$  [6, 7, 8], using the substitutions below and then multiplying by  $(-A^{-3})^{w(D)}$  (where  $A = t^{-1/4}$  and  $w(D)$  is the writhe of the diagram  $D$ ):

$$\begin{aligned} -A^{-3} &\mapsto X_+, & -A^3 &\mapsto X_-, & -A^3 &\mapsto Y_+, & -A^{-3} &\mapsto Y_-, \\ A &\mapsto x_+, & A^{-1} &\mapsto x_-, & A^{-1} &\mapsto y_+, & A &\mapsto y_-. \end{aligned} \quad (1)$$

The result of the theorem then follows from Theorems 3.1 and 4.1 by examining carefully how  $T(F_D)$  (and hence  $J(t)$ ) can be obtained through the repeated variable substitutions given in (1) and Theorem 4.1. Interested reader may refer to [4] for details.

## Acknowledgment

The authors are partially supported by NSF grants DMS-0712958 and DMS-0712997.

## References

- [1] B. Bollobás and O. Riordan, A Tutte Polynomial for Coloured Graph, *Combinatorics, Probability and Computing* **8** (1999), 45–93.
- [2] F. Bonahon and L. Siebenmann, *Geometric splittings of Knots and Conway's algebraic knots*, unpublished manuscript, 1980.
- [3] G. Burde and H. Zieschang, *Knots*, de Gruyter, Berlin, 1986.
- [4] Y. Diao, C. Ernst and U. Ziegler, *Jones Polynomial of Knots formed by Repeated Tangle Replacement Operations*, preprint. Available at [http://www.math.uncc.edu/preprint/2008/2008\\_03.pdf](http://www.math.uncc.edu/preprint/2008/2008_03.pdf).
- [5] Y. Diao, G. Heteyi and K. Hinson, *Invariants of composite networks arising as a tensor product*. Preprint. Available at [http://www.math.uncc.edu/preprint/2007/2007\\_06.pdf](http://www.math.uncc.edu/preprint/2007/2007_06.pdf).
- [6] F. Jaeger, D.L. Vertigan and D.J.A. Welsh, On the Computational Complexity of the Jones and Tutte Polynomials, *Mathematical Proceedings of the Cambridge Philosophical Society* **108** (1990), 35–53.
- [7] L.H. Kauffman, New Invariants in the Theory of Knots, *American Mathematical Monthly* **95**(3) (1988), 195–242.
- [8] L.H. Kauffman, A Tutte Polynomial for Signed Graphs, *Discrete Applied Mathematics* **25** (1989), 105–127.
- [9] M. Murakamia, M. Harab, M. Yamamoto and S. Tani, *Fast algorithms for computing Jones polynomials of certain links*, *Theoretical Computer Science* **374** (2007), 10–24.