TITLE:

Improved models of the piezomagnetic field for the 2011 Mw 9.0 Tohoku-oki earthquake

AUTHOR(S):

Yamazaki, Ken'ichi

CITATION:

Yamazaki, Ken'ichi. Improved models of the piezomagnetic field for the 2011 Mw 9.0 Tohoku-oki earthquake. *Earth and Planetary Science Letters* 2013, 363: 9-15

ISSUE DATE:

2013-02-01

URL:

<http://hdl.handle.net/2433/169695>

RIGHT:

© 2012 Elsevier B.V.; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。; This is not the published version. Please cite only the published version.

- 1 Improved models of the piezomagnetic field for the 2011 Mw 9.0
- 2 Tohoku-oki earthquake
- 3
- 4 Ken'ichi Yamazaki*
- 5 *Miyazaki Observatory, Research Center for Earthquake Prediction, Disaster Prevention*
- 6 *Research Institute, Kyoto University, 3884 Kaeda, Miyazaki 889–2161, Japan*
- 7
- 8 *Corresponding author. Tel.: +81 985651161, Fax: +81 985554005
- 9 *E-mail address:* kenichi@rcep.dpri.kyoto-u.ac.jp (K. Yamazaki)
- 10
- 11

12

13 **ABSTRACT**

14 To assess the feasibility of observing changes in the magnetic field produced by the
15 piezomagnetic effect, an improved model of the piezomagnetic field corresponding to
16 the Mw 9.0 Tohoku-oki earthquake is presented. In contrast to an earlier study, the
17 proposed model explicitly considers the spatial distribution of slip on the seismic fault,
18 and the results from this new model differ significantly from those of the previous
19 model where slip distributions were ignored. Quantitative aspects of the piezomagnetic
20 effect are discussed through comparisons of data and models. One feature clarified is
21 that, because the fault rupture is so far offshore, the expected amplitudes are quite small
22 at onshore existing observation sites; consequently, there would have been little chance
23 of observing sizable piezomagnetic signals at inland sites during the Tohoku-oki
24 earthquake. Nevertheless, piezomagnetic signals were reportedly detected at a few sites,
25 possibly indicating that the stress sensitivity or the initial magnetization was larger (by
26 several factors) than assumed. On the other hand, relatively large variations in the
27 magnetic field of up to 10 nT may have occurred offshore. This means that if
28 ocean-bottom sensors had been installed, larger piezomagnetic signals would have been
29 detected. Moreover, the piezomagnetic field in offshore areas is sensitive to the detailed

30 slip distribution, suggesting that observations of the magnetic field at ocean-bottom

31 sites might provide important constraints on determination of slip models.

32

33 Keywords: piezomagnetic effect; stress sensitivity; 2011 Tohoku-oki earthquake; slip

34 distribution

35

36

37 **1. Introduction**

38

39 The piezomagnetic effect, which describes changes in magnetization caused by
40 mechanical stress, predicts changes in the Earth's magnetic field following a major
41 earthquake. In earlier studies (e.g. Sasai, 1991, 1994, and references therein), a
42 constitutive law of the relation between stress changes and magnetization changes has
43 been proposed, as follows:

44

$$45 \quad \Delta M_i = \frac{3}{2} \beta \Delta T_{ij} M_j, (i, j = x, y, z) \quad (1)$$

46

47 where ΔM_i is the change in remanent and induced magnetization, ΔT_{ij} is the deviatoric
48 stress tensor, M_j is the initial total magnetization (i.e. the sum of induced and remanent
49 magnetization without changes in stress), and β is a proportional coefficient that is
50 usually referred to as the (piezomagnetic) stress sensitivity. Because of the
51 piezomagnetic effect, it should be possible to monitor changes in stress in the Earth's
52 crust with geomagnetic observations. Changes in the magnetic field that arise from the
53 piezomagnetic effect are referred to as piezomagnetic fields. These fields are inverted to

54 changes in magnetization in terms of the magnetic Coulomb's law, and further inverted
55 to changes in stress in terms of eq. (1).

56 However, the usefulness of the piezomagnetic effect as a tool for monitoring
57 changes in stress is still not clear. Whether or not the piezomagnetic field can be
58 observed depends on the spatial distribution of the piezomagnetic field which, in turn,
59 depends on source type, depth and distance. If the piezomagnetic field has a detectable
60 magnitude that is restricted to just a narrow area, then its detection will be difficult with
61 poorly or sparsely spaced arrays of instruments. Numerical examinations of realistic
62 source models need to be performed in order to assess the detectability of the
63 piezomagnetic field.

64 The piezomagnetic stress sensitivity is another uncertain factor that determines
65 whether or not the piezomagnetic field is detectable. While the proportional relation (i.e.
66 eq. 1) is partially inferred from considerations based on thermodynamics (e.g.
67 Nakamura and Nagahama, 1997), and the values of stress sensitivity can be determined
68 by theoretical considerations (Stacey and Johnston, 1972), the actual magnetization
69 fraction and type varies from rock to rock and representative values for a particular
70 region must be determined from magnetic anomaly maps, geology and laboratory
71 experiments. Laboratory experiments (e.g. Nagata and Kinoshita, 1967) suggest that

stress sensitivities are on the order of 10^{-9} Pa^{-1} . A stress sensitivity of this order is usually assumed when the piezomagnetic effect is considered in studies of volcanoes (e.g. Currenti et al., 2005) and earthquakes (e.g. Okubo et al., 2011). However, these values are sometimes too small to explain the observed offsets in the magnetic field associated with stress changes (e.g. Nishida et al., 2004; Oshiman et al., 1990; Zhan, 1989). The effective values of the stress sensitivity on the geophysical scale (i.e. larger than the laboratory scale) should be evaluated by comparing observational and theoretical models.

The 2011 Mw 9.0 Tohoku-chihou Taiheiyou-oki earthquake (herein referred to as the Tohoku-oki earthquake), which occurred on the boundary between the Pacific and Eurasian plates, is one event for which the magnitudes of the piezomagnetic field can be examined. The Tohoku-oki earthquake is the largest seismic event to have been observed with a dense network of modern geophysical instruments. Along with seismological and geodetic data, geomagnetic data were obtained for this extreme event. Utada et al. (2011) presented a prompt and comprehensive report on observed variations in the geomagnetic field associated with the Tohoku-oki earthquake. Together with several types of geomagnetic variations that followed the earthquake, they also reported that magnetic field offsets, which probably arose from the piezomagnetic effect, are

actually observed, but they are only up to 1.0 nT at the observation sites. In their conclusions, Utada et al. (2011) presented a negative view on the detectability of the piezomagnetic field.

Although the observations reported in Utada et al. (2011) provide constraints on the phenomena that actually occurred at the time of the earthquake, their conclusions about the piezomagnetic effect need to be reconsidered because they are based on oversimplified source models that ignore the spatial distribution of slip on the fault. Any reconsideration should incorporate improved piezomagnetic field models in the hope of clarifying the quantitative nature of the piezomagnetic field and evaluating the usefulness of observing it.

The aims of this study are to: (1) provide constraints on the piezomagnetic stress sensitivity around the Tohoku region, near the seismic fault of the Mw 9.0 Tohoku-oki earthquake; and (2) assess the usefulness of the magnetic observations as a tool for detecting stress changes. To these ends, improved models of the piezomagnetic field are presented, and the various models are compared and assessed using the data presented by Utada et al. (2011).

2. Procedures for modeling the piezomagnetic field

This study presents a new model of the piezomagnetic field in which the spatial distribution of slip on the plate-boundary fault is explicitly considered. In many studies of the piezomagnetic field in relation to earthquakes (e.g. Nishida et al., 2007), uniform slip models are employed. However, the slip on a fault is generally heterogeneous, and it is therefore preferable to explicitly consider the spatial distribution of slip. In a uniform slip model, the fault parameters are averaged to obtain a single slip parameter. Such a simplification is valid if the spatial scale of the slip on the fault is smaller than the distance between the observation site and the fault. This criterion is not satisfied in the case of the Tohoku-oki earthquake, for which the spatial scale of the slip and the distance from the surface expression of fault rupture to observational sites are of the order of 100 km. Therefore, the assumption of uniform fault slip used previously by Utada et al. (2011) is likely inadequate in the calculations of the piezomagnetic fields produced by the earthquake.

This study considers the heterogeneous fault slip models of Hayes (2011) and Shao et al. (2011) (herein, referred to as the USGS and UCSB models, respectively), which are derived from seismic wave inversion. Averaged slips from these models were used

125 in the calculations of Utada et al. (2011). These models were published immediately
126 following the Tohoku-oki earthquake and we can expect further refinements as more
127 complete geophysical data sets are taken in account (e.g. Koketsu et al., 2011).
128 Nevertheless, important characteristics of the Tohoku-oki fault slip are apparent in both
129 the USGS and UCSB models. For example, both models indicate large slip near the
130 trench axis. With these slip distributions, it is possible to evaluate the importance of
131 heterogeneous slip distributions and to compare our results with those of Utada et al.
132 (2011).

133 Piezomagnetic fields that correspond to a heterogeneous slip model are calculated
134 according to the following procedure. The fault plane (i.e. plate boundary) is divided
135 into sub-faults, and on each sub-fault the slip is assumed to be uniform. The total
136 piezomagnetic signal at any point is the sum of the contributions from all sub-faults, and
137 each contribution can be calculated by analytical formulae (Utsugi et al., 2000). Using
138 this procedure, we can calculate the piezomagnetic signals that correspond to
139 heterogeneous slip models for the Tohoku-oki earthquake.

140 Because the above procedure involves formulae derived by Utsugi et al. (2000), all
141 the assumptions involved in the formulations of Utsugi et al. (2000) are also used in the
142 present models. The initial total magnetization of the crust is assumed to be uniform

between the ground surface and a constant Curie point depth. The Earth's crust is approximated by elastic half-space, surface of which locates at sea level. These assumptions are not satisfied in reality, thus producing some uncertainty in the models obtained for the piezomagnetic field. Errors should also be included in the slip models, as mentioned above. To estimate the importance of uncertainty in the slip models and Curie point depths, we calculate the piezomagnetic fields that correspond to the two slip models (USGS and UCSB) with two values of H (15 and 30 km). The values for H used here are the same as those used by Utada et al. (2011), and they provide reasonable estimates for the island arc of the Tohoku district and for the subduction zone east of the Tohoku district of Japan (Tanaka et al., 1999). The effects of heterogeneities in the initial magnetization will be discussed separately, later.

3. Features of the new piezomagnetic models

Using the above procedures, and the parameters listed in Table 1, models of the piezomagnetic field have been constructed. The spatial distribution of the expected amplitude of the piezomagnetic effect is shown in Fig. 1. Observations are assumed to have been made at sea level, i.e. the surface of a uniform elastic half-space. Below, I

161 enumerate the features that are commonly observed in the results and which correspond
162 to all sets of parameters. It should be noted that the absolute values given in the results
163 are strongly dependent on the assumed sets of parameters. For this reason, this analysis
164 focuses on relative rather than absolute values.

165 Relatively large signals of the piezomagnetic field are expected to occur in offshore
166 areas in all cases. For the UCSB slip model, piezomagnetic fields larger than 3 nT are
167 predicted in offshore areas. For the USGS slip models, the predicted piezomagnetic
168 fields are smaller than those for the UCSB model, yet changes larger than 1 nT are
169 predicted.

170 In contrast, the expected amplitudes of piezomagnetic signals over the more distant
171 onshore, including the sites of observation, are rather small. Over a vast part of the land
172 area, the predicted amplitudes of the piezomagnetic field are up to 0.4 nT. Precise
173 values of the expected changes at the observation sites are listed in Table 2, together
174 with the observed changes reported by Utada et al. (2011). In some models, the changes
175 predicted at some locations are as large as 0.6 nT. For example, the predicted change at
176 the ESA site is 0.6 nT for model b (i.e. UCSB slip model with $H = 30$ km) whereas the
177 predicted change at the same site is zero for other models. There is no location where all
178 the models predict changes greater than 0.4 nT.

179 It should be pointed out that these models are quite different from those that use
180 uniform slip distribution. Figure 2 shows the calculated piezomagnetic field intensities
181 that correspond to uniform slip models where the slip parameters are averaged over the
182 fault plane. Numerous differences can be observed between Figs. 1 and 2. For example,
183 the amplitudes of the signals predicted with the uniform slip model do not exceed 1.0 nT,
184 except for some localized areas. If we focus on this result, the impression is that the
185 detection of coseismic piezomagnetic signals is a hopeless task, even if the
186 observational area is extended to the seafloor. However, the amplitudes of signals
187 predicted by the heterogeneous slip model are larger than 1–2 nT across a wide area of
188 ocean. In this case, the amplitudes of the coseismic piezomagnetic signals would have
189 been detected, if suitable magnetometers had been installed in the region.

190 The large differences that exist between the uniform and heterogeneous slip models
191 highlight the importance of considering the heterogeneous model for the Tohoku-oki
192 earthquake. The large differences also indicate that many of the conclusions about
193 piezomagnetic signals by Utada et al. (2011) need to be reconsidered and probably
194 changed.

195

4. Discussion

The goals of this study were to provide constraints on stress sensitivity, and to assess the usefulness of magnetic observations as tools for monitoring stress. The former can be accomplished by comparing the data with the models. The latter can be achieved by analyzing the constructed model. These matters are further discussed below.

4.1. Possible values of the piezomagnetic stress sensitivity

To provide constraints on the stress sensitivity using the results of piezomagnetic modeling, I make reference to the data presented by Utada et al. (2011). The idea is as follows. In the proposed model, the stress sensitivity (β) and the intensity of the initial total magnetization ($M = (M_x^2 + M_y^2 + M_z^2)^{1/2}$) are assumed to be given as in Table 1. The assumed values are possibly different from the actual values. As the calculated value of F_p (denoted by $F_p^{\text{calculated}}$) is proportional to the assumed value of βM , $[(\beta M)^{\text{assumed}}]$, the difference between $(\beta M)^{\text{assumed}}$ and the actual value of βM $[(\beta M)^{\text{actual}}]$ yields the disparity between the observed value of F_p (F_p^{observed}) and the calculated F_p .

214 $(F_p^{\text{calculated}})$. The value of $(\beta M)^{\text{actual}}$ is given by

215

$$216 \quad (\beta M)^{\text{actual}} = \frac{F_p^{\text{observed}}}{F_p^{\text{calculated}}} \times (\beta M)^{\text{assumed}}. \quad (2)$$

217

218 If F_p^{observed} and $F_p^{\text{calculated}}$ correlate well, it is possible to determine a plausible value of

219 $(\beta M)^{\text{actual}}$.

220 Regrettably, the correlation between observed and calculated signals of the

221 piezomagnetic field is not good (Fig. 3). This means that assumptions of a uniform

222 Curie point depth, a uniform initial total magnetization, and/or an assumed slip model,

223 are inadequate. In particular, ignoring the heterogeneity of the initial magnetization is

224 possibly problematic because it is known to enhance the piezomagnetic field (e.g.

225 Oshiman, 1990). Aeromagnetic surveys over the Tohoku region have shown that

226 magnetic anomalies in this region are rather strong (i.e. 10–100 nT) (Fig. 4), raising the

227 possibility of a strong heterogeneity in the initial total magnetization. Consideration of

228 the heterogeneity of the initial magnetization is clearly important if we are to calculate

229 the piezomagnetic field accurately. However, an accurate determination of the structure

230 of the initial magnetization is generally laborious and full of possible errors; hence, an

231 accurate determination of a generated piezomagnetic field is difficult in the presence of

a strong heterogeneity in the initial magnetization (e.g. Yamazaki, 2011).

Nevertheless, we can attempt to provide constraints on the possible values of the stress sensitivity using data just from sites KAK and KTR. Around these sites, the gradient of the magnetic anomaly is relatively small (Fig. 4), and we can therefore anticipate that the model with uniform initial magnetization will provide reasonable calculated results. The amplitude of the piezomagnetic signal observed at KTR was -0.8 ± 0.2 nT, whereas those predicted in the theoretical models (Fig. 1a–d) are between -0.2 and -0.3 nT. The piezomagnetic signal observed at KAK was -0.22 , whereas those predicted by theoretical models are between -0.07 and -0.22 . To explain the observations at KAK and KTR, the actual value of βM needs to be larger than the value assumed in the present calculation (i.e. $1.0 \text{ Pa}^{-1} \text{ Am}^{-1}$) by factors of 2–3. Provided that the assumption of $M = 1.0 \text{ A/m}$ (Table 1) is correct, the above result means that the stress sensitivity is about $2.0\text{--}3.0 \times 10^{-9} \text{ Pa}^{-1}$. This value is on the same order as that assumed in many piezomagnetic models (e.g. Johnston et al., 1989).

4.2 Potential usefulness of seafloor magnetic observations

In all the models of the piezomagnetic field shown in Fig. 2, the amplitudes of the

250 piezomagnetic signals are expected to be small on land and relatively large offshore.
251 Because the actual value of βM is possibly larger than assumed, as discussed in the
252 previous subsection, the actual changes in the magnetic field are possibly larger than
253 those shown in Fig. 1. In particular, large offsets in the magnetic field are expected near
254 the trench axis. Although this result is obtained for an assumption that observations are
255 made at sea level, piezomagnetic signals are also expected to be large at seafloor
256 because the seafloor is rather closer to the rupture. If offshore ocean-bottom
257 magnetometers had been installed, they would have detected significant amplitudes of
258 piezomagnetic signals corresponding to the Tohoku-oki earthquake.

259 Observing piezomagnetic signals would not be very useful if the piezomagnetic
260 signals were insensitive to details of the fault parameters, but the results of the
261 piezomagnetic models demonstrate that this is not the case. Indeed, the spatial
262 distributions of the piezomagnetic field are strongly dependent on the slip model that is
263 adopted. It is possible that we could have improved the determination of earthquake
264 source parameters if data from ocean-bottom magnetometers had been available, instead
265 of relying solely on the results of inversions of seismic and geodetic data.

266 In general, it is difficult to measure accurately the distributions of slip for
267 earthquakes that occur on an offshore plate boundary, because geodetic measurements

268 are made mainly on land. In the case of the Tohoku-oki earthquake, extremely large
269 slips near the trench axis have been suggested by inversions of the seismic data (e.g.
270 Hayes, 2011; Shao et al., 2011), but better constraints on slip distributions could have
271 been obtained from seafloor geodetic measurements (Sato et al., 2011). Given that
272 seafloor geodetic equipment is costly and difficult to manage, geomagnetic observations
273 might provide useful additional data for monitoring interplate earthquakes along
274 subduction zones. This solution may still apply, even when we consider magnetic
275 anomalies on the seafloor, because heterogeneities in the magnetization of the crust may
276 possibly enhance the generated piezomagnetic field (e.g. Oshiman, 1990).

277 Regrettably, there are also drawbacks to making seafloor magnetic observations.
278 First, seafloor observations are quite costly. Second, it may be difficult to keep the
279 sensors stably located during quakes, and if a sensor is displaced during a quake, an
280 apparent change in the magnetic field will be recorded. Even if it were possible to
281 obtain accurate data of the geomagnetic field at a certain point, it would be necessary to
282 consider heterogeneities of the initial magnetization and ocean-bottom topography when
283 converting the observed changes in the magnetic field to fault source parameters. For
284 these reasons, the usefulness of observing the piezomagnetic field remains uncertain.
285 However, similar difficulties also exist with respect to making ocean-bottom geodetic

observations. Not only are they extremely costly, but monument stability during earthquakes is also a problem. It is also difficult to process the observations correctly and obtain precise geodetic information. A decision on prioritizing geodetic and geomagnetic techniques should be based on which drawbacks are most easily overcome. If costs allow, an integrated use of both techniques is most desirable because they independently bring useful information to bear on these tectonic phenomena.

5. Conclusions

To calculate the piezomagnetic field that corresponds to the 2011 Mw 9.0 Tohoku-oki earthquake, it is necessary to consider the best representation of the spatial distribution of slip along the fault, and in this paper, I demonstrate the importance of such a consideration, and construct an appropriate slip model. Although this model still cannot entirely explain the observed distribution of piezomagnetic signals, constraints are obtained from data at two onshore sites, where the model seems to provide adequate results. Comparisons between the data and the model show the stress sensitivity to be about $2\text{--}3 \times 10^{-9} \text{ Pa}^{-1}$, which is on the same order as that assumed in many piezomagnetic models. Models of the piezomagnetic field predict that changes in the

304 geomagnetic total forces, due to the piezomagnetic effect, will be relatively large in
305 offshore areas closer to the rupture, and relatively small onshore, far from the rupture.
306 Because the expected magnitudes of the piezomagnetic signals are small at existing sites,
307 stress sensitivity of the piezomagnetic effect is likely to be on the order of 10^{-9} , though
308 this is not tightly constrained. Nevertheless, the possibility of a large piezomagnetic
309 field occurring at ocean-bottom stations is not excluded. Details of the spatial
310 distribution of the piezomagnetic field in oceanic areas are highly dependent on the slip
311 model used. The implication is that detection of the piezomagnetic field with
312 ocean-bottom magnetometers might have provided constraints on the slip models of the
313 Tohoku-oki earthquake, if such observations had been available.

314

315 **Acknowledgments**

316

317 Slip models proposed by Shao et al. (2011) and Hayes (2011), and a program coded
318 by M. Utsugi, were used to calculate the piezomagnetic fields. Numerical data on the
319 geomagnetic anomalies around Japan were provided by the Geospatial Information
320 Authority of Japan through their Web site. Comments and advice from anonymous
321 reviewers and the journal editor (P. Shearer) helped to improve an earlier version of the

322 manuscript. The software Generic Mapping Tools (Wessel and Smith, 1998) was used to

323 prepare some of the figures, including maps.

324

325 **References**

326

327 Currenti, G., Del Negro, C., Nunnari, G., 2005. Inverse modelling of volcanomagnetic

328 fields using a generic algorithm technique. *Geophys. J. Int.* 163, 403–418. doi:

329 10.1111/j.1365-246X.2005.02730.x

330 Hayes, G.P., 2011. Rapid source characterization of the 2011 Mw 9.0 off the Pacific

331 coast of Tohoku Earthquake. *Earth Planets Space* 63, 529–534.

332 Johnston, M. J. S., 1989. Review of magnetic and electric field effects near active faults

333 and volcanoes in the U.S.A. *Phys. Earth Planet. Interiors*, **57**, 47–63.

334 Koketsu, K., Yokota, Y., Nishimura, N., Yagi, Y., Miyazaki, S., Satake, K., Fujii, Y.,

335 Miyake, H., Yamanaka, Y., Sakai, S., Okada, T., 2011. A unified source model for

336 the 2011 Tohoku earthquake. *Earth Planet. Sci. Lett.* 310, 480–487.

337 Nagata, T., Kinoshita, H., 1967. Effect of hydrostatic pressure on magnetostriction and

338 magnetocrystalline anisotropy of magnetite. *Phys. Earth Planet. Int.* 1, 44–48.

339 Nakamura, N., Nagahama, H., 1997. Magnetic susceptibility and plastic strain of rocks

340 by the differential geometric theory of the physical interaction field. *Phys. Chem.*341 *Earth* 22, 167–173.

342 Nishida, Y., Sugisaki, Y., Takahashi, K., Utsugi, M., Oshima, H., 2004. Tectonomagnetic

- 343 study in the eastern part of Hokkaido, NE Japan: discrepancy between observed
344 and calculated results. *Earth Planets Space* 56, 1049–1058.
- 345 Nishida, Y., Utsugi, M., Mogi, T., 2007. Tectonomagnetic study in the eastern part of
346 Hokkaido, NE Japan (II): Magnetic fields related with the 2003 Tokachi-oki
347 earthquake and the 2004 Kushiro-oki earthquake. *Earth Planets Space* 59,
348 1181–1186.
- 349 Okubo, K., Takeuchi, N., Utsugi, M., Yumoto, K., Sasai, Y., 2011. Direct magnetic
350 signals from earthquake rupturing: Iwate–Miyagi earthquake of M 7.2, Japan.
351 *Earth Planet. Sci. Lett.* 305, 65–72.
- 352 Oshiman, N., 1990. Enhancement of tectonomagnetic change due to non-uniform
353 magnetization in the Earth's crust — two dimensional case studies. *J. Geomag.*
354 *Geoelectr.* 42, 607–619.
- 355 Oshiman, N., Sasai, Y., Miyakoshi, J., Nishida, R., Shiozaki, I., 1990. Continuous
356 observation of piezomagnetic changes due to ground loading by Lake Nichinan,
357 Tottori, Japan. *Proceedings of Conductivity Anomaly Symposium (1991)*, 137–148.
358 (In Japanese)
- 359 Sasai, Y., 1991. Tectonomagnetic modeling on the basis of the linear piezomagnetic
360 effect. *Bull. Earthq. Res. Inst., Univ. Tokyo* 66, 585–722.

- 361 Sasai, Y., 1994. Piezomagnetic fields produced by dislocation sources. *Surv. Geophys.*
362 15, 363–382.
- 363 Sato, M., Ishikawa, T., Ujihara, N., Yoshida, S., Fujita, M., Mochizuki, M., Asada, A.,
364 2011. Displacement Above the Hypocenter of the 2011 Tohoku-Oki Earthquake.
365 *Science* 332, 1395.
- 366 Shao, G., Li, X., Ji, C., Maeda, T., 2011. Focal mechanism and slip history of 2011 Mw
367 9.1 off the Pacific coast of Tohoku earthquake, constrained with teleseismic body
368 and surface waves. *Earth Planets Space* 63, 559–564.
- 369 Stacey, F.D., Johnston, M.J.S., 1972. The theory of the piezomagnetic effect. *Pure Appl.*
370 *Geophys.*, 97, 146–155.
- 371 Tanaka, A., Okubo, Y., Matsubayashi, O., 1999. Curie point depth based on spectrum
372 analysis of the magnetic anomaly data in east and southeast Asia. *Tectonophys.* 306,
373 461–470.
- 374 Utada, H., Shimizu, H., Ogawa, T., Maeda, T., Furumura, T., Yamamoto, T., Yamazaki,
375 N., Yoshitake, Y., Nagamachi, S., 2011. Geomagnetic field changes in response to
376 the 2011 off the Pacific Coast of Tohoku Earthquake and Tsunami. *Earth Planet.*
377 *Sci. Lett.* 311, 11–27.
- 378 Utsugi, M., Nishida, Y., Sasai, Y., 2000. Piezomagnetic potentials due to an inclined

- 379 rectangular fault in a semi-infinite medium. *Geophys. J. Int.* 140, 479–492.
- 380 Wessel, P., Smith, W.H.F., 1998. New, improved version of Generic Mapping Tools
- 381 released. *EOS Trans. Am. Geophys.* 79, 579.
- 382 Yamazaki, K., 2011. Calculation of the piezomagnetic field arising from uniform
- 383 regional stress in inhomogeneously magnetized crust (II): Limitation in general
- 384 cases. *Earth Planets Space* 63, 1217–1220.
- 385 Zhan, Z., 1989. Investigations of tectonomagnetic phenomena in China. *Phys. Earth*
- 386 *Planet. Inter.* 57, 11–22.
- 387

Figure captions

Fig. 1. Models of the piezomagnetic field corresponding to the UCSB and USGS slip

models. (a) UCSB slip model with $H = 15$ km. (b) UCSB slip model with $H = 30$

km. (c) USGS slip model with $H = 15$ km. (d) USGS slip model with $H = 30$ km.

The rectangle represents the fault plane on which heterogeneous slip is considered.

Fig. 2. Models of the piezomagnetic field corresponding to averaged versions of the

UCSB and USGS slip models. (a) UCSB slip model with $H = 15$ km. (b) UCSB

slip model with $H = 30$ km. (c) USGS slip model with $H = 15$ km. (d) USGS slip

model with $H = 30$ km. The rectangle represents the fault plane on which uniform

slip is considered.

Fig. 3. Comparisons between observed and calculated piezomagnetic signals.

Calculated values in this figure are the averages of four piezomagnetic models.

Error bars in calculated values represent maximum and minimum values for the

four models. Error bars in observed values are from Utada et al. (2011). Open

circles indicate the results at sites KAK and KTR, where the magnetic anomalies

are rather small, while solid circles indicate the results at other sites.

Fig. 4. The magnetic anomaly over the Tohoku region as observed by an aeromagnetic

survey at a height of 5000 m. Contour intervals are 10 nT. Observations were

conducted by the Geographical Survey Institute (predecessor of the Geospatial

Information Authority) of Japan in 1990, and the data are available on their Web

site, in Japanese

(http://vldb.gsi.go.jp/sokuchi/geomag/menu_03/aeromag_data.html; last access: 26

October 2012).

Figure 1 Models of the piezomagnetic field corresponding to the UCSB and USGS slip models. (a) UCSB slip model with $H = 15$ km. (b) UCSB slip model with $H = 30$ km. (c) USGS slip model with $H = 15$ km. (d) USGS slip model with $H = 30$ km. The rectangle represents the fault plane on which heterogeneous slip is considered.

Figure 2 Models of the piezomagnetic field corresponding to averaged versions of the UCSB and USGS slip models. (a) UCSB slip model with $H = 15$ km. (b) UCSB slip model with $H = 30$ km. (c) USGS slip model with $H = 15$ km. (d) USGS slip model with $H = 30$ km. The rectangle represents the fault plane on which uniform slip is considered.

Figure 3 Comparisons between observed and calculated piezomagnetic signals. Calculated values in this figure are the averages of four piezomagnetic models. Error bars in calculated values represent maximum and minimum values for the four models. Error bars in observed values are from Utada et al. (2011). Solid circles indicate the results at sites KAK and KTR, where the magnetic anomalies are rather small, while open circles indicate the results at other sites.

Figure 4 The magnetic anomaly over the Tohoku region as observed by an aeromagnetic survey at a height of 5000 m. Contour intervals are 10 nT. Observations were conducted by the Geographical Survey Institute (predecessor of the Geospatial Information Authority) of Japan in 1990, and the data are available on their Web site, in Japanese (http://vldb.gsi.go.jp/sokuchi/geomag/menu_03/aeromag_data.html; last access: 26 October 2012).

Table 1

Parameters assumed in the modeling of the piezomagnetic field.

Parameter	Value
Rigidity	$57(53) \times 10^9 \text{ Pa}$
Poisson's ratio	0.25
Magnetization	1.0 A m^{-1}
Piezomagnetic stress sensitivity	$1 \times 10^{-9} \text{ Pa}^{-1}$
Curie point depth	15 and 30 km
Inclination of the ambient geomagnetic field	51.0 degree
Declination of the ambient geomagnetic field	-7.5 degree
Observation altitude	0 m

Table 2

Comparisons of coseismic changes in the geomagnetic total intensity, as predicted by the piezomagnetic models versus those calculated from data reported in Utada et al. (2011). Piezomagnetic field models are determined for Curie point depths of 30 and 15 km together with two slip models (USGS and UCSB).

Station code	USGS slip model		UCSB slip model		Observed (error)	
	30 km	15 km	30 km	15 km		
ESA	+0.05	+0.05	+0.55	+0.49	−0.09	(0.46)
HAR	−0.56	−0.37	−0.40	−0.06	−0.43	(0.36)
IWK	−0.48	−0.28	−0.44	−0.14	+0.21	(0.42)
KTR	−0.18	−0.10	−0.29	−0.18	−0.83	(0.20)
OTA	+0.14	+0.16	−0.08	−0.03	−0.01	(0.28)
KAK	−0.22	−0.14	−0.19	−0.07	−0.22	(0.10)